

Sub Theme : Adaptation and Production
ISBN : 978-602-8915-93-9

Oral Presentation

CHARACTERIZATION OF EGCG COMPOUND USE ^1H NMR SPECTRUM ON CAMELLIA SINENSIS (L.) CALLUS

Sutini ¹, Tatik W², Sutiman B³, R. Verpoorte ⁴

¹Agrotechnology Department of Agriculture Faculty UPN 'Veteran', Surabaya-East Java.

² Agronomy Department of Agriculture Faculty, Brawijaya University, Malang-East Java

³Biology Department of FMIPA, Brawijaya University, Malang-East Java

⁴Plant Metabolomics Department - Leiden University. Netherlands

Email : tien_basuki@yahoo.com

ABSTRACT

Epigallocatechin gallate (EGCG) are secondary metabolite on *Camellia sinensis* L as obesity preventing agent. The characterisation of this plant use ^1H NMR spectroscopy often have been done, however characterisation on callus both drying with open air and without drying is rare. The purpose of this research is characterize EGCG of tea callus via process both drying with open air and vacuum. Tis method use ^1H NMR spectroscopy. The result show that EGCG character of tea callus via process both drying with open air and vacuum are significantly different.

Key note: Epigallocatechin gallate, ^1H NMR, Camellia sinensis L callus

INTRODUCTION

Epigallocatechin gallate (EGCG) bioactive is available on tea (*Camellia sinensis* L). The advantage of this are anti obesity, anti cancer, anti diabetic, anti cholesterol, anti bakterial, kardiovaskuler disease and osteoporosis prevention agent. Many function of tea on industry that is beverage, cosmetic, pharmacy, and food (Hartoyo, 2003). EGCG bioactive compound structure as figure 1, is one of flavonoid derivat of phenol on tea (*Camellia sinensis* L.). Structure that have many hydroxyl/ OH easy to bond free radical so EGCG identified have multi function in health.




Figure 1. EGCG structure (Thomson, 2004).

Peter W.L. (2000) say that EGCG harvested on winter and summer season different on EGCG value. This is relevant with Caffin, N., D'Arcy, B., Yao L., Rintou, N. (2004) note that EGCG amount of tea leaves is increasing harvested on summer (May), however decreasing on winter (November).

In general, this research is aimed at developing production of EGCG technology *in vitro* by callus culture technique. Characteristics of EGCG are: binding with several biologic matrix and heavy metals, catalyzing electron transportation, and trapping free radicals. Four characteristics above made it a bioactive agent. Hence secondary metabolite of EGCG must be characterized with ^1H NMR both on drying and undrying process to improve the product quality. :

Purpose :

Characterize EGCG of tea callus with process both drying in open air and with vacuum.

MATERIALS AND METHOD

^1H NMR spectroscopy 500 MHz (Bruker, Jerman), 1.5 mL-ependorff tube mL -2 mL, 5 mm NMR tube, centrifuge, Ultrasonic, vortex, vacuum dry, metanol-deuterium ($\text{CH}_3\text{OH-d}_4$), buffer KH_2PO_4 in D_2O (pH 6,0) containing 0,01% (b / b) TSP, aqua bidestilata.

Extraction

Preparing metanol-deuterium ($\text{CH}_3\text{OH-d}_4$) without add standar internal, buffer KH_2PO_4 in D_2O (pH 6,0) containing 0,01% (b / b) TSP. Measure gentle powder of 25-50 mg tea callus that both with drying in open air and vacuum. Then add $\text{CH}_3\text{OH-d}_4$ (without any internal standard), KH_2PO_4 buffer in D_2O (pH 6.0) containing 0.1% (w/w) TSP, in to 2 mL-ependorff tube. This solution vortex for 1 minute at room temperature and then ultrasonication for 5-20 minute at room temperature. This solution centrifuge at room temperature for 5 – 20 minute using microtube centrifugator (13000 rpm, room temperature). Transfer supernatant (more than 1 mL) to 1.5 mL-ependorff tube.

If more centrifugation is necessary centrifugator using microtube centrifugator (13000 rpm, 1 minute, room temperature). Then transfer 800 mL of supernatant to 5 mm NMR tube.

Characterisation Use ^1H Nmr

The study done use 500 spectrometer MHz ^1H NMR (Bruker, Jerman) completed by cryoprobes. Chemical shift (δ) is measured on ppm, with standart referency use tetrametil silen zero ppm, with chemical shift range between 4.52-7.08.

RESULTS AND DISCUSSION

The liquid of green brownly pure extraction, then spectrum observed. EGCG spectra of tea callus proceed both with drying in open air and wet tea callus in vacuum and standar as Figure 1.

Spectrum ^1H NMR 500 MHz on methanol deuterium solvent, (Table 1) showed that chemical shift (δ ppm) and space between two spin/kopling constanta (J in Hz). Proton position are structure from EGCG resonance on H-6, H-8, H-2', H-5', H-6' (Markam, et al, 1994). Chemical shift and coupling constanta on proton position resonance for EGCG show that tea callus with drying in vacuum almostly same with standart. This show that tea callus characterization with drying vacuum can identify EGCG character, that not happen in tea callus with drying in open air. Proton position of EGCG resonance on chemical shift (δ) and coupling constanta (J), tea callus with open air drying can not show the character because that compound oxydated by air.

Based on Nathalie V, G. research (2001) that flavonoid oxydation caused by temperatur, UV light, and ion Cu^{2+} then change to be unstable quinon into sulfonat. Using ^1H NMR spectroscopy, can characterize proton of EGCG. This is relevant to Moco research (2007) show that ^1H NMR can identify flavonoid compound on tomato plant. Then, Tarachiwin L. Et al., 2007 note that ^1H NMR spectroscopy combined with multivariat analysis can descript secondary metabolit profile.

However using ^1H NMR spectroscopy, have disadvantage that is: 1). Relatively low sensitivity than using other analysis technique such as MS, 2). Can produce more than one ambiguous spectra, 3). Chemical shift influenced by the surrounding chemical environment. There is many ways to solve that is: 1). Combine 2D spectrum (two dimension) NMR, 2). This research use standart data comparison refer to the same sovent material.




Figure 1. Spektrum ^1H NMR 500 MHz on $\text{CH}_3\text{OH-d}_4$ solvent from: (A) tea callus with vacuum drying, (B), standart , (C) tea callus with drying in open air.

Table 1. Proton position δ and J EGCG , tea callus drying in open air, tea callus Drying with vacuum and standart

Proton Position	δ EGCG tea callus drying in open air, (J in Hz)	δ EGCG tea callus drying with vacuum, (J in Hz)	δ EGCG with standart (J in Hz)
H-2	-	4.90 (s)	4.90 (s)
H-3	-	5.50 (s)	5.51 (s)
H-4 α	-	2.97 (dd)	2.97 (dd)
H-4 β	-	2.83 (dd)	2.83 (dd)
H-6	-	5.94(s,1.79)	5.93(s,1.79)
H-8	-	5.94(s,1.79)	5.93(s,1.79)
H-2'	6.34(s)	6.49(s,1.80)	6.48(s,1.80)
H-5'	-	-	-
H-6'	6.40(s,)	6.49(s,1.80)	6.48(s,1.80)
H-2''	-	6.95(s)	6.93 (s)

CONCLUSION

Achieve character from EGCG body that is ^1H NMR as H-6, H-8, H-2', H-5', H-6' (Markam, et al, 1994). Observation on chemical shift located between 5,94 – 6,49, this value based on existence range of EGCG compound. This relevant with study by McLeod (2010) that aromatic bonding area located on 5,8 – 8,8

ACKNOWLEDGEMENT :

Thanks to Prof. Dr. Rob. Verpoorte and staff of Department Plant Metabolomics, Leiden University, Netherlands for help and attention.

REFERENCES

- Caffin, N., dkk., 2004. Developing an index of quality for Australian tea, Rural Industries Research and Development Corporation, <http://www.rirdc.gov.au>
- Hartoyo Arif. 2003. Teh dan kasiatnya bagi kesehatan. Sebuah tinjauan ilmiah, Kanisius. Yogyakarta.
- Markam, dkk., 1994. H. Nuclear magnetic resonance spectroscopy of flavonoids and their glycosides in hexamethylsulfoxide. Di dalam Harborn J.B., The flavonoid in research Science, Chapman and Hall limited London.
- McLeod. 2010. New Trend Research on Eco-friendly technology for medicine and Agriculture. Basic science national seminar. Universitas Brawijaya Malang.
- Moço Sofia. 2007. Metabolomics Technologies applied to the Identification of Compounds in Plants. Dissertation. Wageningen University. Netherlands.
- Nathalie V, G. dkk., 2001. Study and quantification of monomeric flavan-3-ol and dimeric oxidation J. Sci Food Agric (81): p 1172-1179
- Peter W.L. 2002. Biochemical analysis for identification of quality in black tea (Camellia sinensis), Disertasi. Faculty of Natural and Agricultural Sciences. Department of Biochemistry. University of Pretoria. Pretoria. South Africa. <http://upetd.up.ac.za/thesis/available/etd-03012005084935/unrestricted/00dissertation.pdf>
- Tarachiwin L. dkk., 2007. ^1H NMR Based Metabolic Profiling in the Evaluation of Japanese Green Tea Quality. *J. Agric. Food Chem.*(55): p. 9330–9336.
- Thomson Leonor. 2004. Enrichment of Biologically Active Compounds from Selected Plants Using Adsorptive Bubble Separation, Dissertation Fakultät Wissenschaftszentrum Weihenstephan für Ernährung, Landnutzung und Umwelt Technischen Universität München. <http://www.google.co.id/search?hl=id&q=production+catechin++from+callus+camellia+sinensis+leaf+pdf+%2Bfree+purchase+research+&meta>.

Question :

1. What the structure resonance in EGCG ?

Answer :

1. Aquired characteristic from EGCG structure resonance on H that H – 6, H – 8, H - 2, H – 5, H – 7, etc.