FINAL VERSION

Synchronization of Stochastic Genetic Oscillator Networks with Time Delays and Markovian Jumping Parameters

Yao Wang^a, Zidong Wang^{a,b,*}, Jinling Liang^c, Yurong Li^d and Min Du^d

Abstract

Genetic Oscillator networks (GONs) are inherently coupled complex systems where the nodes indicate the biochemicals and the couplings represent the biochemical interactions. This paper is concerned with the synchronization problem of a general class of stochastic GONs with time delays and Markovian jumping parameters, where the GONs are subject to both the stochastic disturbances and the Markovian parameter switching. The regulatory functions of the addressed GONs are described by the sector-like nonlinear functions. By applying up-to-date 'delay-fractioning' approach for achieving delay-dependent conditions, we construct novel matrix functional to derive the synchronization criteria for the GONs that are formulated in terms of linear matrix inequalities (LMIs). Note that LMIs are easily solvable by the Matlab toolbox. A simulation example is used to demonstrate the synchronization phenomena within biological organisms of a given GON and therefore shows the applicability of the obtained results.

Keywords

Genetic oscillator networks; system biology; stochastic synchrony; Markovian switching; random perturbation; linear matrix inequality (LMI).

I. INTRODUCTION

The oscillatory behavior of genetic networks, as a fundamental challenge in the research field of systems biology, has recently attracted an increasing attention, see e.g. [1,5,6,10–12,14,29]. Generally speaking, the genetic networks are a class of complex dynamical networks since the genetic oscillators can be expressed in terms of complicated biological functions [9, 16, 17, 23]. In such kind of genetic oscillator networks (GONs), the nodes represent the genetic oscillators, while the inner or outer couplings denote the interactions. In order to research into the intrinsic biological organisms of GONs, it is of great importance to investigate the collective dynamics of coupled genetic oscillators with hope to understand the rhythmic behavior of living organisms. Synchronization, as a universal phenomenon, occurs typically in genetic networks. For example,

This work was supported in part by the Biotechnology and Biological Sciences Research Council (BBSRC) of the U.K. under Grants BB/C506264/1 and 100/EGM17735, the Royal Society of the U.K., the National Natural Science Foundation of China under Grant 60804028, the Teaching and Research Fund for Excellent Young Teachers at Southeast University of China, the International Science and Technology Cooperation Project of China under Grant 2009DFA32050, and the Alexander von Humboldt Foundation of Germany.

^aSchool of Information Science and Technology, Donghua University, Shanghai 200051, China.

^bDepartment of Information Systems and Computing, Brunel University, Uxbridge, Middlesex, UB8 3PH, United Kingdom. ^cDepartment of Mathematics, Southeast University, Nanjing 210096, China.

^dDepartment of Electrical Engineering, Fuzhou University, Fuzhou 350002, China.

^{*}Corresponding author. E-mail: Zidong.Wang@brunel.ac.uk

 $\mathbf{2}$

in [19], a synthetic gene network in Escherichia coli has been shown to have two features: the system acts as a relaxation oscillator and uses an intercell signaling mechanism to couple the oscillators and induce synchronous oscillations. A coupling scheme has been proposed that leads to synchronous behavior across a population of cells, and an analytical treatment of the synchronization process has been conducted. Up to now, the synchronization motion analysis problem for genetic oscillator networks has attracted considerate research attention. In [11,14,16,17], the synchronization problem in genetic networks has been thoroughly investigated via experiments (e.g. synchronization of cellular clock in the suprachiasmatic nucleus in genetic networks), numerical simulation (e.g. biological networks of identical genetic oscillators) as well as theoretical analysis (i.e., synchronizability of coupled nonidentical genetic oscillators).

It has been demonstrated experimentally that the networks states or oscillatory expression are significantly affected by the inherent state delay due primarily to the slow processes of transcription, translation, and translocation or the finite switching speed of amplifiers. From the synthetic biology viewpoint, it is necessary to address the time-delay effects in the mathematical models, and then a more accurate state values of the biological oscillators could be obtained from oscillatory expression measurements [22, 26, 30]. Note that the stability analysis issue of genetic regulatory networks with either constant or time-varying delays has recently been a research focus, see [30] and references therein. It is noter to achieve less conservative delay-dependence conditions, see e.g. [20, 27, 28, 35].

Biological data promises to enhance the fundamental understanding of life at the molecular level, from regulation of gene expression and gene function to cellular mechanisms, and may prove useful in medical diagnosis, treatment, and drug design. Substantial effort is being made to build models to analyze microarray data. It is evident that genetic networks are always affected by the random fluctuations [1,3,13,21,25,26,29,31]. Therefore, to have an accurate prediction of the dynamical behaviors of genetic oscillators, it is important to consider the random effects including intrinsic and intrinsic noise perturbations [1,3,25,26,29]. Also, as shown in [7,8,13], in gene regulatory networks, the transition from one state to the next usually takes place in accordance with certain transition probabilities, which forms a homogeneous Markov chain with finite state space. Subsequently, the dynamics of the so-called Markovian genetic regulatory networks, which are subject to mode switching (or jumping), has been thoroughly investigated in [7,8]. It should be pointed out that, up to now, the control and filtering problems for Markovian jumping systems have already been widely studied [4, 18, 33, 34]. Recently, the stochastic synchrony study has been carried out for genetic networks in [32], where an adaptive filtering approach is elegantly developed to estimate uncertain delayed genetic regulatory networks. However, the stochastic synchrony problem for *Markovian* delayed genetic networks of specific structures has not gained adequate research attention yet, and this constitutes the main focus of this paper.

In this paper, we aim to make one of the first attempts to investigate the synchronization problem for stochastic GONs with Markovian jumping parameters and time delays so as to exhibit more realistic characteristics of the GONs, where the regulation functions are assumed to be sector-like, and the intrinsically stochastic fluctuation is a scalar Brownian motion. The main results obtained are illustrated through a numerical simulation example. The rest of this paper is organized as follows. Section II introduces the model formulation and some preliminary works. In Section III, by utilizing the approach of 'delay-fractioning' and a novel matrix functional method, stochastic analysis is conducted to obtain delay-dependent sufficient criteria described by linear matrix inequalities (LMIs) [2] that can be easily checked by using standard numerical software. Section IV illustrates the obtained results and Section V concludes the paper.

Notations: Throughout this paper, \mathfrak{R}^n and $\mathfrak{R}^{n\times m}$ denote, respectively, the *n* dimensional Euclidean space and the set of all $n \times m$ real matrices. P > 0 means that matrix *P* is real, symmetric and positive definite. *I* and 0 denote the identity matrix and the zero matrix with compatible dimensions, respectively; and diag $\{\cdots\}$ stands for a block-diagonal matrix, $\operatorname{col}\{\cdots\}$ denotes a matrix column with blocks given by the matrices in $\{\cdots\}$. If *A* is a matrix, the notation $\lambda_{max}(A)$ means the largest eigenvalue of *A*. The superscript "*T*" stands for matrix transposition and the asterisk "*" in a matrix is used to represent the term which is induced by symmetry. The Kronecker product of matrices $Q \in \mathfrak{R}^{m \times n}$ and $R \in \mathfrak{R}^{p \times q}$ is a matrix in $\mathfrak{R}^{mp \times nq}$ and denoted as $Q \otimes R$. We let $\mathcal{C}([-h, 0]; \mathfrak{R}^n)$ denote the family of all continuous functions φ from [-h, 0] to \mathfrak{R}^n with the norm $|\varphi| = \sup_{-h \leq \theta \leq 0} ||\varphi(\theta)||$, where $|| \cdot ||$ is the Euclidean norm on \mathfrak{R}^n . Moreover, let $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t\geq 0}, \mathcal{P})$ be a complete probability space with a filtration $\{\mathcal{F}_t\}_{t\geq 0}$ satisfying the usual conditions (i.e., the filtration contains all \mathcal{P} -null sets and is right continuous). Denote by $L^p_{\mathcal{F}_0}([-h, 0]; \mathfrak{R}^n)$ the family of all \mathcal{F}_0 -measurable $\mathcal{C}([-h, 0]; \mathfrak{R}^n)$ -valued random variables $\xi = \{\xi(\theta) : -h \leq \theta \leq 0\}$ such that $\sup_{-h \leq \theta \leq 0} \mathcal{E}\{|\xi(\theta)|^p\} < \infty$, where $\mathcal{E}\{\cdot\}$ stands for the mathematical expectation operator with respect to the given probability measure \mathcal{P} . Sometimes, the arguments of a function will be omitted in the analysis when no confusion arises.

II. PROBLEM FORMULATION AND PRELIMINARIES

Let r(t) $(t \ge 0)$ be a right-continuous Markovian chain on a probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t\ge 0}, P)$ taking values in a finite state space $\mathbb{S} = \{1, 2, ..., m\}$ with generator $\Pi = \{\pi_{ij}\}$ given by

$$P\{r(t+\Delta) = j \mid r(t) = i\} = \begin{cases} \pi_{ij}\Delta + o(\Delta), & \text{if } i \neq j, \\ 1 + \pi_{ij}\Delta + o(\Delta), & \text{if } i = j. \end{cases}$$

Here $\Delta > 0$, and $\pi_{ij} \ge 0$ is the transition rate from *i* to *j* if $j \ne i$ while

$$\pi_{ii} = -\sum_{j \neq i} \pi_{ij}.$$

Among many models of genetic networks, the differential equation model is one of the mostly adopted ones. A general delayed genetic oscillator network could be described by the following vector form [16, 17]:

$$\frac{dy(t)}{dt} = Ay(t) + \sum_{i=1}^{l} B_i f_i(y(t)) + \sum_{i=1}^{l} C_i g_i(y(t-\tau)),$$
(1)

where l is a positive integer and $y(t) = col\{y_1(t), y_2(t), \ldots, y_n(t)\} \in \mathfrak{R}^n$ represents the concentrations of proteins, mRNAs and chemical complexes; A, B_i, C_i $(i = 1, 2, \ldots, l)$ are matrices in $\mathfrak{R}^{n \times n}$; $f_i(y(t)) = col\{f_{i1}(y_1(t)), f_{i2}(y_2(t)), \ldots, f_{in}(y_n(t))\} \in \mathfrak{R}^n$ and $g_i(y(t-\tau)) = col\{g_{i1}(y_1(t-\tau)), g_{i2}(y_2(t-\tau)), \ldots, g_{in}(y_n(t-\tau))\} \in \mathfrak{R}^n$ are monotonic genetic regulatory functions which are usually taken as the Hill form. The scalar $\tau > 0$ denotes the translation time delay in the translation process.

As discussed in the introduction, the genetic oscillators in biological networks are tightly coupled between each other, and both the stochastic perturbations [1, 3, 13, 21, 25, 26, 29] and Markovian jumping parameters

$$dx_{k}(t) = \left[A(r(t))x_{k}(t) + B(r(t))f(x_{k}(t)) + C(r(t))g(x_{k}(t-\tau)) + \sum_{l=1}^{N} w_{kl}\Gamma_{r(t)}x_{l}(t)\right]dt + \sigma_{k}(x_{k}(t), x_{k}(t-\tau), t, r(t))d\omega(t),$$

$$x_{k}(t) = \phi_{k}(t), \quad r(t)|_{t=0} = r_{0} \in \mathbb{S}; \quad t \in [-\tau, 0], \quad k = 1, 2, \dots, N$$
(2)

where $x_k(t) = \operatorname{col}\{x_{k1}(t), x_{k2}(t), \cdots, x_{kn}(t)\} \in \mathfrak{R}^n$ is the state vector of the *k*th genetic oscillator representing the concentrations of proteins, mRNAs and chemical complexes, which are of limited values; For $r(t) = i \in \mathbb{S}$, A(i) includes the degradation terms and all the other linear terms of the *k*th genetic oscillator; B(i), C(i)are known matrices in $\mathfrak{R}^{n \times n}$; $f(x_k(t)) = \operatorname{col}\{f_1(x_{k1}(t)), f_2(x_{k2}(t)), \cdots, f_n(x_{kn}(t))\} \in \mathfrak{R}^n$ and $g(x_k(t-\tau)) =$ $\operatorname{col}\{g_1(x_{k1}(t-\tau)), g_2(x_{k2}(t-\tau)), \cdots, g_n(x_{kn}(t-\tau))\} \in \mathfrak{R}^n$ are usually monotonic functions satisfying the sector-bounded conditions that will be given later; $\phi_k(t) \in L^p_{\mathcal{F}_0}([-h, 0]; \mathfrak{R}^n)$ is the initial condition of $x_k(t)$.

The matrix $\Gamma_{r(t)} = [\gamma_{kl,r(t)}]_{n \times n}$ is a matrix linking the state variable of the *l*th genetic oscillator in the genetic network mode r(t) if $\gamma_{kl,r(t)} \neq 0$; and $W = [w_{kl}]_{N \times N}$ is the coupling matrix that represents the coupling topology, direction, as well as the coupling strength of the genetic network. The definition is given as follows: if there is a link from the *k*th oscillator to the *l*th oscillator $(k \neq l)$, then w_{kl} equals to a positive constant denoting the coupling strength of this link; otherwise $w_{kl} = 0$; $w_{kk} = -\sum_{k=1,k\neq l}^{N} w_{kl}$. Note that many real-world genetic oscillator networks are sparse, and therefore reducible near a tree topological structure. In this paper, however, we are looking into the synchronization problem for a genetic oscillator network from a theoretical viewpoint. In such as case, we suggest that the presence of irreducible dependencies among the genes is likely to identify direct regulatory interactions mediated by a transcription factor binding to a target genes promoter region, although other types of interactions may also be identified.

In the system (2), $\omega(t)$ is a scalar Wiener process (Brownian Motion) on $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t\geq 0}, \mathcal{P})$, which is independent of the Markov chain $r(\cdot)$ and satisfies

$$\mathcal{E}\{d\omega(t)\} = 0, \qquad \mathcal{E}\{[d\omega(t)]^2\} = dt.$$
(3)

The noise intensity function vector $\sigma_k(\cdot, \cdot, \cdot, \cdot) : \mathfrak{R}^n \times \mathfrak{R}^n \times \mathfrak{R}^n \times \mathfrak{R}^n \times \mathfrak{R}^n$ is Borel measurable and is assumed to satisfy the following Lipschitz condition.

Assumption 1: There exist constant matrices Σ_1 and Σ_2 of appropriate dimensions such that the following inequality:

$$[\sigma_k(u_1, v_1, t, i) - \sigma_l(u_2, v_2, t, i)]^T [\sigma_k(u_1, v_1, t, i) - \sigma_l(u_2, v_2, t, i)] \le \|\Sigma_1(u_1 - u_2)\|^2 + \|\Sigma_2(v_1 - v_2)\|^2$$
(4)

holds for all t > 0; u_q, v_q $(q = 1, 2) \in \Re^n$; $k, l = 1, 2, \cdots, N$; $i \in \mathbb{S}$.

Assumption 2: The nonlinear functions $f(\cdot)$ and $g(\cdot)$ satisfy the following sector-like conditions:

$$0 \le \frac{f_i(s) - f_i(t)}{s - t} < \kappa_{1i}, \quad 0 \le \frac{g_i(s) - g_i(t)}{s - t} < \kappa_{2i}, \qquad i = 1, 2, \cdots, n.$$

Remark 1: From Assumption 2, it is not difficult to see that

$$f^{T}(x)(f(x) - K_{1}x) \le 0, \qquad g^{T}(x)(g(x) - K_{2}x) \le 0$$

Remark 2: Notice that the sector-like description of the nonlinearities has been used to model the structure and regulation mechanism of the genetic regulatory networks in many papers, see e.g. [17,29]. Traditionally, as monotonic regulation functions, $f(\cdot)$ or $g(\cdot)$ usually takes the Hill form or the Michaelis-Menten form, which is a special case of the sector-like functions.

For the sake of notation simplicity, we use the matrix Kronecker product " \otimes " to rewrite (2) in the following compact form:

$$dx(t) = \left[(I_N \otimes A(r(t)) + W \otimes \Gamma_{r(t)}) x(t) + (I_N \otimes B(r(t))) F(x(t)) + (I_N \otimes C(r(t))) G(x(t-\tau)) \right] dt + \sigma(x(t), x(t-\tau), t, r(t)) d\omega(t),$$
(5)

where

$$\begin{aligned} x(t) &= \operatorname{col}\{x_1(t), x_2(t), \dots, x_N(t)\}, \quad F(x(t)) &= \operatorname{col}\{f(x_1(t)), f(x_2(t)), \dots, f(x_N(t))\}, \\ G(x(t)) &= \operatorname{col}\{g(x_1(t-\tau)), g(x_2(t-\tau)), \dots, g(x_N(t-\tau))\}, \\ \sigma(x(t), x(t-\tau), t, r(t)) &= \operatorname{col}\{\sigma_1(x_1(t), x_1(t-\tau), t, r(t)), \dots, \sigma_N(x_N(t), x_N(t-\tau), t, r(t))\} \end{aligned}$$

Before stating the main results, a definition and some lemmas are introduced here.

Definition 1: The genetic oscillator network (2) is said to be globally asymptotically synchronized in the mean square sense if

$$\mathcal{E}\left\{\|x_k(t,\phi_k,r_0) - x_l(t,\phi_l,r_0)\|^2\right\} \longrightarrow 0, \text{ as } t \to +\infty$$

holds for any $\phi_k(\cdot)$, $\phi_l(\cdot) \in L^p_{\mathcal{F}_0}([-h, 0], \Re^n)$; $k, l \in \{1, 2, ..., N\}$ and $r_0 \in \mathbb{S}$.

Lemma 1: [15] The Kronecker product has the following properties:

- (1) $(\alpha A) \otimes B = A \otimes (\alpha B);$
- (2) $(A+B) \otimes C = A \otimes C + B \otimes C;$
- (3) $(A \otimes B)(C \otimes D) = (AC) \otimes (BD);$
- $(4) \ (A \otimes B)^T = A^T \otimes B^T.$

Lemma 2: Let $\mathcal{U} = (\alpha_{ij})_{N \times N}$, $P \in \mathbb{R}^{n \times n}$, $x = \operatorname{col}\{x_1, x_2, ..., x_N\}$ where $x_i = \operatorname{col}\{x_{i1}, x_{i2}, ..., x_{in}\} \in \mathbb{R}^n$ and $y = \operatorname{col}\{y_1, y_2, ..., y_N\}$ where $y_i = \operatorname{col}\{y_{i1}, y_{i2}, ..., y_{in}\} \in \mathbb{R}^n$ (k = 1, 2, ..., N). If $\mathcal{U} = \mathcal{U}^T$ and each row sum of \mathcal{U} is zero, then

$$x^{T}(\mathcal{U} \otimes P)y = -\sum_{1 \leq i < j \leq N} \alpha_{ij}(x_{i} - x_{j})^{T} P(y_{i} - y_{j}).$$

Lemma 3: [24] Let f be a nonnegative function defined on $[0, +\infty)$. If f is Lebesgue integrable and is uniformly continuous on $[0, +\infty)$, then $\lim_{t\to +\infty} f(t) = 0$.

The main aim of this paper to deal with the exponential mean-square synchronization problem of the genetic oscillator networks (2) or (5), and derive LMI-based sufficient conditions that guarantee the network to be exponentially synchronous in the mean-square sense for all admissible time delays, nonlinearities and stochastic disturbances.

III. MAIN RESULTS

Letting

$$y(t, r(t)) := [I_N \otimes A(r(t)) + W \otimes \Gamma_{r(t)}]x(t) + (I_N \otimes B(r(t)))F(x(t)) + (I_N \otimes C(r(t)))G(x(t-\tau)), \quad (6)$$

then the network (5) can be recast into the following form

$$dx(t) = y(t, r(t))dt + \sigma(x(t), x(t-\tau), t, r(t))d\omega(t).$$
(7)

For each $i \in \mathbb{S}$, Eq. (7) becomes

$$dx(t) = y^{(i)}(t)dt + \sigma(x(t), x(t-\tau), t, i)d\omega(t),$$
(8)

where $y^{(i)}(t) = y(t, r(t))$ as r(t) = i.

Theorem 1: For a fixed integer $d \ge 1$, the genetic oscillator networks (2) or (5) with time-delay $\tau \in (0, h]$ is globally exponentially synchronous in the mean square if there exist $n \times n$ matrices $P_i > 0$, $Q_k > 0$, R > 0, a positive-define matrix $X^{(i)} \in \Re^{(dn+4n)\times(dn+4n)}$, positive scalars λ^* , e_l and matrices $H^{(i)}$, $M_1^{(i)}$, $M_2^{(i)}$, ..., $M_{d+4}^{(i)}$ with appropriate dimensions such that

$$P_i \le \lambda^* I,\tag{9}$$

$$\Xi_{kl}^{(i)} = \Pi_{kl}(i) + hX^{(i)} < 0, \tag{10}$$

$$\Omega_{kl}^{(i)} = \begin{bmatrix} X^{(i)} & M^{(i)} \\ * & R \end{bmatrix} \ge 0,$$
(11)

where $k = 1, 2, \ldots, d, l = 1, 2, i \in \mathbb{S}$ and

$$\begin{split} \Pi_{kl}(i) &= W_Q^T \tilde{Q} W_Q + \sum_{q=1}^2 (W_M^T M^{(i)} W_M^{(q)} + (W_M^T M^{(i)} W_M^{(q)})^T) + W_H^T \tilde{H}_{kl}^{(i)} W_H, \\ W_Q &= \begin{bmatrix} I_{dn \times dn} & 0_{dn \times n} & 0_{dn \times 3n} \\ 0_{dn \times n} & I_{dn \times dn} & 0_{dn \times 3n} \end{bmatrix}, \quad \tilde{Q} = \begin{bmatrix} \mathbb{Q} & 0 \\ 0 & \mathbb{Q} \end{bmatrix}, \quad M^{(i)} = \operatorname{col}\{M_1^i, M_2^i, \cdots, M_{d+4}^i\}, \\ W_M &= I_{d+4} \otimes I_n, \quad W_M^{(1)} = [I_n \ 0_{n \times (d+3)n}], \quad W_M^{(2)} = [0_{n \times dn} \ -I_n \ 0_{n \times 3n}], \\ W_H &= \begin{bmatrix} I_n \ 0_{n \times (dn+3n)} \\ 0_{n \times (dn+n)} \ I_n \ 0_{n \times 2n} \\ 0_{n \times (dn+2n)} \ I_n \ 0_{n \times n} \\ 0_{n \times (dn+3n)} \ I_n \end{bmatrix}, \\ \tilde{H}_{kl}^{(i)} &= \begin{bmatrix} (1,1) \ 0 \ e_1 K_1^T + P_i B(i) \ P_i C(i) \ (1,5) \\ * \ \lambda^* \Sigma_2^T \Sigma_2 \ 0 \ e_2 K_2^T \ 0 \\ * \ * \ -2e_1 I \ 0 \ B^T(i) H^{(i)^T} \\ * \ * \ * \ -2e_2 I \ C^T(i) H^{(i)^T} \\ * \ * \ * \ hR - H^{(i)} - H^{(i)^T} \end{bmatrix}, \\ (1,1) &= \sum_{1 \le j \le m} \pi_{ij} P_j + \lambda^* \Sigma_1^T \Sigma_1 + P_i A(i) + A^T(i) P_i - N w_{kl} (P_i \Gamma_i + \Gamma_i^T P_i), \\ (1,5) &= -N w_{kl} \Gamma_i^T H^{(i)^T} + A^T(i) H^{(i)^T}. \end{split}$$

Proof: Since $\{(x(t), r(t)), t \ge 0\}$ is not a Markov process, in order to cast our model into the framework for a Markov system, let us define a new Markov process $\{x_t, r(t), t \ge 0\}$ with

$$x_t(s) = x(t+s), \ -\tau \le t \le 0.$$

Based on the idea of 'delay-fractioning', we consider the following novel matrix functional candidate for the genetic oscillator networks (2) or (5):

$$V(x_t, t, r(t)) = x^T(t)(U \otimes P(r(t)))x(t) + \int_{t-\tau}^t \int_{\beta}^t y^T(\alpha, r(\alpha))(U \otimes R)y(\alpha, r(\alpha))d\alpha d\beta + \sum_{k=1}^d \int_{t-\frac{k}{d}\tau}^{t-\frac{k-1}{d}\tau} x^T(\theta)(U \otimes Q_k)x(\theta)d\theta$$
(12)

with $d \ge 1$ (number of fractions) being an integer and $U = [u_{kl}]_{N \times N}$ with $u_{kl} = \begin{cases} -1, & k \ne l \\ N-1, & k = l \end{cases}$. For $r(t) = i \in \mathbb{S}, P(r(t)) = P_i > 0, R > 0, Q_k > 0.$

For the function $V : L^p_{\mathcal{F}_0}([-h, 0]; \mathfrak{R}^{Nn}) \times \mathfrak{R}^+ \times \mathbb{S} \to \mathfrak{R}^{Nn}$ given above, its infinitesimal operator \mathcal{L} [4] is defined as

$$\mathcal{L}V(x_t, t, i) = \lim_{\Delta \to 0^+} \frac{1}{\Delta} \bigg[\mathcal{E} \bigg\{ V(x_{t+\Delta}, t+\Delta, r(t+\Delta)) | x_t, r(t) = i \bigg\} - V(x_t, t, i) \bigg].$$
(13)

Then, for the stochastic process $\{x_t, r(t), t \ge 0\}$ along the network (2) or (5), it is easy to obtain that

$$\mathcal{L}V(x_{t},t,i) = 2x^{T}(t)(U \otimes P_{i})y^{(i)}(t) + \sum_{j=1}^{m} \pi_{ij}x^{T}(t)(U \otimes P_{j})x(t) + \operatorname{trace}[\sigma^{T}(x(t), x(t-\tau), t, i)(U \otimes P_{i})\sigma(x(t), x(t-\tau), t, i)] + \tau y^{(i)}(t)^{T}(U \otimes R)y^{(i)}(t) - \int_{t-\tau}^{t} y^{T}(\beta, r(\beta))(U \otimes R)y(\beta, r(\beta))d\beta + x^{T}(t)(U \otimes Q_{1})x(t) - x^{T}(t-\tau)(U \otimes Q_{d})x(t-\tau) - \sum_{l=1}^{d-1} \left\{ x^{T}\left(t - \frac{l}{d}\tau\right) [U \otimes (Q_{l} - Q_{l+1})]x\left(t - \frac{l}{d}\tau\right) \right\}.$$
(14)

With the following formula

$$x(t) - x(t - \tau) = \int_{t-\tau}^{t} dx(\beta),$$

we have for $r(t) = i \in \mathbb{S}$ that

$$2(\xi^{(i)}(t))^T M_U^{(i)} \left[x(t) - x(t-\tau) - \int_{t-\tau}^t y(\beta, r(\beta)) d\beta - \int_{t-\tau}^t \sigma(x(\beta), x(\beta-\tau), \beta, r(\beta)) d\omega(\beta) \right] = 0,$$
(15)

where $M_U^{(i)} = \operatorname{col}\{U \otimes M_1^{(i)}, U \otimes M_2^{(i)}, \cdots, U \otimes M_{d+4}^{(i)}\}$ is of appropriate dimensions and

$$\xi^{(i)}(t) = \operatorname{col}\left\{x(t), \ x\left(t - \frac{1}{d}\tau\right), \cdots, \ x\left(t - \frac{d-1}{d}\tau\right), \ x(t-\tau), \ F(x(t)), \ G(x(t-\tau)), \ y^{(i)}(t)\right\}.$$

Considering condition (3), one has

$$\mathcal{E}\left\{2(\xi^{(i)}(t))^T M_U^{(i)} \int_{t-\tau}^t \sigma(x(\beta), x(\beta-\tau), \beta, r(\beta)) d\omega(\beta)\right\} = 0, \quad i = 1, 2, \dots, m.$$

$$\tag{16}$$

Also, by (6), for any matrix $H^{(i)}$ with appropriate dimensions, we have

$$2(y^{(i)}(t))^{T}(U \otimes H^{(i)}) \left[(I_{N} \otimes A(i) + W \otimes \Gamma_{i})x(t) + (I_{N} \otimes B(i))F(x(t)) + (I_{N} \otimes C(i))G(x(t-\tau)) - y^{(i)}(t) \right] = 0.$$
(17)

Noticing the fact of UW = NW, it follows from Lemma 1 that

$$(U \otimes P_i)(W \otimes \Gamma_i) = (UW) \otimes (P_i \Gamma_i) = (NW) \otimes (P_i \Gamma_i), \ \forall i \in \mathbb{S}.$$
(18)

Substituting (6), (15)-(16) into (14) and utilizing Lemma 2, one obtains

$$\begin{split} \mathcal{L}V(x_{l},t,i) \\ &\leq \sum_{1 \leq k < l \leq N} \left\{ (x_{k}(t) - x_{l}(t))^{T} \Big[\Big(\sum_{1 \leq j \leq m} \pi_{ij} P_{j} + P_{i} A(i) + A^{T}(i) P_{i} - Nw_{kl}(P_{i}\Gamma_{i} + \Gamma_{i}^{T}P_{i}) \Big) \\ &\times (x_{k}(t) - x_{l}(t)) + 2P_{i} B(i)(f(x_{k}(t)) - f(x_{l}(t))) + 2P_{i} C(i)(g(x_{k}(t - \tau))) - g(x_{l}(t - \tau)))) \Big] \\ &+ (\sigma_{k}(x_{k}(t), x_{k}(t - \tau), t, i) - \sigma_{l}(x_{l}(t), x_{l}(t - \tau), t, i))^{T} P_{i} \\ &\times (\sigma_{k}(x_{k}(t), x_{k}(t - \tau), t, i) - \sigma_{l}(x_{l}(t), x_{l}(t - \tau), t, i)) \\ &+ h(y_{k}^{(i)}(t) - y_{l}^{(i)}(t))^{T} R(y_{k}^{(i)}(t) - y_{l}^{(i)}(t)) \\ &- \int_{t-\tau}^{t} (y_{k}(\beta, r(\beta)) - y_{l}(\beta, r(\beta)))^{T} R(y_{k}(\beta, r(\beta)) - y_{l}(\beta, r(\beta))) d\beta \\ &+ (\Upsilon_{k}(t) - \Upsilon_{l}(t))^{T} \mathbb{Q}(\Upsilon_{k}(t) - \Upsilon_{l}(t)) - \Big(\Upsilon_{k}(t - \frac{\tau}{d}) - \Upsilon_{l}(t - \frac{\tau}{d}) \Big)^{T} \mathbb{Q}\Big(\Upsilon_{k}(t - \frac{\tau}{d}) - \Upsilon_{l}(t - \frac{\tau}{d}) \Big) \\ &+ 2(\xi_{k}^{(i)}(t) - \xi_{l}^{(i)}(t))^{T} M^{(i)} \Big[(x_{k}(t) - x_{l}(t)) - (x_{k}(t - \tau) - x_{l}(k - \tau)) - \int_{t-\tau}^{t} (y_{k}(\beta, r(\beta)) - y_{l}(\beta, r(\beta))) d\beta \\ &- \int_{t-\tau}^{t} (\sigma_{k}(x_{k}(\beta), x_{k}(\beta - \tau), \beta, i) - \sigma_{l}(x_{l}(\beta), x_{l}(\beta - \tau), \beta, i)) d\omega(\beta) \Big] \\ &+ 2(y_{k}^{(i)}(t) - y_{l}^{(i)}(t))^{T} \Big[(H^{(i)}A(i) - Nw_{kl}H^{(i)}\Gamma_{i})(x_{k}(t) - x_{l}(t)) + H^{(i)}B(i)(f(x_{k}(t)) - f(x_{l}(t))) \\ &+ H^{(i)}C(i)(g(x_{k}(t - \tau)) - g(x_{l}(t - \tau))) - H^{(i)}(y_{k}^{(i)}(t) - y_{l}^{(i)}(t)) \\ &+ h(\xi_{k}^{(i)}(t) - \xi_{l}^{(i)}(t))^{T} X^{(i)}(\xi_{k}^{(i)}(t) - \xi_{l}^{(i)}(t)) - \int_{t-\tau}^{t} (\xi_{k}^{(i)}(t) - \xi_{l}^{(i)}(t))^{T} X^{(i)}(\xi_{k}^{(i)}(t) - \xi_{l}^{(i)}(t)) ds \Big] \Big\},$$

where $M^{(i)} = \operatorname{col}\{M_1^{(i)}, M_2^{(i)}, \dots, M_{d+4}^{(i)}\}, \mathbb{Q} = \operatorname{diag}\{Q_1, Q_2, \cdots, Q_d\}, \Upsilon_k(t) = \operatorname{col}\{x_k(t), x_k(t - \frac{\tau}{d}), \cdots, x_k(t - \frac{d-1}{d}\tau)\}.$

From conditions (4) and (9), we have

$$[\sigma_{k}(x_{k}(t), x_{k}(t-\tau), t, i) - \sigma_{l}(x_{l}(t), x_{l}(t-\tau), t, i)]^{T} P_{i}[\sigma_{k}(x_{k}(t), x_{k}(t-\tau), t, i) - \sigma_{l}(x_{l}(t), x_{l}(t-\tau), t, i)]$$

$$\leq \lambda_{\max}(P_{i}) \left[\sigma_{k}(x_{k}(t), x_{k}(t-\tau), t, i) - \sigma_{l}(x_{l}(t), x_{l}(t-\tau), t, i) \right]^{T} \\ \times \left[\sigma_{k}(x_{k}(t), x_{k}(t-\tau), t, i) - \sigma_{l}(x_{l}(t), x_{l}(t-\tau), t, i) \right]$$

$$\leq \lambda^{*} \left[(x_{k}(t) - x_{l}(t))^{T} \Sigma_{1}^{T} \Sigma_{1}(x_{k}(t) - x_{l}(t))) + (x_{k}(t-\tau) - x_{l}(t-\tau))^{T} \Sigma_{2}^{T} \Sigma_{2}(x_{k}(t-\tau) - x_{l}(t-\tau)) \right].$$
(20)

Moreover, it implies from Assumption 2 and Remark 1 that

$$e_{1} \begin{bmatrix} x_{k}(t) - x_{l}(t) \\ f(x_{k}(t)) - f(x_{l}(t)) \end{bmatrix}^{T} \begin{bmatrix} 0 & -K_{1}^{T} \\ -K_{1} & 2I \end{bmatrix} \begin{bmatrix} x_{k}(t) - x_{l}(t) \\ f(x_{k}(t)) - f(x_{l}(t)) \end{bmatrix} \leq 0,$$
(21)

$$e_{2} \begin{bmatrix} x_{k}(t-\tau) - x_{l}(t-\tau) \\ g(x_{k}(t-\tau)) - g(x_{l}(t-\tau)) \end{bmatrix}^{T} \begin{bmatrix} 0 & -K_{2}^{T} \\ -K_{2} & 2I \end{bmatrix} \begin{bmatrix} x_{k}(t-\tau) - x_{l}(t-\tau) \\ g(x_{k}(t-\tau)) - g(x_{l}(t-\tau)) \end{bmatrix} \leq 0$$
(22)

where $e_j > 0$ (j = 1, 2) and $k, l = 1, 2, \dots, N$.

Combining (20) with (22), it can be concluded that

$$\mathcal{E}\{\mathcal{L}V(x_{t},t,i)\} \leq \mathcal{E}\left\{\sum_{1\leq k< l\leq N} (\xi_{k}^{(i)}(t) - \xi_{l}^{(i)}(t))^{T} \Xi_{kl}^{(i)}(\xi_{k}^{(i)}(t) - \xi_{l}^{(i)}(t))\right\} \\ - \int_{t-\tau}^{t} \mathcal{E}\left\{(\xi_{k}^{(i)}(t,\beta) - \xi_{l}^{(i)}(t,\beta))^{T} \Omega_{kl}^{(i)}(\xi_{k}^{(i)}(t,\beta) - \xi_{l}^{(i)}(t,\beta))\right\} d\beta,$$

$$(23)$$

where $\Xi_{kl}^{(i)}$ and $\Omega_{kl}^{(i)}$ are defined in (10)–(11) and $\xi_k^{(i)}(t,\beta) - \xi_l^{(i)}(t,\beta) = \operatorname{col}\{(\xi_k^{(i)}(t) - \xi_l^{(i)}(t)), (y_k(\beta, r(\beta)) - y_l(\beta, r(\beta)))\}$.

From condition (11), we can see that

$$\mathcal{E}\{\mathcal{L}V(x_t, t, i)\} \le \lambda(\Xi) \mathcal{E}\left\{\sum_{1 \le k < l \le N} (\xi_k^{(i)}(t) - \xi_l^{(i)}(t))^T (\xi_k^{(i)}(t) - \xi_l^{(i)}(t))\right\},\tag{24}$$

where $\lambda(\Xi) = \max_{i \in \mathbb{S}} \{\lambda_{\max}(\Xi_{kl}^{(i)})\}.$

Under condition (10), it follows readily that $\lambda(\Xi)$ is a negative constant. Therefore, we have

$$\mathcal{E}\{V(x_t, t, i)\} - \mathcal{E}\{V(x_0, 0, r_0)\} = \int_0^t \mathcal{E}\{\mathcal{L}V(x_s, s, i)\} ds$$

$$\leq \lambda(\Xi) \int_0^t \mathcal{E}\left\{\sum_{1 \le k < l \le N} (\xi_k^{(i)}(s) - \xi_l^{(i)}(s))^T (\xi_k^{(i)}(s) - \xi_l^{(i)}(s))\right\} ds,$$
(25)

which implies that

$$\int_{0}^{t} \mathcal{E}\left\{\sum_{1 \le k < l \le N} \|\xi_{k}^{(i)}(s) - \xi_{l}^{(i)}(s)\|^{2}\right\} ds \le -\{\lambda(\Xi)\}^{-1} \mathcal{E}\{V(x_{0}, 0, r_{0})\}.$$
(26)

Moreover, under Assumption 2, we can obtain $V(x_0, 0, r_0) < +\infty$ and, subsequently, $\sum_{1 \le k \le l \le N} \|\xi_k^i(s) - \xi_l^i(s)\|^2$ is uniformly continuous on $[0, +\infty)$. Then, by using Lemma 3, we obtain

$$\mathcal{E}\left\{\sum_{1\leq k\leq l\leq N} \|\xi_k^i(s) - \xi_l^i(s)\|^2\right\} \to 0, \text{ as } t \to +\infty.$$

$$(27)$$

In other words, all the subsystem in genetic oscillator networks (2) are asymptotically synchronous in the mean square sense. This completes the proof of Theorem 1.

Remark 3: By taking advantage of a novel matrix functional and linear matrix inequality (LMI) techniques, the stochastic synchrony criteria have been derived in the form of LMIs for the genetic oscillator networks with time delays, Markovian switching parameters as well as stochastic disturbances. The LMI-based conditions can be readily checked by using the LMI toolbox in Matlab or other standard numerical software. An important feature of the reported results lies in that all the conditions are dependent on the upper bound of the timedelays, which is made possible by utilizing the most updated techniques for achieving delay dependence. Note that the main criteria involve 1) the matrix Γ_i that links the state variable of the *l*th genetic oscillator in the genetic network mode *i*; 2) the coupling matrix *W* that represents the coupling topology, direction, as well as the coupling strength of the genetic network. Therefore, the network topology does affect the dynamics of the synchronization.

IV. NUMERICAL EXAMPLE

In this section, we present a numerical example to illustrate the usefulness and applicability of the developed approach in this paper.

Consider the genetic oscillator network (2) or (5) with N gene oscillators and Markovian switching between two modes. The parameters are given as follows:

$$\begin{aligned} A(1) &= \begin{bmatrix} -1.0 & 0.6 \\ 0.5 & -1.5 \end{bmatrix}, \ A(2) &= \begin{bmatrix} -1.5 & 0.5 \\ 0.4 & -2.0 \end{bmatrix}, \ B(1) &= \begin{bmatrix} 1.8 & 1.0 \\ 1.2 & 1.4 \end{bmatrix}, \\ B(2) &= \begin{bmatrix} 1.4 & 1.2 \\ 1.0 & 1.0 \end{bmatrix}, \ C(1) &= \begin{bmatrix} 2.8 & 0.5 \\ 0.3 & 1.6 \end{bmatrix}, \ C(2) &= \begin{bmatrix} 2.5 & 0.35 \\ 0.25 & 0.9 \end{bmatrix}, \\ \Sigma_1 &= \begin{bmatrix} 0.1 & 0 \\ 0.1 & 0.1 \end{bmatrix}, \ \Sigma_2 &= \begin{bmatrix} 0.1 & -0.1 \\ 0 & 0.1 \end{bmatrix}, \ \Gamma_1 &= 1.5I, \ \Gamma_2 &= I, \\ \tau &= 2.5, \ \Pi &= \begin{bmatrix} -3 & 3 \\ 4 & -4 \end{bmatrix}, \ W &= [w_{kl}]_{6 \times 6} &= \begin{cases} 1, & k \neq l; \\ -5, & k = l. \end{cases} \end{aligned}$$

The nonlinear regulatory functions are of the following form

$$f_l(x_{kl}(s)) = \frac{x_{kl}^2(s)}{1 + x_{kl}^2(s)}, \quad g_l(x_{kl}(s-\tau)) = \frac{x_{kl}^2(s-\tau)}{2 + x_{kl}^2(s-\tau)},$$
(28)

and it can then be verified that $K_1 = 0.65I$ and $K_2 = 0.48I$.

By using the Matlab LMI toolbox, we solve the LMIs (9)-(11) to obtain a feasible solution as follows:

$$P_{1} = \begin{bmatrix} 8.7634 & -0.7823 \\ -0.7823 & 10.0816 \end{bmatrix}, P_{2} = \begin{bmatrix} 10.1496 & -0.9153 \\ -0.9153 & 11.5092 \end{bmatrix}, Q_{1} = \begin{bmatrix} 45.2497 & -10.3339 \\ -10.3339 & 57.1284 \end{bmatrix}, R = \begin{bmatrix} 0.1812 & 0.0154 \\ 0.0154 & 0.1775 \end{bmatrix}, \lambda^{*} = 34.0188, e_{1} = 26.4168, e_{2} = 30.9079.$$

Therefore, it follows from the Theorem 1 that genetic oscillator networks (2) or (5) with given parameters is globally synchronized in the mean square sense.

In order to confirm the theoretical results, the simulation study is carried out. Figs. 1–2 are plotted to show the evolution dynamics of mRNA concentrations of all the coupled gene oscillators with different random initial values. It can be seen clearly from both Figs. 1–2 and Fig. 3 that the time evolution of the stochastic synchronization error between all the coupled gene oscillators indeed approaches zero, and the expected stochastic synchrony is well achieved.



Fig. 1. Oscillation dynamics of the mRNA concentrations of $x_{k1}(t)$ $(k = 1, 2, \dots, 6)$ of all coupled oscillators with different random initial values.

V. Conclusions

In this paper, the stochastic synchronous analysis problem has been studied for a general array of genetic oscillator networks with Markovian switching, random perturbations as well as constant time-delays. We have established easily verifiable conditions under which the addressed delayed genetic oscillator network is asymptotically synchronization in the mean square sense in the presence of both Markovian jumping parameters and time-delays. By constructing a novel matrix functional based on the idea of 'delay-fractioning' and combining the stochastic analysis with the linear matrix inequality (LMI) technique, the delay-dependent synchrony criterion has been derived in the form of LMIs. A numerical example has been provided to demonstrate the effectiveness and applicability of the proposed testing conditions.

References

- M. Barrio, K. Burrage, A. Leier and T. Tian, Oscillatory regulation of Hes1: discrete stochastic delay modelling and simulation. PLoS Comput Biol 2(9): 1017–1030, 2006.
- [2] S. Boyd, L. E. Ghaoui, E. Feron and V. Balakrishnan, Linear Matrix Inequalities in System and Control Theory. Philadelphia: SIAM, 1994.
- [3] X. Cai and X. Wang, Stochastic modeling and simulation of gene networks. IEEE Signal Processing, 1: 27–36, 2007.



Fig. 2. Oscillation dynamics of the mRNA concentrations of $x_{k2}(t)$ $(k = 1, 2, \dots, 6)$ of all coupled oscillators with different random initial values.

- [4] Y. Cao, J. Lam and L. Hu, Delay-dependent stochastic stability and H_{∞} analysis for time-delay systems with Markovian jumping parameters. Journal of the Franklin Institute, 340: 423–434, 2003.
- [5] G. Chesi, Y.S. Hung, Stability analysis of uncertain genetic sum regulatory networks. Automatica, 44: 2298–2305, 2008.
- [6] B. Chen and P. Chen, Robust engineered circuit design principles for stochastic biochemical networks with parameter uncertainties and disturbances. IEEE Tran. on Biomedical Circuits and Systems, 2(2): 114–133, 2008.
- [7] A. Datta, A. Choudhary, M.L. Bittner and E.R. Dougherty, External control in Markovian genetic regulatory networks, Machine Learning, 52(1-2): 169-181, 2003
- [8] A. Datta, A. Choudhary, M.L. Bittner and E.R. Dougherty, External control in Markovian genetic regulatory networks: the imperfect information case, Bioinformatics, 20(6): 924-930, 2004.
- H. de Jong and R. Lima, Modeling the dynamics of genetic regulatory networks: continuous and discrete approaches. Lect. Notes Phys., 671: 307–340, 2005.
- [10] M.B. Elowitz and S. Leibler, A synthetic oscillatory network of transcriptional regulators. Nature, 403: 335–338, 2000.
- R.E. Mirollo and S.H. Strogatz, Synchronization of pulse-coupled biological oscillators. SIAM J. APPL. MATH., 50(6):1645– 1662, 1990.
- [12] M. Hecker, S. Lambeck, S. Toepfer, E. Someren, R. Guthke, Gene regulatory network inference: Data integration in dynamic models-A review. BioSystems, 96(1): 86–103, 2009.
- [13] I. Ivanov and E. R. Dougherty, Modeling genetic regulatory networks: continuous or disctete? Journal of Biological Systems, 14(2): 219–229, 2006.
- [14] A. Kuznetsov, M. Kaern and N. Kopell, Synchrony in a population of hystersis-based genetic oscillators. SIAM J. Appl.



Fig. 3. The time evolution of the stochastic synchronization error between all the coupled gene oscillators.

Math, 65(2): 392-425, 2004.

- [15] A. N. Langville and W. J. Stewart, The Kronecker product and stochastic automata networks. Journal of Computational and Applied Mathematics, 167: 429-447, 2004.
- [16] C. Li, L. Chen and K. Aihara, Synchronization of coupled nonidentical genetic oscillators. Phys. Biol. 3: 37–44, 2006.
- [17] C.Li, L. Chen and K. Aihara, Stochastic synchronization of genetic oscillator networks. BMC Systems Biology, 1: Article No. 6, 2007.
- [18] Y. Liu, Z. Wang and X. Liu, Exponential synchronization of complex networks with Markovian jump and mixed delays. Physics Letters A, 372: 3986–3998, 2008.
- [19] D. McMillen, N. Kopell, J. Hasty and J. J. Collins, Synchronizing genetic relaxation oscillators by intercell signaling, PNAS, 99(2): 679–684, 2002.
- [20] S. Mou, H. Gao, W. Qiang and K. Chen, New delay-dependent exponential stability for neural networks with time delay. IEEE Trans. Systems, Man, and Cybernetics - Part B, 38(2): 571-576, 2008.
- [21] J. Paulsson, Summing up the noise in gene networks. Nature, 427(29):415-418, 2004.
- [22] F. Ren, J. Cao, Asymptotic and robust stability of genetic regulatory networks with time-varying delays. Neurocomputing, 71: 834–842, 2008.
- [23] E. Someren, L. Backer and M. Reinders, Genetic network modeling. Lect. Notes Phys., 671: 307–340, 2005.
- [24] J.E. Slotine and W. Li, Applied Nonlinear Control. Prentice-Hall, New Jersy, 1991.
- [25] V. Shahrezaei and P. Swain, The stochastic nature of biochemical networks. Biotechnology, 19:369–374, 2008.
- [26] T. Tian, K. Burrage, Pamela M. Burrage and Margherita Carletti, Stochastic delay differential equations for genetic regulatory networks. Journal of Computational and Applied Mathematics, 205:696–707, 2007.
- [27] Y. Wang, Z. Wang and J. Liang, Global synchronization for delayed complex networks with randomly occurring nonlinearities and multiple stochastic disturbances. J. Phys. A: Math. Theor. 42(13), art. No. 135101, 2009.
- [28] Y. Wang, Z. Wang and J. Liang, A delay fractioning approach to global synchronization of delayed complex networks with stochastic disturbances. Physics Letters A, 372(9): 6066–6073, 2008.
- [29] Z. Wang, J. Lam, G. Wei, K. Fraser, and X. Liu, Filtering for nonlinear genetic regulatory networks with stochastic disturbances. IEEE Tran. on Automatic Control, 53(10): 2448–2458, 2008.

- [30] Z. Wang, H. Gao, J. Cao and X. Liu, On delayed genetic regulatory networks with polytopic uncertainties: robust stability analysis, *IEEE Trans. NanoBioscience*, 7(2): 154-163, 2008.
- [31] Z. Wang, F. Yang, D.W.C. Ho, S. Swift, A. Tucker and X. Liu, Stochastic dynamic modeling of short gene expression time series data, *IEEE Trans. NanoBioscience*, 7(1): 44–55, 2008.
- [32] W. Yu, J. Lü, G. Chen, Z. Duan and Q. Zhou, Estimating uncertain delayed genetic regulatory networks: an adaptive filtering approach, IEEE Trans. Automatic Control, 54(4): 892–897, 2009.
- [33] D. Yue and Q. Han, Delay-Dependent exponential stability of stochastic systems with time-varying delay, nonlinearity, and Markovian switching. IEEE Trans. Automat. Control, 50(2): 217–222, 2005.
- [34] D. Yue, J. Fang and S. Won, Delay-dependent robust stability of stochastic uncertain systems with time delay and Markovian jump parameters. Circuits, Systems and Signal Processing, 22(4):351–365, 2003.
- [35] Y. Zhao, H. Gao, J. Lam and B. Du, Stability and stabilization of delayed T-S fuzzy systems: a delay partitioning approach. IEEE Transactions on Fuzzy Systems, 17(4):750–762, 2009.

Yao Wang received his B.Sc. degree and M.Sc. degree in applied mathematics from Department of Mathematics, Yangzhou University, China, in 2004 and 2007, respectively. He is currently pursuing the Ph.D. degree in the School of Information Science and Technology at Donghua University, Shanghai, China. His research interests include dynamical systems, complex networks, systems biology, stochastic systems and time-delay systems. He is a very active reviewer for many international journals.

Zidong Wang was born in Jiangsu, China, in 1966. He received his B.Sc. degree in Mathematics in 1986 from Suzhou University, Suzhou, China, the M.Sc. degree in Applied Mathematics in 1990 and the Ph.D. degree in Electrical and Computer Engineering in 1994, both from Nanjing University of Science and Technology, Nanjing, China.

Currently, he is a Professor of Dynamical Systems and Computing at Brunel University in the UK. His research interests include dynamical systems, signal processing, bioinformatics, control theory and applications. He has published more than 120 papers in refereed international journals.

Prof. Wang is currently serving as an Associate Editor for 12 international journals including *IEEE Transactions on* Automatic Control, *IEEE Transactions on Neural Networks*, *IEEE Transactions on Signal Processing*, *IEEE Transac*tions on Systems, Man, and Cybernetics - Part C and *IEEE Transactions on Control Systems Technology*.

Jinling Liang received her B.Sc. degree in Mathematics in 1997 and M.Sc. degree in Mathematics in 1999, both from Northwest University, Xi'an, China, and the Ph.D. degree in 2006 in Applied Mathematics from Southeast University, Nanjing, China.

Dr. Liang is now an Associate Professor in the Department of Mathematics at Southeast University, Nanjing, China. From January 2004 to March 2004, she was a research assistant in the University of Hong Kong. From March to April 2004, she was a research assistant in the City University of Hong Kong. From April 2007 to March 2008, she was a postdoctoral research fellow with the Department of Information Systems and Computing at Brunel University in the UK sponsored by the Royal Society Sino-British Fellowship Trust Award of the UK.

Dr. Liang's research interests include neural networks, complex networks, nonlinear systems and bioinformatics. She has published around 20 papers in refereed international journals. She is a member of program committee for some international conferences and serves as a very active reviewer for many international journals.

Yurong Li received her master degree in industry automation and PhD in control theory and control engineering from Zhejiang University in 1997 and 2001, respectively. Now she is an assistant professor at Fuzhou University. And since 2007, she is the member of Fujian Key Laboratory of Medical Instrumentation & Pharmaceutical Technology. Her research interests include biomedical instrument and intelligent information processing.

Min Du received her PhD in electrical engineering from Fuzhou University in 2005. Now she is a professor and doctorial supervisor at Fuzhou University. And since 2007, she is the associate director of Fujian Key Laboratory of Medical Instrumentation & Pharmaceutical Technology. Her research interests include smart instrument and photoelectrical system.