University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

MEDI 9331 Scholarly Activities Clinical Years

School of Medicine

Fall 10-23-2020

Long non-coding RNA (IncRNA) as a new biomarker for hepatocellular carcinoma (HCC) progression and drug resistance

Areeb Masood

The University of Texas Rio Grande Valley, areeb.masood01@utrgv.edu

Sanjaya Satapathy

Northwell Health/ North Shore University Hospital

Manish Tripathi

The University of Texas Rio Grande Valley, manish.tripathi@utrgv.edu

Follow this and additional works at: https://scholarworks.utrgv.edu/som9331



Part of the Hepatology Commons, Neoplasms Commons, and the Oncology Commons

Recommended Citation

Masood, Areeb; Satapathy, Sanjaya; and Tripathi, Manish, "Long non-coding RNA (IncRNA) as a new biomarker for hepatocellular carcinoma (HCC) progression and drug resistance" (2020). MEDI 9331 Scholarly Activities Clinical Years. 7.

https://scholarworks.utrgv.edu/som9331/7

This Article is brought to you for free and open access by the School of Medicine at ScholarWorks @ UTRGV. It has been accepted for inclusion in MEDI 9331 Scholarly Activities Clinical Years by an authorized administrator of ScholarWorks @ UTRGV. For more information, please contact justin.white@utrgv.edu, william.flores01@utrgv.edu.

Long non-coding RNA (IncRNA) as a new biomarker for hepatocellular carcinoma (HCC) progression and drug resistance

Areeb Masood¹, Manish Tripathi, PhD¹, Sanjaya Satapathy, MD²

Abstract

Hepatocellular carcinoma (HCC) is a primary malignant liver tumor that commonly occurs as a progression of chronic liver inflammation due to long-standing viral hepatitis infection, toxins, metabolic conditions, and congenital disorders. In 2018, HCC was the 4th leading cause of cancer-related deaths worldwide, accounting for approximately 780,000 deaths in total.

HCC's prognosis is directly correlated with early detection. Unfortunately, HCC has an asymptomatic pattern of growth in the early stages of the disease which makes early detection challenging. When HCC progresses to advanced stages, it leaves clinicians with limited therapeutic and curative options, leading to high rates of morbidity and mortality. In general, one major hurdle in treating advanced cancer is that cancer cells develop drug resistance.

Research studies have elucidated the role of long non-coding RNAs (IncRNAs) in progression and drug resistance in various cancers such as bladder cancer, prostate cancer, and non-small cell lung cancer. Long non-coding RNAs (IncRNAs) are non-coding RNA segments that are >200 nucleotides long that play various roles in biological processes regarding DNA, RNA and protein regulation. In carcinogenesis, IncRNAs can promote growth and differentiation of cancer stem cell populations, promoting chemoresistance and tumor recurrence. The role of IncRNAs in chemoresistance of HCC has not been well-studied. There is a need to uncover novel IncRNA biomarkers for both the early detection of HCC and to create drug strategies for clinicians when predicting chemoresistance.

The goal of our review is three-fold. Firstly, we aim to describe HCC risk factors, the pathologic progression of HCC from benign pathologies such as viral hepatitis and non-alcoholic fatty liver disease (NAFLD), current diagnostic standards and challenges, and approved treatment options for HCC. Secondly, we aim to explain the molecular role of lncRNAs in cancer progression and drug resistance. Lastly, we aim to compile a comprehensive list of studied lncRNAs involved in HCC progression and drug resistance, their mechanisms of action, their binding partners, and the impact these lncRNAs had on metastatic potential and tumor proliferation. Additionally, we aim to include a list of proteins involved in HCC and their interactions with lncRNAs.

¹ Department of Immunology and Microbiology, University of Texas Rio Grande Valley School of Medicine, Edinburg, TX

² Department of Medicine, Northwell Health/ North Shore University Hospital, Manhasset, NY