

Measure, Topology and Probabilistic Reasoning in Cosmology[†]

Erik Curiel[‡]

October 18, 2014

A man said to the universe:
“Sir, I exist!”
“However,” replied the universe,
“The fact has not created in me
A sense of obligation.”

— Stephen Crane

ABSTRACT

I explain the difficulty of making various concepts of and relating to probability precise, rigorous and physically significant when attempting to apply them in reasoning about objects (*e.g.*, spacetimes) living in infinite-dimensional spaces, working through many examples from cosmology. I focus on the relation of topological to measure-theoretic notions of and relating to probability, how they diverge in unpleasant ways in the infinite-dimensional case, and are even difficult to work with on their own. Even in cases where an appropriate family of spacetimes is finite-dimensional, and so admits a measure of the relevant sort, however, it is always the case that the family is not a compact topological space, and so does not admit a physically significant, well behaved probability measure. Problems of a different but still deeply troubling sort plague arguments about likelihood in that context, which I also discuss. I conclude that most standard forms of argument used in cosmology to estimate the likelihood of the occurrence of various properties or behaviors of spacetimes have serious mathematical, physical and conceptual problems.

[†]I thank the members of the working group on Cosmology and Probability at the 2014 MCMP/Lausanne Summer School on Probability in Physics for invaluable feedback on a presentation of an earlier version of this paper, especially Jeremy Butterfield, Juliusz Doboszewski, Sam Fletcher, Karim Thébeault and Jos Uffink. I particularly thank Juliusz for catching errors in an earlier draft, and Sam for not letting me getting away with anything shady.

[‡]**Author’s address:** Munich Center for Mathematical Philosophy, Ludwigstraße 31, Ludwig-Maximilians-Universität, 80539 München, Germany; **email:** erik@strangebeautiful.com

Contents

1	Probabilistic Reasoning in Cosmology	2
2	Topology, Measure, Probability	5
2.1	Topological Spaces	5
2.2	Measure Spaces	8
2.3	Probability	10
3	Topology and Measure in Infinite-Dimensional Spaces	12
4	Topologies and Measures on Families of Spacetimes	13
4.1	Topologies	13
4.2	Measures	16
5	Genericity, Stability, and Prediction	19

1 Probabilistic Reasoning in Cosmology

There is, by any standard measure, exactly one actual cosmos, and its evolution cannot be repeated. It is, therefore, perhaps surprising when one first learns that probabilistic reasoning of various kinds pervades cosmology as a science—reasoning not just about the statistics of repeated and repeatable subsystems of the cosmos, but reasoning that purports to assign probabilities to uniquely global properties and structures of the cosmos itself. It should, therefore, perhaps not be surprising that problems arise for probabilistic reasoning in this context peculiar to it.

Physicists and philosophers have tended to focus on problems with probabilistic reasoning in cosmology that, in the end, boil down to one of the following two forms.

1. What can probability mean, when there is only one physical system of the type at issue to observe?
2. How can one justify attributions of definite values of probability when one cannot measure frequencies (because one cannot repeat experiments), which is to ask, what kinds of evidence may be available to try to substantiate attributions of probability?

I shall not address these sorts of questions and problems in this paper.¹ I shall rather address the relationship between topological and measure-theoretic methods in probabilistic reasoning and the problems that arise for it in the case of infinite-dimensional spaces, as naturally occur in cosmology.

Although it is far more common to associate the mathematical theory of measure spaces with probabilistic notions and reasoning, if one takes a broad-minded view of what counts as “probabilistic” reasoning, then, in many areas of physics, topological concepts and methods ground much of what it is reasonable to think of as probabilistic reasoning. This is particularly true in a science such as cosmology, in which well defined probability measures over families of systems are few and far between. In such situations, physicists often argue that a property or behavior of interest is typical or generic or stable in a family of possible systems, or is scarce or meagre or rigid, and so

¹See [Ellis \(2007\)](#) and [Smeenk \(2012a, 2012b\)](#) for excellent reviews and discussion of these questions and problems.

on, with no serious attempt to make those ideas quantitatively precise, though they clearly are intended to have probabilistic import. Often, the arguments are grounded on topological considerations with gestures at interpreting the conclusions in measure-theoretic terms so as to justify the intended probabilistic import.

Say we are interested in the likelihood of the appearance of a particular feature (having a singularity, *e.g.*) in a given family of spacetimes satisfying some fixed condition (say, being spatially open). If one can convincingly argue that spacetimes with that feature form a “large” open set in some appropriate, physically motivated topology on the family, then one concludes that such spacetimes are generic in the family, and so have high prior probability of occurring. If one can similarly show that such spacetimes form a meagre or nowhere-dense set in the family, one concludes they have essentially zero probability. The intuition underlying the conclusions always seems to be that, though we may not be able to define it in the current state of knowledge, there should be a physically significant measure consonant with the topology in the sense that it will assign large measure to “large” open sets and essentially zero measure to meagre or nowhere-dense sets. Similarly for stability and rigidity: if one can show that a given feature is topologically stable under “small” perturbations, one can conclude that the probability is very high that a spacetime approximately satisfying the relevant conditions will still have the feature; if the feature is topologically rigid under “small” perturbations, one can conclude that the probability is essentially zero that a spacetime approximately satisfying the relevant conditions will still have the feature. In order to justify the probabilistic nature of the conclusion, one again assumes the existence of an appropriate measure consonant with the topology in the sense that the smallness of the perturbation is to be judged by the fact that the resulting spacetime is in a neighborhood of the initial spacetime, of “small” measure.

In cosmology, reasoning of this form occurs ubiquitously, in the context of the following of kinds problem:

1. characterizing the likelihood of observing certain kinds of events, given the situation of possible observers in a spacetime, *i.e.*, the fact that observers are limited in observations to what lies in their past light-cone, by the sensitivity of their apparatus, by the amount of time a process will emit energy of a given magnitude or greater, and by how far to the past of the observers such processes may occur
2. characterizing the likelihood that the value of a universal constant lies within a fixed range
3. characterizing the likelihood that cosmological initial conditions of a particular kind or form, or having a particular property or characteristic, occurred
4. characterizing the likelihood that large-scale structure of a particular kind would form
5. characterizing the likelihood that a spacetime has a particular global (causal, topological, projective, conformal, affine, metrical) property

Common specific examples of such problems are:

1. characterizing the likelihood that observers such as ourselves would come to exist in the sort of spatiotemporal region we occupy in a spacetime of this sort

2. characterizing the likelihood that we are “typical” observers in the universe
3. characterizing the likelihood that the cosmological constant has any non-zero value, and has, moreover, a value near that actually observed
4. characterizing the likelihood of various “fine-tuning coincidences”: the seeming equality of densities of dark energy and dark matter in the current epoch, the approximate flatness of the observed universe, the approximate isotropy and spatial homogeneity of the observed universe, the seemingly required special entropic state of the very early universe, *etc.*
5. characterizing the likelihood that a spatially open spacetime is singular

In most branches of physics, one would address such problems by fixing an appropriate reference class of physical systems and a physically significant probability-measure on that class. When one cannot rigorously define such a measure, or one is not that interested in quantitative exactness, one will often rest content with arguing (or just stipulating) that a physically significant measure exists whose distribution of weight harmonizes in a particular way with a natural topology on the class of systems, to wit, one assumes that non-trivial positivity of measure is at least strongly correlated with openness of sets and likewise that smallness or nullness of measure is correlated with topological meagreness of sets. In this case, one will base one’s estimates of likelihood on the topological properties of the families of systems at issue.

Even in cases where one does have a well defined measure to give quantitative exactness to estimates of genericity or typicality, however, one still needs the measure to harmonize with an underlying topology in the appropriate way. The point is simple, though it does not seem to be widely appreciated or even recognized, either in the physics or the philosophy literature: genericity and typicality, roughly speaking, mean something like “most systems are similar in this respect” (and *mutatis mutandis* for meagreness and scarcity); “most”, however, is a measure-theoretic notion, whereas “similar in this respect” is a topological notion.² Most systems satisfy a property if the family of such systems forms a set of large measure; a given family of physical systems are similar in a given respect if the topological neighborhood-systems of the elements of the space representing the physical systems stand in some appropriate relation to each other, which often will be as simple as the fact that the family of elements representing the physical systems forms an open set.

In cosmology, however, the systems one most often focuses on are entire spacetimes, and families of spacetimes usually form infinite-dimensional spaces of a particular kind. And now one comes to the heart of the problem: it is a theorem (as I discuss in some detail in §3 below) that infinite-dimensional spaces of that kind do not admit non-trivial measures that harmonize in the right way with any underlying topology. It follows that one simply does not have available the kinds of reasoning normally employed to draw even qualitative conclusions about the likelihoods of properties or features or behaviors of spacetimes. To be clear, I do not claim that it is not possible to draw well grounded conclusions about such likelihoods, only that arguments of the standard forms cannot, not even in principle, be made rigorous, and so conclusions based on them are *prima*

²One can of course quantify similarity using a metric as well, but in this case the metric will give rise to a topology. The measure will still have to harmonize with the metric and so will automatically harmonize (or not) with the induced topology.

facie suspect, and should be treated with far more caution and skepticism than is common in the physics and philosophy literature. It is exactly the standard forms of argument, however, that cosmologists make when reasoning about likelihoods.

In §2, I quickly review the basics of topology, measure theory and probability theory, emphasizing technical and interpretative points that the rest of the paper relies on. Cognoscenti may want to skip that section, though I do discuss some issues (such as the character of topologies on spaces of functions, and the topological character of the uniqueness of the Lebesgue measure on \mathbb{R}^n) sometimes unfamiliar even to those with a solid grounding in topology and measure theory. I also present the basic facts about topology in a somewhat unusual way, based on the idea of an accumulation point, which is particularly suited to the goals of this paper. In my presentation of the basics of probability, moreover, I focus on those foundational problems most relevant to the kinds of cosmological argument I examine. In §3, I briefly rehearse the relevant aspects of topology and measure theory in the context of infinite-dimensional Fréchet spaces, and conclude with a statement of the fundamental theorem relevant to this paper and explain its import. In §4.1, I discuss the few well defined topologies on families of spacetimes commonly used in cosmology, and show that they have severe problems of physical interpretation on their own. In §4.2, I do the same for the only known example of a well defined measure on a finite-dimensional family of spacetimes of real physical interest. I conclude in §5 with a discussion of several standard cosmological arguments about likelihood in the context of infinite-dimensional spaces of spacetimes, and show how the reasoning runs afoul of the mismatch between topology and measure in such spaces.

2 Topology, Measure, Probability

2.1 Topological Spaces

A topology \mathfrak{T} is a family of sets, including the null set \emptyset , closed under arbitrary unions and finite intersections.³ In particular, the union \mathcal{T} of all elements of \mathfrak{T} itself belongs to \mathfrak{T} , and is called the topological space with topology \mathfrak{T} . The elements of \mathfrak{T} are its open sets; a neighborhood of a point of \mathcal{T} is a subset of \mathcal{T} , not necessarily in \mathfrak{T} , that contains an open set containing that point. In general, one can associate many different topologies with the same set of points \mathcal{T} . (We will, however, still abuse notation and terminology in the usual way when no ambiguity can arise, sometimes referring to a topological space simply by its associated set without specifying which topology on it we mean.)

As is always the case with mathematical fields of study, there are many ways to think about the subject of topology, both in the sense of intuitive visualization and in the sense of rigorous formalization. For our purposes, the sense in which topology captures the idea of the study of “continuity”—what remains invariant under deformations of a space that don’t rip or puncture it and don’t glue different parts together—is the most important.⁴ The neighborhoods of a topology capture an idea of relative proximity relevant to the idea of continuity: two points of the underlying set are in proximity (relative to the fixed topology) if the family of neighborhoods of one stands in one of a number of relations to the family of neighborhoods of the other. Intuitively speaking,

³All the material I cover in this section is developed with thoroughness and illuminating insight in [Kelley \(1955\)](#).

⁴A topologist is a person who doesn’t know the difference between a coffee-cup and a doughnut.

a neighborhood is a region of the space in which, at the point of which it is a neighborhood, “arbitrarily small perturbations” don’t take one out of the region. If one thinks of the topology as capturing something like a similarity relation among entities, then a neighborhood of an entity is a collection of other entities similar to the first to some degree.

For our purposes, one of the most important of these relations among families of neighborhoods is grounded on the idea of an accumulation point. Given any subset $O \subset \mathcal{T}$ (whether an open set or not), an *accumulation point* of O is a point p such that every neighborhood of p has a non-trivial intersection with $O - \{p\}$. In agreeably suggestive language, one may say that an accumulation point is “arbitrarily close” to its associated set. Much information about the topology of a topological space is encoded in the behavior of infinite sequences of points, and in particular by the situation of any accumulation points they may have. (Indeed, under mild restrictions, which all the examples we consider here satisfy, a topology can be fully characterized by the behavior of the accumulation points of all infinite sequences.) A set is *closed* if it contains all its accumulation points.

Given a sequence $P = \{p_i\}$ ($i \in \mathbb{I}^\uparrow$, the non-negative integers), we say P is *eventually in* a set O if there is an $m \in \mathbb{I}^\uparrow$ such that $p_n \in O$ for all $n > m$. Clearly, if there is a p such that a sequence P is eventually in every one of its neighborhoods, then p is an accumulation point of P . In this case, we say P *converges* to p .⁵ If a sequence converges at all, there may be more than one point the sequence converges to, depending on global properties of the topology. A topology is *Hausdorff* if every two distinct points have disjoint neighborhoods. In a Hausdorff space, if a sequence converges, its convergence point is unique.

A function from one topological space to another is *continuous* if the inverse image of an open set in the range is an open set in the domain: if you tell me how proximate you want to be to a point in the range, under the mapping, I’ll tell you how proximate you need to be to its pre-image in the domain. Under mild conditions on the topology, the continuity of a function can be characterized by the behavior of infinite sequences of points: roughly speaking, a function f is continuous if, for every sequence P in the domain that accumulates at a point p , the sequence $f[P]$ in the range accumulates at $f(p)$.

Whether or not a given mapping between two point-sets is continuous depends sensitively on the topologies one imposes on the sets. As a general rule, the fewer open sets a topology has, the easier it is for a function having the space as its range to be continuous; contrarily, the fewer open sets a topology has, the harder it is for a function having it as its domain to be continuous. The intuition behind this rough claim is easy to grasp: the more open sets there are, the harder it is for a sequence to have an accumulation point. Of two topologies on a given set, one is *finer* than the other if every one of its open sets is also an open set of the other. (One also says that the other is *coarser* than the one.) Finer topologies have more continuous functions from them; coarser topologies have more continuous functions to them.

One of the most central and important ideas in topology is compactness. The motivation behind

⁵A sequence may have an accumulation point it does not converge to. A sequence is *frequently in* a set O if, for every $m \in \mathbb{I}^\uparrow$, there is an $n > m$ such that $p_n \in O$. If a sequence is frequently in every neighborhood of a point, that point is a *cluster point* of a sequence. A cluster point is an accumulation point; a sequence may, but does not necessarily, converge to a cluster point. Roughly speaking, a sequence may ceaselessly approach arbitrarily close to and then recede from a cluster point, but never come to remain permanently near it.

the idea comes from the classic Heine-Borel Theorem. To state it, we need two more definitions. An *open cover* of a subset of a topological space is a family of open sets whose union contains the subset. A *subcover* of a cover is a proper subset of a cover that is also itself a cover.

Theorem 2.1.1 (Heine-Borel) *Every open cover of a closed, bounded interval of \mathbb{R} (under its standard topology) has a finite subcover.*

This is remarkable. No matter how large and fiendishly Baroque one makes an open cover of a closed, bounded interval, one can *always* select a finite number of elements from it that will still cover the interval. As with all the best theorems, the conclusion of the Heine-Borel Theorem has become a definition of fundamental importance: a subset of a topological space is *compact* if every one of its open covers has a finite subcover. (Of course, the entire space itself may be compact.)

Compact sets have particularly pleasant properties for our purposes, perhaps the two most important of which are that, first, under mild restrictions on the topology, every infinite sequence in a compact set has at least one accumulation point, and, second, under no restrictions at all on the topology, the Cartesian product of any family of compact spaces is itself compact under the natural product topology. (The latter statement is known as Tychonov’s Theorem.) Intuitively speaking, then, compact sets don’t “extend out to infinity”, and they also contain “every point they could possibly have had in the first place”—in a natural sense, they are bounded, and they don’t have any gaps or holes. An important weakening of the notion of compactness retains almost all its nice properties: a topological space is *locally compact* if every point has a compact neighborhood.

Finally, we record a few definitions and propositions that will play an important role in what follows. A subset D of a topological space is *dense* if every point of the space is an accumulation point of some sequence of points in D . Intuitively, D extends arbitrarily closely to every point of the space. The rational numbers, for example, form a dense subset of \mathbb{R} (indeed, a countable one). A topological space is *separable* if it has a countable dense subset. A subset of a topological space is *nowhere dense* if the union of the set and all its accumulation points do not contain an open set. If a subset N is nowhere dense, then, given any point not in N , one can find a neighborhood around that point such that no sequence in N accumulates on the neighborhood.

The case of most interest for us will be topologies on the family of continuous, differentiable or smooth functions between two topological spaces—in particular, the family of cross-sections on the fiber bundle of Lorentz metrics over a candidate spacetime manifold (connected, paracompact, Hausdorff, four-dimensional).⁶ Consider two topological spaces \mathcal{T}_1 and \mathcal{T}_2 , and the family of continuous functions \mathfrak{F} from the former to the latter. Define $\mathcal{N}(K, O) := \{f \in \mathfrak{F} : f[K] \subset O, \text{ for } K \subset \mathcal{T}_1 \text{ compact and } O \subset \mathcal{T}_2 \text{ open}\}$. The family of all such collections, for all such K and O , forms a subbase for the *compact-open topology* on \mathfrak{F} .⁷ A topology on \mathfrak{F} is said to be *jointly continuous* if the mapping $P : \mathfrak{F} \times \mathcal{T}_1 \rightarrow \mathcal{T}_2$ that takes (f, p) to $f(p)$ is itself continuous, in the product topology

⁶A topological space is *connected* if it is not the union of two open, nonempty, disjoint sets. The exact definition of paracompactness is too involved to give here; suffice it to say that it means the space is not “too big”. Indeed, one has to work hard to construct a topological space that is not paracompact (Hocking and Young 1988). In any event, a theorem due to Geroch (1969) shows that a manifold has a Lorentz metric only if it is paracompact, so we lose nothing by restricting attention to such manifolds.

⁷A *base* for a topology is a collection of open sets such that every other open set can be formed from a union of sets in the base. A *subbase* is a collection of open sets such that one can form a base by taking finite intersections of them.

on $\mathfrak{F} \times \mathcal{T}_1$.⁸ Say a topological space \mathcal{T} is *regular* if for every $p \in \mathcal{T}$ and every neighborhood N of p , there is a closed neighborhood U of p such that $U \subset N$. Then the following proposition captures the sense in which the compact-open topology is the coarsest mathematically reasonable topology to impose on a function space, so long as one wants that topology to respect the structure of the elements of the space *as functions*.

Proposition 2.1.2 *If the topological space \mathcal{T} is locally compact and regular, then the compact-open topology is the coarsest jointly continuous topology one can impose on the family of continuous functions from \mathcal{T} to any other topological space.*

Most relatively well behaved topologies—and all the ones we will consider here—are Hausdorff, separable, regular, and locally compact.

2.2 Measure Spaces

A σ -algebra is an ordered pair (\mathcal{S}, Σ) consisting of a set \mathcal{S} , and Σ , a non-empty collection of subsets of \mathcal{S} closed under the operations of finite set-differences and countable unions.⁹ Where no confusion can arise, we will often abuse notation in the standard way and refer to Σ itself as the σ -algebra. Write ‘ \mathbb{R}^\uparrow ’ for the set of non-negative real numbers.

Definition 2.2.1 *A measure on a σ -algebra Σ is a function $\mu : \Sigma \rightarrow \mathbb{R}^\uparrow \cup \{\infty\}$ such that*

1. $\mu(S) < \infty$ for at least one $S \in \Sigma$
2. for any countable, pairwise-disjoint family $\{S_i\} \subset \Sigma$,

$$\mu\left(\bigcup_i S_i\right) = \sum_i \mu(S_i)$$

A *measure space* is an ordered pair consisting of a σ -algebra and a measure on it; the elements of the σ -algebra are called *measurable sets*. It follows from the definitions that the null set \emptyset is always measurable, and any measure assigns value zero to it.¹⁰ In general, however, the null set will not be the only set assigned a measure of zero. We say a property that holds for all points of a measure space except for a subset of measure zero holds *almost everywhere*.

A σ -algebra has much the same feel about it as a topology, naturally giving rise to the question whether one can construct measures on topological spaces that relate in a natural way to the topology.

Definition 2.2.2 *Let \mathcal{T} be a Hausdorff, locally compact topological space. The Borel sets \mathcal{B} of \mathcal{T} consist of the smallest σ -algebra containing all its open sets. A Borel measure is a measure μ on the Borel sets such that $\mu(C) < \infty$ for every compact set C .*

⁸The product topology for the Cartesian product of two topological spaces is exactly what one would expect: all sets of the form $O_1 \times O_2$, where O_1 is an open set in the first factor and O_2 open in the second, constitute a base for the product topology.

⁹All the material I cover in this section is developed with thoroughness and illuminating insight in [Halmos \(1950\)](#).

¹⁰While it is not unusual, it is also not entirely standard to demand condition 1 for a measure. I do it because it simplifies matter greatly, in particular guaranteeing that the null set is measurable, of measure zero, without having to require it as a separate axiom. Also, it seems to me a quite reasonable bare minimum one should require of something one wants to call a measure, if it is to be useful in physics at all.

A Borel measure, in an obvious and natural sense, respects the topology of the underlying topological space.¹¹ It is a simple matter to construct a natural Borel measure on \mathbb{R} : one is uniquely picked out by the requirement that $\mu([a, b]) = b - a$ for every real interval $[a, b]$. This measure, suitably generalized to \mathbb{R}^n , is not however unique, and its multiplicity can be traced to the fact that it lacks one feature, completeness, that it is convenient to have: we say a measure is *complete* if every subset of every set of measure zero is itself measurable (and so necessarily of measure zero).

An *extension* of a measure space $(\mathcal{S}, \Sigma, \mu)$ is another measure space $(\mathcal{S}, \Sigma', \mu')$ such that $\Sigma \subset \Sigma'$ and $\mu'(A) = \mu(A)$ for all $A \in \Sigma$. The following is easily proven.

Proposition 2.2.3 *There is exactly one complete extension of the natural Borel measure on \mathbb{R}^n (for any $n \in \mathbb{I}^+$).*

Lebesgue measure μ_L on \mathbb{R}^n is the unique complete extension of the natural Borel measure. Any countable subset of \mathbb{R}^n has Lebesgue measure zero, but uncountable subsets also can. The Cantor Set is an example.

Let A be a subset of \mathbb{R}^n ; then, for any $p \in \mathbb{R}^n$, denote by ' $A + p$ ' the set that results by adding p to every element of A , sometimes called the p -translate of A . Now, the following proposition shows the most important properties of Lebesgue measure in relation to the natural topology and linear structure on \mathbb{R}^n .

Proposition 2.2.4 *Lebesgue measure is locally finite, strictly positive and translation invariant, i.e.:*

1. every $p \in \mathbb{R}^n$ has an open neighborhood O such that $\mu_L(O) < \infty$
2. $\mu_L(O) > 0$ for every non-empty open set O
3. for every measurable set A and every $p \in \mathbb{R}^n$, $\mu_L(A + p) = \mu_L(A)$

(Lebesgue measure is obviously not the unique measure satisfying these conditions, because there are non-complete Borel measures that also satisfy them.) The translation invariance of Lebesgue measure is commonly taken to be its most characteristic feature, to the point that any translation-invariant measure on any linear space is often referred to as a Lebesgue measure. The analogous property is particularly important in a measure that would be used to define a probability space over a family of events that itself has an appropriate algebraic structure, for reasons I discuss in §2.3 below.

The following theorem captures the precise sense in which Lebesgue measure is the unique measure that respects both the topology and the linear structure of \mathbb{R}^n .

Theorem 2.2.5 *Lebesgue measure is the unique complete, translation-invariant measure on the Borel sets in \mathbb{R}^n .*

¹¹Again, while not unusual, it is not wholly standard to demand that a Borel measure assigns finite measure to all compact sets. And, again, this seems to me the minimum one should require of such a thing for it to be usefully applicable in physics.

2.3 Probability

Measures allow for one elegant and important way to formalize the notion of probability: a *probability space* is an ordered pair consisting of a σ -algebra (\mathcal{P}, Π) and a measure μ on it such that $\mu(\mathcal{P}) = 1$. Intuitively speaking, the elements of \mathcal{P} represent the totality of possible outcomes for some family of phenomena we are interested in, those of Π the collections of outcomes to which it makes sense to assign probabilities, and the value assigned by μ to an element of Π the probability of that collection of outcomes. It is trivial to show that a probability space satisfies the standard Kolmogorov axioms of probability theory.

It is of fundamental importance to recognize that, when one wants to be precise and rigorous, it never makes sense to ask for “the probability” *simpliciter* of some event or collection of events. One must have in hand a probability space that includes the event or collection of events in its σ -algebra (or, at least, some structure formally equivalent to one). In general, for any given event or collection of events, there will be many, many, many such probability spaces, with different σ -algebras and with different measures. Picking the most appropriate σ -algebra for the question or investigation at hand is known as the *reference-class problem*. I am not aware of any standard name for the problem of picking the most appropriate measure, but it is equally as important and difficult in general as the reference-class problem. The kinds of consideration that should bear on those choices will depend on the nature of the subject matter one is treating, and on the nature of the problem concerning that subject matter. In physics, of course, when using measures to assign probabilities to collections of events, one wants to find a σ -algebra that represents “all appropriately similar events”, where the similarity has manifest physical significance for the problem at issue, and to fix a measure on it that captures a property of real physical significance shared by the events that relates in a clear, direct, determinate way to the probabilities one wants to characterize.¹² Without making sure that the measure latches on to and respects a physically significant feature of the problem space with manifest relevance to the determination of probabilities, there will be no reason to think of the values the measure assigns as representing real physical probabilities. All these issues play a crucial role in attempts to evaluate the soundness of many kinds of argument in cosmology.

Although measures on their own can be used to define probability spaces, it is often the case that topological considerations play an important role in probabilistic reasoning. It is almost always desirable, for instance, especially in physics, for an appropriate probability measure to be a Borel measure, and in particular to assign non-zero probability to any collection of outcomes that forms an open set in a physically natural topology. This captures the idea that, if an event has non-zero probability (measure greater than zero), then “arbitrarily small” perturbations of it shouldn’t render the result impossible, *i.e.*, send it into a set of measure zero. One can guarantee this by having the original event lie in an open set, which, because the measure is Borel, will have non-zero measure. If this were not the case, then, given the necessarily limited precision of observations in physics, we would find ourselves in the position of predicting outcomes with non-zero probability that we could never in principle observe.

Topology plays other important roles in probabilistic reasoning as well. In many cases, the

¹²Peirce (1878a, 1878b) gives a particularly beautiful discussion of these issues, although of course he does not use the language of measures.

quantitative exactness delivered by a measure is either not feasible or not desirable. Sometimes it is enough merely to know that an event is very likely or not likely at all, without attaching a quantitatively exact probability to it. Let's say that we make a prediction for the dynamical evolution of a system starting from a set of exact initial conditions. We want to know how likely it is that the system, if prepared with approximately those initial conditions will evolve in approximately the predicted way. (Roughly speaking, this is known as the Hadamard stability problem for the initial-value formulation of the system.) One natural way to make the question precise is to find appropriate topologies for the space of initial conditions and the space of dynamical evolutions, define the mapping taking initial conditions to dynamical evolutions, and determine whether it is jointly continuous. ("Do arbitrarily small perturbations of the initial conditions leave the later dynamical behavior essentially unchanged?") If so, then, if the prediction is sound and if we have good reason to believe that the system starts with initial conditions close enough to the exact initial conditions used to generate the prediction, it is very likely that we will get the expected behavior even though we know that, due to the finite exactness of measurement and preparation, the system almost certainly did not start with those exact initial conditions. If the mapping is not jointly continuous, then it may be very unlikely that we will get the expected behavior, no matter how close to the exact initial conditions the system starts evolving from. (This is one of the reasons why it is almost always desirable, from a physical point of view, to have one's function-space topology be jointly continuous.)

In a closely related vein, say that we have found an appropriate topology for the space representing the possible states of a type of system, and that, moreover, the points of that space representing the system as possessing a certain property with values in a fixed range form an open dense subset.¹³ Then there is a natural sense in which it is overwhelmingly likely that an appropriately random sampling of such systems will all evince values for the given property falling within the fixed range. If the subset is nowhere dense, it is very unlikely to find a system having the property with value in the fixed range in an appropriately random sample.¹⁴ Of course, one must bear in mind that such conclusions depend not only on the physical propriety of the topology one has fixed on the space of states, but, at least as importantly, they depend on the propriety of the mechanism one has chosen to construct the random sample. If one's sampling mechanism is biased in some way, then it doesn't matter how "appropriate" one's measure is: one will not get physically reliable results. Again, these issues play a crucial role in the evaluation of many kinds of cosmological argument.

Finally, one can use measures in similar ways to draw qualitative judgments about the likelihood of an event or kind of event: a property that holds almost everywhere in an appropriate measure space will be very likely to occur, and one that holds in a set of measure zero will be very unlikely, even if the measure is not a probability measure, so long as one has been able to demonstrate an appropriate relation between the measure and the relevant physical properties of the system at issue. It is only in the latter case that the value the measure attributes to a set may reasonably

¹³If the topology arises from or is compatible with a complete metric, one can use the more general criterion that the set be a G_δ -set, *i.e.*, that it be a countable intersection of open dense subsets, in so far as the complement of a G_δ -set in this case is nowhere dense.

¹⁴If the topology arises from or is compatible with a complete metric, one can use the more general criterion that the set be *meagre*, *i.e.*, that it be a countable union of nowhere dense subsets, in so far as the complement of a meagre set in this case is dense. (This is known as the Baire Category Theorem.)

be thought of as a representation of a real physical probability for the class of events in the set.

3 Topology and Measure in Infinite-Dimensional Spaces

Now, as we have seen, there is a natural sense in which, in \mathbb{R}^n , Lebesgue measure respects both the topology and the linear structure, both of which are desirable features in a measure one wants to found probabilistic reasoning on, for the reasons discussed in §2.3, *inter alia*. The situation in infinite-dimensional spaces, therefore, as we will see, poses considerable problems for the hopeful physicist. This matters because, for our purposes—the application of probabilistic reasoning to families of spacetimes—the natural spaces one works with are spaces of functions (Lorentzian metrics on differential manifolds). These spaces tend to be infinite-dimensional spaces with natural algebraic structures accruing to them.

To apply probabilistic reasoning to families of spacetimes, one must first choose what sort of spacetime metric one is going to work with, C^n or C^∞ . Each has virtues and demerits. To see what is at issue in a simpler setting, consider the set of functions on the unit disk. For C^n functions, we have the norm

$$\|f\| = \sup |f| + \dots + \sup |\nabla^{(n)} f|$$

resulting in a Banach space, since this norm is complete for any finite n —because the disk is compact and the functions are continuous, the supremum is always finite. (A *Banach space* is a normed vector space, complete with respect to the metric the norm induces.) For C^∞ , we do not get a Banach space because the resulting infinite sum may not converge. Instead, we define

$$\langle\langle f \rangle\rangle = \frac{\sup |f|}{1 + \sup |f|} + \dots + \frac{1}{2^n} \frac{\sup |\nabla^{(n)} f|}{1 + \sup |\nabla^{(n)} f|} + \dots$$

which manifestly converges. This operation defines a metric in the obvious way,

$$(f, g) = \langle\langle f - g \rangle\rangle$$

and this metric is complete in the sense that all its Cauchy sequences converge to a point in the space. This operation, however, does not define a norm, since it is not the case that

$$\langle\langle a f \rangle\rangle = |a| \langle\langle f \rangle\rangle$$

for scalar a . (There is no norm for C^∞ functions that both accounts for all their derivatives and, in a natural sense, extends the norm for C^n functions.) The metric is, however, manifestly invariant with respect to translations. The resulting space is a *Fréchet space*, a metrizable, locally convex vector space, complete with respect to a translation-invariant metric.

Is it true in such a large space that every well-behaved vector field has unique integral curves (a necessity, *e.g.*, for certain forms of local stability analysis)? In a Banach space, yes, but in a Fréchet space, no. (Sometimes there are no integral curves, sometimes they are not unique). As an example of the way things can go awry, consider a map f from functions on the disk to functions on the disk:¹⁵ fix a vector field ξ^a on the disk; then define

$$\phi(f) = \xi^n \nabla_n f$$

¹⁵I thank Bob Geroch for conversations in which we worked out the details of this example.

ϕ wants to be a linear functional on our space of functions, *i.e.*, to be a vector field on the vector space of functions on the disk, but, while it is in fact a vector field on the Fréchet space of C^∞ functions, it is not one on the Banach space of C^n functions, since for $f \in C^n$ it is not necessarily the case that $\nabla_a f \in C^n$. On the Fréchet space, however, the vector field resulting from this mapping does not yield unique solutions—when one slides a function along the disk, as the mapping in effect asks one to do, one gets to make up whatever one wants to fill up the “back” part of the disk, as the “front” part of the function slides off the disk, *i.e.*,

$$\frac{d}{dt}f(t, 0) = \xi^n \nabla_n f(t, 0)$$

has no unique integral curves. So one can have unique integral curves, but no vector field (on Banach space), or a vector field but no unique integral curves (on Fréchet space).

In general, however, there may be difficulties with defining tangent vectors to curves on a Fréchet space. In a Banach space B , $\gamma : \mathbb{R} \rightarrow B$ has derivative $v \in B$ if, $\forall \epsilon > 0$, $\exists c > 0$ such that

$$\|\gamma(t) - \gamma(t_0) - (t - t_0)v\| \leq c|t - t_0|^2$$

for $|t - t_0| < \epsilon$, but in a Fréchet space one has no norm, only a metric, and if one uses the metric to try to define derivatives, nice properties like “sum of two differentiable vector fields is a differentiable vector field” will likely fail since one does not have the nice norm properties.

Now, once one has bitten the bullet and chosen to work with the type of space one considers the lesser of two evils, one might have thought that all one’s travail would be behind one. Sadly, no.

Theorem 3.1 *The only locally finite, translation-invariant Borel measure on an infinite-dimensional, separable Fréchet space is the trivial measure (viz., the one that assigns measure zero to every measurable set).*

Thus, since a Banach space is automatically a Fréchet space, any translation-invariant measure on any reasonably well behaved infinite-dimensional space assigns infinite measure to all open sets, unless the measure is the trivial measure.

4 Topologies and Measures on Families of Spacetimes

Even though, as should now be clear, we cannot hope for the most satisfactory framework—a well behaved Borel measure—on which to found probabilistic reasoning about families of spacetimes, we may still hope to find topologies or measures on their own appropriate for addressing specific sorts of problems. Perhaps, the hope goes, we can find a well behaved, physically significant measure that will return probabilities in such a way that its lack of relation to a topology will not necessarily lead to conundrums or implausibility. Or perhaps we can find a topology that, though not related to a measure, will still allow us to reason qualitatively about likelihoods in a physically significant way. Alas, in the event, things do not look good.

4.1 Topologies

Fix a candidate spacetime manifold M , and consider the family \mathfrak{G} of all Lorentz metrics on it, *i.e.*, the family of all cross-sections of the fiber bundle of Lorentz metrics over M . There are two

standard topologies relativists impose on \mathfrak{G} when addressing problems related to likelihoods. The first is a standard compact-open topology on a function space; the second is a standard Whitney topology on a function space.¹⁶ (The compact-open is strictly coarser than the Whitney, unless M itself is compact, in which case the two coincide; we will not consider that case.) The idea behind each is to fix a standard of “distance” between Lorentz metrics by fixing an arbitrary positive-definite metric on M and using it to assign magnitudes to the algebraic differences of Lorentz metrics. As we shall see, both have severe problems of physical interpretation, which can in large part be traced to the fact that the positive-definite metrics themselves used to fix the similarity relations among Lorentz metrics have no physical significance.¹⁷

Roughly speaking, the compact-open topology cares only whether or not metrics are similar on bounded regions in the interior of the spacetime manifold; it does not care about their relative asymptotic behavior. To characterize it, we must define the neighborhoods of a given Lorentz metric g_{ab} . A neighborhood $\mathcal{N}(h_{ab}, K, \epsilon; g_{ab})$ is determined by a positive-definite metric h_{ab} on M , a compact subset K of M , and a real number $\epsilon > 0$. A Lorentz metric g'_{ab} is in the neighborhood if and only if $h^{mn}h^{rs}(g_{mr} - g'_{mr})(g_{ns} - g'_{ns}) < \epsilon$ everywhere in K . The family of all such neighborhoods forms a subbase for the compact-open topology.¹⁸ The compact-open topology has the pleasant properties of being locally compact, Hausdorff and regular (because the fiber bundle of Lorentz metrics over M is). It also, according to proposition 2.1.2, is the coarsest mathematically reasonable topology to use on \mathfrak{G} .

An example from Geroch (1971) shows, however, that its physical significance is questionable at best. Consider the sequence of metrics on \mathbb{R}^4 of the form $\text{diag}(t_m, -1, -1, -1)$, for $m \in \mathbb{I}^+$ (the strictly positive integers), where $t_m := 1 + \frac{m}{1 + (x - m)^{1/2}}$, x being a global Cartesian spacelike coordinate function. Roughly speaking, each of these metrics is essentially flat almost everywhere except for a sharp peak of curvature around the t - y - z -hypersurface defined by $x = m$. As m increases, moreover, this peak of curvature becomes higher and sharper, as it moves further out along the x -axis. It does not seem physically reasonable that such a sequence should converge to Minkowski spacetime, since the spacetimes in it have curvature that in a sense one can make precise grows without bound, and yet that is what it does under the compact-open topology.¹⁹ The problem is that the compact-open topology is too coarse—it does not have enough open sets to stop pathological sequences from converging.

Roughly speaking, the Whitney topology cares whether or not metrics are similar on the entire spacetime manifold, including their relative asymptotic behavior. A neighborhood $\mathcal{N}(h_{ab}, \epsilon; g_{ab})$ of a given Lorentz metric g_{ab} is determined by a positive-definite metric h_{ab} on M , and a real number

¹⁶Geroch (1971) calls the former the coarse and the latter the fine topology. Hawking (1971) calls the latter the open topology, and calls a yet finer topology the fine topology. We shall not consider here the fine topology of Hawking (1971), for, as we shall soon see, the Whitney topology already has “too many open sets”. Another common class of topologies used in relativity theory are the Sobolev topologies, which play an important role in the analysis of the Cauchy problem in general relativity (Ringström 2009). Since these are even finer than the finest one Hawking (1971) considers, and since we shall not discuss the Cauchy problem, we shall, again, not worry about them.

¹⁷See Geroch (1967, 1971) and Fletcher (2015) for insightful discussions of these problems.

¹⁸One can also form compact-open topologies that account for derivatives of the Lorentz metrics and how they differ, but we will not need to do so.

¹⁹This sequence of metrics does not converge at all under the Whitney topology, which one may perhaps think of as the “correct” or “naturally expected” result.

$\epsilon > 0$. A Lorentz metric g'_{ab} is in the neighborhood if and only if $h^{mn}h^{rs}(g_{mr} - g'_{mr})(g_{ns} - g'_{ns}) < \epsilon$ everywhere in M . The family of all such neighborhoods forms a subbase for the topology.²⁰ The Whitney topology also has the pleasant properties of being locally compact, Hausdorff and regular (because the fiber bundle of Lorentz metrics over M is).

The Whitney topology fairs even worse than the compact-open one with regard to physical significance, as examples from Geroch (1970, 1971) again show. Consider the sequence of metrics on \mathbb{R}^4 of the form $\text{diag}(t_m, -1, -1, -1)$, for $m \in \mathbb{I}^+$, where now $t_m := 1 + \frac{1}{m^2 + x^2 + y^2 + z^2}$. Each metric in this family is essentially flat almost everywhere except for a spherically symmetric bump of curvature centered on the origin; this bump, moreover, decreases smoothly to zero as m increases. This sequence, however, does *not* converge to Minkowski spacetime under the Whitney topology. Even more egregiously, the one-parameter family of metrics $\{\lambda g_{ab}\}$, for $\lambda \in \mathbb{R}^+$ (the set of strictly positive real numbers), where g_{ab} is *any* Lorentz metric on *any* non-compact M , fails to be a continuous curve under the Whitney metric. But each metric in the family represents the same physical spacetime! Multiplying a spacetime metric by a constant does nothing other than change the effective units one (implicitly) uses to quantify physical magnitudes such as mass and acceleration.²¹ The problem now is that the Whitney topology is too fine—it has so many open sets that almost no reasonable sequence will converge.

One could perhaps argue with some justice that Geroch’s example speaking against the compact-open topology is not so bad as to preclude its usefulness in many cases and for many purposes, and I would not necessarily disagree. The problem arises in the example because the compact-open topology, roughly speaking, does not contain enough open sets to control the similarity relations between metrics with respect to their asymptotic behavior. In other words, it does not care about global similarity, only local similarity. For the sorts of problems for which one would want to use a topology on a family of spacetime metrics to ground qualitative probabilistic reasoning in the context of cosmology, however, it is exactly the global similarity of metrics that will in general be at issue. We will see physically important examples of this in §5 below. The Whitney topology rules unhelpfully in such simple and fundamental cases as to make it, to my mind, never a viable option for addressing global issues.²²

It will be useful to conclude the section by considering problems with these topologies in the context of a more physically interesting example, to substantiate my claims. Consider the question of the stability of the occurrence of singularities in the family of metrics over a given manifold. One wants to show that the occurrence of a singularity is (topologically) stable “under small perturbations”. (I discuss this question, and the problems facing attempts to address it, in more detail in §5 below.) In this case, the impropriety of the Whitney topology can be easily illustrated

²⁰One can also form Whitney topologies that account for derivatives of the Lorentz metrics and how they differ, but again we will not need to do so.

²¹The compact-open topology seems to get both of these examples right: under it, the sequence converges to Minkowski spacetime, and the one-parameter family forms a continuous curve.

²²One may thus wonder about the usefulness of the Sobolev measures used in analyzing the stability of the Cauchy problem in general relativity, for those topologies are always strictly finer even than the Whitney topology. In fact, though, in this case the promiscuity of the topologies is a virtue—one *wants* to show stability under as difficult conditions as possible. Even though one may not be able to elucidate the physical significance of the Sobolev topologies, they are surely finer than any topology one will be able so to elucidate, and so stability under the Sobolev topologies ensures stability under more restrained, more physically plausible ones, whatever they may be.

by noting that any reasonable sense of “small perturbation” will yield an operation discontinuous with respect to it. For example, given a spacetime (\mathcal{M}, g_{ab}) , one might define a small perturbation as follows. Consider a one-parameter family of spacetimes $\mathfrak{M}_\epsilon := \{(\mathcal{M}, (1 + \phi_\lambda)g_{ab}) : \lambda \in [0, \epsilon)\}$, for some small ϵ , where each ϕ_λ is a non-negative smooth function on \mathcal{M} such that $0 \geq \sup_{\mathcal{M}} \phi_{\lambda'} < \lambda' < \sup_{\mathcal{M}} \phi_\lambda < \lambda$, for all $\lambda', \lambda \in [0, \epsilon)$, and the family of functions $\{\phi_\lambda\}$ varies smoothly with respect to λ in the supremum norm, and the supremum approaches zero “slowly”.²³ Then

$$\left(1 + \frac{d\phi_\lambda}{d\lambda} \Big|_{\lambda=0}\right) g_{ab}$$

is a small perturbation off g_{ab} . It is easy to see by construction that \mathfrak{M}_ϵ forms an everywhere discontinuous curve on the family of metrics with respect to the Whitney topology, and so *any* property of (\mathcal{M}, g_{ab}) one may want to consider is trivially stable under such small perturbations. (The only physically reasonable “small perturbation” continuous with respect to the Whitney topology is the identity operation.)

Non-trivial small perturbations defined in this way can easily be constructed so as to be continuous with respect to the compact-open topology, so this looks initially more promising. For the treatment of singularities, though, the compact-open topology is not physically appropriate. If one characterizes a singularity by the existence of incomplete, inextendible causal geodesics, then the compact-open topology will never be able to discriminate singular from non-singular metrics: it is only in highly pathological cases that incomplete, inextendible geodesics are contained in compact subsets of a spacetime (Curiel 1999). Every open neighborhood of a singular metric in the compact-open topology contains non-singular metrics, and vice-versa. There are no other well defined topologies on the family of Lorentz metrics standardly used by physicists.²⁴

4.2 Measures

The space of Lorentz metrics over a manifold—the family of possible spacetime models having that as its underlying manifold—is an infinite-dimensional Fréchet space, and so by theorem 3.1 it has no non-trivial Borel measure. The only rigorously defined measure on a reasonably interesting family of spacetimes is the Gibbons-Hawking-Stewart (GHS) measure μ_{GHS} on “minisuperspace” Γ (Gibbons, Hawking, and Stewart 1987). Roughly speaking, Γ comprises the family of initial data for FLRW spacetimes with compact Cauchy surfaces, sourced by a minimally coupled homogeneous scalar field. A little more precisely, one constructs the constraint-reduced phase space for an appropriately gauge-fixed Hamiltonian formulation of general relativity; restricting attention to compact 3-geometries sourced by homogeneous, minimally coupled scalar fields, one finds that the resulting space, remarkably, simplifies to the point that it itself is only four-dimensional. In essence, the reduced phase space is fully parametrized by the field-intensity ϕ of the scalar field and the Hubble expansion factor a on a Cauchy surface, and their “time-derivatives”, $\dot{\phi}$ and \dot{a} , off

²³Since the family of Lorentz metrics is a locally convex space, all the $(1 + \phi_\lambda)g_{ab}$ will also be Lorentzian for small enough ϵ .

²⁴In order to try to address such problems with the compact-open and the Whitney topologies, Fletcher (2014) constructs a novel topology, in some ways similar to the compact-open topology but which yields the “natural” answers for Geroch’s examples I discussed above. It would be of great interest to determine whether Fletcher’s topology is appropriate for the characterization of the stability of singularities not confined to compact subsets of spacetime.

the Cauchy surface; these quantities are constant on the Cauchy surface by homogeneity of the spacetime. The standardly defined Liouville measure on this phase space (modulo a few technical difficulties that do not concern us) is the GHS measure. All of a sudden, things are looking up for our eternally hopeful cosmologist who would engage in probabilistic reasoning, at least with regard to this (admittedly quite restricted, but still physically important) family of spacetimes—we have a rigorously defined Borel measure on a finite-dimensional space. (A Liouville measure is always a Borel measure.²⁵) It is not long, however, before a bucket of cold water is dashed in her face with the realization that, even though the space is finite-dimensional and the measure is Borel, it cannot be turned into a probability measure, for the measure it assigns the entire phase space is infinity— Γ is not compact.

Still, let us see whether we may not salvage something useful from this mess. We want to see whether the GHS measure can support any, even weak, form of probabilistic reasoning. Say we want to determine whether we can meaningfully attribute a probability to the occurrence of a physical property X , given the fixed reference class Γ . Let $P_X \subset \Gamma$ be the family of spacetimes evincing X . There are four cases to consider.

1. P_X is not measurable
2. $\mu_{\text{GHS}}(P_X) < \infty$
3. $\mu_{\text{GHS}}(\Gamma \setminus P_X) < \infty$
4. $\mu_{\text{GHS}}(P_X) = \infty$ and $\mu_{\text{GHS}}(\Gamma \setminus P_X) = \infty$

In the first case, we can say nothing at all, but one assumes or stipulates or hopes or demands or pleads or dreams that physically significant properties will not manifest such topological pathology in their distribution across spacetimes. In the second case, one can unambiguously attribute a probability of zero to it, and in the third a probability of one. In the fourth, one can say nothing simple or straightforward, without ambiguity, but now one does not even have the solace of yelling at the property and demanding that it not be pathological, as in the first case, for there is nothing pathological about such topological behavior at all.

There is, however, a “natural” schema for regularization procedures that one can use to try to derive a finite probability in such cases.²⁶ One approximates P_X by a nested sequence of finite-measure subsets of Γ , such that the union of the sequence is P_X and the sequence of measures of the subsets, appropriately weighted, converge to a finite value in $[0, 1]$:

1. assume Γ is σ -finite (*i.e.*, is a countable union of subsets of finite measure)
2. find “physically appropriate” nested sequence of subsets of Γ , $\{S_i\}$ ($i \in \mathbb{I}^+$), such that $\Gamma = \bigcup_i S_i$ and $\mu_{\text{GHS}}(S_i) < \infty$
3. define $\text{Pr}(P_X) = \lim_{i \rightarrow \infty} \frac{\mu_{\text{GHS}}(P_X \cap S_i)}{\mu_{\text{GHS}}(S_i)}$

²⁵Since the final, four-dimensional space does not naturally have the structure of a cotangent bundle, and so the symplectic structure one uses is not perspicuously related to the underlying topology, it is actually not clear to me that the standardly constructed Liouville measure ends up in this case being a Borel measure. It strikes me as plausible that it is, so I will assume so for the sake of argument. It would be an interesting exercise, which I leave to the reader, to verify this.

²⁶I follow here the exposition of [Schiffrin and Wald \(2012\)](#).

Minisuperspace is σ -finite, so we're off to a good start. The serious problem arises with the second condition: one can get pretty much any answer one wants by judicious choice of $\{S_i\}$, *i.e.*, different regularization procedures can yield wildly different results.

A simple example illustrates the general form of the problem. What is the probability that a randomly chosen natural number is even? *Prima facie*, the question makes no sense. Let's fix a regularization procedure to attempt to address it. Let S_i be the subset consisting of the first i natural numbers, in their normal ordering; then the regularization procedure yields the well defined probability $\frac{1}{2}$ for a natural number's being even. Now, however, order the natural numbers as follows, $\{1, 3, 2, 5, 7, 4, \dots\}$, and again let S_i be the subset consisting of the first i numbers. This yields a well defined probability, but now it is $\frac{1}{3}$.

In cosmology, the problem is nicely illustrated by attempts to calculate the probability of inflation for spacetimes in Γ . Using regularization procedures derived from arguments based on (topological) stability of initial conditions yielding "slow-roll" inflation, Gibbons and Turok (2008) deduced extremely low probability for $N \gg 1$ e -foldings of inflation, whereas Carroll and Tam (2010) deduced extremely high probability for $N \gg 1$ e -foldings of inflation. Both analyses, moreover, have strong, physically plausible justifications for the regularization procedures they employ (*i.e.*, their choice of $\{S_i\}$). The resolution to this seemingly paradoxical state of affairs is that they each used a different criterion for topological stability for initial conditions yielding inflation in fixing the choice of $\{S_i\}$. Roughly speaking, Carroll and Tam (2010) characterized topological stability based on the behavior of spacetimes entering an inflationary phase, whereas Gibbons and Turok (2008) did so based on spacetimes leaving the inflationary phase. This difference naturally leads them to consider the weight the GHS measure assigns to physically quite different open sets in Γ . It should therefore be no surprise that those open sets get assigned divergent weights.²⁷

Even in the cases where one can unambiguously attribute a probability to the occurrence of a property based on μ_{GHS} , one must ask about the physical significance of that probability, which, if it indeed has any, must come from the physical significance of the GHS measure itself, if it indeed has any. Schiffrin and Wald (2012) argue persuasively on multiple grounds that, at best, much work must be done to justify the physical significance of the GHS measure, and, at worst, it has none. In particular, they note that the standard justifications for the use of a Liouville measure are given by arguments based on special properties of the dynamical evolution of the system at issue and in particular on how it equilibrates. In particular, the arguments rely on the fact that the amount of time the system spends in a portion of phase space is proportional to its Liouville measure. Those arguments, however, are not available when:

1. the system is not ergodic
2. OR one has not waited a time much greater than the equilibration time after the system was prepared
3. OR the system has a time-dependent Hamiltonian that varies on a timescale that is small or comparable to the equilibration time

²⁷My diagnosis of the conflict between the two conclusions is in some ways similar to that of Schiffrin and Wald (2012), but also differs in one important way, *viz.*, my emphasis on the role topological stability plays in their arguments in fixing the open sets whose measures are relevant to the problem.

All of those conditions hold, however, for the canonical “dynamics” of that sector of general relativity represented by minisuperspace and its Hamiltonian. The system is not ergodic because the phase space has infinite measure—the dynamics cannot adequately explore the whole space in any finite time. For the same reason, there is no finite time in which the system can explore enough of the phase space in order for one to be able to conclude that it has satisfactorily equilibrated: the system always “remembers” its initial state, which precludes true statistical equilibration. That the time-dependent Hamiltonian in this case varies over timescales small compared to the equilibration time follows for the same reason.

In the face of these problems, [Hollands and Wald \(2002\)](#) and [Schiffrin and Wald \(2012\)](#) conclude that the only justification for the use of a Liouville measure in cosmology, in our current state of knowledge, is the bare assumption of the conceit of [Penrose \(1979\)](#), to wit, that the universe’s initial conditions were, by some appropriate process, randomly selected from a probability distribution fixed by the Liouville measure—the “creator” blindly threw a dart at a dartboard whose values are distributed according to it. [Schiffrin and Wald \(2012, p. 20\)](#) drily observe that this “has the status of an unsupported hypothesis.” I demure. There is no tongue long enough and no cheek deep enough to sufficiently warrant endowing this assumption with the honorific ‘hypothesis’. There is *no* known physical justification for the use of the Liouville measure in cosmology.

5 Genericity, Stability, and Prediction

As I already remarked above, the space of Lorentz metrics over a manifold—the family of possible spacetime models having that as its underlying manifold—is an infinite-dimensional Fréchet space, and so by [theorem 3.1](#) it has no non-trivial Borel measure. Standard probabilistic forms of argument in cosmology, however, mix topological and measure-theoretic concepts and methods in a way that depends on relations between topology and measure that are guaranteed to obtain only for Borel measures. In particular, those standard forms (always implicitly) assume at least one of the following propositions.

- Fix a “randomly selected” spacetime with a given property; if “small perturbations” (in a topological sense) destroy that property, then the collection of spacetimes with that property has zero measure. (The property is “scarce”; theorems showing the existence of the property are “rigid”.)
- Fix a “randomly selected” spacetime with a given property; if “small perturbations” (in a topological sense) preserve that property, then the collection of spacetimes with that property has large (or at least discernibly non-zero) measure. (The property is “generic”; theorems showing the existence of the property are not “rigid”.)
- If the collection of spacetimes with a given property has large (or at least discernibly non-zero) measure (“generic”), then that property is topologically stable under “small perturbations” (not “rigid”).
- If the collection of spacetimes with a given property has zero measure (“scarce”), then that property is topologically unstable under “small perturbations” (“rigid”).

The probabilistic element of the conclusions can be expressed using the idea of likelihood (in a non-technical sense). Standard arguments then take the following form. Assume that the property is generic and that observations we make indicate that the actual spacetime approximately satisfies the conditions of an existence theorem for that property; then the topological stability under small perturbations entailed by genericity guarantees that the inevitable inaccuracies and inexactitudes in the observations cannot block the inference that the likelihood that the property obtains in the actual universe is high; and so we conclude that the likelihood is in fact high. Because one does not have a Borel measure in infinite-dimensional Fréchet spaces, however, none of these propositions hold in general for the space of Lorentz metrics over a fixed manifold.²⁸

A good example of a powerful probabilistic conclusion based on topological reasoning dressed up in measure-theoretic clothing pertains to the likelihood of finding singularities in a certain class of spacetimes. Geroch (1966) conjectured that essentially all spatially closed spacetimes either have singularities or do not satisfy the SEC, or, somewhat more precisely, that singularities are generic and their occurrence is stable in the family of spatially closed spacetimes.²⁹ One compelling way to make Geroch’s conjecture precise is given by the so-called Lorentzian splitting theorems.³⁰ These theorems may be thought of as rigidity meta-theorems for singularity theorems invoking the strong energy condition, for the splitting theorems show that, under certain other assumptions, there will be no singularities only when the spacetime is static and globally hyperbolic. The reasoning then runs, static and globally hyperbolic spacetimes are “of measure zero” in the space of all spacetimes, and so being free of singularities is, under the ancillary conditions, unstable under arbitrarily small perturbations; thus, the likelihood of a “randomly selected” spatially closed spacetime being singularity-free is very low.³¹ These conclusions, however, are simply not justified in the absence of a Borel measure, even if one had a physically appropriate topology to use for the rigorous characterization of stability in the first place.

²⁸Hawking (1971) is particularly clear and explicit in sketching what I just proposed as a typical scheme for this sort of argument, though he does not note the mathematical issues I focus on.

²⁹The strong energy condition requires that for any timelike vector ξ^a , $R_{mn}\xi^m\xi^n \geq 0$, where R_{ab} is the Ricci tensor associated with the spacetime metric.

³⁰In order to state the most relevant splitting theorem, we need two definitions. First, the *edge* of an achronal, closed set Σ is the set of points $p \in \Sigma$ such that every open neighborhood of p contains a point $q \in I^-(p)$, a point $r \in I^+(p)$ and a timelike curve from q to r that does not intersect Σ . Second, let Σ be a non-empty subset of spacetime; then a future inextendible causal curve is a *future Σ -ray* if it realizes the supremal Lorentzian distance between Σ and any of its points lying to the future of Σ (Galloway and Horta 1996); *mutatis mutandis* for a *past Σ -ray*. (If γ is a Σ -ray, it necessarily intersects Σ .)

Theorem 5.1 (Lorentzian splitting theorem) *(\mathcal{M}, g_{ab}) be a spacetime that contains a compact, acausal space-like hypersurface Σ without edge and obeys the SEC; if it is timelike geodesically complete and contains a future Σ -ray γ and a past Σ -ray η such that $I^-(\gamma) \cap I^+(\eta) \neq \emptyset$, then it is isometric to $(\mathbb{R} \times \Sigma, t_a t_b - h_{ab})$, where (Σ, h_{ab}) is a compact Riemannian manifold and t^a is a timelike vector-field in \mathcal{M} .*

In particular, (\mathcal{M}, g_{ab}) must be globally hyperbolic and static. See Galloway and Horta (1996) for a proof.

³¹See, e.g., Hawking (1971), Penrose (1979), and Senovilla (1998) for examples of physicists explicitly using such measure-theoretic language to characterize the genericity of the occurrence of singularities in these families of spacetimes, based on topological stability of the occurrence of singularities. Those same physicists also offer similar arguments for the genericity of singularities in spatially open spacetimes. One can make the conjecture in this case precise by using a variation of the Lorentzian splitting theorem given in footnote 30 (Galloway and Horta 1996); see, e.g., Ringström (2009), for arguments of the sort I criticize based on the Lorentzian theorem for the spatially open case.

An example of a different sort is provided by the anthropic argument of [Weinberg \(1987\)](#) predicting “the most likely” range of values for the cosmological constant Λ . His argument runs as follows:

1. work with a family of near-FLRW spacetimes (*i.e.*, ones derived by allowing small perturbations off FLRW spacetimes, introducing small inhomogeneities);
2. then the existence of large, gravitationally bound systems places upper and lower bounds on possible values of Λ —if Λ is too positive, then potentially bound systems would be pulled apart, and if it is too negative, then the universe would recollapse before they can form;
3. argue for the topological stability of the formation of such bound systems under small changes in the value of Λ ;
4. use an anthropic argument (the presence of conscious observers as a selection effect, assuming we are typical observers, *i.e.*, that the value of Λ in our spacetime is typical of spacetimes with such observers) to fix the shape and peak of an appropriate measure on the family of near-FLRW spacetimes;
5. predict that the probability of the occurrence of a cosmological constant with a value lying in the range fixed in the second step is high, according to the posited measure.

The inadmissibility of the reasoning should, again, be clear. The argument assumes that there exists a measure and a topology that harmonize in such a way as to allow one both to characterize topological stability under small perturbations and to characterize typicality of a class of observers in a consistent way. On any reasonable family of near-FLRW spacetimes, however, there will be no such measure and topology, for the inhomogeneities ensure that the family will form an infinite-dimensional space.

My arguments do not show that the conclusions of the sorts of arguments I have considered in this section are necessarily wrong, only that the arguments currently given for those conclusions, in their present form, have serious mathematical, physical and conceptual problems that must be addressed before any real confidence can be had in those conclusions.

References

- Carroll, S. and H. Tam (2010). Unitary evolution and cosmological fine-tuning. Available online at [arXiv:1007.1417 \[hep-th\]](#).
- Curiel, E. (1999). The analysis of singular spacetimes. *Philosophy of Science* 66(S1), 119–145. A more recent version, with corrections and emendations, is available at <http://strangebeautiful.com/phil-phys.html>.
- Ellis, G. (2007). Issues in the philosophy of cosmology. In J. Butterfield and J. Earman (Eds.), *Handbook of Philosophy of Physics, Part B*, pp. 1183–1286. Dordrecht: North Holland. Preprint available at [arXiv:astro-ph/0602280v2](#).
- Fletcher, S. (2014). Global spacetime similarity. Unpublished manuscript.

- Fletcher, S. (2015). Similarity, topology, and physical significance in relativity theory. *British Journal for the Philosophy of Science*. Forthcoming.
- Galloway, G. and A. Horta (1996, May). Regularity of Lorentzian Busemann functions. *Transactions of the American Mathematical Society* 348(5), 2063–2084.
- Geroch, R. (1966). Singularities in closed universes. *Physical Review Letters* 17, 445–447. doi:10.1103/PhysRevLett.17.445.
- Geroch, R. (1967). Topology in general relativity. *Journal of Mathematical Physics* 8(4), 782–786. doi:10.1063/1.1705276.
- Geroch, R. (1969). Spinor structure of space-times in general relativity I. *Journal of Mathematical Physics* 9, 1739–1744.
- Geroch, R. (1970). Singularities. In M. Carmeli, S. Fickler, and L. Witten (Eds.), *Relativity*, pp. 259–291. New York: Plenum Press.
- Geroch, R. (1971). General relativity in the large. *General Relativity and Gravitation* 2(1), 61–74.
- Gibbons, G., S. Hawking, and J. Stewart (1987, February). A natural measure on the set of all universes. *Nuclear Physics B* 281(3–4), 736–751. doi:10.1016/0550-3213(88)90008-9.
- Gibbons, G. and N. Turok (2008). Measure problem in cosmology. *Physical Review D* 77, 063516. doi:10.1103/PhysRevD.77.063516. Preprint available online at [arXiv:hep-th/0609095](https://arxiv.org/abs/hep-th/0609095).
- Halmos, P. (1950). *Measure Theory*. New York: Van Nostrand and Co.
- Hawking, S. (1971). Stable and generic properties in general relativity. *General Relativity and Gravitation* 1(4), 393–400.
- Hocking, J. and G. Young (1988). *Topology*. New York: Dover Publications, Inc. Originally published in 1961 by Addison-Wesley Publishing Co.
- Hollands, S. and R. Wald (2002). An alternative to inflation. *General Relativity and Gravitation* 34, 2043–2055. doi:10.1023/A:1021175216055. Preprint available at [arXiv:gr-qc/0205058](https://arxiv.org/abs/gr-qc/0205058).
- Kelley, J. (1955). *General Topology*. The University Series in Higher Mathematics. Princeton: D. Van Nostrand Company, Inc.
- Peirce, C. S. (1878a). The doctrine of chances. See [Peirce \(1992\)](#), Chapter 10, pp. 142–154. Originally published in *Popular Science Monthly* 12(March, 1878):604–615.
- Peirce, C. S. (1878b). The probability of induction. See [Peirce \(1992\)](#), Chapter 10, pp. 155–69. Originally published in *Popular Science Monthly* 12(April, 1878):705–18.
- Peirce, C. S. (1992). *The Essential Peirce: Selected Philosophical Writings*, Volume 1 (1867–1893). Bloomington, IN: Indiana University Press.
- Penrose, R. (1979). Singularities and time-asymmetry. In S. Hawking and W. Israel (Eds.), *General Relativity: An Einstein Centenary Survey*, pp. 581–638. Cambridge University Press.
- Ringström, H. (2009). *The Cauchy Problem in General Relativity*. ESI Lectures in Mathematics and Physics. Zürich: European Mathematical Society Publishing House.

- Schiffrin, J. and R. Wald (2012). Measure and probability in cosmology. [arXiv:1202.1818 \[gr-qc\]](#).
- Senovilla, J. (1998). Singularity theorems and their consequences. *General Relativity and Gravitation* 30(5), 701–848. [doi:10.1023/A:1018801101244](#). *N.b.*: on the article itself is written that the paper appears in volume 29, 1997; this is erroneous, as the author of the paper has assured me.
- Smeenk, C. (2012a). The logic of cosmology revisited. Unpublished manuscript. Available on request from the author.
- Smeenk, C. (2012b). Philosophy of cosmology. Unpublished manuscript. Available on request from the author.
- Weinberg, S. (1987). Anthropic bound on the cosmological constant. *Physical Review Letters* 59(November), 2607. [doi:10.1103/PhysRevLett.59.2607](#).