

This is a repository copy of *The structure of sperm Izumo1 reveals unexpected similarities with Plasmodium invasion proteins.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/166881/>

Version: Published Version

Article:

Nishimura, Kaoru, Han, Ling, Bianchi, Enrica et al. (3 more authors) (2016) The structure of sperm Izumo1 reveals unexpected similarities with Plasmodium invasion proteins. *Current Biology*. R661-R662. ISSN 0960-9822

<https://doi.org/10.1016/j.cub.2016.06.028>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Correspondence

The structure of sperm Izumo1 reveals unexpected similarities with *Plasmodium* invasion proteins

Kaoru Nishimura^{1,4}, Ling Han^{1,4},
Enrica Bianchi², Gavin J. Wright²,
Daniele de Sanctis³,
and Luca Jovine^{1,*}

Fertilization, the culminating event in sexual reproduction, occurs when haploid sperm and egg recognize each other and fuse to form a diploid zygote. In mammals this process critically depends on the interaction between Izumo1, a protein exposed on the equatorial segment of acrosome-reacted sperm, and the egg plasma-membrane-anchored receptor Juno [1,2]. The molecular mechanism triggering gamete fusion is unresolved because both Izumo1 and Juno lack sequence similarity to known membrane fusogens. Here we report the crystal structure of Izumo1, which reveals a membrane distal region composed of a four-helix bundle connected to a carboxy-terminal immunoglobulin (Ig)-like domain through a β -hairpin stabilized by disulfide bonds. Remarkably, different regions of Izumo1 display significant structural similarities to two proteins expressed by the invasive sporozoite stage of *Plasmodium* parasites: SPECT1, which is essential for host cell traversal and hepatocyte invasion [3]; and TRAP, which is necessary for gliding motility and invasion [4]. These observations suggest a link between the molecular mechanisms underlying host cell invasion by the malaria parasite and gamete membrane fusion at fertilization.

Juno and Izumo are structurally unrelated: Juno adopts a modified folate receptor family fold [5], whereas Izumo1 is a type I transmembrane protein consisting of an extracellular region of about 300 residues and a short cytoplasmic tail [2]. The biological activity of Izumo1 depends

on the amino-terminal half of its ectodomain — a region named the ‘Izumo domain’ because it is shared with the paralogous sperm proteins Izumo2, 3 and 4 [6]; this region is followed by a glycosylated Ig-like domain [2] (Figure 1A).

To investigate how Izumo1 might trigger gamete fusion, we expressed the entire ectodomain in mammalian cells and determined a crystal structure to 2.5 Å resolution (Figure S1 in Supplemental Information and PDB ID 5B5K). The structure shows that the Izumo domain consists of a four-helix bundle with a flexible α -helical hook/ β -sheet insertion, followed by a β -hairpin that connects the bundle to a seven-stranded Ig-like domain (Figure 1A,B). These features result in an elongated architecture stabilized by five intramolecular disulfide bonds that are conserved in the three Izumo paralogs, and are clearly discernible within the sequence of Spaca6, another sperm surface protein recently reported to be required for gamete fusion [7] (Figure S2A,B). This suggests that the similarity between Izumo1 and Spaca6 is not limited to the carboxy-terminal Ig-like domain, which can be superimposed onto the canonical V-set Ig-like domain of human CD2 (PDB ID 1HNF) — a molecule mediating the adhesion of lymphocytes to antigen-presenting cells — with a root mean square deviation (RMSD) of 2.1 Å over 86 residues.

The helical character of the carboxy-terminal half of the Izumo domain, which corresponds to helices α 3 and α 4 in our structure (Figure 1A,B), was shown to be important for the function of Izumo1 [6]. The structure readily explains why mutation of leucine residues to proline interferes with this biological activity by disrupting the fold of the four-helix bundle (Figure S2A,C). However, binding assays using peptides representing α 3 and α 4 suggest that this region of Izumo1 is either not involved in binding to Juno or not sufficient for this interaction (Figure 1C).

Strikingly, structural homology searches revealed a high confidence match between the four-helix bundle of Izumo1 and that of SPECT1, a secreted *Plasmodium* protein that is essential for host cell traversal by

the infective sporozoite form of the parasite (Dali Z-score 6.9, RMSD 3.3 Å over 98 residues; Figure 1D). The nearly parallel/antiparallel arrangement of the SPECT1 helical bundle, which also contains a helical hook (albeit inserted at a different position than in Izumo1), has been suggested to adopt a metastable structure which favors the transition from a solvent-exposed to membrane-associated form [8]. In addition, the β -hairpin of Izumo1 is structurally similar to the extensible β -ribbon of TRAP, another *Plasmodium* protein that mediates sporozoite gliding and host cell invasion [4] (RMSD 1.4 Å over 19 residues; Figure 1E). The TRAP β -ribbon is thought to undergo conformational changes upon attachment to host cells and acts as an elbow-like flexible spacer between the amino-terminal α -helical von Willebrand factor A (VWA) domain of the protein (which interacts with the host cell) and its carboxy-terminal β -rich thrombospondin type I repeat (TSR) domain.

What might be the functional implications of the structural similarities to *Plasmodium* cell traversal and invasion proteins with Izumo1? Our findings suggest that the four-helix bundle of the Izumo domain and SPECT1 may orchestrate analogous fusion-related molecular recognition events at the sperm-egg and parasite-host cell interfaces. Izumo1 is not considered a fusogen per se because it cannot induce membrane fusion when heterologously expressed in commonly used cell lines [1]. The presence of solvent-exposed bulk aromatic residues in the Izumo1 four-helix bundle (Figure S2A; a feature also observed in SPECT1) and the report that some monoclonal antibodies that target the Izumo domain do not block binding of sperm to the egg but hinder gamete fusion [6] are consistent with the possibility that an additional, as yet unknown, component is required to form a functional fusogenic complex [9]. Whether SPECT1 is also part of a larger protein complex that could organize proteins that have membrane-disrupting functions, such as SPECT2, is currently unclear [10].

Both SPECT1 and TRAP are expressed by the sporozoite stage






Figure 1. Experimental results and structural comparisons with *Plasmodium* invasion proteins. (A) Domain architecture of the extracellular region of Izumo1. The Izumo domain, which consists of a four-helix bundle ($\alpha 1$ – $\alpha 4$), an insertion (ins: $\alpha 2/3$ hook and $\beta 0$) and a β -hairpin ($\beta 1$ – $\beta 2$) — see panel (B) — is boxed. A protease-sensitive carboxy-terminal region that was included in the expression construct used for this study, but is not defined in the electron density map, is indicated by a dashed line. (B) Crystal structure of mouse Izumo1 (amino acids C22–K256), shown in cartoon representation with different regions of the molecule colored as in (A). Amino/carboxyl termini and secondary structure elements are marked. Cysteine residues and the N-acetylglucosamine (NAG) residue attached to N204 are represented in ball-and-stick notation, with circled pink numbers indicating the five disulfide bonds (see also Figure S2A). (C) Microscale thermophoresis experiments show that Juno directly binds to the Izumo1 ectodomain, but does not interact with peptides corresponding to Izumo1 helices $\alpha 3$ and $\alpha 4$ (or an equimolar mixture thereof), which span a region reported to be sufficient for binding to the egg membrane [6]. Vertical error bars represent s.d. of the mean ($n \geq 3$). (D) Structural alignment of the four-helix bundles of Izumo1 and *Plasmodium* SPECT1 (PDB ID 4U5A [8]), in side (left) and top (right) views. Izumo1 helices are colored as in (A,B), SPECT1 is grey. By optimizing the match found by Dali, 84 residues can be superimposed with a RMSD of 3.0 Å. (E) Superposition of the β -hairpin of Izumo1 (red) and the extensible β -ribbon of *Plasmodium* protein TRAP (black; PDB ID 4HQO [4]). Both elements separate amino-terminal α -helical domains from carboxy-terminal β -rich regions in the respective proteins.

of *Plasmodium* parasites and are similarly sequestered within specialized intracellular secretory organelles called micronemes [3,4]. We do not yet know if these proteins, like Izumo1, have a specific binding partner that mediates cell-specific recognition events, or, as has been suggested for sp18 (a five-helical bundle sperm acrosomal protein from the mollusk abalone), are disrupting membrane structure by interacting directly with phospholipids. Nevertheless, it is intriguing that, similar to SPECT1 and TRAP, Izumo1 is sequestered in the acrosome and that its release is also highly regulated by localized secretion.

The structure of Izumo1 provides new mechanistic insights into a fascinating basic biological process and, also considering the protein's recent implication in

human immunofertility, presents new opportunities in the rational design of new fertility treatments and contraceptives.

SUPPLEMENTAL INFORMATION

Supplemental Information includes two figures and experimental procedures and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2016.06.028>.

ACKNOWLEDGEMENTS

We thank R. Aricescu and Y. Zhao (University of Oxford) for the mammalian expression vector pHLsec, and are grateful to T. Shimizu (University of Tokyo) for sharing the coordinates of human IZUMO1 before publication. Access to structural biology beamlines at ESRF is also gratefully acknowledged. Research was supported by Karolinska Institutet; the Center for

Innovative Medicine; Swedish Research Council grant 2012-5093; the Göran Gustafsson Foundation for Research in Natural Sciences and Medicine; the Sven and Ebba-Christina Hagberg foundation; an EMBO Young Investigator award; and the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement 260759 (to L.J.); the Wellcome Trust grant number 098051 and the UK Medical Research Council grant MR/M012468/1 (to E.B. and G.J.W.).

REFERENCES

- Bianchi, E., Doe, B., Goulding, D., and Wright, G.J. (2014). Juno is the egg Izumo receptor and is essential for mammalian fertilization. *Nature* 508, 483–487.
- Inoue, N., Ikawa, M., Isotani, A., and Okabe, M. (2005). The immunoglobulin superfamily protein Izumo is required for sperm to fuse with eggs. *Nature* 434, 234–238.
- Ishino, T., Yano, K., Chinzei, Y., and Yuda, M. (2004). Cell-passage activity is required for the malarial parasite to cross the liver sinusoidal cell layer. *PLoS Biol.* 2, E4.
- Song, G., Koksai, A.C., Lu, C., and Springer, T.A. (2012). Shape change in the receptor for gliding motility in *Plasmodium* sporozoites. *Proc. Natl. Acad. Sci. USA* 109, 21420–21425.
- Han, L., Nishimura, K., Sadat Al Hosseini, H., Bianchi, E., Wright, G.J., and Jovine, L. (2016). Divergent evolution of vitamin B₃ binding underlies Juno-mediated adhesion of mammalian gametes. *Curr. Biol.* 26, R100–R101.
- Inoue, N., Hamada, D., Kamikubo, H., Hirata, K., Kataoka, M., Yamamoto, M., Ikawa, M., Okabe, M., and Hagihara, Y. (2013). Molecular dissection of IZUMO1, a sperm protein essential for sperm-egg fusion. *Development* 140, 3221–3229.
- Lorenzetti, D., Poirier, C., Zhao, M., Overbeek, P.A., Harrison, W., and Bishop, C.E. (2014). A transgenic insertion on mouse chromosome 17 inactivates a novel immunoglobulin superfamily gene potentially involved in sperm-egg fusion. *Mamm. Genome* 25, 141–148.
- Hamaoka, B.Y., and Ghosh, P. (2014). Structure of the essential *Plasmodium* host cell traversal protein SPECT1. *PLoS ONE* 9, e114685.
- Inoue, N., Hagihara, Y., Wright, D., Suzuki, T., and Wada I. (2015). Oocyte-triggered dimerization of sperm IZUMO1 promotes sperm-egg fusion in mice. *Nat. Commun.* 6, 8858.
- Ishino, T., Chinzei, Y., and Yuda, M. (2005). A *Plasmodium* sporozoite protein with a membrane attack complex domain is required for breaching the liver sinusoidal cell layer prior to hepatocyte infection. *Cell Microbiol.* 7, 199–208.

¹Department of Biosciences and Nutrition & Center for Innovative Medicine, Karolinska Institutet, Huddinge, SE-141 83, Sweden.

²Cell Surface Signalling Laboratory, Wellcome Trust Sanger Institute, Hinxton, Cambridge, CB10 1SA, UK. ³ESRF - The European Synchrotron, Grenoble 38000, France.

⁴Co-first author.

*E-mail: luca.jovine@ki.se