

PUBLICATIONS

2006

SOIL DEPTH DISTRIBUTION OF NITRATE-NITROGEN IN COASTAL BERMUDAGRASS PASTURES SUBJECTED TO LONG-TERM STOCKING RATES AND FERTILITY REGIMENS

M.L. Silveira, F.M. Rouquette, Jr., J.L. Kerby, G.H. Nimr, V.A. Haby, and G.R. Smith

Background. A detailed description of stocking rates and fertility regimens from 1969 through 2005 are presented in a companion 2006 Field Day Report by Rouquette et al. The objective of this study was to investigate nitrate-nitrogen ($\text{NO}_3\text{-N}$) distribution within the soil profile in Coastal bermudagrass pastures continuously stocked for 19 years and subjected to different fertility management regimens.

Research Findings. After 5 years (1985-1989) of annual N fertilizer rates > 400 lbs/ac, soil $\text{NO}_3\text{-N}$ concentrations in 1989 were predictably greater in N-fertilized bermudagrass pastures overseeded with ryegrass than in bermudagrass/clover pastures that received no N fertilizer (Figure 1). At both low and high stocking rates, $\text{NO}_3\text{-N}$ concentrations in Coastal bermudagrass pastures plus N were considerably greater (~ 20 ppm) within the top 36 inches (Figure 1a). Deeper soil depths in the N-fertilized pastures showed a significant decrease in $\text{NO}_3\text{-N}$ levels (~ 5 ppm). In contrast, Coastal bermudagrass pastures receiving no N fertilizer exhibited more homogeneous $\text{NO}_3\text{-N}$ distribution within the soil profile compared to bermudagrass pastures receiving N.


In 1994, after 5 years of lower N application rates (238 lbs N/ac year), soil $\text{NO}_3\text{-N}$ concentrations were dramatically decreased in the bermudagrass + N pastures (Fig. 1b). This trend was more evident in the top 36 inches. Similar results were observed for both stocking rates.

Soil $\text{NO}_3\text{-N}$ concentrations in 2004 (Fig. 1c) were comparable to that in 1994. Bermudagrass without N fertilizer showed a slight increase in soil $\text{NO}_3\text{-N}$ levels in 2004 compared to 1994, which was probably related to the better P nutrient status in 2004 (soil test P = 21 ppm) compared to 1994 (soil test P = 12 ppm); thus, greater forage production was promoted. Bermudagrass pastures, with no N-fertilizer application, sustained nearly constant soil $\text{NO}_3\text{-N}$ concentrations over 19 years (1985-2004) of continuous stocking. Long-term stocking rates showed no detrimental effect on soil $\text{NO}_3\text{-N}$ concentrations. Rather, nutrient cycling via animal excreta was apparently playing a major role maintaining adequate soil $\text{NO}_3\text{-N}$ levels.


Applications. Nitrogen fertilization can tremendously affect soil $\text{NO}_3\text{-N}$ concentrations in Coastal bermudagrass pastures and in certain soil types. Fertilization rates can be a much more important factor affecting soil $\text{NO}_3\text{-N}$ concentrations than stocking rates. Clover proved to be a

reasonable alternative to supply satisfactory amounts of N via animal excreta for Coastal bermudagrass production on sandy, acid soils in East Texas.

a. 1989 Coastal Bermudagrass



b. 1994 Coastal Bermudagrass



c. 2004 Coastal Bermudagrass




Figure 1. Nitrate-nitrogen ($\text{NO}_3\text{-N}$) distribution within the soil profile as a function of fertility regimens.