



# University of Chester



This work has been submitted to ChesterRep – the University of Chester's  
online research repository

<http://chesterrep.openrepository.com>

Author(s): Gavin J Phillips ; Jim Thieser ; Mingjin Tang ; Nicolas Sobanski ;  
Johannes Fachinger ; Frank Drewnick ; Jos Lelieveld ; John N Crowley

Title: CINO<sub>2</sub> and nitrate formation via N<sub>2</sub>O<sub>5</sub> uptake to particles: Derivation of N<sub>2</sub>O<sub>5</sub>  
uptake coefficients from ambient datasets

Date: 2015

Originally published in: Geophysical Research Abstracts

Example citation: Phillips, G. J., Thieser, J., Tang, M., Sobanski, N., Fachinger, J.,  
Drewnick, F., Lelieveld, J., & Crowley, J. N. (2015). CINO<sub>2</sub> and nitrate formation via  
N<sub>2</sub>O<sub>5</sub> uptake to particles: Derivation of N<sub>2</sub>O<sub>5</sub> uptake coefficients from ambient  
datasets. *Geophysical Research Abstracts*, 17, EGU2015-10435.

Version of item: Published version

Available at: <http://hdl.handle.net/10034/345822>



## **ClNO<sub>2</sub> and nitrate (NO<sub>3</sub><sup>-</sup>) formation via N<sub>2</sub>O<sub>5</sub> uptake to particles: Derivation of N<sub>2</sub>O<sub>5</sub> uptake coefficients from ambient datasets**

Gavin J. Phillips (1,2), Jim Thieser (1), Ming J. Tang (1), Nicolas Sobanski (1), Johannes Fachinger (3), Frank Drewnick (3), Jos Lelieveld (1), and John N. Crowley (1)

(1) Max Planck Institute for Chemistry, Division of Atmospheric Chemistry, Mainz, Germany, (2) Department of Natural Sciences, University of Chester, Thornton Science Park, CH2 4NU, UK, (3) Max Planck Institute for Chemistry, Division of Particle Chemistry, Mainz, Germany

We present estimates of the uptake coefficient of N<sub>2</sub>O<sub>5</sub>,  $\gamma(\text{N}_2\text{O}_5)$ , using ambient measurements of the trace gases N<sub>2</sub>O<sub>5</sub> and ClNO<sub>2</sub> and particle composition and surface area at the Kleiner Feldberg observatory, near Frankfurt, SW Germany, during the PARADE campaign (summer 2011). Three methods used to extract  $\gamma(\text{N}_2\text{O}_5)$  from the datasets were found to be in reasonable agreement, generating values between 0.001 and 0.4.  $\gamma(\text{N}_2\text{O}_5)$  displayed a significant dependence on relative humidity (RH), the largest values obtained, as expected, at high RH. No significant dependence of  $\gamma(\text{N}_2\text{O}_5)$  on particle organic content or sulphate-to-organic ratio was observed. The variability in  $\gamma(\text{N}_2\text{O}_5)$  is however large, indicating that humidity is not the sole factor determining the uptake coefficient. There is also an indication that the yield of ClNO<sub>2</sub> with respect to N<sub>2</sub>O<sub>5</sub> uptake is larger with lower concentrations of PM1 total organics. Our results will be compared to existing uptake coefficients from laboratory studies and those derived from field-observations.