

## Energy Intake and Expenditure Assessed ‘In-Season’ in an Elite European Rugby Union Squad

Warren Bradley<sup>1</sup>, Bryce Cavanagh<sup>2</sup>, William Douglas<sup>2</sup>, Timothy F, Donovan<sup>3</sup>, Craig Twist<sup>4</sup> James P Morton<sup>1</sup>  
and Graeme L Close<sup>1</sup>

<sup>1</sup>Research Institute for Sport and Exercise Sciences  
Liverpool John Moores University  
Tom Reilly Building  
Liverpool  
L3 3AF  
UK

<sup>2</sup>Munster Rugby  
Tyco Building  
Cork Institute of Technology  
Cork  
Ireland

<sup>3</sup>Sport and Exercise Sciences,  
Glyndwr University,  
Plas Coch Campus,

Wrexham  
UK

<sup>4</sup>Department of Sport and Exercise Sciences  
University of Chester  
Parkgate Road  
Chester  
CH1 4BJ

### Address for Correspondence:

Dr Graeme L. Close  
Research Institute for Sport and Exercise Sciences,  
Tom Reilly Building  
Byrom St Campus  
Liverpool John Moores University,  
Liverpool,  
UK  
L3 3AF  
0151 904 6266  
[g.l.close@ljmu.ac.uk](mailto:g.l.close@ljmu.ac.uk)

## 1Abstract

2

3Rugby Union is a complex, high-intensity intermittent collision sport with emphasis placed on players  
4possessing high lean body-mass and low body-fat. After an 8-12 week pre-season focused on physiological  
5adaptations, emphasis shifts towards competitive performance. However, there are no objective data on the  
6physiological demands or energy intake (EI) and expenditure (EE) for elite players during this period.  
7Accordingly, in-season training load using GPS and session RPE (sRPE), alongside six-day assessments of  
8EE and EI were measured in 44 elite Rugby Union players. Mean weekly distance covered was  $7827 \pm 954$   
9m and  $9572 \pm 1233$  m with a total mean weekly sRPE of  $1776 \pm 355$  and  $1523 \pm 434$  AU for forwards and  
10backs, respectively. Mean weekly EI was  $16.6 \pm 1.5$  and  $14.2 \pm 1.2$  MJ, and EE was  $15.9 \pm 0.5$  and  $14 \pm 0.5$   
11MJ. Mean carbohydrate intake was  $3.5 \pm 0.8$  and  $3.4 \pm 0.7$  g.kg<sup>-1</sup> body mass, protein intake was  $2.7 \pm 0.3$  and  
12 $2.7 \pm 0.5$  g.kg<sup>-1</sup> body mass, and fat intake was  $1.4 \pm 0.2$  and  $1.4 \pm 0.3$  g.kg<sup>-1</sup> body mass. All players who  
13completed the food diary self-selected a 'low' carbohydrate 'high' protein diet during the early part of the  
14week, with carbohydrate intake increasing in the days leading up to a match, resulting in the mean EI  
15matching EE. Based on EE and training load data, the EI and composition seems appropriate, although  
16further research is required to evaluate if this diet is optimal for match day performance.

17Keywords: Rugby, Pre-Season, GPS, Physiology, Nutrition

18

## 19Introduction

20 Rugby Union (RU) is a high impact collision sport played over 80 minutes, which is split into two  
21 forty-minute halves. RU is characterized by frequent bouts of high intensity exercise such as sprinting,  
22 accelerations, tackling, scrummaging, rucking and mauling (Roberts, Trewartha, Higgitt, El-Abd, & Stokes,  
23 2008) and is predominantly aerobic in nature. Players are classified as either forwards or backs, with the  
24 forwards tending to be heavier and stronger compared with the backs who tend to be leaner and faster  
25 (Duthie, Pyne, & Hooper, 2003). During a typical rugby union game, players will cover  $68 \text{ m} \cdot \text{min}^{-1}$  (Cahill,  
26 Lamb, Worsfold, Headey, & Murray, 2013), which is much lower than locomotive rates described for  
27 Australian football,  $\sim 123 \text{ m} \cdot \text{min}^{-1}$ ; (Kempton, Sullivan, Bilsborough, Cordy, & Coutts, 2015), rugby league,  
28  $\sim 85 \text{ m} \cdot \text{min}^{-1}$  (Waldron, Twist, Highton, Worsfold, & Daniels, 2011) and soccer,  $\sim 104 \text{ m} \cdot \text{min}^{-1}$  (Varley,  
29 Gabbett, & Aughey, 2014). Activities such as rucking, mauling, scrummaging and lineouts are likely  
30 explanations for the lower locomotive rates observed between rugby union and other football codes.

31 A typical in-season in rugby union lasts approximately 34-36 weeks followed by 3-6 weeks of rest  
32 time depending on whether play off stages are reached. The central focus of the in-season is peak  
33 performance during competition. Strategies to prepare for, and optimally recover from competition are  
34 therefore the objectives of this period with emphasis also placed on the maintenance of body composition to  
35 values attained at the end of pre-season (Bradley et al., 2014). An understanding of players' day-to-day  
36 training and energy requirements is essential to avoid residual fatigue (Gamble, 2006), in the identification of  
37 appropriate recovery strategies, and to determine appropriate training loads to maximise performance  
38 (Fowles, 2006). While researchers have investigated the movement patterns and physiological demands of  
39 matches (Cunniffe, Proctor, Baker, & Davies, 2009; Duthie et al., 2003), there are no data currently available  
40 on the internal and external demands, or the energy intakes and expenditures of elite rugby union players  
41 during the in-season period.

42 The daily nutritional intake of an athlete should meet the fuel requirements of high training  
43 intensities and competition, promote optimal recovery, and provide essential micronutrients for general  
44 health and well being. Data have previously been published on the nutritional intakes of elite rugby union

45 players during training (Bradley et al., 2014); however, these data looked specifically at the pre-season with  
46 training and nutrition tailored towards physiological adaptation. Transition from the pre-season to in-season  
47 shifts focus from physiological adaptation to competition preparation and recovery, with training  
48 programmes modified to reflect this transition and consequently nutritional intakes must also be modified to  
49 meet training and competition requirements. To the authors' knowledge, no data evaluating the nutritional  
50 intake of elite rugby union players during in-season training is available and, as such, evidence-based  
51 recommendations regarding the energy requirements to fuel a rugby player's training plan in-season are  
52 currently lacking.

53 For team sports such as Rugby Union, daily carbohydrate (CHO) intakes have traditionally been  
54 high. However, in recent years many rugby players have adopted a lower CHO diet during the beginning of a  
55 training week in attempts to maintain or reduce body fat (Morton, Robertson, Sutton, & MacLaren, 2010)  
56 maximize adaptations to training (Morton et al., 2009). Thereafter, intakes are increased in the day(s) leading  
57 up to a match to maximize glycogen stores. For example, Bradley et al. (2014) reported carbohydrate intakes  
58 of  $3.3 \pm 0.7$  and  $4.1 \pm 0.4$  g·kg<sup>-1</sup> for forwards and backs, respectively, during a rugby union pre-season. These  
59 data are similar to those reported in professional soccer players ( $3.4$  g·kg<sup>-1</sup> (Maughan, 1997)), but lower than intakes  
60 generally suggested for team sports engaged in moderate exercise programmes where values of 5-7 g·kg<sup>-1</sup>  
61 have been recommended (Burke, Hawley, Wong, & Jeukendrup, 2011). To date however there are no data  
62 on typical macronutrient intakes of elite rugby players during in-season training.

63 To implement a valid nutritional plan it is important to understand the day-to-day energy  
64 requirements of an athlete. Due to the physicality of rugby, the measurement of energy expenditure (EE) is  
65 somewhat difficult given that many of the tools available would not be suitable either through danger to the  
66 athlete or to the equipment. Currently the doubly labelled water (DLW) stable isotope method is considered  
67 the gold standard for measuring EE (Ekelund, Yngve, Westerterp, & Sjostrom, 2002), despite not allowing  
68 day-to-day comparisons to be made. Multi-sensor, wearable body monitoring technology might therefore  
69 provide an effective means of assessing daily EE in rugby players.

70 Although Bradley et al., (2014) has reported the training demands and nutritional intakes of an elite  
71 rugby union pre-season, to date there are no studies showing the training demands and energy intakes and  
72 expenditures during the competitive season. Due to the importance of competitive performance, these data  
73 would be of great significance to the strength and conditioning professional to allow informed decisions to  
74 be made with regards to players' diets during this competitive period. Therefore the aim of this study was to  
75 1) characterize the weekly external and internal training demands of a rugby union in-season using GPS  
76 technology and session RPE (sRPE); 2) evaluate the typical energy intakes and macro- and micronutrient  
77 intakes, and 3) analyse the energy expenditures of elite rugby union players during the in-season period.

78

## 79 **Methods**

### 80 *Study design*

81 Players began in-season training at the rugby club after a 12-week pre-season period. The first week of in-  
82 season training started in early October and this was classed as Week 13. Players then began 3 x 12-week in-  
83 season training macrocycles as prescribed by the club. During the 'in season', running activity was  
84 monitored at every training session using GPS technology and session RPE (sRPE) was used to quantify the  
85 overall training load. Body composition assessment and food diaries were completed as part of the club's  
86 normal in-season training regime and were routinely performed by all of the players who were therefore  
87 familiar with each test. During weeks 32 (n = 5), 33 (n = 5) and 34 (n = 4) of the season 14 players wore  
88 Senseware armbands and completed a detailed seven-day food diary to assess energy expenditure and  
89 nutrient intake. A typical in-season training week is depicted in Table 1.

90

### 91 *Participants*

92 Forty-four elite rugby union players currently playing in the European Rabo Direct Pro 12 league  
93 volunteered for this study. Based on playing position, these were divided in sub-groups of forwards (n= 24)  
94 and backs (n = 20). The sample population was collected on the first team squad that included 12 current

95international players and four British & Irish Lions. Of the 44 players that completed the ‘in-season’ season,  
96all completed sRPE and anthropometric assessments every 8-weeks. All 44 players also trained wearing the  
97GPS units at some stage during the competitive season although only 17 players wore units during any one  
98training session due to the availability of equipment. Only 14 players (seven forwards and seven backs) from  
99the squad of 44 players completed the energy expenditure and dietary analysis due to time constraints on the  
100players and limited equipment. A summary of the participant characteristics can be seen in Table 2. The local  
101ethics committee of Liverpool John Moores University granted ethical approval for the study. All  
102participants provided written informed consent before commencement of the study and all participants were  
103greater than 18 years of age (age range 21-34 years old).

#### 104**Procedures**

##### 105*Quantification of weekly external and internal training load*

106 Distances covered by forwards and backs during field sessions over four ‘typical’ in-season weeks  
107were assessed using GPS technology. Seventeen GPS units were rotated around the team ensuring that all  
108positions were accounted for during each training session. Movements were recorded using a Minimax S4  
109GPS unit (Catapult Innovations, Melbourne, Australia) sampling at a frequency of 10 Hz. A recent review  
110has demonstrated that 10 Hz units provide more accurate and reliable data compared with lower sampling  
111frequency devices (Cummins, Orr, O'Connor, & West, 2013). Indeed, the 10 Hz units used in this study are  
112two to three times more accurate at detecting changes in velocity, and up to six-fold more reliable than  
113devices sampling at 5 Hz (Varley, Fairweather, & Aughey, 2012). The CV of these units across a range of  
114speeds have been reported as 3.1 to 8.3% at a constant velocity, 3.6 to 5.9% for accelerations and 3.6 to  
11511.3% for decelerations (Varley et al., 2012). GPS units were used to collect data on total distance (m) and  
116relative distance covered in standing ( $0-2.0 \text{ m}\cdot\text{s}^{-1}$ ), walking ( $2.0-4.4 \text{ m}\cdot\text{s}^{-1}$ ), jogging ( $4.4-5.6 \text{ m}\cdot\text{s}^{-1}$ ), high-  
117speed running ( $5.6-7.5 \text{ m}\cdot\text{s}^{-1}$ ) and sprinting ( $7.5 + \text{ m}\cdot\text{s}^{-1}$ ) based on the clubs in-house classification of speed  
118zones. Tri-axial accelerometers and gyroscopes sampling at 100 Hz, also provided data on the number of  
119maximal accelerations ( $>5 \text{ m}\cdot\text{s}^{-2}$ ), physical collisions, and repeated high-intensity efforts (RHIE). A RHIE  
120was defined as three consecutive efforts (sprint, contact or acceleration) each separated by less than 21 s

121(Gabbett, Wiig, & Spencer, 2013). The unit was worn in a fitted neoprene vest, on the upper back of the  
122players. Quantification of gym and pitch session training loads was also assessed using the session rating of  
123perceived exertion (sRPE), (Foster et al., 2001). Using a modified 10-point Borg Scale (Borg, Hassmen, &  
124Lagerstrom, 1987) individual RPEs were provided by each player ~20 minutes after a training session from  
125which sRPE (AU) was calculated by multiplying RPE by total training time or total number of repetitions x  
126RPE for field or gym sessions, respectively.

127

### 128**Energy intake (EI)**

129 A six-day food diary was used to analyze the macronutrient and micronutrient and reported as days  
130away from a match (Game day -5, -4, -3, -2, -1 and game day +1) in megajoules (MJ). This time period is  
131believed to provide reasonably accurate and precise estimations of habitual energy and macronutrient  
132consumption (Braakhuis, Meredith, Cox, Hopkins, & Burke, 2003). Players were instructed to document a  
133complete account of all foods and fluids ingested over a six-day period, with careful attention to detail such  
134as timing of intakes, volumes and quantities, and specific brand names where possible. The nutrient intakes  
135were calculated using Nutritics professional diet analysis software (Nutritics LTD, Ireland) to obtain energy  
136and macro- and micronutrient composition. Each athlete's individual physical activity was known from the  
137weekly training schedule.

138

### 139**Energy expenditure (EE)**

140SenseWear Pro2 wearable armband (SWA; BodyMedia, USA) was used to assess the energy expenditure of  
141the players. Five armbands were rotated between the athletes over a three-week period during the same  
142macrocycle. Athletes wore the armband 24-hours a day for six consecutive days, except during water or  
143heavy contact based activities. The SWA were removed on match day to avoid disruption during match  
144preparations and also due to contacts sustained during competition. Studies have demonstrated that the SWA  
145provides accurate results for energy expenditure during low-to-moderate intensity physical exercise with a

146threshold for accurate measurements at intensities of around ten METs (Drenowatz & Eisenmann, 2011).  
147However, given that the compendium of physical activities indicates an intensity of 8.3 METs for rugby  
148union competition (Ainsworth et al., 2011), the use of SWA for rugby union appears appropriate. The  
149armband was worn on the back of the upper right arm and utilized a two-axis accelerometer, heat flux sensor,  
150galvanic skin response sensor, skin temperature sensor, and a near-body ambient temperature sensor to  
151capture data leading to the calculation of energy expenditure. SenseWear computer software (BodyMedia,  
152USA) was used to analyze player energy expenditure and reported as days away from competition (Game  
153day -5, -4, -3, -2, -1 and game day +1) in MJ. 07:00 was chosen as the 24-hour start point determined by  
154average player wake-up time according to the clubs daily monitoring.

### 155Statistical analysis

156 Statistical tests were performed using the Statistical Package for the Social Sciences (SPSS, Version  
15718). All data were initially checked for normality. Differences between positional groups in mean weekly  
158external (GPS) and internal (sRPE) training load measures were assessed using separate independent *t*-tests.  
159Differences between EE and EI for forwards and backs were analyzed using a two-way mixed design  
160analysis of variance (ANOVA). Differences between macronutrient intakes across time were analyzed using  
161a one-way repeated measures ANOVA. If Mauchley's test of sphericity indicated a minimum level of  
162violation, as assessed by a Greenhouse Geisser epsilon ( $\epsilon$ ) of  $\geq 0.75$ , data were corrected using the Huynh-  
163Feldt  $\epsilon$ . If Mauchley's test of sphericity was violated (Greenhouse Geisser  $\epsilon$  of  $\leq 0.75$ ) data was corrected  
164using the Greenhouse Geisser  $\epsilon$  (Field, 2007). If any significant *F* values were observed least significant  
165difference (LSD) tests were performed *post hoc* to determine where any significant differences occurred. An  
166alpha value of  $P \leq 0.05$  was utilized for all tests. All data are expressed as mean (SD).

167

### 168Results

169

170Weekly external and internal training load



171The backs had significantly higher mean weekly total distances ( $P<0.0005$ ) than forwards, comprising more  
172standing ( $P=0.002$ ), walking ( $P=0.002$ ), jogging ( $P<0.0005$ ) high-speed running ( $P<0.0005$ ) and sprinting  
173( $P=0.004$ ). The backs performed more accelerations ( $P=0.0005$ ) but less contacts ( $P<0.0005$ ) than forwards  
174during a training week. However, there was no difference in the weekly number of RHIE during training  
175between the forwards and backs ( $P=0.15$ ). The total weekly sRPE was not different between positions  
176( $P=0.13$ ), with values of 1776 (355) AU and 1523 (434) AU for forwards and backs, respectively. All  
177external and internal training load data are presented in Table 3.

178

### 179Energy intake and expenditure

180 Energy intake (EI) and expenditure (EE) over the six assessment days, presented in megajoules  
181(MJ), are shown in Figure 1. Mean EI and EE was 16.6 (1.25) MJ and 15.9 (0.53) MJ, and 14.2 (1.2) MJ and  
18214 (0.47) MJ for forwards and backs, respectively. There was a significant time x condition interaction for EI  
183and EE intake over the course of the training week ( $P<0.0005$ ). *Post hoc* analysis confirmed that there was a  
184significant change in EI ( $P<0.0005$ ) and EE ( $P<0.0005$ ) intake over the week. EI was significantly lower  
185than EE on GD-5 ( $P=0.022$ ) and GD-4 ( $P=0.002$ ), and significantly higher on GD-1 ( $P=0.002$ ) and GD+1  
186( $P=0.014$ ) for the forwards. EI was significantly lower than EE on GD-5 ( $P=0.01$ ), GD-4 ( $P<0.0005$ ) and  
187GD-3 ( $P=0.042$ ), and significantly higher on GD-1 ( $P<0.0005$ ) and GD+1 ( $P<0.0005$ ) for the backs. EE  
188( $P=0.024$ ) and EI ( $P=0.046$ ) were significantly different from GD-5 for forwards. EE ( $P=0.033$ ) and EI  
189( $P=0.045$ ) were significantly different from GD-5 for the backs on GD-2 and GD-1 respectively. The  
190forwards elicited significantly higher weekly EI ( $P=0.0006$ ) and EE ( $P=0.002$ ) than the backs.

191

### 192Macronutrient profile

193 Macronutrient intakes from six-day food diaries (presented in  $\text{g}\cdot\text{kg}^{-1}$  body mass) can be seen in  
194Figure 2. There was no difference ( $P=0.53$ ) in mean weekly CHO intake between the forwards and backs,  
195with values of 3.5 (0.8)  $\text{g}\cdot\text{kg}^{-1}$  (38% total calories) and 3.4 (0.7)  $\text{g}\cdot\text{kg}^{-1}$  (37% total calories), respectively.  
196Similarly, mean weekly protein intake between the forwards was 2.7 (0.5)  $\text{g}\cdot\text{kg}^{-1}$  (30% total calories)

197 compared to backs 2.7 (0.3) g·kg<sup>-1</sup> (30% of total calories) and was not different ( $P=0.97$ ). Mean fat intake  
198 was also similar between positions ( $P=0.8$ ), with values of 1.4 (0.2) and 1.4 (0.3) g·kg<sup>-1</sup> (32 and 33% of total  
199 calories) for forwards and backs, respectively.

200 For forwards CHO intake changed during the week ( $P=0.004$ ), with *post hoc* analysis confirming a  
201 higher CHO intake on GD-4 ( $P=0.036$ ) and GD-1 ( $P=0.008$ ) compared with GD-5 (the first day of the  
202 week). This coincided with changes in protein intake during the same period ( $P=0.008$ ), which comprised an  
203 increase in protein intake on GD-1 ( $P<0.0005$ ) and GD+1 ( $P=0.049$ ) compared with GD-5. Fat intake for  
204 forwards did not change during the week ( $P=0.093$ ). CHO intake also changed during the week for backs  
205 ( $P=0.003$ ), with a higher CHO intake on GD-2 ( $P=0.003$ ) and GD-1 ( $P=0.014$ ) compared with GD-5.  
206 Similarly, there was a significant difference in protein intake across the week ( $P<0.0005$ ), with *post hoc*  
207 analysis confirming an increase in protein intake on GD-4 ( $P=0.045$ ) and GD-2 ( $P=0.026$ ), GD-1 ( $P<0.0005$ )  
208 and GD+1 ( $P=0.004$ ) compared with GD-5. Fat intake did change during the week ( $P=0.003$ ), whereby  
209 values increased on GD-1 ( $P=0.0005$ ) and GD+1 ( $P=0.014$ ) compared with GD-5.

210

### 211 **Micronutrient profile**

212 Daily average micronutrient intakes for the squad taken from six-day food diaries can be seen in  
213 Table 3. Mean micronutrient intakes met and exceeded RDA's for physical activity for all minerals and  
214 vitamins apart from vitamin K which fell 24µg under this RDA but met and exceeded the RDA for general  
215 health.

## 216 Discussion

217 The aims of the present study were 1) to quantify the external and internal training loads during a  
218 typical in-season training week for elite rugby union players and 2), evaluate the EI and EE of elite rugby  
219 players during the competitive season. It is reported for the first time that distances of ~8-10 km are covered  
220 by elite rugby union forwards and backs during a typical in-season week, which equates to a total weekly  
221 internal load of ~1500 - 1800 AU. We also report that the daily EE and EI of elite rugby union players  
222 during this same training period are 14-16 and 14-17 MJ, respectively. Our data also demonstrate  
223 considerable variation in the day-to-day energy EE with peak EE occurring early in the week and tapering  
224 down in preparation for competition. Interestingly, although EI also varied on a day-to-day basis, the  
225 temporal pattern did not match EE with EI being the lowest when EI was the highest and EI increasing in  
226 preparation for game-day. This inverse pattern may be essential to allow players to load with carbohydrate in  
227 preparation for game day without excessive total energy intake during the week which over the course of the  
228 season may lead to unwanted gains in body fat.

229 GPS analysis of the training sessions revealed that weekly total distances of  $7.8 \pm 1$  km and  $9.6 \pm 1.2$   
230 km were covered by forwards and backs, respectively. These distances were achieved over a five-day period  
231 as players were given rest days before and after game day. Backs covered more distance in all speed zones,  
232 along with a greater number of maximal accelerations but less collisions than forwards. These differences  
233 probably reflect the contrasting training regimes between rugby union forwards and backs. For example,  
234 forwards engage in more activities that involve tackling, rucking, mauling and line-outs, while the backs  
235 perform more acceleration and ball-in-hand running play. Interestingly, the frequency of RHIE was similar  
236 for positional groups despite clear differences in the movement characteristics of forwards and backs. This is  
237 probably explained by how the GPS software detects RHIE, which is defined as three consecutive efforts  
238 (sprint, contact or acceleration) each separated by  $<21$  s (Gabbett et al., 2013). So, while both positional  
239 groups perform a similar number of high intensity bouts, the movement actions that determine the RHIE are  
240 likely to be different between forwards and backs.

241 Weekly sRPE of ~1778 and ~1522 AU were observed for forwards and backs. These values are

242 lower than those seen in an elite rugby union players during pre-season (~2900-3400 AU; (Bradley et al.,  
243 2014) and reflects the periodization of a rugby training programme. Indeed, lower training loads during the  
244 competitive season are deliberately administered to allow optimal recovery and for players to peak around  
245 games, whereas higher training loads are used in the pre-season when physiological adaptation is key and  
246 competition not a priority. Despite experiencing a higher external load during a training week, forwards did  
247 exhibit a higher internal load (sRPE) than backs. While sRPE is an appropriate measure of training load in  
248 rugby players (Lovell, Sirotic, Impellizzeri & Coutts, 2013), variances in perceptual responses will be  
249 influenced by several internal and external factors. Here, differences in perceived weekly load between  
250 positions is probably explained by higher number of collisions experienced in training by forwards. Indeed,  
251 collisions have been purported to contribute significantly to the variance in sRPE between players during  
252 rugby training (Lovell et al., 2013). Our findings reaffirm the complexity of factors influencing perceptual  
253 measures of training load and the necessity to adopt both internal and external measures to monitor training  
254 in rugby.

255

256 Energy expenditure changed during the training week for both forwards and backs, with higher EE  
257 elicited during the first four days of the training week and significantly reducing around competition (Figure  
258 1). The six-day food diary also revealed changes in EI during the training week for both forwards and backs,  
259 following an inverse trend to EE (Figure 1). Fluctuations in EI represent lower intakes during the first 4-  
260 days of the training week concurrent with higher EE. This is likely attributed to rugby players attempting to  
261 reduce or maintain body fat before significantly increasing EI by increasing carbohydrate intake leading up  
262 to competition in an attempt to increase muscle glycogen stores. It is possible that players might have  
263 intentionally (Bingham, 1987; Deakin, 2000) underreported their total energy intake. However, since  
264 approximately half of the daily nutrition consumed was observed by the authors, including a meal provided  
265 on arrival at the club, whey protein before and after training, and a lunch provided post-training, this seems  
266 unlikely.

267 Although our data indicate lower training loads and total distances compared to those of rugby union  
268 players in pre-season (Bradley et al., 2014), mean EI was slightly higher in-season for both forwards (15.8

269cf. 14.8 MJ) and backs (14.1 cf. 13.3 MJ). We attribute this to players increasing total EI in the days leading  
270up to competition. It must be stressed, however, that the pre-season study used a 24-hour dietary recall,  
271which might compromise the comparison between the two studies. Interestingly, while EE and EI differ on a  
272day-to-day basis, mean EE and EI were surprisingly similar for forwards (16.6 and 15.8 MJ) and backs (14.2  
273and 14.1 MJ). This suggests that although athletes might fail to meet energy requirements on some training  
274days, light training days or rest days before a game correspond with players increasing EI (mainly through  
275CHO increases) to maximise muscle glycogen stores. The lower EI early in the week may be necessary to  
276prevent a positive energy balance that over the course of a season could result in unwanted gain in body-fat.

277         Interestingly, all the players in the present study self-selected what could be classed as a low CHO /  
278high protein diet for the first four days of the training week, and increased CHO intake the day before game  
279day. This practice contravenes earlier recommendations for CHO intakes of 8-12 g·kg<sup>-1</sup> (Burke, Kiens, &  
280Ivy, 2004), as well as more recent guidelines that state values of 6-10 g·kg<sup>-1</sup> (Burke et al., 2011) for athletes  
281engaged in moderate to high intensity exercise lasting 1-3 hours. Therefore it could still be argued that  
282players in the current study failed to meet the daily recommended CHO requirements. However, current  
283guidelines also clearly state that CHO intakes should be designed to meet the fuel requirements of the  
284training programme (Burke et al., 2011). We therefore propose that players are attempting to match  
285carbohydrate intakes with training demands such that CHO intakes of 4-6 g·kg<sup>-1</sup> body mass are not 'low' and  
286are in fact 'appropriate' for this group of athletes providing CHO intake is increased in the day before and  
287after a game. Given that CHO intake altered significantly over the week and was the main macronutrient  
288contributor in the daily EI fluctuations, we suggest that the players are indeed following the recent guidelines  
289and matching their CHO intakes to the fuel requirements of the training programme. Players could be using  
290some periods of lower CHO intake to enhance training adaptations (Morton et al., 2009) and for the  
291maintenance of low body fat (Morton et al., 2010), yet still increasing glycogen stores in preparation for  
292competition (Hawley, Schabort, Noakes, & Dennis, 1997). Interestingly, backs utilized a two-day load  
293compared with a single day by the forwards. This might reflect 1) a lower CHO intake in the first three days  
294to reduce body fat or 2) a purposeful attempt to increase glycogen more aggressively than the forwards due  
295to the varying physiological challenges of the positions.

296

297 This cycling of CHO intake reported in the present study might be a suitable way of maintaining  
298 weekly energy balance yet still allowing sufficient CHO intake as to increase muscle glycogen and thus  
299 enhance match day performance. Playing performance is however unquestionably improved with a high  
300 CHO diet leading up to team sport based games (Hawley et al., 1997; Jardine, Wiggins, Myburgh, & Noakes,  
301 1988), and although we have reported a significant increase in CHO intake in the days leading up to  
302 competition, the intakes reported in this study are still below recommended CHO intake for elite athletes  
303 (Burke et al., 2011). It is still possible that such intakes are not optimal for match day performance and future  
304 studies should now attempt to measure pre and post game muscle glycogen demands in elite rugby.

305 Protein intakes of  $2.7 \text{ g}\cdot\text{kg}^{-1}$  reported in the present study was similar to values reported in an elite  
306 rugby union pre-season ( $2.5$  and  $2.6 \text{ g}\cdot\text{kg}^{-1}$ ; (Bradley et al., 2014). These intakes are much higher than the  
307  $1.4 \text{ g}\cdot\text{kg}^{-1}$  reported in soccer (Maughan, 1997) and  $1.8 \text{ g}\cdot\text{kg}^{-1}$  described for strength based athletes (Tipton &  
308 Wolfe, 2004). However, to maintain muscle mass whilst decreasing body fat, protein intakes of  $2.5 \text{ g}\cdot\text{kg}^{-1}$   
309 have been recommended (Mettler, Mitchell, & Tipton, 2010) suggesting that the protein intakes in this study  
310 might in fact have been appropriate. Moreover, the athletes in the present study would have deliberately  
311 timed protein intakes around training in an attempt to maximise muscle protein synthesis, which might  
312 explain in these higher protein intakes. The backs significantly increased protein intake from four days  
313 before the match. However, this higher protein intake early in the week could simply be the lower CHO  
314 intake in this group early in the week with protein being used as a CHO substitute. Dietary fat intakes in the  
315 present study were approximately  $1.4 \text{ g}\cdot\text{kg}^{-1}$  body mass, slightly higher than the current recommendations  
316 (Bishop, Blannin, Walsh, Robson, & Gleeson, 1999) but similar to those seen in elite Australian athletes  
317 (Burke et al., 2003). Consumption of oily fish, meats, and the use of cooking oils accounts for most of the fat  
318 intake, and although intakes were high, given the importance of healthy fats for performance it would be  
319 unwise to suggest a reduction in dietary fat intake.

320 Micronutrient intakes met and exceeded the RDAs for physical activity (Whiting & Barabash, 2006)  
321 for all minerals and vitamins apart from vitamin K, which fell slightly below the guidelines for physical

322activity (-24 µg, less than 1 small stem of broccoli, see Table 3). These values did, however, meet and  
323exceeded the RDA for general health. Although supplement use is common practice in sport with 40 to  
324100% of athletes using supplements (Baume, Hellemans, & Saugy, 2007), it seems inappropriate to  
325supplement the athletes in this study with a multi-vitamin or a mega dose single vitamin supplement given  
326the lack of any micronutrient deficiencies (Whiting & Barabash, 2006). The exception to this could be  
327vitamin D with recent data suggesting the current RDA for general health is too low (Holick & Chen, 2008)  
328and deficiencies are commonplace in many athletes (Close et al., 2013). Future studies might wish to  
329investigate blood vitamin D concentrations in elite rugby players to further investigate this hypothesis.

330        Although this study provides novel data for the literature it is not without limitations, many of which  
331are a direct consequence of collecting data from elite athletes outside of the controlled laboratory  
332environment. These data were collected on a single professional rugby team, which may not accurately  
333represent every rugby club, and therefore future studies might choose to collect data from a variety of teams.  
334Finally, energy expenditure was not assessed on game day due to dangers associated with the use of the  
335utilized EE technology in contact sports where physical collisions could cause damage to both player and  
336equipment. The use of DLW over the course of a seven-day training week should now be performed. Energy  
337intake was not assessed on game day to avoid adding to player's game day stresses while performing at an  
338elite standard.

339

340        To conclude, for the first time this study has attempted to quantify the training demands and assess  
341energy expenditure, intake and micronutrient intakes of elite rugby players during the in-season. We report  
342that mean energy intake and expenditure followed an inverse trend, with expenditure exceeding intake during  
343the first four-days of the training week and then reversed in the day leading up to competition with intake  
344exceeding expenditure. This is likely due to a heavier training load and players desire to maintain body fat  
345during the beginning of the training week, followed by a decrease in training load and increase in CHO  
346intake leading up to competition in order to maximise glycogen stores. Interestingly, mean energy intake  
347exceeded expenditure for both forwards and backs despite CHO consumption falling short of recommended

348 guidelines. This is likely attributable to relatively low training loads and running distances that attempt to  
 349 provide sufficient stimulus to maintain player strength and fitness during the in-season, while reducing  
 350 residual fatigue and promoting competition preparation. Alongside no micronutrient deficiencies, the current  
 351 dietary practices of these elite rugby players are sufficient to fuel training during the in-season, providing  
 352 energy intake and CHO are increased leading up to a match. However, whether this intake is optimal for  
 353 game day performance remains unknown.

## 354 References

- 355  
 356
- 357 Ainsworth, B. E., Haskell, W. L., Herrmann, S. D., Meckes, N., Bassett, D. R., Jr., Tudor-Locke,  
 358 C., . . . Leon, A. S. (2011). 2011 Compendium of Physical Activities: a second update of  
 359 codes and MET values. *Med Sci Sports Exerc*, 43(8), 1575-1581. doi:  
 360 10.1249/MSS.0b013e31821ece12
- 361 Baume, B., Hellemans, L., & Saugy, M. (2007). Guide to over-the-counter sports supplements for  
 362 athletes *International Sports Medicine Journal*, 8, 2-10
- 363 Bingham, S. (1987). Definitions and intakes of dietary fiber. *Am J Clin Nutr*, 45(5 Suppl), 1226-  
 364 1231
- 365 Bishop, N. C., Blannin, A. K., Walsh, N. P., Robson, P. J., & Gleeson, M. (1999). Nutritional  
 366 aspects of immunosuppression in athletes. *Sports Med*, 28(3), 151-176
- 367 Borg, G., Hassmen, P., & Lagerstrom, M. (1987). Perceived exertion related to heart rate and blood  
 368 lactate during arm and leg exercise. *Eur J Appl Physiol Occup Physiol*, 56(6), 679-685
- 369 Braakhuis, A. J., Meredith, K., Cox, G. R., Hopkins, W. G., & Burke, L. M. (2003). Variability in  
 370 estimation of self-reported dietary intake data from elite athletes resulting from coding by  
 371 different sports dietitians. *Int J Sport Nutr Exerc Metab*, 13(2), 152-165
- 372 Bradley, W., Cavanagh, B., Douglas, W., Donovan, T. F., Morton, J. P., & Close, G. L. (2014).  
 373 Quantification of training load, energy intake and physiological adaptations during a Rugby  
 374 pre season: A case study from an Elite European Rugby Union Squad. *J Strength Cond Res*.  
 375 doi: 10.1519/JSC.0000000000000631
- 376 Burke, L., Kiens, B., & Ivy, J. (2004). Carbohydrates and fat for training and recovery. *Journal of*  
 377 *Sports Sciences*, 22(1), 15-30. doi: 10.1080/0264041031000140527
- 378 Burke, L. M., Hawley, J. A., Wong, S. H., & Jeukendrup, A. E. (2011). Carbohydrates for training  
 379 and competition. *J Sports Sci*, 1-11. doi: 938533953 [pii]
- 380 10.1080/02640414.2011.585473
- 381 Burke, L. M., Slater, G., Broad, E. M., Haukka, J., Modulon, S., & Hopkins, W. G. (2003). Eating  
 382 patterns and meal frequency of elite Australian athletes. *Int J Sport Nutr Exerc Metab*,  
 383 13(4), 521-538
- 384 Cahill, N., Lamb, K., Worsfold, P., Headey, R., & Murray, S. (2013). The movement characteristics  
 385 of English Premiership rugby union players. *J Sports Sci*, 31(3), 229-237. doi:  
 386 10.1080/02640414.2012.727456
- 387 Close, G. L., Russell, J., Copley, J. N., Owens, D. J., Wilson, G., Gregson, W., . . . Morton, J. P.  
 388 (2013). Assessment of vitamin D concentration in non-supplemented professional athletes



- 389 and healthy adults during the winter months in the UK: implications for skeletal muscle  
390 function. *J Sports Sci*, 31(4), 344-353. doi: 10.1080/02640414.2012.733822
- 391 Cummins, C., Orr, R., O'Connor, H., & West, C. (2013). Global positioning systems (GPS) and  
392 microtechnology sensors in team sports: a systematic review. *Sports Med*, 43(10), 1025-  
393 1042. doi: 10.1007/s40279-013-0069-2
- 394 Cunniffe, B., Proctor, W., Baker, J. S., & Davies, B. (2009). An evaluation of the physiological  
395 demands of elite rugby union using Global Positioning System tracking software. *J Strength  
396 Cond Res*, 23(4), 1195-1203. doi: 10.1519/JSC.0b013e3181a3928b
- 397 Deakin, V. (Ed.). (2000). *Measuring nutritional status of athletes: clinical and research  
398 perspectives*. Sydney, Australia: McGraw- Hill Book Company.
- 399 Drenowatz, C., & Eisenmann, J. C. (2011). Validation of the SenseWear Armband at high intensity  
400 exercise. *Eur J Appl Physiol*, 111(5), 883-887. doi: 10.1007/s00421-010-1695-0
- 401 Duthie, G., Pyne, D., & Hooper, S. (2003). Applied physiology and game analysis of rugby union.  
402 *Sports Med*, 33(13), 973-991
- 403 Ekelund, U., Yngve, A., Westerterp, K., & Sjostrom, M. (2002). Energy expenditure assessed by  
404 heart rate and doubly labeled water in young athletes. *Med Sci Sports Exerc*, 34(8), 1360-  
405 1366
- 406 Field, A. (2007). *Discovering statistics using SPSS* (Third ed.). London, UK: Sage Publications.
- 407 Foster, C., Florhaug, J. A., Franklin, J., Gottschall, L., Hrovatin, L. A., Parker, S., . . . Dodge, C.  
408 (2001). A new approach to monitoring exercise training. *J Strength Cond Res*, 15(1), 109-  
409 115
- 410 Fowles, J. R. (2006). Technical issues in quantifying low-frequency fatigue in athletes. *Int J Sports  
411 Physiol Perform*, 1(2), 169-171
- 412 Gabbett, T. J., Wiig, H., & Spencer, M. (2013). Repeated high-intensity running and sprinting in  
413 elite women's soccer competition. *Int J Sports Physiol Perform*, 8(2), 130-138
- 414 Gamble, P. (2006). Periodization of training for team sports athletes. *Strength and Conditioning  
415 Journal*, 28, 56-66
- 416 Hawley, J. A., Schabort, E. J., Noakes, T. D., & Dennis, S. C. (1997). Carbohydrate-loading and  
417 exercise performance. An update. *Sports Med*, 24(2), 73-81
- 418 Holick, M. F., & Chen, T. C. (2008). Vitamin D deficiency: a worldwide problem with health  
419 consequences. *The American Journal of Clinical Nutrition*, 87(4), 1080S-1086S. doi:  
420 87/4/1080S [pii]
- 421 Jardine, M. A., Wiggins, T. M., Myburgh, K. H., & Noakes, T. D. (1988). Physiological  
422 characteristics of rugby players including muscle glycogen content and muscle fibre  
423 composition. *S Afr Med J*, 73(9), 529-532
- 424 Kempton, T., Sullivan, C., Bilsborough, J. C., Cordy, J., & Coutts, A. J. (2015). Match-to-match  
425 variation in physical activity and technical skill measures in professional Australian  
426 Football. *J Sci Med Sport*, 18(1), 109-113. doi: 10.1016/j.jsams.2013.12.006
- 427 Maughan, R. J. (1997). Energy and macronutrient intakes of professional football (soccer) players.  
428 *Br J Sports Med*, 31(1), 45-47
- 429 Mettler, S., Mitchell, N., & Tipton, K. D. (2010). Increased protein intake reduces lean body mass  
430 loss during weight loss in athletes. *Med Sci Sports Exerc*, 42(2), 326-337. doi:  
431 10.1249/MSS.0b013e3181b2ef8e
- 432 Morton, J. P., Croft, L., Bartlett, J. D., Maclaren, D. P., Reilly, T., Evans, L., . . . Drust, B. (2009).  
433 Reduced carbohydrate availability does not modulate training-induced heat shock protein  
434 adaptations but does upregulate oxidative enzyme activity in human skeletal muscle. *J Appl  
435 Physiol*, 106(5), 1513-1521. doi: 00003.2009 [pii]
- 436 10.1152/jappphysiol.00003.2009

- 437Morton, J. P., Robertson, C., Sutton, L., & MacLaren, D. P. (2010). Making the weight: a case  
438 study from professional boxing. *Int J Sport Nutr Exerc Metab*, 20(1), 80-85
- 439Roberts, S. P., Trewartha, G., Higgitt, R. J., El-Abd, J., & Stokes, K. A. (2008). The physical  
440 demands of elite English rugby union. *J Sports Sci*, 26(8), 825-833. doi:  
441 10.1080/02640410801942122
- 442Tipton, K. D., & Wolfe, R. R. (2004). Protein and amino acids for athletes. *J Sports Sci*, 22(1), 65-  
443 79. doi: 10.1080/0264041031000140554
- 444Varley, M. C., Fairweather, I. H., & Aughey, R. J. (2012). Validity and reliability of GPS for  
445 measuring instantaneous velocity during acceleration, deceleration, and constant motion. *J*  
446 *Sports Sci*, 30(2), 121-127. doi: 10.1080/02640414.2011.627941
- 447Varley, M. C., Gabbett, T., & Aughey, R. J. (2014). Activity profiles of professional soccer, rugby  
448 league and Australian football match play. *J Sports Sci*, 32(20), 1858-1866. doi:  
449 10.1080/02640414.2013.823227
- 450Waldron, M., Twist, C., Highton, J., Worsfold, P., & Daniels, M. (2011). Movement and  
451 physiological match demands of elite rugby league using portable global positioning  
452 systems. *J Sports Sci*, 29(11), 1223-1230. doi: 10.1080/02640414.2011.587445
- 453Whiting, S. J., & Barabash, W. A. (2006). Dietary Reference Intakes for the micronutrients:  
454 considerations for physical activity. *Appl Physiol Nutr Metab*, 31(1), 80-85. doi:  
455 10.1139/h05-021
- 456
- 457
- 458
- 459