

DIMENSION OF THE LISBON VOTING RULES IN THE EU COUNCIL: A CHALLENGE AND NEW WORLD RECORD

SASCHA KURZ AND STEFAN NAPEL

ABSTRACT. The new voting system of the Council of the European Union cannot be represented as the intersection of six or fewer weighted games, i.e., its dimension is at least 7. This sets a new record for real-world voting bodies. A heuristic combination of different discrete optimization methods yields a representation as the intersection of 13 368 weighted games. Determination of the exact dimension is posed as a challenge to the community. The system's Boolean dimension is proven to be 3.

Keywords: simple games, weighted games, dimension, real-world voting systems, set covering problem, computational challenges

MSC: 90C06, 05B40, 91B12, 91A12

JEL: C71, C63, D72

1. INTRODUCTION

Consider a group or committee whose members jointly decide whether to accept or reject a proposal (or, more generally, any system which outputs 1 if a minimal set of binary conditions are true and 0 otherwise). The mapping of given configurations of approving members to a collective “yes” (1) or “no” (0) defines a so-called *simple game*. It can often be described by a weighted voting rule: each member i gets a non-negative weight w_i ; a proposal is accepted iff the weight sum of its supporters meets a given quota q . The simple game is then known as a *weighted game*.

Many real-world decision rules can be represented as weighted games, but not all. It is sometimes necessary to consider the intersection of multiple weighted games, or their union, in order to correctly delineate all acceptance and rejection configurations. The minimal number of weighted games whose intersection represents a given simple game is known as its *dimension* [13]; the corresponding number in the disjunctive case is its *co-dimension* [6]. The (co-)dimension of a rule which involves finitely many decision makers is finite, but can grow exponentially in the group size [14, Thm. 1.7.5]. It is NP-hard to determine the exact dimension of a given game [3].

Taylor [12] remarked in 1995 that he did not know of any real-world voting system of dimension 3 or higher. Amendment of the Canadian constitution [9] and the US federal legislative system [13] are classical examples of dimension 2. More recently, systems of dimension 3 have been adopted by the Legislative Council of Hong Kong [2] and the Council of the European Union (EU Council) under its Treaty of Nice rules [5]: until late 2014, each EU member implicitly wielded a 3-dimensional vector-valued weight and proposals were accepted iff their supporters met a 3-dimensional quota. Real-world cases with dimension 4 or more, however, have not been discovered yet (at least to our knowledge). This suggests that determining the dimension of a given simple game might be a hard problem in theory but not in practice.

We establish that the situation is changed by the new voting rules of the EU Council, which were agreed to apply from Nov. 2014 on in the Treaty of Lisbon (with a transition period). They involve a dual majority requirement where (i) at least 55% of the EU member states need to support a motion and (ii) these supporters shall represent at least 65% of the total EU population. However, (iii) the “no”-votes of at least four EU member states are needed in order to block a proposal. A coalition of the 25 smallest among the 28 EU members fails to meet provision (ii) but is still winning due to (iii). We show that representing these rules as the union of one weighted game with the intersection of two more involves no redundancy, even for moderate changes of the current populations. So the *Boolean dimension* (see Def. 1) of (i)–(iii) is 3, and robustly so. Restricting representations to pure intersections or pure unions, however, increases the minimal number of weighted constituent games significantly.

We can prove that the dimension of the EU28's new voting rules is an integer between 7 and 13 368; its co-dimension lies above 2000. This makes the EU28 a new record holder among real-world institutions.

The *determination of the exact dimension* of voting rules in the EU Council is an open computational challenge, which we here wish to present to a wider audience. It is related to the classical set covering problem in combinatorics and computer science.

The EU voting rules aside, the paper provides a general algorithmic approach for determining the dimension of simple games. We combine combinatorial and algebraic techniques, exact and heuristic optimization methods in ways that are open to other applications and further refinements. This contrasts with previously mostly tailor-made arguments for specific group decision rules.

2. NOTATION AND DEFINITIONS

We first introduce notation and some selected results on simple games; [14] is recommended for a detailed treatment. Given a finite set $N = \{1, \dots, n\}$ of *players*, a *simple (voting) game* v is a mapping $2^N \rightarrow \{0, 1\}$ from the subsets of N , called *coalitions*, to $\{0, 1\}$ (interpreted as a collective “no” and “yes”) which satisfies $v(\emptyset) = 0$, $v(N) = 1$, and $v(S) \leq v(T)$ for all $\emptyset \subseteq S \subseteq T \subseteq N$. Coalition $S \subseteq N$ is called *winning* if $v(S) = 1$ and *losing* otherwise. If S is winning but all of its proper subsets are losing, then S is called a *minimal winning coalition*. Similarly, a losing coalition T whose proper supersets are winning is called a *maximal losing coalition*. A simple game is more compactly characterized by its set \mathcal{W}^m of minimal winning coalitions than by the corresponding set \mathcal{W} of winning coalitions (or, equivalently, by its set \mathcal{L}^M of maximal losing coalitions rather than the set \mathcal{L} of all losing coalitions).

Players of a simple game can often be ranked according to their ‘influence’ or ‘desirability’. Namely, if $v(S \cup \{i\}) \geq v(S \cup \{j\})$ for players $i, j \in N$ and all $S \subseteq N \setminus \{i, j\}$ then we write $i \sqsupseteq j$ (or $j \sqsubseteq i$) and say that player i is *at least as influential* as player j . The case $i \sqsupseteq j$ and $j \sqsupseteq i$ is denoted as $i \sqsubseteq j$; we then say that both players are *equivalent*. The \sqsubseteq -relation partitions the set of players into equivalence classes. It is possible that neither $i \sqsupseteq j$ nor $j \sqsupseteq i$ holds, i.e., players may be incomparable. A simple game v is called *complete* if the binary relation \sqsubseteq is complete, i.e., $i \sqsupseteq j$ or $j \sqsupseteq i$ for all $i, j \in N$. Complete simple games form a proper subclass of simple games.

Given a complete simple game v , a minimal winning coalition S is called *shift-minimal winning* if $S \setminus \{i\} \cup \{j\}$ is losing for all $i \in S$ and all $j \in N \setminus S$ with $i \sqsupseteq j$ but not $i \sqsubseteq j$, i.e., S would become losing if any of its players i were replaced by a strictly less influential player j . Similarly, a maximal losing coalition T is called *shift-maximal losing* if $T \setminus \{i\} \cup \{j\}$ is winning for all $i \in T$ and $j \in N \setminus T$ with $j \sqsupseteq i$ but not $i \sqsubseteq j$. A complete simple game is most compactly characterized by the partition of the players into equivalence classes and a description of either the shift-minimal winning or shift-maximal losing coalitions.

If there exist weights $w_i \in \mathbb{R}_{\geq 0}$ for all $i \in N$ and a quota $q \in \mathbb{R}_{> 0}$ such that $v(S) = 1$ iff $w(S) := \sum_{i \in S} w_i \geq q$ for all coalitions $S \subseteq N$ then we call the simple game v *weighted*. Every weighted game is complete but the converse is false. We call the vector (q, w_1, \dots, w_n) a *representation* of v and write $v = [q; w_1, \dots, w_n]$. If v is weighted, there also exist representations such that all weights and the quota are integers. If $\sum_{i=1}^n w_i$ is minimal with respect to the integrality constraint, we speak of a *minimum sum integer representation* (see, e.g., [10]).

If v_1, v_2 are weighted games with identical player set N and respective sets of winning coalitions \mathcal{W}_1 and \mathcal{W}_2 then the winning coalitions of $v_1 \wedge v_2$ are given by $\mathcal{W}_1 \cap \mathcal{W}_2$. The smallest number k such that a simple game v coincides with the intersection $v_1 \wedge \dots \wedge v_k$ of k weighted games with identical player set is called the *dimension* of v . Similarly, the winning coalitions of $v_1 \vee v_2$ are $\mathcal{W}_1 \cup \mathcal{W}_2$, and the smallest number of weighted games whose union $v_1 \vee \dots \vee v_k$ coincides with a simple game v is the *co-dimension* of v . Freixas and Puente have shown that there exists a complete simple game with dimension k for every integer k [7]. It is not known yet whether the dimension of a complete simple game is polynomially bounded in the number of its players or can grow exponentially (like for general simple games).

Lemma 1. (cf. [14, Theorem 1.7.2]) *The dimension of a simple game v is bounded above by $|\mathcal{L}^M|$ and the co-dimension is bounded above by $|\mathcal{W}^m|$.*

Proof. For each coalition $S \in \mathcal{L}^M$ we set $q^S = 1$, $w_i^S = 0$ for all $i \in S$ and $w_i^S = 1$ otherwise. Note that $\mathcal{L}^M \neq \emptyset$ since \emptyset is a losing coalition. With this $w^S(S) = 0 < q^S$. However, for all $T \subseteq N$ with $T \not\subseteq S$ we have $w(T) \geq 1 = q^S$. Thus, we have $v = \bigwedge_{S \in \mathcal{L}^M} [q^S; w_1^S, \dots, w_n^S]$. Similarly, for each $S \in \mathcal{W}^m$ we set $\tilde{q}^S = |S|$, $\tilde{w}_i^S = 1$ for all $i \in S$ and $w_i^S = 0$ otherwise. Note that $\mathcal{W}^m \neq \emptyset$ since N is a winning

#	Member state	Population	w_2	#	Member state	Population	w_2
1	Germany	80 780 000	4 659 052	16	Bulgaria	7 245 677	417 900
2	France	65 856 609	3 798 333	17	Denmark	5 627 235	324 556
3	United Kingdom	64 308 261	3 709 031	18	Finland	5 451 270	314 406
4	Italy	60 782 668	3 505 689	19	Slovakia	5 415 949	312 369
5	Spain	46 507 760	2 682 373	20	Ireland	4 604 029	265 541
6	Poland	38 495 659	2 220 268	21	Croatia	4 246 700	244 932
7	Romania	19 942 642	1 150 208	22	Lithuania	2 943 472	169 767
8	Netherlands	16 829 289	970 643	23	Slovenia	2 061 085	118 875
9	Belgium	11 203 992	646 199	24	Latvia	2 001 468	115 436
10	Greece	10 992 589	634 006	25	Estonia	1 315 819	75 890
11	Czech Republic	10 512 419	606 312	26	Cyprus	858 000	49 486
12	Portugal	10 427 301	601 403	27	Luxembourg	549 680	31 703
13	Hungary	9 879 000	569 780	28	Malta	425 384	24 535
14	Sweden	9 644 864	556 276				
15	Austria	8 507 786	490 693		Total	507 416 607	2 9265 662

TABLE 1. EU population 01.01.2014 (<http://ec.europa.eu/eurostat>); minimum sum integer weights of v_2

coalition. With this $\tilde{w}^S(S) = \tilde{q}^S$. However, for all $T \subseteq N$ with $S \not\subseteq T$ we have $w(T) < \tilde{q}^S$. Thus, we have $v = \bigvee_{S \in \mathcal{W}^m} [\tilde{q}^S; \tilde{w}_1^S, \dots, \tilde{w}_n^S]$. \square

Let $\Phi = \{u_1, \dots, u_k\}$ be a set of weighted games, interpreted as Boolean variables, and let φ be a *monotone Boolean formula* over Φ , i.e., a well-formed formula of propositional logic over Φ which uses parentheses and the operators \wedge and \vee only. The *size* $|\varphi|$ of formula φ is the number of variable occurrences, i.e., the number of \wedge and \vee operators plus one. For instance, the size of $u_1 \vee (u_1 \wedge u_2)$ is 3.

Definition 1. *The Boolean dimension of a simple game v is the smallest integer m such that there exist $k \leq m$ weighted games u_1, \dots, u_k and a monotone Boolean formula φ of size $|\varphi| = m$ satisfying $\varphi(u_1, \dots, u_k) = v$.*

Clearly, the Boolean dimension of v is at most the minimum of v 's dimension and co-dimension. Because combinations of \wedge with \vee have a size of at least 3, the Boolean dimension must exceed 2 whenever the dimension and co-dimension do. The dimension can be exponential in the Boolean dimension of a simple game [4, Thm. 4]; the Boolean dimension of a simple game can be exponential in the number of players [4, Cor. 2].

3. LISBON VOTING RULES IN EU COUNCIL

We now formalize the provisions (i)–(iii) for decision making by the EU Council (see Sec. 1). The membership requirement (i) – approval of at least $16 = \lceil 0.55 \cdot 28 \rceil$ member states – is easily reflected by the weighted game $v_1 = [16; 1, \dots, 1]$. The population requirement (ii) could be represented by using the official population counts as weights and 65% of the total population as quota (see Table 1). Its computationally more convenient minimum sum integer representation is given by $v_2 = [q; \mathbf{w}_2]$ with $q = 19\,022\,681$ and the weights indicated in the w_2 -columns of Table 1.¹ The additional minimal blocking requirement (iii) can be described as $v_3 = [25; 1, \dots, 1]$, since $28 - 4 + 1 = 25$ member states suffice to pass a proposal. The Lisbon voting rule of the EU Council is then formally characterized as $v_{\text{EU28}} = (v_1 \wedge v_2) \vee v_3$ or $v_{\text{EU28}} = v_1 \wedge (v_2 \vee v_3)$.

The 268 435 456 coalitions of v_{EU28} are partitioned into 30 340 718 winning and 238 094 738 losing coalitions. Of these, 8 248 125 are minimal winning and 7 179 233 maximal losing. So the dimension of v_{EU28} must be below 7.18 millions.

¹We remark that *rounding* populations to, say, thousands is common in applied work because this simplifies computations, e.g., of the voting power distribution in the EU Council. Rounding, however, leads to a different set of winning coalitions, i.e., is analyzing ‘wrong’ rules.

The influence partition of the Boolean combination of weighted games generally corresponds to the coarsest common refinement of the respective partitions in the constituent games. Here, there is only a single equivalence class of players in v_1 and v_3 , respectively, while v_2 has 28 equivalence classes (all minimum sum weights differ by at least 2). So each player forms its own equivalence class in v_{EU28} . There are only 60 607 shift-minimal winning and 60 691 shift-maximal losing coalitions in v_{EU28} .²

4. WEIGHTEDNESS AND BOUNDING STRATEGY

Determining whether a given simple game is weighted or not will be crucial for our analysis of v_{EU28} . Answers can be given by combinatorial, algebraic or geometric methods (see [14, Ch. 2]). We will draw on the first two.

Combinatorial techniques usually invoke so-called ‘trades’. A *trading transform* for a simple game v is a collection of coalitions $J = \langle S_1, \dots, S_j; T_1, \dots, T_j \rangle$ such that $|\{h: i \in S_h\}| = |\{h: i \in T_h\}|$ for all $i \in N$. An m -trade for v is a trading transform with $j \leq m$ such that all S_h are winning and all T_h are losing coalitions. Existence of, say, a 2-trade $\langle S_1, S_2; T_1, T_2 \rangle$ implies that the game cannot be weighted: $w(S_1), w(S_2) \geq q$ and $w(T_1), w(T_2) < q$ would contradict $w(S_1) + w(S_2) = w(T_1) + w(T_2)$. The simple game v is called *m-trade robust* if no m -trade exists for it. Taylor and Zwicker have shown that a simple game is weighted iff it is $m = 2^{2^n}$ -trade robust (see, e.g., [14, Thm. 2.4.2]). Sharper bounds for m have been provided by [8], but the lower one is still linear and the upper exponential in n .

Example 1. Consider the complete simple game v with $N = \{1, 2, 3, 4, 5, 6\}$ and

$$\mathcal{L}^M = \{\{1, 3, 5\}, \{1, 3, 6\}, \{1, 4, 5\}, \{1, 4, 6\}, \{2, 3, 5\}, \{2, 3, 6\}, \{2, 4, 5\}, \{2, 4, 6\}\}.$$

All coalitions in \mathcal{L}^M are also shift-maximal losing, but only coalitions $\{1, 2\}$, $\{1, 3, 4\}$, $\{2, 3, 4\}$ and $\{3, 4, 5, 6\}$ of

$$\mathcal{W}^m = \{\{1, 2\}, \{1, 3, 4\}, \{2, 3, 4\}, \{3, 4, 5, 6\}, \{1, 3, 5, 6\}, \{1, 4, 5, 6\}, \{2, 3, 5, 6\}, \{2, 4, 5, 6\}\}$$

are also shift-minimal winning. Since

$$\langle \{1, 2\}, \{3, 4, 5, 6\}; \{1, 3, 5\}, \{2, 4, 6\} \rangle$$

is a 2-trade, v is not weighted.³

Algebraic methods exploit that a simple game v is weighted iff the inequality system $\sum_{i \in S} w_i \geq q \forall S \in \mathcal{W}^m$, $\sum_{i \in T} w_i \leq q - 1 \forall T \in \mathcal{L}^M$, $w_i \in \mathbb{R}_{\geq 0} \forall i \in N$, and $q \in \mathbb{R}_{\geq 1}$ admits a solution. Linear programming (LP) techniques can be applied. In case that no solution exists, the dual multipliers provide a certificate of non-weightedness. A suitable subset of the constraints – those for the minimal winning and some maximal losing coalitions, say – often suffice to conclude infeasibility and thus non-weightedness.

For a complete simple game v with sets \mathcal{W}^{sm} and \mathcal{L}^{sM} of shift-minimal winning and shift-maximal losing coalitions, the linear inequality system can further be simplified. Namely, v is weighted iff

$$(1) \quad \begin{aligned} & \sum_{i \in S} w_i \geq q \forall S \in \mathcal{W}^{sm}, \quad \sum_{i \in T} w_i \leq q - 1 \forall T \in \mathcal{L}^{sM}, \\ & w_i \geq w_j \in \mathbb{R}_{\geq 0} \forall i, j \in N \text{ with } i \sqsupset j, \quad w_i \in \mathbb{R}_{\geq 0} \forall i \in N \text{ and } q \in \mathbb{R}_{\geq 1} \end{aligned}$$

admits a solution. Note that non-weightedness of v says no more about v 's dimension than that it exceeds 1.

One might hope that it is possible to construct a representation of a complete simple game v as the intersection of $|\mathcal{L}^{sM}|$ weighted games as follows: look at one coalition $T_l \in \mathcal{L}^{sM}$ at a time; find a weighted game v_l such that (a) $v_l(T_l) = 0$ and (b) $v_l(S) = 1$ for every $S \in \mathcal{W}^{sm}$ by ignoring all constraints $\sum_{i \in T'} w_i \leq q - 1$ in system (1) for $T' \in \mathcal{L}^{sM} \setminus T_l$; finally obtain $v_1 \wedge \dots \wedge v_{|\mathcal{L}^{sM}|}$ as a representation of v . Unfortunately, this does not work in general. For instance, we can infer from infeasibility of $w_1 + w_2 \geq q$, $w_3 + w_4 + w_5 + w_6 \geq q$, $w_1 + w_3 + w_5 \leq q - 1$, $w_1 = w_2$, $w_3 = w_4$ and $w_5 = w_6$ that there exists no weighted game v_1 which respects the ordering condition $w_i \geq w_j \iff i \sqsupset j$ and in which $T_1 = \{1, 3, 5\} \in \mathcal{L}^{sM}$ is losing and (at least) $\{1, 2\}$ and $\{3, 4, 5, 6\}$ are winning (see Example 1). Counterexamples exist also when no two players are equivalent. The basic idea of this heuristic construction is still

²For example, every 16-member winning coalition is minimal but few are also shift-minimal.

³The example is the smallest possible: all complete simple games with $n \leq 5$ are weighted.

useful, and will be applied in order to provide an *upper bound* on v_{EU28} 's dimension. In order to establish a *lower bound*, we will use

Observation 1. *Given a simple game v with winning coalitions \mathcal{W} and losing coalitions \mathcal{L} , let $\mathcal{L}' = \{T_1, \dots, T_k\} \subseteq \mathcal{L}$ be a set of losing coalitions with the following ‘pairwise incompatibility property’: for each pair $\{T_i, T_j\}$ with $T_i \neq T_j \in \mathcal{L}'$ there exists no weighted game in which all coalitions in \mathcal{W} are winning while T_i and T_j are both losing. Then if $v = \bigwedge_{1 \leq l \leq m} v_l$ is the intersection of m weighted games, we must have $m \geq k$, i.e., v 's dimension is at least k .*

The observation generalizes the construction used in [5]. A quick way to establish that there is no weighted game with T_i and T_j losing and all $S \in \mathcal{W}$ winning is to find a 2-trade $\langle S_1, S_2; T_i, T_j \rangle$ for some $S_1, S_2 \in \mathcal{W}$. Not finding a 2-trade does not guarantee that such weighted game exists; and checking for 3-trades, 4-trades, etc. gets computationally demanding. However, in order to provide a lower bound k for v_{EU28} 's dimension, it suffices to provide *any* set \mathcal{L}' of k pairwise incompatible losing coalitions. So one can focus on sets in which 2-trades are easily obtained for all $\binom{k}{2}$ pairs, and improve the resulting bound by extending \mathcal{L}' if needed.

We remark that it is possible to formulate the *exact determination* of the dimension of a simple game as a discrete optimization problem. To this end let \mathcal{C} collect all subsets $\mathcal{S} \subseteq \mathcal{L}^M$ with the property that there exists a weighted game where all elements of \mathcal{W} are winning and all elements of \mathcal{S} are losing. In particular, all singleton subsets of \mathcal{L}^M are contained in \mathcal{C} (cf. proof of Lemma 1); so is, e.g., $\{\{1, 3, 5\}, \{1, 3, 6\}, \{1, 4, 5\}, \{1, 4, 6\}\}$ in Example 1, but not $\{\{1, 3, 5\}, \{2, 4, 6\}\}$.

Having constructed \mathcal{C} , the dimension of v can be determined by finding a *minimal covering* of \mathcal{L}^M , using the elements of \mathcal{C} . Specifically, v 's dimension is the optimal value of $\min \sum_{\mathcal{S} \in \mathcal{C}} x_{\mathcal{S}}$ subject to the constraints $\sum_{\mathcal{S} \in \mathcal{C}: T \in \mathcal{S}} x_{\mathcal{S}} \geq 1$ for all $T \in \mathcal{L}^M$ and $x_{\mathcal{S}} \in \{0, 1\}$ for all $\mathcal{S} \in \mathcal{C}$. However, this set covering formulation is, in general, computationally intractable. For v_{EU28} , already the construction of \mathcal{C} is out of reach because \mathcal{L}^M has more than $2^{7.1 \cdot 10^6}$ subsets. We hence have to contend ourselves with lower and upper bounds which may be brought to identity at some point in the future.

5. BOUNDS FOR v_{EU28} 'S DIMENSION

Since v_{EU28} has so many maximal losing coalitions we have focused our search for a suitable set \mathcal{L}' of pairwise incompatible losing coalitions on the subset $\mathcal{L}_{23,24} \subset \mathcal{L}$ of losing coalitions with 23 or 24 members. They fail the 65% population and 25 member thresholds. For each pair of these 4533 coalitions we have performed a greedy search for a 2-trade. Specifically, let two such losing coalitions $T_i \neq T_j \in \mathcal{L}_{23,24}$ be given, set $I = T_i \cap T_j$, and then extend I to a winning coalition S_1 with 25 members by choosing the least populous elements of $(T_i \cup T_j) \setminus I$. Coalition S_2 is then defined by $((T_i \cup T_j) \setminus S_1) \cup I$. If S_2 is winning, we have found a 2-trade, i.e., pair $\{T_i, T_j\}$ satisfies the incompatibility criterion. Marking this occurrence as an edge in a graph \mathcal{G} with vertex set $\mathcal{L}_{23,24}$, we can perform a clique search on \mathcal{G} . It turns out that \mathcal{G} contains 24 452 800 cliques of size 6 but no larger clique. One of the 6-cliques corresponds to $\mathcal{L}' =$

$$\left\{ \begin{array}{l} \{1, 4, 5, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28\}, \\ \{3, 4, 5, 6, 7, 8, 10, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28\}, \\ \{2, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28\}, \\ \{2, 3, 5, 6, 8, 9, 10, 11, 13, 14, 15, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28\}, \\ \{2, 3, 4, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28\}, \\ \{1, 3, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 21, 22, 23, 24, 25, 26, 27, 28\} \end{array} \right\}^4$$

This 6-clique is actually the most robust one regarding changes of the relative population distribution in the EU: it is not upset by moves between states, births, or deaths as long as the new relative population vector pop' and the old one, pop , based on Table 1, have a $\|\cdot\|_1$ -distance less than 0.0095. This distance could accommodate arbitrary moves of up to 2.5 million EU citizens. The robustness is noteworthy because high numbers in the minimum sum representation of v_2 indicate that v_{EU28} is rather sensitive to population changes.

⁴Just to give an example, $\{\{4, \dots, 28\}, \{1, 3, 4, 5, 7, 8, 10, 12, 15, \dots, 28\}; \{1, 4, 5, 7, \dots, 12, 14, \dots, 28\}, \{3, \dots, 8, 10, 12, 13, 15, \dots, 28\}\}$ is a 2-trade for the first two losing coalitions.

The above set \mathcal{L}' can be extended, without affecting robustness, by adding the maximal losing coalition $\{1, \dots, 15\}$ of the 15 largest member states, which was excluded by the initial focus on $\mathcal{L}_{23,24}$. This establishes:

Proposition 1. *Let v be the simple game arising from v_{EU28} by replacing the underlying relative population vector pop by the relative population vector pop' . If $\|pop - pop'\|_1 \leq 0.95\%$ then v has dimension at least 7.*

An alternative for establishing a lower bound d for v_{EU28} 's dimension is to replace the graph-theoretic search for 2-trades by a straightforward integer linear program (ILP) such as⁵

$$\begin{aligned} \max \Delta \quad \text{s.t.} \quad & \sum_{i=1}^{28} l_i^j \leq 24 \quad \forall 1 \leq j \leq d, \quad \sum_{i=1}^{28} pop_i \cdot l_i^j \leq 0.65 - \Delta \quad \forall 1 \leq j \leq d \\ & \sum_{i=1}^{28} w_i^{j,h,1} \geq 25 \quad \forall 1 \leq j < h \leq d, \quad \sum_{i=1}^{28} pop_i \cdot w_i^{j,h,2} \geq 0.65 + \Delta \quad \forall 1 \leq j < h \leq d \\ & l_i^j + l_i^h = w_i^{j,h,1} + w_i^{j,h,2} \quad \forall 1 \leq i \leq 28, 1 \leq j < h \leq d, l_i^j \in \{0, 1\} \quad \forall 1 \leq i \leq 28, 1 \leq j \leq d \\ & \sum_{i=1}^{28} w_i^{j,h,2} \geq 16 \quad \forall 1 \leq j < h \leq d, w_i^{j,h,k} \in \{0, 1\} \quad \forall 1 \leq i \leq 28, 1 \leq j < h \leq d, k \in \{1, 2\}. \end{aligned}$$

This turned out to be impractical for $d > 6$ but has yielded a simple, robust certificate for $d = 3$, which will be useful for obtaining Corollary 1 below:

Proposition 2. *Let v be the simple game arising from v_{EU28} by replacing the underlying population vector pop by the relative population vector pop' . If $\|pop - pop'\|_1 \leq 2.19\%$ then v has dimension at least 3.*

Proof. Three losing coalitions whose pairs can be completed to a 2-trade are:

$$\{1, 4, 5, 7, 8, 9, 11, \dots, 14, 16, \dots, 26, 28\}, \quad \{3, \dots, 6, 8, 9, 10, 12, 14, \dots, 24, 26, 27\}, \quad \text{and} \\ \{2, 4, \dots, 8, 10, 11, 13, 15, 17, \dots, 20, 22, \dots, 25, 27, 28\}.$$

□

In order to bring down the baseline upper bound of $|\mathcal{L}^M| \approx 7.18$ mio. for v_{EU28} 's dimension (Lemma 1), we draw on LP formulation (1) and the indicated idea to check for each $T_l \in \mathcal{L}^{sM}$ whether inequality system (1) with \mathcal{L}^{sM} replaced by $\{T_l\}$ has a feasible solution. This yields weighted games for 57 869 out of $|\mathcal{L}^{sM}| = 60 691$ coalitions. The remaining 2 822 *stubborn* shift-maximal losing coalitions correspond to exactly 17 003 maximal losing coalitions, which are not yet covered by the identified weighted games. We could apply the construction in the proof of Lemma 1 to these and would obtain an upper bound of 74 872.

This, however, is easily improved by the following procedure: (I) try to greedily cover many shift-maximal losing coalitions with a few selected weighted games; (II) find a weighted game v_j for each still uncovered and non-stubborn $T_j \in \mathcal{L}^{sM}$; (III) deal with the maximal losing coalitions related to all stubborn T_k . We utilized the following ILP in order to iteratively find helpful games in step (I)

$$\begin{aligned} \max \sum_{T \in \mathcal{L}''} x_T \quad \text{s.t.} \quad & x_T \in \{0, 1\} \quad \forall T \in \mathcal{L}'', w_i \geq w_{i+1} \quad \forall 1 \leq i \leq 27, \sum_{i=1}^{28} w_i \leq M, \\ & \sum_{i \in S} w_i \geq q \quad \forall S \in \mathcal{W}^{sm}, \sum_{i \in T} w_i \leq q - 1 + (1 - x_T)M, w_i, q \in \mathbb{N} \quad \forall 1 \leq i \leq 28. \end{aligned}$$

This ILP exploits that $1 \sqsupset \dots \sqsupset 28$ in v_{EU28} , the constant M is chosen so as to give integer weights with suitable magnitude (e.g., thousands), and \mathcal{L}'' is the part of \mathcal{L}^{sM} which is still uncovered or a subset thereof. It is possible, for instance, to cover 34 323 shift-maximal losing coalitions in step (I) with just 10 weighted games. Adding more weighted games to these, the *lowest upper bound* which we have obtained so far is 13 368. The games and a checking tool can be obtained from the authors.

⁵For the general ILP modeling of weighted games we refer to [11].

All of these considerations can easily be translated to the *co-dimension*. There, we have to consider unions of weighted games, where all coalitions in \mathcal{L}^M are losing and the winning coalitions in \mathcal{W}^m end up being covered by a suitable selection of constituent games. We skip the details for space reasons.

Proposition 3. *Let v be the simple game arising from v_{EU28} by replacing the underlying relative population vector pop by the relative population vector pop' . If $\|pop - pop'\|_1 \leq 5\%$ then v has co-dimension at least 7.*

Proof. Seven winning coalitions whose pairs can be completed to a 2-trade are:

$\{2, \dots, 5, 7, 8, 9, 11, \dots, 15, 17, \dots, 20\}$, $\{1, 2, 3, 6, 8, \dots, 15, 17, 18, 19, 25\}$,
 $\{1, 3, 5, \dots, 16, 19, 20\}$ $\{1, 2, 5, \dots, 17, 22\}$, $\{1, 2, 4, 5, 7, 9, \dots, 15, 19, 23, 24, 26\}$,
 $\{1, 2, 4, 6, 8, \dots, 16, 18, 20, 21\}$, and $\{1, 2, 3, 5, 7, 10, \dots, 15, 18, 20, 21, 22, 28\}$. \square

The combination of Propositions 2 and 3 yields:

Corollary 1. *Let v be the simple game arising from v_{EU28} by replacing the underlying relative population vector pop by the relative population vector pop' . If $\|pop - pop'\|_1 \leq 2.19\%$ then v has Boolean dimension exactly 3.*

We remark that is not too hard to determine 2 000 winning coalitions such that each pair can be completed to a 2-trade. So the co-dimension of v_{EU28} with populations exactly as in Table 1 is at least 2 000.

6. CONCLUDING REMARKS

Simple game v_3 rules out that three of the EU's "Big Four" (see Table 1) can cast a veto in the Council. This has very minor consequences for the mapping of different voting configurations to a collective "yes" or "no": the disjunction with v_3 adds a mere 10 to the 30 340 708 coalitions which are already winning in $v_1 \wedge v_2$. Prima facie, provision (iii) should therefore have only symbolic influence on the distribution of voting power in the Council.⁶ Quite surprisingly, however, provision (iii) has tremendous effect on the conjunctive dimensionality of the rules. Namely, the EU Council sets a new world record, among the political institutions that we know of: the dimension of its decision rule is at least 7.

The link to classical set covering problems in optimization which we have identified and partly exploited in Sections 4 and 5 implies that there exist algorithms which should – at least in theory – terminate with an answer to the simple question: what is the dimension of v_{EU28} ? In practice, heuristic methods which establish and improve bounds are needed. The suggested mix of combinatorial and algebraic techniques, integer linear programming and graph-theoretic methods has rather general applicability. It also lends itself to robustness considerations, which we hope will become more popular in the literature. (A potentially negative referendum on EU membership in the UK and a consequent exit, for instance, would leave our lower bounds intact.)

The drawback of our relatively general approach is that the resultant upper bound of 13 368 is still pretty high; the record lower bound of 7 may not be the final word either. Alternative approaches, which might use unexploited specifics of v_{EU28} , will potentially lead to much sharper boundaries in the future.

The certification of better dimension bounds is a problem which we would here like to advertise to the optimization community. The application of meta-heuristics, such as simulated annealing and genetic algorithms, or column generation techniques could be promising. The ultimate challenge is, of course, to determine the *exact dimension* of the group decision rule in the EU Council.

REFERENCES

- [1] J. Alonso-Mejjide, M. Álvarez-Mozos, and M. Fiestras-Janeiro. The least square nucleolus is a normalized Banzhaf value. *Optim. Lett.*, pages 1–7, to appear.
- [2] W.-S. Cheung and T.-W. Ng. A three-dimensional voting system in Hong Kong. *European J. Oper. Res.*, 236(1):292–297, 2014.
- [3] V. G. Deĭneko and G. J. Woeginger. On the dimension of simple monotonic games. *European J. Oper. Res.*, 170(1):315–218, 2006.

⁶In order to check this intuition, we have computed the difference $\|\mathcal{P}(v_{EU28}) - \mathcal{P}(v_1 \wedge v_2)\|_1$ for four different power measures \mathcal{P} (cf. [1]): it is only around $7 \cdot 10^{-7}$ for the least square nucleolus and $9 \cdot 10^{-7}$ for the normalized Banzhaf index, but 0.00537 for the Shapley-Shubik index and 0.167 for the nucleolus.

- [4] P. Faliszewski, E. Elkind, and M. Wooldridge. Boolean combinations of weighted voting games. In *Proceedings of the 8th International Conference on Autonomous Agents and Multiagent Systems*, pages 185–192. International Foundation for Autonomous Agents and Multiagent Systems, 2009.
- [5] J. Freixas. The dimension for the European Union Council under Nice rules. *European J. Oper. Res.*, 156(2):415–419, 2004.
- [6] J. Freixas and D. Marciniak. On the notion of dimension and codimension. In L. A. Petrosyan and N. A. Zenkevich, editors, *Proceedings of the 3rd International Conference Game Theory and Management*, pages 67–81. The International Society of Dynamic Games (Russian Chapter), 2010.
- [7] J. Freixas and M. A. Puente. Dimension of complete simple games with minimum. *European J. Oper. Res.*, 188(2):555–568, 2008.
- [8] T. Gvozdeva and A. Slinko. Weighted and roughly weighted simple games. *Math. Social Sci.*, 61(1):20–30, 2011.
- [9] M. D. Kilgour. A formal analysis of the amending formula of Canada’s Constitution Act. *Canad. J. Pol. Sci.*, 16:771–777, 1983.
- [10] S. Kurz. On minimum sum representations for weighted voting games. *Ann. Oper. Res.*, 196(1):361–369, 2012.
- [11] S. Kurz. On the inverse power index problem. *Optimization*, 61(8):989–1011, 2012.
- [12] A. D. Taylor. *Mathematics and Politics*. Springer, Berlin, 1995.
- [13] A. D. Taylor and W. S. Zwicker. Weighted voting, multicameral representation, and power. *Games Econ. Behav.*, 5(1):170–181, 1993.
- [14] A. D. Taylor and W. S. Zwicker. *Simple Games: Desirability Relations, Trading and Pseudoweightings*. Princeton University Press, 1999.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BAYREUTH, 95440 BAYREUTH, GERMANY., TEL.: +49-921-557353,
FAX: +49-921-557352, SASCHA.KURZ@UNI-BAYREUTH.DE

DEPARTMENT OF ECONOMICS, UNIVERSITY OF BAYREUTH, AND PCRC, UNIVERSITY OF TURKU