

1 **HYPOXIA DOES NOT INFLUENCE THE RESPONSE OF FISH TO A**
2 **MIXTURE OF ESTROGENIC CHEMICALS**

3

4

5 Jayne V. Brian^{1*}, Nicola Beresford¹, Julie Walker¹, Giulio Pojana² Andrea Fantinati²,
6 Antonio Marcomini² and John P. Sumpter¹.

7

8

9 ¹ Institute for the Environment, Brunel University, Middlesex, UK.

10 ² Department of Environmental Sciences, University of Venice, Italy.

11

12

13 * Corresponding author

14 Tel. 01895 266 267

15 Fax. 01895 269 761

16 jayne.brian@brunel.ac.uk

17

18 Institute for the Environment,

19 Brunel University,

20 Uxbridge,

21 Middlesex,

22 UB8 2NG,

23 UK

24

25 **ABSTRACT**

26 Chemical risk assessment procedures assign a major role to standardised toxicity tests,
27 in which the response of a particular organism to a single test substance is determined
28 under otherwise constant and favourable conditions in the laboratory. This approach
29 fails to consider the potential for chemical interactions, as well as failing to consider
30 how the toxicological response varies, depending on the conditions of exposure. As
31 yet, the issue of confounding factors on chemically-mediated effects in wildlife has
32 received little attention, despite the fact that a range of physicochemical parameters,
33 including temperature, water quality and pH, are known to modify chemical toxicity.
34 Here, we consider how the estrogenic response of fish varies with regard to hypoxia.
35 Fathead minnows (*Pimephales promelas*) were exposed to a mixture of estrogenic
36 chemicals under hypoxic or normoxic conditions. Their estrogenic response was
37 characterised using an *in vivo* assay, involving the analysis of the egg yolk protein,
38 vitellogenin (VTG). The results revealed that there was no effect of hypoxia on the
39 VTG response in either treatment group at the end of the exposure period. This
40 suggests that this endpoint is robust and relatively insensitive to the effects of any
41 physiological changes that arise as a result of hypoxia. The implications of these
42 negative findings are discussed in terms of their relevance with regard to the
43 development of risk assessment policy.

44 **KEY WORDS:**

45 Fathead minnow, multiple stress, hypoxia, endocrine disruption, estrogen, mixture,
46 vitellogenin.

47 **1. INTRODUCTION**

48 Hypoxia is a phenomenon that occurs in both marine and freshwater environments,
49 affecting many thousands of km², worldwide. In this context, it is defined as
50 dissolved oxygen (DO) levels of less than 2.8 mg/l (1). These conditions are
51 generally of detriment to the survival of aquatic organisms, having been associated
52 with mass mortalities, benthic defaunation and declining fisheries production (2).
53 Although hypoxia can occur as a result of natural stratification in some systems,
54 through the formation of haloclines and thermoclines, the incidence and extent of this
55 phenomenon has increased in recent decades as a result of excessive inputs of
56 nutrients and organic matter into water bodies with poor circulation. An example is
57 provided by the situation in the northern Gulf of Mexico, where the hypoxic region
58 has averaged over 15,600 km² in size since 1993, as a result of the increased use of
59 nitrate fertilisers (3). Situations such as this are likely to be exacerbated in the future
60 due to the increase in the intensity of agricultural practices and the rate of human
61 population growth in coastal areas, combined with the impacts of global climate
62 change (4).

63 Fish have developed two main strategies for coping with hypoxia. The first is to
64 invoke various behavioural and physiological responses that increase oxygen delivery.
65 For example, ventilation rates are increased and glycolysis, with lactic acid as an end
66 product, is induced to resist the effects of hypoxia in the short term (5). A second
67 strategy, which may be invoked following prolonged exposure, is to conserve energy
68 by metabolic suppression (2). This is apparent from the analysis of gene expression
69 patterns in the mudsucker, *Gillichthys mirabili*, which revealed that cellular growth is
70 suppressed under hypoxic conditions in order to allow energy to be channelled into

71 essential metabolic processes (6). However, in contrast with their capacity to protect
72 against hypoxia, it would appear that these responses are associated with a reduction
73 in the tolerance of fish to simultaneous chemical challenge. Increased toxicity under
74 low oxygen conditions has been demonstrated for a range of micro-pollutants (e.g.
75 cyanide, ammonia) and for some metals (e.g. copper, cadmium), leading to reduced
76 survival of fish in multiple stress exposure situations (7-11). This phenomenon may
77 be linked to the enhanced uptake of toxicants under hypoxic conditions; there is an
78 apparent link between DO, ventilation rate and toxicity (9). However, the evidence
79 surrounding this issue remains equivocal (8).

80 Currently, little is known about the potential influence of hypoxia on the response of
81 fish to endocrine disrupting chemicals (EDCs), such as the environmental estrogens.
82 However, this issue is pertinent for two reasons. Firstly, the input of nutrients into the
83 environment from anthropogenic sources often coincides with the presence of EDCs
84 (e.g. in sewage treatment works effluent), creating multiple stress exposure situations.
85 Secondly, there is growing evidence that hypoxia can, on its own, cause endocrine-
86 mediated disturbances in fish. Whilst the mechanism(s) responsible are still under
87 investigation, it would appear that changes in the hormonal balance of the common
88 carp, *Cyprinus carpio*, that occur in response to hypoxia are associated with retarded
89 gonadal development, reduced spawning success, sperm motility, fertilisation success,
90 hatching rate and larval survival (12). Subsequent research has revealed effects on
91 sex differentiation and development in the zebrafish, *Danio rerio*, leading to male-
92 dominated populations (13). In view of this evidence, it seems likely that hypoxia
93 may act as a confounding factor in determining the way in which fish respond to
94 chemical challenges mediated via the endocrine system.

95 The environmental literature rarely considers the influence of physicochemical factors
96 on endocrine mediated effects, which is probably due, at least in part, to the difficulty
97 in designing appropriate experiments for detecting these highly complex interactions.
98 However, some insight into influence of hypoxia on the estrogenic response can be
99 garnered from the biomedical field. For example, cancer research, using microarray
100 technology, has revealed that hypoxia and estrogen interact to modulate gene
101 expression in an *in vitro* study, involving human breast cancer cells (14).

102 In the present study, we compare the response of fathead minnows (FHM; *Pimephales*
103 *promelas*) exposed to a mixture of estrogenic chemicals under hypoxic vs. normoxic
104 conditions, using an *in vivo* assay that is based on the induction of egg yolk protein
105 (vitellogenin; VTG) synthesis in male fish (15). There is already evidence to suggest
106 that hypoxia is associated with altered VTG levels in wild estuarine fish, as well as
107 those maintained under laboratory conditions (16). However, here, we will consider
108 the influence of hypoxia on the VTG response of male fish stimulated by exposure to
109 a defined mixture of estrogenic chemicals. The data generated will contribute to our
110 understanding of the risks that exist in multiple stress exposure situations, which is of
111 relevance with regard to the development of risk assessment methodology.

112 **2. MATERIALS AND METHODS**

113 *2.1 Experimental Design*

114 The design of this investigation is based on that of a previous study by Brian et al.
115 (17), in which the response of male FHM to a defined mixture of estrogenic chemicals
116 was characterised, using the induction of plasma VTG as an endpoint, following an
117 exposure period of two weeks. The mixture consisted of the endogenous steroidal

118 estrogen, 17β -estradiol (E2) and the synthetic steroidal estrogen, 17α -ethinylestradiol
119 (EE2), as well as three other environmentally relevant chemicals that have the
120 capacity to mimic the actions of estrogen; namely 4-*tert*-nonylphenol (NP), 4-*tert*-
121 octylphenol (OP) and bisphenol-A (BPA). Stocks of E2 (98% purity), EE2 (98%
122 purity), OP (97% purity) and BPA (99% purity) were purchased from Sigma Aldrich,
123 Dorset, UK. NP (99% purity) was obtained from ACROS Organics, Leicestershire,
124 UK. Each of the chemicals was combined at a fixed ratio, based on their potency with
125 regard to the induction of VTG. The joint action of these chemicals is known to be
126 consistent with predictions based on concentration additivity (CA; *17*).

127 A master stock, containing each component of the mixture at a concentration that was
128 known to elicit a 50% response with regard to the induction of VTG (i.e. its EC₅₀),
129 was prepared in dimethylformamide (DMF; VWR International, Leicestershire, UK).
130 This master stock, which comprised 13.5 $\mu\text{g/l}$ EE2, 375 $\mu\text{g/l}$ E2, 105 mg/l NP, 675
131 mg/l OP and 2.25 g/l BPA, was then diluted in DMF to produce five further stocks
132 that were 0.5, 0.3, 0.2, 0.1 and 0.05 of the original mixture concentration. The stock
133 solutions were diluted by 1:15000 with de-chlorinated tap water (pre-heated to 25 °C)
134 prior to delivery to the experimental tanks. This flow-through exposure system is
135 described in more detail in an earlier publication (*17*).

136 The resulting mixture concentrations in the fish tanks were sufficient to cover the full
137 extent of the concentration response curve in fish maintained under normal oxygen
138 conditions (7 mg/l DO \pm 1 mg/l; *17*). A solvent control tank was run alongside those
139 containing each of the various dilutions of the mixture. This was dosed with a stock
140 of pure DMF, which was delivered at the same rate as the mixture-treated tanks.

141 The chemical dosing commenced one week before the start of each exposure study.
142 This conditioning process ensured that the chemical concentrations in the tanks were
143 accurate. Analytical chemistry was used to verify the water concentrations in samples
144 collected immediately prior to the addition of the fish and after one week of exposure.
145 A third and final set of samples was collected on the day that the exposure study was
146 terminated. The phenolic compounds (NP, OP and BPA) were measured by direct
147 injection onto a reverse phase HPLC column, according to the methods described by
148 Pojana et al. (18). The steroids (E2 and EE2) were analysed by RIA, using the
149 technique outlined by Länge et al. (19).

150 *2.2 Protocol*

151 Two exposure studies were set up, in parallel, according to the design outlined above.
152 One set of tanks was maintained under hypoxic conditions (<2 mgO₂/l). This was
153 achieved by bubbling nitrogen gas through the tanks, which displaced the oxygen in
154 the water. Each tank was supplied with a close fitting glass lid with a hole at either
155 end to allow the delivery of water and pressurised nitrogen to the tanks, via silicone
156 tubing. The nitrogen, which was fed by a series of nitrogen cylinders with individual
157 flow controls to each tank, was then diffused into the water using a 15 cm ceramic air
158 stone (PlanetRena, Charlotte, NC, USA). The second set of tanks was set up in an
159 identical manner, except that they were supplied with pressurised air, as opposed to
160 nitrogen. It was expected that the oxygen conditions in these tanks would be close to
161 100% saturation (7-8 mg/l in our system), thereby representing normoxic conditions.
162 One week prior to exposure, whilst the experimental tanks were being conditioned,
163 male fathead minnows were selected from our laboratory-reared stocks. These fish
164 were split into two groups before being transferred to two sets of holding tanks. In

165 one set of holding tanks, the fish were equilibrated to hypoxic conditions. This was
166 achieved by increasing the flow of nitrogen into the tanks such that the oxygen levels
167 were reduced by approximately 1mg/l each day. In the other set of tanks, the flow
168 rate of air was increased in a similar manner. At the end of the week the fish in each
169 group were randomly allocated to the treated tanks (8 per tank).

170 During the equilibration period and the experiment itself, the fish in each treatment
171 group were fed twice daily: once with frozen brine shrimp and once with flaked fish
172 food. The photoperiod was maintained on a 16 hr light/8 hr dark cycle with 20 minute
173 dawn and dusk transition periods. The DO concentrations in each tank were recorded
174 several times daily using an Oxi 340i digital meter and Cellox® 325 probe (WTW;
175 Weilheim, Germany). Water temperature was also measured daily. Various other
176 water quality parameters (i.e. ammonia, nitrite and nitrate) were analysed at routine
177 intervals to ensure that there were no differences between the two sets of fish tanks,
178 aside from the oxygen availability.

179 *2.3 Sampling and Analysis*

180 At the end of the experiment, the fish were sacrificed by overdose with anaesthetic
181 (MS222; Sigma Aldrich). Their lengths and weights were recorded. Blood samples
182 were then collected from the caudal peduncle using heparinised capillary tubes. The
183 blood samples were centrifuged at 4000 g for 5 minutes and the plasma drawn off and
184 snap frozen on dry ice. The plasma samples were then stored at -20 °C until required
185 for the determination of VTG protein levels. This was carried out using a FHM VTG
186 ELISA kit, which was supplied by Biosense Laboratories AS (Bergen, Norway).

187 *2.4 Statistical Analysis*

188 A series of statistical tests were performed on the physicochemical and biological data
189 sets. Whilst there was a clear difference between the DO levels in each of the parallel
190 exposures, the levels measured within each set of tanks were compared statistically to
191 determine their variability. This was achieved using the ANOVA procedure, followed
192 by Tukey's pairwise comparisons. The chemical concentrations measured in each set
193 of tanks at the start of the experiment were also compared statistically to ensure that
194 there were no differences between the exposure levels. The measurements were first
195 converted into proportions by dividing by the nominal values and comparisons were
196 made between tanks with the same nominal exposure levels using paired t-tests. The
197 VTG levels were also analysed, using t-tests, to compare the mean response of fish in
198 each treatment group across the parallel exposures. Where necessary, these data were
199 log transformed prior to analysis in order to achieve normality. The statistical testing
200 was carried out using Minitab version 13.1 (Minitab Inc. State College, PA, USA).

201 **3. RESULTS**

202 *3.1. Oxygen Conditions*

203 There was a clear difference between the oxygen conditions in the parallel exposure
204 studies (Figure 1). The mean daily DO concentration in the normoxic tanks ranged
205 between 6.58 and 7.18 mg/l. Some of these values were slightly lower than the target
206 of 100% oxygen saturation. This can be attributed to the fact that these tanks suffered
207 from a slight bacterial build up towards the end of the exposure period. The bacterial
208 levels tended to be higher in the tanks that contained more of the mixture, which is
209 reflected by the trend of reducing DO with increasing level of exposure. However,
210 despite of this, there were no significant differences between the DO levels in this set

211 of tanks and the DO concentrations were well within the range of encountered under
212 normoxic conditions.

213 In contrast, the mean daily DO levels in the hypoxic tanks during the exposure were
214 between 1.44 and 1.75 mg/l. These levels were less variable than those recorded in
215 the normoxic tanks, probably due to the lower levels of bacteria in the hypoxia tanks
216 (these factors may have been related). However, there was a statistically significant
217 difference ($p < 0.01$) between the levels measured in the tanks containing the mixture
218 at a 0.05 and 1.0 dilution, which had the lowest and highest mean DO concentrations,
219 respectively. Nevertheless, the oxygen conditions within this set of tanks fell below
220 the hypoxic threshold of 2 mg/l throughout the period of exposure, thereby satisfying
221 the experimental criteria.

222 The differential growth of bacteria in each of the parallel exposures, which became
223 apparent at the beginning of the second week of exposure, raised concerns regarding
224 the chemical concentrations in each set of tanks. This was based on prior experience
225 indicating that bacterial blooms may be associated with increased rates of chemical
226 biodegradation, potentially lead to a reduction in the exposure levels in the affected
227 tanks (20). Hence, as a precaution, the decision was taken to terminate the experiment
228 early and, as a result, the duration of the exposure was reduced from 14 to 10 days.

229 *3.2. Chemical Concentrations*

230 The analysis of the chemical concentrations in each fish tank revealed that, in general,
231 there was good agreement between the nominal and actual exposure concentrations
232 measured at the start of the experiment (Table 1). There was also good agreement
233 between the concentrations measured across each of the parallel exposures. However,

234 BPA was an exception to this rule. Whilst the actual concentrations of this chemical
235 were close to nominal in the hypoxic tanks, these measurements were considerably
236 lower in the tanks maintained under normoxic conditions and, hence, a significant
237 difference ($p < 0.05$) between the levels measured in the parallel studies was detected.
238 However, the agreement between the nominal and actual exposure concentrations in
239 the normoxic tanks improved at the second and third time points (see the supporting
240 information, S1 and S2, respectively), suggesting that the inconsistency between the
241 levels measured at the start of the experiment may have been an analytical anomaly,
242 although the values were consistently lower than those measured in the hypoxic tanks.
243 This may reflect the ease with which BPA is biodegraded in the presence of bacteria,
244 which was more prevalent in the normoxic tanks. In contrast, the other components
245 of the mixture (E2, EE2, NP and OP) appeared to be unaffected by the presence of the
246 bacteria, with their concentrations remaining consistent throughout the experimental
247 period.

248 *3.2. VTG Protein Induction*

249 The analysis of VTG induction levels at the end of the experiment (Figure 2) revealed
250 that there was a clear and consistent concentration-response to the mixture in each of
251 the parallel exposures. The potency was similar to that reported in previous work by
252 the same authors (17, 20) in that a 50% VTG response was induced by the 0.2 mixture
253 dilution, which contained each chemical at a fifth of its individual EC50. There was
254 no evidence that the VTG response of fish differed between the normoxic and hypoxic
255 conditions, as reflected by the fact that there were no significant differences detected
256 between the mean VTG levels within each treatment group.

257 **4. DISCUSSION**

258 The results of this study clearly demonstrate that the VTG response of FHM exposed
259 to a mixture of estrogenic chemicals is similar under hypoxic vs. normoxic conditions.
260 Hence, the data refute the hypothesis that the estrogenic response may be elevated at
261 low oxygen levels, either as a result of increased chemical uptake or due to changes in
262 the rate of physiological processing under varying physicochemical conditions. These
263 findings contrast with evidence from similar studies involving the exposure of fish to
264 micro-pollutants and metals. For example, recent research by Hattink et al. (8)
265 revealed that common carp (*Cyprinus carpio*) are around three times more sensitive to
266 the effects of cadmium under hypoxia (at 25% oxygen saturation) in relation to those
267 maintained under normal oxygen conditions, although it was not possible to identify
268 the mechanism responsible. The lack of response to hypoxia in the present study
269 indicates that EDCs may not behave in the same way as other toxicants under hypoxic
270 conditions and that the rate at which they are taken up and metabolised remains
271 constant, regardless of oxygen availability. However, this theory is not consistent
272 with *in vitro* data, which shows that hypoxia and estrogen treatment act together to
273 affect molecular-level responses, leading to significant effects on gene expression
274 profiles (14).

275 Whilst we would normally expect molecular responses, such as those reported by
276 Seifeddine et al. (14), to be reflected at higher levels of biological organisation, it is
277 possible that effects on VTG induction were not observed *in vivo* due to the influence
278 of negative feedback processes. As a result, we cannot exclude the possibility that
279 hypoxia was associated with effects on rates of chemical uptake and metabolism:
280 these alterations may have acted against one another, thereby countering any overall
281 effect. The potential for physiological interactions of this nature is highlighted by
282 recent evidence that the expression of the estrogen receptor, ER α , is three-fold lower

283 in wild fish inhabiting hypoxic sites, compared to those at normoxic locations (16).
284 Presumably, differences in receptor activity have the capacity to affect the rate and
285 efficiency with which molecular responses are transcribed and subsequently translated
286 at the proteomic level, thereby altering the magnitude of the response to estrogenic
287 stimuli. It is therefore possible that changes in the expression of ER α could,
288 potentially, have masked any effects arising as a result of changes in the rate of
289 chemical uptake, although this hypothesis requires further investigation.

290 In addition, whilst there was no effect of hypoxia on the VTG response measured after
291 10 days of exposure, it is possible that differences may have been detected at earlier
292 time points (e.g. after 24 hours or 7 days). This response pattern has previously been
293 reported for FHM exposed to the same estrogenic mixture at different temperatures
294 (21). The transient nature of this response was attributed to an increase in the rate of
295 induction of the VTG response at higher temperatures, which was mediated via both
296 transcriptional and translational effects. After two weeks, however, these differences
297 were no longer apparent and the VTG response was identical for fish maintained at 20
298 and 30 °C. The same temporal pattern may have been apparent in the present study,
299 with fish maintained under hypoxic conditions exhibiting an elevated response to the
300 mixture at earlier time points due to an increased rate of chemical uptake under these
301 conditions. However, the potential for short-term effects on the estrogenic response
302 were not considered because the relevance of such transient alterations with regard to
303 chemical risk assessment remains unclear (21).

304 To conclude, the results of this study reiterate that mixtures of estrogenic chemicals,
305 or indeed any chemicals that act via a common mechanism, have the capacity to act
306 together to exert combined effects *in vivo*, as previously reported by Brian et al. (17,

307 20), thereby highlighting the need to take account of their joint effects. However,
308 there was no evidence to suggest that the estrogenic response was confounded by the
309 effects of an additional physicochemical variable, which was, in this case, represented
310 by hypoxia, contrary to expectations based on previous studies. The lack of response
311 indicates that the VTG response is robust and relatively insensitive to the effects of
312 additional challenges that arise in multiple stress exposure situations. Hence, it would
313 appear that existing safety factors are sufficient to protect against the effects of inter-
314 actions with confounding factors, such as low oxygen conditions. Nevertheless, this
315 conclusion should be interpreted with caution, as the response may vary, depending
316 both on the nature of the physicochemical challenge and the characteristics of the
317 toxicant in question. It is also possible that the response becomes more plastic at
318 higher levels of biological organisation, which means that confounding factors may
319 have a greater impact on endpoints that relate to survival and reproduction. Hence,
320 despite the negative conclusion of the present study, there may be a need for greater
321 stringency when assessing the risk posed by chemicals in the “real world”, in which
322 multiple stress exposure situations are the norm.

323 **5. ACKNOWLEDGEMENTS**

324 This work was funded by a grant from the Natural Environment Research Council
325 (NE/D00389X/1).

326 **6. SUPPORTING INFORMATION**

327 The nominal and actual exposure concentrations in each set of tanks at time points 1
328 (after 7 days) and 2 (on the final day of exposure) are presented in tabular form in S1
329 and S2.

330 **7. REFERENCES**

- 331 (1) Diaz, R.J.; Rosenberg, R. Marine benthic hypoxia: a review of its ecological
332 effects and the behavioural responses of benthic macrofauna. *Oceanogr. Mar. Biol.*
333 *Annu. Rev.* **1995**, 33, 245-303.
- 334 (2) Wu, R.S.S. Hypoxia: from molecular responses to ecosystem responses. *Mar.*
335 *Poll. Bull.* **2002**, 45, 35-45.
- 336 (3) Scavia, D.; Donnelly, K.A. Reassessing hypoxia forecasts for the Gulf of Mexico.
337 *Environ. Sci. Technol.* **2007**, 41, 8111-8117.
- 338 (4) Justić, D.; Rabalais, N.N.; Turner, R.E. Coupling between climate variability and
339 coastal eutrophication: evidence and outlook for the Gulf of Mexico. *J. Sea Res.* **2005**,
340 54, 25-35.
- 341 (5) Heath, A.G. *Water pollution and fish physiology*; CRC Press: Cleveland, 1987.
- 342 (6) Gracey, A.Y.; Troll, J.V.; Somero, G.N. Hypoxia-induced gene expression
343 profiling in the euryoxic fish *Gillichthys mirabilis*. *PNAS* **2001**, 98, 1993-1998.
- 344 (7) Alabaster, J.S.; Shurben, D.G.; Mallet, M.J. The acute lethal toxicity of mixtures
345 of cyanide and ammonia to smolts of salmon, *Salmo salar* L. at low concentrations
346 of dissolved oxygen. *J. Fish Biol.* **1983** 22, 215-222.
- 347 (8) Hattink, J.; De Boeck, G.; Blust R. The toxicokinetics of cadmium in carp under
348 normoxic and hypoxic conditions. *Aquat. Toxicol.* **2005**, 75, 1-15.
- 349 (9) Lloyd, R. Effects of dissolved oxygen concentration on the toxicity of several
350 poisons to Rainbow trout (*Salmo gairdneri* Richardson) *J. Exp. Biol.* **1961**, 38, 447-
351 455.

- 352 (10) Magaud, H.; Migeon, B.; Morfin, P.; Garric, J.; Vindimian, E. Modelling fish
353 mortality due to urban storm run off: interacting effects of hypoxia and un-ionised
354 ammonia. *Water Research* **1997**, 31, 211-218.
- 355 (11) Sorensen, E.M. *Metal poisoning in fish*; CRC Press Inc.: Boca Raton, 1991.
- 356 (12) Wu, R.S.S.; Zhou, B.S.; Randall, D.J.; Woo, N.Y.S.; Lam, P.K.S. Aquatic
357 hypoxia is an endocrine disruptor and impairs fish reproduction. *Environ. Sci.*
358 *Technol.* **2003**, 37, 1137-1141.
- 359 (13) Shang, E.H.H.; Yu, R.M.K.; Wu, R.S.S. Hypoxia affects sex differentiation and
360 development, leading to a male-dominated population in zebrafish (*Danio rerio*).
361 *Environ. Sci. Technol.* **2006**, 40, 3118-3122.
- 362 (14) Seifeddine, R.; Dreiem, A.; Tomkiewicz, C.; Fulchignoni-Lataud, M.; Brito, I.;
363 Danan, J.; Favaudon, V.; Barounki, R.; Massaad-Massade, L. Hypoxia and estrogen
364 co-operate to regulate gene expression in T-47D human breast cancer cells. *J. Steroid*
365 *Biochem. Mol. Biol.* **2007**, 104, 169-179.
- 366 (15) Sumpter, J.P.; Jobling, S. Vitellogenesis as a biomarker of oestrogenic
367 contamination of the aquatic environment. *Environ. Health. Persp.* **1995**, 103,
368 Supplement 7, 173-17.
- 369 (16) Thomas, P.; Saydur Rahman, M.; Khan, I.A.; Kummer, J.A. Widespread
370 endocrine disruption and reproductive impairment in an estuarine fish population
371 exposed to seasonal hypoxia. *Proc. R Soc. B* **2007**, 247, 2693-2701.
- 372 (17) Brian, J.V.; Harris, C.A.; Scholze, M.; Backhaus, T.; Booy, P.; Lamoree, M.;
373 Pojana, G.; Jonkers, N.; Runnalls, T.; Bonfa, A.; Marcomini, A.; Sumpter, JP.
374 Accurate prediction of the response of freshwater fish to a mixture of estrogenic
375 chemicals. *Environ. Health Perspect.* **2005**, 113, 721-728.

- 376 (18) Pojana, G.; Bonfà, A.; Buseti, F.; Collarin, A.; Marcomini, A. Determination of
377 natural and synthetic estrogenic compounds in coastal lagoon waters by HPLC-
378 electrospray-mass spectrometry. *Intern. J. Environ. Anal. Chem.* **2004**, 84, 717-727.
- 379 (19) Lange, R.; Hutchinson, T. H.; Croudace, C. P.; Siegmund, F.; Schweinfurth, H.;
380 Hampe, P.; Panter, G. H.; Sumpter, J. P. Effects of the synthetic estrogen 17 alpha-
381 ethinylestradiol on the life-cycle of the fathead minnow (*Pimephales promelas*).
382 *Environ. Toxicol. Chem.* **2001**, 20, 1216–1227.
- 383 (20) Brian, J.V.; Harris, C.A.; Scholze, M.; Kortenkamp, A.; Booy, P.; Lamoree, M.;
384 Pojana, G.; Jonkers, N.; Bonfa, A.; Marcomini, A.; Sumpter, J.P. Evidence of
385 estrogenic mixture effects on the reproductive performance of fish. *Environ. Sci.*
386 *Technol.* **2007**, 41, 337-344.
- 387 (21) Brian, J.V.; Harris, C.A.; Runnalls, T.J.; Fantinati, A.; Pojana, G.; Marcomini,
388 A.; Booy, P.; Lamoree, M.; Kortenkamp, A.; Sumpter, J.P. Evidence of temperature-
389 dependent effects on the estrogenic response of fish: Implications with regard to
390 climate change. *Sci. Total Environ.* **2008**, 397, 72-81.

391 Table 1: Nominal and actual chemical concentrations in each tank at the beginning of the parallel exposure studies. The water samples were
 392 collected immediately prior to the addition of the fish.

393


394		Nominal					Hypoxic					Normoxic				
395	Treatment	EE2	E2	NP	OP	BPA	EE2	E2	NP	OP	BPA	EE2	E2	NP	OP	BPA
396		ng/l	ng/l	μg/l	μg/l	μg/l	ng/l	ng/l	μg/l	μg/l	μg/l	ng/l	ng/l	μg/l	μg/l	μg/l
397	Tank 1: Control	0	0	0	0	0	0.0	0.3	0.3	0.0	0.0	0.0	0.1	0.3	0.0	0.2
398	Tank 2: 0.05 dilution	0.05	1.25	0.35	2.25	7.5	0.07	1.19	0.46	5.22	0.5	0.06	0.93	0.50	1.34	1.0
399	Tank 3: 0.1 dilution	0.09	2.5	0.7	4.5	15	0.14	2.39	0.65	2.38	16.6	0.13	1.82	0.69	2.55	9.8
400	Tank 4: 0.2 dilution	0.18	5	1.4	9	30	0.24	3.94	1.12	14.2	36.1	0.24	4.17	1.51	8.80	0.0
401	Tank 5: 0.3 dilution	0.27	7.5	2.7	13.5	45	0.39	6.36	1.59	13.6	52.3	0.38	6.86	1.86	15.0	0.3
402	Tank 6: 0.5 dilution	0.45	12.5	3.7	22.5	75	0.63	9.34	3.36	32.9	87.7	0.69	10.3	3.38	27.8	0.0
403	Tank 7: 1:0 dilution	0.9	25	7	45	150	1.68	19.2	5.15	57.6	169	1.26	21.5	6.12	59.4	12

404 **LEGENDS**

405 Figure 1: Mean of the daily dissolved oxygen (DO) concentrations in each tank
406 throughout each of the parallel exposure studies. Daily DO concentrations were taken
407 to be the average value recorded based on 6-10 measurements that were made on each
408 day. Error bars represent one standard error of the mean. The letters (a and b) denote
409 that there was a significant difference between the mean DO level in tanks containing
410 the 0.05 and 1.0 mixture dilution under hypoxia. No differences were detected within
411 the remaining tanks maintained under each set of oxygen conditions.

412 Figure 2: Mean of the plasma VTG concentrations in each tank following each of the
413 parallel exposure studies. Error bars represent one standard error of the mean, which
414 was calculated on the basis of measurements made from eight fish in each tank.


415 Figure 1:



416

417

418 Figure 2:



419

420

421 Brief:

422 The response of fish to a mixture of estrogenic chemicals is not affected by

423 concomitant exposure to hypoxia.