



A survey of psychological practitioner workplace wellbeing.

Journal:	<i>Clinical Psychology & Psychotherapy</i>
Manuscript ID	CPP-2316.R2
Wiley - Manuscript type:	Assessment
Date Submitted by the Author:	n/a
Complete List of Authors:	Summers, Elisabeth; Cardiff University, Clinical Psychology; Aneurin Bevan University Health Board Morris, Reg; Cardiff University, clinical psychology Bhutani, Gita; Lancashire Care NHS Foundation Trust, Psychological Services; University of Liverpool, Roa, Amra ; Psychological Horizons Clarke, Jeremy; New Savoy Partnership
Keywords:	survey, psychological practitioner, psychological therapist, wellbeing, workplace wellbeing, psychologists

SCHOLARONE™
Manuscripts

Abstract

The wellbeing of the psychological workforce is an area of concern. However, it has been sparsely studied in an holistic manner encompassing workplace wellbeing as well as burnout. This study reports a survey of 1,678 psychological practitioners accessed through professional networks.

The short Warwick Edinburgh Mental Wellbeing Scale (SWEMWBS) and the Psychological Practitioner Workplace Wellbeing Measure (PPWWM) were administered with a demographic questionnaire. The mean for the SWEMWBS was below that of a national population survey. The inter-correlation of these tests was .61. Subgroup analyses showed significant differences: assistant psychologists, counsellors and psychological wellbeing practitioners demonstrated better than average workplace wellbeing. But for general wellbeing (SWEMWBS) trainee clinical psychologists and assistant psychologists showed lower than average wellbeing while psychological wellbeing practitioners were higher than average.

Other factors associated with wellbeing were: contract type –both measures (higher workplace wellbeing in those with temporary contracts and the self-employed); employment sector –for PPWWM only (private organisation/independent workers and third sector/charitable organisation workers scored above the PPWWM mean); ethnicity – for both measures (Asian groups except Chinese had higher wellbeing than average for the PPWWM and SWEMWBS); disability was strongly associated with lower wellbeing on both measures. Harassment, feeling depressed or a failure and wanting to leave the NHS were associated with lower wellbeing. Greater age, pay and years of service were *negatively* correlated with wellbeing.

A five-factor structure was obtained with this sample.

The results confirmed psychological practitioners as an at-risk group and identified a number of factors associated with workplace wellbeing.

Practitioner Points

- Workplace wellbeing is related to, but distinct from, general wellbeing and should be measure by a specialist instrument.
- The general wellbeing of psychological practitioners is below the national average and this requires consideration by policymakers and workforce planners.
- The association of gender and sexual orientation with wellbeing do not appear to be strong, but disability has a strong generalised association requiring further research and action.

- The associations between ethnicity and temporary contracts and wellbeing are complex and require further study.

Introduction

The mental health of psychological practitioners has become a focus of attention in recent years (Dattilio, 2015). Workplace wellbeing is a vital consideration if health services are to retain staff and deliver an effective and safe service (Health Education England, 2019; Hall et al., 2016).

Research in healthcare professional populations has focused particularly on the assessment of burnout. Burnout can be thought of as a specific form of chronic workplace psychological stress (Ruotsalainen, Verbeek, Mariné & Serra, 2015; World Health Organization, 2018). Although burnout is correlated with anxiety, depression and fatigue, burnout is thought to be distinct from mental health, from general stress and from other work phenomena such as job dissatisfaction (Maslach, Schaufeli & Leiter, 2001). In healthcare professional populations, burnout has been defined by Maslach and colleagues and is often measured by the Maslach Burnout Inventory (Maslach, Jackson, & Leiter, 1996). It is characterised by three simultaneously existing dimensions: (1) *emotional exhaustion*, the depletion of emotional reserves, energy loss and feeling less able to experience emotion related to work; (2) *depersonalisation*, the distancing from patients and negative or cynical attitudes and feelings towards patients; (3) *lack of personal accomplishment*, negative self-evaluation, particularly in relation to patient work, feelings of dissatisfaction with work accomplishments and achievements (Maslach et al., 1996; Maslach & Jackson, 1981). The Copenhagen Burnout Inventory (CBI), a further measure of burnout, has also been used in healthcare populations (Di Benedetto & Swadling, 2014; D'Souza, Egan & Rees, 2011). It was designed to have utility across different professional domains beyond healthcare (Kristensen, Borritz, Villadsen & Christensen, 2005). The authors propose fatigue and exhaustion are at the core of burnout, with an additional key feature being “the *attribution* of fatigue and exhaustion to specific *domains* or spheres in the person’s life” (Kristensen et al., 2005, p. 197).

Table 1 shows the results of studies of burnout in psychological practitioners. The majority found evidence for higher than average burnout levels in psychological practitioners. This is concerning for the profession, but not all these studies used the established thresholds for burnout to identify ‘average’ levels (Maslach et al., 2017). Concerning results for the UK were reported by Hacker-Hughes, Rao, Dosanjh, Cohen-Tovée, Clarke & Bhutani (2016). These studies demonstrated increasingly poor wellbeing amongst psychological practitioners across successive annual surveys until 48% felt depressed, almost 50% felt like

1 a failure and 92% found their job stressful at least some of the time. Of additional concern
2 is the lack of priority given to self-care in psychologists and mental health professionals. It
3 has been proposed that despite the potential increased personal challenges associated with
4 being a mental health professional, they are prone to avoiding treatment or support
5 (Dattilio, 2015).
6
7
8

9
10 *Table 1 about here*
11

12 There is a little research comparing the rates of burnout across professions to allow
13 comparison of the wellbeing of psychological practitioners with the wider mental health
14 workforce. Most studies appear to either focus on levels of burnout in single professional
15 groups, or levels of burnout are assessed within a single service type. However, Johnson et
16 al. (2012) conducted a large sample of mental health professionals in England (n= 2,258)
17 including social workers, occupational therapists, nurses, clinical psychologists, nursing
18 assistants, psychiatrists, and service managers. They reported significant differences across
19 the professional groups with the mean for social workers, nurses and occupational
20 therapists reaching the threshold for high burnout while the other professions were below
21 the threshold. In contrast, in Israel, no differences in stress or burnout were observed
22 between 249 female psychologists, nurses and social workers, except for the
23 depersonalization outcome of burnout, which was significantly lower among psychologists
24 than among nurses or social workers (Ben-Zur & Michael, 2007). However, this sample size
25 was small for detecting statistically significant subgroup differences.
26
27
28
29
30
31
32
33

34 Two issues with workplace wellbeing research to date are the lack of a validated measure
35 specific to psychological practitioners and the focus on burnout and the negative facets of
36 work experience rather than wellbeing. Maben et al., (2012) made a convincing case for
37 considering both positive and negative aspects of workplace experience to reach an overall
38 appraisal of employee wellbeing. This approach is echoed by models of workplace
39 wellbeing that identify resources as positive aspects and demands or threats to resources
40 as negative aspects (Bakker & Demerouti, 2007; Hobfoll & Shiron, 2001). Summers et al.
41 (2020) addressed the dual problems of the lack of a validated measure of workplace
42 wellbeing for psychological practitioners and the need to consider both the positive and
43 negative facets of wellbeing. This research developed the Psychological Practitioner
44 Workplace Wellbeing Measure (PPWWM), a brief 26-item measure based on items derived
45 from a qualitative study of practitioners.
46
47
48
49
50
51
52

53 The present study used existing data collected for the annual national survey delivered by
54 The New Savoy Partnership and Leadership and Management Faculty of the British
55 Psychological Society, Division of Clinical Psychology. This survey was predicated by
56 concerns for the wellbeing of the psychological workforce and was designed to inform
57 policy and guidelines. It was sent to psychological practitioners from a range of
58 psychological sub-professions with a focus on those delivering talking-based therapies. The
59
60

1 survey used the PPWWM, a generic wellbeing measure (The Short Form Warwick-
2 Edinburgh Mental Wellbeing Scale – SWEMWBS—Stewart-Brown et al., 2009) and some
3 specific questions from the National Health Service (UK) staff survey (NHS staff survey
4 coordination centre, 2018).
5
6
7

8 Specific research questions included: the determination of averages for workplace
9 wellbeing of psychological practitioner sub-groups in the UK; the exploration of subgroup
10 differences (including profession, gender and disability); association of wellbeing with
11 demographic variables; the correlation of the PPWWM with the SWEMWBS and its
12 association with specific questions from the NHS staff survey relating to harassment and
13 bullying. Since gender differences in the workforce are currently an area of interest and
14 concern (Office for National Statistics, 2019) key results were reported separately for males
15 and females.
16
17
18
19
20

21 Additionally, the study was designed to augment existing psychometric data for the
22 distribution and norms of the PPWWM and its factor structure.
23
24
25

26 27 **Method**

28 29 30 **Consent and Data Management**

31
32 The study was based on a secondary analysis of an anonymised data base produced by a
33 Division of Clinical Psychology/New Savoy Partnership survey of the UK psychology
34 profession. In common with earlier annual surveys conducted by the Division of Clinical
35 Psychology/New Savoy Partnership, the purpose of the original survey was to assess the
36 state of the profession and was not construed as a research study requiring research ethical
37 approval. The survey was conducted under the terms of the European General Data
38 Protection Regulations (2016/679). It was anonymous from the point of data collection and
39 no identifiable information was collected (e.g . location, IP address, etc.). Participation was
40 voluntary and consents for use of the data were given within the survey. Participants
41 agreed for their data to be used *“in future reports or publications.”*
42
43
44
45
46
47

48 **Procedure**

49
50 The first author was involved in the collection of data. A web-based Qualtrics® survey was
51 created and participants were sent a web-based link for completion. The questionnaire is
52 available as supplementary material to this paper. The survey was disseminated via
53 psychological practitioner mailing lists, professional organisations and networks and a
54 closed social media group for clinical psychologists. Members of the professional groups
55 were emailed a web-based link to the survey. Individuals were encouraged to share the
56 invitation for participation with their colleagues and professional groups (a ‘snowball’
57 sample). Response rate cannot be determined in such samples. The inclusion criterion was
58
59
60

1 those currently employed in a UK health/social care organisation as a psychological
2 practitioner.
3

4 **Measures**

5
6
7
8 The Qualtrics® survey included a demographic survey, the PPWWM (Summers, et al., 2020),
9 The Short Warwick-Edinburgh Mental Well-being Scale (Stewart-Brown et al., 2009) and a
10 number of items taken from the NHS staff survey relating to harassment and bullying (NHS
11 staff survey coordination centre, 2018). The questionnaire required a response to all
12 questions therefore questionnaires did not have missing items and incomplete
13 questionnaires were screened out.
14
15
16

17 **Demographic Survey**

18
19
20 The demographic questionnaire collected the following information: profession, type of
21 contract, years since professional qualification, years worked since qualification, years in
22 current post, contracted hours, additional paid and unpaid hours, type of organisation, pay
23 scale, gender, sexual orientation, ethnicity, illness/disability and age. The illness/disability
24 question asked 'Do you have a long-standing illness or disability?'
25
26
27

28 **The Psychological Practitioner Workplace Wellbeing Measure (PPWWM)**

29
30
31 The PPWWM is a 26-item Likert scale (1 *strongly disagree* to 5 *strongly agree*) with a score
32 range of 26 to 130. Higher scores indicate greater wellbeing. It was validated with a sample
33 of 400 psychological practitioners (Summers et al., 2020) and showed good construct
34 validity against: a generic workplace wellbeing measure, The Health and Safety Executive
35 Management Standards Indicator Tool (Health and Safety Executive, 2004) $r=.88$; The
36 Satisfaction with Life Scale (Diener et al., 1985) $r=.50$; The General Health Questionnaire
37 (12) (Goldberg, 1992) $r=-.31$. Test-retest reliability and internal consistency were $r=.94$ and
38 $\alpha=.92$, respectively. The PPWWM produced a six-factor structure: 'professional and
39 organisational satisfaction', 'support and flexibility', 'professional role', 'physical
40 environment', 'clinical supervision' and 'external personal' support.
41
42
43
44
45

46 **Short Warwick-Edinburgh Mental Well-being Scale**

47
48
49 The Short Warwick-Edinburgh Mental Well-being Scale (SWEMWEBS) (Stewart-Brown et al.,
50 2009) is a unidimensional self-report, 7-item scale using a 5-point rating system (1 *none of*
51 *the time* to 5 *all of the time*) to assess mental-wellbeing. It was derived from the original 14-
52 item Warwick-Edinburgh Mental Well-being Scale (WEMWEBS) (Tennant et al., 2007). Its
53 score range is 7 to 35 with higher scores indicating greater wellbeing. The WEMWEBS was
54 developed using Rasch analysis (Stewart-Brown et al., 2009) to measure wellbeing in the
55 general population and for the evaluation of projects and policies that aim to improve
56 mental wellbeing. It has been widely used within the UK in population surveys (for
57 example, Bellis, Lowey, Leckenby, Hughes & Harrison, 2013) and for the assessment of
58
59
60

1 interventions to improve mental wellbeing (for example, Shah, Cader, Andrews, Wijesekera,
2 & Stewart-Brown, 2018). Cronbach's α for the SWEMWBS was .84 (Fat, Scholes, Boniface,
3 Mindell & Stewart-Brown, 2017), indicating high internal consistency (DeVellis, 2017). The
4 test-retest reliability has not been evaluated. The SWEMWBS is negatively correlated with
5 the GHQ-12 and positively correlated with happiness and positive health state (Ng Fat et al.,
6 2017).
7
8
9

10 ***Additional items***

11
12
13
14 Nine additional quantitative items from the National Health Service (UK) staff survey
15 (NHS staff survey coordination centre, 2018) were included. These tapped harassment,
16 bullying and discrimination at work, organisational change, staffing levels and overall mood.
17 Participants were also given the opportunity to provide additional comments within a 'free-
18 text' format relating to their wellbeing and the survey questions.
19
20

21 **Data Cleaning and Analysis**

22
23
24
25 As noted above, the forced choice nature of the questionnaire did not generate missing
26 questions. Data were cleaned by removing any incomplete questionnaires and responses
27 from participants who did not meet the inclusion criteria.
28
29

30 Data analysis used IBM SPSS® version 25. All correlations and ANOVA analyses used
31 bootstrapping (2000 iterations) to give a more robust test with respect to deviation from
32 assumptions. Bonferroni corrections for analyses involving demographic variables were not
33 made because they are highly conservative and can fail to identify significant relationships
34 in survey research (Bland & Altman, 1996). Commentaries suggest there is a lack of
35 consensus as to how Bonferroni should be applied (Drezner & Drezner, 2016) and that their
36 routine use should be avoided (Armstrong, 2014). To counter Type 1 errors comparisons
37 were made with Summers et al. (2020) to establish replicability.
38
39
40
41
42

43 The comparisons of the SWEMWBS with the survey of Ng Fat et al (2017) used a one-
44 sample t-test. The significance of PPWWM distribution skew and kurtosis was evaluated
45 using their standard errors to obtain a z-statistic. One-way ANOVA was used to compare
46 the PPWWM and SWEMWBS scores across sub-groups and follow-up tests were two-tailed
47 one-sample t-tests comparing subgroup means with the overall mean. Pearson correlation
48 with bootstrapping was used to measure associations between interval variables (e.g.
49 PPWWM and SWEMWBS).
50
51
52
53

54 Key demographic variables such as age and salary were reported separately for males and
55 females.
56
57

58 The Factor analysis used a Varimax rotation of the principal components matrix with
59 rotation of factors with Eigen values >1.
60

Results

Key demographic variables are presented separately for males and females in Table 2 together with statistical comparisons and further demographic data appears in Table 3.

Table 2 about here

For the total sample of 1,678 the means for the PPWWM and the SWEMWBS were 91.88 ($SD = 17.55$) and 22.21 ($SD = 3.92$) respectively and the inter-correlation was $r = .61$; $p < .0001$. The mean wellbeing score for the SWEMWBS in this sample were 22.09 for men and 22.24 for women compared with 23.70 and 23.20 in a national UK survey (Ng Fat et al., 2017). These scores were significantly below those of the national survey; $t = -7.22$; $df = 319$; $p < .001$ and $t = -9.05$; $df = 1333$; $p < .001$, respectively.

The PPWWM distribution was approximately normal with non-significant skew (-0.05) but there was negative kurtosis (-0.45 , $p < 0.01$). (Figure 1.) Significance of deviation from normality is not uncommon in very large samples that can detect small effects, and a Q-Q plot showed good conformity to the normal distribution apart from lower than expected values at the extreme lower end of the scale.

Figure 1 about here

Means of the PPWWM and SWEMWBS for subgroups with 15 or more respondents are given in Table 3. Post-hoc subgroup follow-up tests comparing groups to the overall mean were completed when the overall ANOVA was significant. All post-hoc follow-up t-tests were two-tailed.

Table 3 about here

For professional groups a one-way ANOVA showed that there were significant differences across all groups for the PPWWM ($F_{(10,1667)} = 2.89$; $p < .001$) and the SWEMWBS ($F_{(10,1667)} = 2.90$; $p < .001$). For the PPWWM assistant psychologists, psychological wellbeing practitioner wellbeing and counsellors scored significantly higher than the overall mean ($t = -2.16$, $df = 23$; $p = .042$, $t = 2.47$, $df = 225$; $p = .018$ and $t = 2.22$, $df = 94$; $p = .029$). But for the SWEMWBS trainee clinical psychologists and assistant psychologists showed lower than average general wellbeing ($t = -3.10$, $df = 27$; $p = .007$ and $t = -2.12$, $df = 23$; $p = .039$). While psychological wellbeing practitioners had higher than average general wellbeing ($t = 2.91$, $df = 225$; $p = .006$).

There was also a significant effect of type of contract (PPWWM, $F_{(3,1674)} = 7.27$; $p < .0001$: SWEMWBS, $F_{(3,1674)} = 6.99$; $p < .0001$). Self-employed practitioners and those on non-

1 permanent contracts demonstrated above average PPWWM scores ($t = 2.91, df = 18; p$
2 $= .009$, two-tailed; $t = 3.70, df = 188; p < .001$, two-tailed) while those on permanent
3 contracts were below the mean, but not significantly so.
4

5
6 Only the self-employed/private practice/independent group differed significantly from the
7 mean on the SWEMBS ($t = 3.85, df = 18; p = .003$).
8
9

10 For sector of employment the differences were significant for the PPWWM ($F_{(5,1672)} = 4.79; p$
11 $< .0001$) but not for the SWEMWBS ($F_{(5,1672)} = 2.05; p = .069$). Follow-up tests for the
12 PPWWM showed that only private organisation/independent workers ($t = 2.39, df = 55; p =$
13 $.025$) and third sector /charitable organisation workers ($t = 4.43, df = 71; p < .001$) scored
14 above the overall mean.
15
16
17
18

19
20 Gender differences for the PPWWM and SWEMWBS were not statistically significant.
21 Similarly, there were no statistical differences for sexual orientation.
22
23

24
25 The overall differences between ethnic groups were statistically significant (PPWWM,
26 $F_{(4,1617)} = 2.89; p < .021$; SWEMWBS, $F_{(4,1617)} = 3.82; p < .004$). For the PPWWM and SWEMBS
27 Asian groups were above the mean ($t = 2.18, df = 65; p = .042$ and $t = 2.0, df = 65; p = .048$).
28 No other group differences were significant.
29
30

31
32 Those reporting disabilities (17.1% of the sample) had significantly lower wellbeing on the
33 PPWWM and the SWEMWBS (PPWWM, $F_{(1,1630)} = 16.13; p < .0001$; SWEMWBS, $F_{(1,1630)} =$
34 $19.55; p < .0001$).
35
36
37
38

39
40 Correlations for key variables with the PPWWM are shown in Table 4. Noteworthy
41 associations are those with the questions about experiences of harassment, especially by
42 managers (19.0% reported experiencing harassment or bullying), feeling depressed (43.3%
43 reported feeling depressed 'some of the time' or 'often') or a failure (42.3% reported feeling
44 a failure 'some of the time' or 'often') and wanting to leave the NHS (74.7% reported
45 wanting to leave 'at least once or twice a year'). Although not large, there were significant
46 *negative* correlations with age and pay scale and with years served in the current post.
47
48
49
50
51
52

53 *Table 4 about here*
54

55 Table 5 depicts the Varimax structure matrix for the factor analysis of the PPWWM. Factors
56 with Eigen vales >1 gave a five factor solution.
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 5 about here

Discussion

The current sample PPWWM mean and standard deviation of 91.88 (SD=17.55) and Cronbach's Alpha of .92 correspond with the results of the validation study (Summers et al., 2020) which were 93.47(17.67) and .92. The means did not differ significantly between the two studies. The PPWWM distribution had significant negative kurtosis, as also found by Summers et al. (2020) but in this case there was no significant skew. The large sample size made it possible to detect statistical significance of minor deviations from normality. But the Q-Q plot showed good conformity to the normal distribution suggesting that the deviation is trivial. The mean scores for the SWEMWBS were below the UK norms for men and women (Ng Fat et al., 2017) suggesting that this professional group experience reduced wellbeing compared to the general population. This corroborates the previous literature on burnout and distress (Table 1).

The small, but significant, *negative* correlations with age, years since qualification and pay scale (Table 4) were not expected. In this profession at least, it appears that higher pay is not associated with improved workplace wellbeing. This small negative association with age and age-related variables replicates the results of Summers et al. (2020). While it is important not to over-interpret these small negative correlations, these findings do contrast with results from the USA and Australia demonstrating significantly *higher* scores on indices of burnout in younger, less experienced psychologists (Dorociak, Patricia, Rupert and Zahniser, 2017; Ackerley, Burnell, Holder & Kurdek, 1988; Rupert & Kent, 2007; Rupert & Morgan, 2005; Rupert, Stevanovic & Hunley, 2009; D'Souza, Egan & Rees. 2011; Di Benedetto & Swadling, 2014). There may be two reasons for this. First, the results may be culture-specific. All these studies were cross-sectional. The specific experiences of older cohorts in the UK may have detruded their workplace well-being, possibly as a result of the re-structuring of public health service pay and conditions between 2004 and 2007 and the erosion of pensions from 2008 onwards. Second, the PPWWM measured positive aspects of workplace well-being as well as the negative aspects (burnout) measured in the USA and Australian studies. Possibly the positive and negative aspects of wellbeing have different longitudinal trajectories.

The PPWWM and the SWEMWBS scores differed significantly between professional subgroups. This effect for the PPWWM was not observed by Summers et al. (2020), but their sample was much less heterogeneous with over 73% being clinical psychologists compared with 49% in this sample. Psychological wellbeing practitioners were above the mean on both measure and counsellors were above the mean on the PPWWM. Meanwhile trainee psychologists and assistant psychologists reported the highest group average scores on the PPWWM and the lowest average scores on the SWEMWBS. For assistant psychologists differences from the overall mean were significant on both measures; for

1 trainee psychologists the differences were significant only for the SWEMWBS. These
2 groups are both on the usual pathway to qualifying as clinical psychologists. Perhaps
3 hopeful career anticipation elevated the workplace wellbeing score while other factors,
4 such as the need to relocate and make personal sacrifices for career, detruded general
5 wellbeing. This decoupling between general wellbeing and workplace wellbeing highlights
6 the importance of using specific measures to evaluate workplace wellbeing.
7
8
9
10

11 Staff on non-permanent contracts were above the mean for the PPWWM and self-employed
12 and private practitioners were above the mean on both the PPWWM and the SWEMWBS. It
13 was unexpected that those on fixed term and temporary contracts and the self-
14 employed/private group would demonstrate higher workplace wellbeing scores than those
15 on permanent organisational contracts. A review of studies of workers on temporary
16 contracts (De Kuyper & De Witte, 2009) also concluded temporary contracts were not
17 associated with poor outcomes and there was some evidence for better wellbeing in
18 temporary workers. This result also replicates Summers et al. (2020) and seems to be a
19 robust finding. It may be because the temporary contract group contains more early career
20 staff who exhibit high workplace wellbeing.
21
22
23
24
25
26

27 The private sector/independent sector group and the third sector/charitable group both
28 had averages above the overall mean for the PPWWM. This was not found by Summers et
29 al. (2020) where nearly 90% of the sample were wholly employed by the NHS. It is
30 concerning that NHS employees were not in this elevated category and this concern is
31 reflected by reports of poor staff wellbeing in large healthcare organisations such as the UK
32 NHS (Health Education England, 2019).
33
34
35
36

37 Wellbeing did not differ across genders nor across those with different sexual orientations.
38 This absence of gender difference is consistent with the finding of Summers et al. (2020).
39
40

41 Recently there has been considerable interest in gender difference in pay and career
42 progression (Office for National Statistics, 2019). This sample did show a small gender
43 difference in favour of males in Agenda for Change banding of NHS workers (Table 2).
44 However, male workers in this sample were on average older and had worked more years
45 since qualification than the females and both these factors were highly associated with
46 salary. This is not to say that female workers do not experience disadvantage in the
47 workforce since 'the motherhood penalty', for example, has the potential to create
48 inequalities in both years worked and age when leaving the workforce (Chung, Graham,
49 Downs, Sandler & Sienkiewicz 2017).
50
51
52
53
54

55 Ethnicity showed an overall statistical difference for PPWWM scores with those identifying
56 as Asian (other than Chinese) having a significantly higher than average score on both
57 measures and those identifying as mixed race having the lowest scores on both measures
58 (but not significantly so). Lower wellbeing in black and minority ethnic people is well
59 documented (Stevenson & Rao, 2014), but the current study revealed a mixed picture for
60

1 psychological practitioners with white practitioners demonstrating somewhat lower
2 wellbeing than all groups other than those of mixed race. However, sample sizes for the
3 non-white groups are small, so differences should be treated with caution.
4
5

6 The percentage of psychological practitioners self-identifying as disabled (over 17%) was
7 unexpected; it was only 12% in Summers et al.'s (2020) mainly clinical psychologist sample.
8 However, this figure is close to that of the 2018 NHS staff survey figure of 18% (NHS staff
9 survey coordination centre, 2018). There was a very clear and highly significant difference
10 between those reporting disabilities and those who did not in terms of workplace wellbeing
11 and general wellbeing in favour of those without reported disability. This reflects the
12 negative relationship between job satisfaction and long-term health problems/disabilities
13 (Gazioglu & Tansel, 2006; Clark, 1996) and indicates the need for improved understanding
14 of workplace support for workers with disabilities (Nevala, Pehkonen, Koskela, Ruusuvaori,
15 & Anttila, 2015).
16
17
18
19
20
21
22
23
24

25 Self-reported '*feeling depressed*' (43.3%) and '*a failure*' (42.3%) was comparable with the
26 survey of Hacker-Hughes et al., (2016) who reported 48%. The correlations of these indices
27 with the PPWWM are indicative of the negative aspects of workplace wellbeing. This
28 relationship with depression was expected and reflects the findings of copious previous
29 research (Glass & McKnight, 1996). However, the functional relationship between
30 depression and burnout remains unclear, especially the question of whether they are
31 aspects of the same phenomenon or different stages in a developmental sequence (Bianchi,
32 Shonfeld & Laurent, 2015).
33
34
35
36
37

38 Self-reported harassment and bullying also correlated with the PPWWM. This finding
39 corroborates previous cross-sectional and longitudinal work on the relationship between
40 bullying and harassment and psychological and physical symptoms (Beswick, Gore,
41 Palferman, 2006).
42
43
44

45 Finally, the negative association between the PPWWM and wanting to leave the NHS is
46 indicative of the potential for factors associated with poor workforce wellbeing to engender
47 high staff turnover rates (Boorman, 2009; Griffeth, Hom & Gaertner, 2000). This highlights
48 the importance of interventions to promote wellbeing as a means of reducing unplanned
49 absences and the costs associated with high turnover (Michie & Williams, 2003).
50
51
52
53
54

55 The factor structure of the PPWWM in this study differed somewhat from that of the
56 validation study. There may be several reasons for this. The sample was larger and more
57 diverse in terms of professional groups. Also, the PPWWM is designed to be a brief
58 questionnaire which necessarily reduces the number of items per factor and this in turn
59 reduces factorial stability (Costello & Osborne, 2005; Field, 2018). However, the differences
60

1 that did emerge are explicable. For example, the Supervision Factor, a separate two-item
2 factor in the validation study, was subsumed under the Support and Flexibility Factor in the
3 present study which affirms the supportive nature of supervision. Two items (21 & 22,
4 about organisational support for personal development) from the Professional and
5 Organisational Satisfaction Factor of the validation study loaded equally on the satisfaction
6 and support factors in this study. Since they referred to support, they were included under
7 this factor, but clearly also have an association with organisational satisfaction. The
8 Professional Role Factor in the present study encompassed three items (Item 2 & 6,
9 referencing support from colleagues and Item 4, referencing sense of belonging) from the
10 Support and Flexibility Factor in the validation study. Again, colleague support and
11 belonging could equally be perceived as an aspect of professional role. Finally, Item 1,
12 which referred to having someone to turn to, was originally under the External Personal
13 Factor, but in the current study was under the Support and Flexibility Factor.

22 **Limitations and Further Research**

24 Although large, the sample may not have been representative. It was a volunteer sample
25 and included less than 10% of the total number of qualified clinical psychologists and
26 probably about the same percentage of the total psychological practitioner workforce of
27 the UK. Moreover, numbers in some of the subgroups were relatively small which limited
28 the dependability of inter-group comparisons.

32 The PPWWM measures positive and negative aspects of workplace wellbeing. The positive
33 aspects explain the correlation with the SWEMWBS measure of general wellbeing. But it
34 would have been useful to also have a general measure of distress to confirm the
35 relationship with the negative aspects of workplace wellbeing. The temporal stability (test-
36 re-test reliability) of the SWEMWS has yet to be established. Until this is established, the
37 effect on correlations of error due to temporal factors cannot be quantified.

42 Surveys of this kind require large numbers of comparisons that inflate Type 1 error rates.
43 Bonferroni correction can correct for this, but as noted earlier, they are highly conservative
44 and inflate Type 2 error rates. An alternative approach is to demonstrate dependability of
45 significant results through replication over successive surveys, as here.

49 These results suggest several areas of future research. It would strengthen confidence in
50 the associations between factors and workplace wellbeing if they were evaluated
51 longitudinally. Longitudinal research could also shed light on the functional relationship
52 between positive and negative aspects of workplace wellbeing (including depression and
53 burnout) and whether they exert mutual influence over time. It could also be helpful in
54 explaining the cross-cultural variation in the relationship between workplace wellbeing and
55 age. Another question that requires clarification stems from by the inverse relationship
56 between workplace wellbeing and general wellbeing in some early career groups. Finally,
57
58
59
60

1 the particularly low wellbeing people who report disabilities urgently requires further
2 investigation.
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

References

- Ackerley, G. D., Burnell, J., Holder, D. C., & Kurdek, L. A. (1988). Burnout among licensed psychologists. *Professional psychology: Research and practice*, 19(6), 624. doi: 0735-7028/88
- Armstrong, R. A. (2014). When to use the Bonferroni correction. *Ophthalmic and Physiological Optics*, 34(5), 502-508. doi:10.1111/opo.12131
- Bakker, A. B., & Demerouti, E. (2007). The job demands-resources model: State of the art. *Journal of managerial psychology*, 22(3), 309-328. doi: 10.1108/02683940
- Ben-Zur, H., & Michael, K. (2007). Burnout, social support, and coping at work among social workers, psychologists, and nurses: The role of challenge/control appraisals. *Social work in health care*, 45(4), 63-82.
- Beswick, J., Gore, J., & Palerman, D. (2006). Bullying at work: a review of the literature WPS/06/04. *Health and safety laboratory*, 14. Buxton Derbyshire: HSE.
- Bellis, M. A., Lowey, H., Leckenby, N., Hughes, K., & Harrison, D. (2013). Adverse childhood experiences: retrospective study to determine their impact on adult health behaviours and health outcomes in a UK population. *Journal of public health*, 36(1), 81-91. doi:10.1093/pubmed/fdt038
- Berjot, S., Altintas, E., Grebot, E., & Lesage, F. X. (2017). Burnout risk profiles among French psychologists. *Burnout research*, 7, 10-20. doi: 10.1016/j.burn.2017.10.001
- Bianchi, R., Schonfeld, I. S., & Laurent, E. (2015). Burnout–depression overlap: A review. *Clinical psychology review*, 36, 28-41. doi: 10.1016/j.cpr.2015.01.004
- Bland, J. M. & Altman, D. G. (1996). Measurement error. *British Medical Journal*, 313, 744. doi: 10.1136/bmj.313.7059.744
- Boorman, S. (2009). *NHS health and well-being: Final report*. London: Department of Health.
- Chung, Y., Graham, R., Downs, B., Sandler, D.H, Sienkiewicz, R. (2017). The parental gender earnings gap in the United States. Center for Economic Studies, CES 17-68. Washington DC: US Census Bureau
- Clark, A. E. (1996). Job satisfaction in Britain. *British journal of industrial relations*, 34(2), 189-217. doi:10.1111/j.1467-8543.1996.tb00648.x
- Costello, A. B., & Osborne, J. W. (2005). Best practices in exploratory factor analysis: Four recommendations for getting the most from your analysis. *Practical Assessment, Research and Evaluation*, 10(7), 1-9. doi: 10.1.1.110.9154

- 1 D'Souza, F., Egan, S. J., & Rees, C. S. (2011). The relationship between perfectionism, stress and burnout in
2 clinical psychologists. *Behaviour Change*, 28(1), 17-28. doi:10.1375/behc.28.1.17
- 3
4
5 Dattilio, F. M. (2015). The self-care of psychologists and mental health professionals: A Review and
6 practitioner guide. *Australian Psychologist*, 50(6), 393-399. doi: 10.1111/ap.12157
- 7
8
9 De Cuyper, N., & De Witte, H. (2009). Temporary employment: Associations with employees' attitudes,
10 well-being and behaviour. A review of Belgian research. *Psychologica Belgica*, 49(4), 249.
11
12
13
14
15 doi:10.5334/pb-49-4-249
- 16 DeVellis, R. F. (2017). *Scale development: Theory and applications* (4th ed.). Los Angeles: Sage Publications.
- 17
18 Diener, E., Emmons, R. A., Larsen, R. J., & Griffin, S. (1985). The satisfaction with life scale. *Journal of*
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Di Benedetto, M., & Swadling, M. (2014). Burnout in Australian psychologists: Correlations with work-
setting, mindfulness and self-care behaviours. *Psychology, Health and Medicine*, 19(6), 705-715.
doi:10.1080/13548506.2013.861602
- Dorociak, K. E., Rupert, P. A., & Zahniser, E. (2017). Work life, well-being, and self-care across the
professional lifespan of psychologists. *Professional Psychology: Research and Practice*, 48(6), 429-
437. doi: 10.1037/pro0000160
- Drezner, Z. & Drezner, T.D. (2016). A Remedy for the Overzealous Bonferroni Technique for Multiple
Statistical Tests. *Bulletin of the Ecological Society of America*, 97(1), 91-98. doi:10.1002/bes2.1214
- Emery, S., Wade, T. D., & McLean, S. (2009). Associations among therapist beliefs, personal resources and
burnout in clinical psychologists. *Behaviour Change*, 26(2), 83-96. doi: 10.1375/behc.26.2.83.
- Field, A. (2018). *Discovering statistics using IBM SPSS Statistics* (5th ed.) London: Sage Publications.
- Gazioglu, S., & Tansel, A. (2006). Job satisfaction in Britain: individual and job related factors. *Applied*
economics, 38(10), 1163-1171. doi: 10.1080/00036840500392987
- Glass, D. C., & McKnight, J. D. (1996). Perceived control, depressive symptomatology, and professional
burnout: A review of the evidence. *Psychology and health*, 11(1), 23-48. doi:
10.1080/08870449608401975
- Goldberg, D. (1992). *General Health Questionnaire (GHQ-12)*. Windsor: NFER-NELSON

- 1 Griffeth, R. W., Hom, P. W., & Gaertner, S. (2000). A meta-analysis of antecedents and correlates of
2 employee turnover: Update, moderator tests, and research implications for the next millennium.
3 Journal of management, 26(3), 463-488. doi: 10.1177/014920630002600305
4
5
6
7 Hacker-Hughes, J. H., Rao, A. S., Dosanjh, N., Cohen-Tovée, E., Clarke, J., & Bhutani, G. (2016). Physician
8 heal thyself (Luke 4: 23). *The British Journal of Psychiatry*, 209(6), 447-448.
9 doi:10.1192/bjp.bp.116.185355
10
11
12
13 Hall, L. H., Johnson, J., Watt, I., Tsipa, A., & O'Connor, D. B. (2016). Healthcare staff wellbeing, burnout, and
14 patient safety: a systematic review. *PloS one*, 11(7), e0159015. doi:10.1371/.0159015.
15
16
17 Health and Safety Executive (HSE). (2004). Management Standards Indicator Tool. Retrieved from
18 <https://www.hse.gov.uk/stress/assets/docs/indicatortool.pdf>
19
20
21
22 Health Education England (2019). *NHS Staff and Learners' mental wellbeing commission*. Birmingham:
23 Department of Health.
24
25
26
27 Hobfoll, S. E., & Shirom, A. (2001). Conservation of resources theory: Applications to stress and
28 management in the workplace. In R. T. Golembiewski (Ed.), *Handbook of organizational*
29 *behavior* (pp. 57-80). New York, US: Marcel Dekker.
30
31
32
33
34 Johnson, S., Osborn, D. P., Araya, R., Wearn, E., Paul, M., Stafford, M., ... & Anderson, E. (2012). Morale in
35 the English mental health workforce: questionnaire survey. *The British Journal of Psychiatry*, 201(3),
36 239-246.
37
38
39
40 Kristensen, T.S., Borritz, M., Villadsen, E., & Christensen, K.B. (2005). The Copenhagen Burnout Inventory: A
41 new tool for the assessment of burnout. *Work and Stress*, 19(3), 192-207.
42 doi:10.1080/02678370500297720
43
44
45
46 Maben, J., Peccei, R., Adams, M., Robert, G., Richardson, A., Murrells, T., & Morrow, E. (2012). Exploring
47 the relationship between patients' experiences of care and the influence of staff motivation, affect
48 and wellbeing. Final report. Southampton: NIHR Service Delivery and Organization Programme.
49
50
51
52 Maslach, C., & Jackson, S. E. (1981). The measurement of experienced burnout. *Journal of Organizational*
53 *Behavior*, 2(2), 99-113. doi: 10.1002/job.4030020205
54
55
56
57 Maslach, C., Jackson, S.E., & Leiter, M.P. (1996). *Maslach Burnout Inventory manual* (3rd ed.). Palo Alto:
58 Consulting Psychologists Press.
59
60

- 1 Maslach, C., Jackson, S. E., & Leiter, M. P. (2017) Maslach burnout inventory manual. (4th ed.) USA: Mind
2 Garden, Inc.
3
4
- 5 Maslach, C., Schaufeli, W. B., & Leiter, M. P. (2001). Job burnout. *Annual Review of Psychology*, 52, 397-
6 422. doi: 10.1146/annurev.psych.52.1.397
7
8
- 9
10 Michie, S & Williams, S. (2003). Reducing work related psychological ill health and sickness absence: A
11 systematic literature review. *Occupational and Environmental Medicine*, 60(1), 3-9.
12 doi.org/10.1136/oem.60.1.3.
13
14
- 15
16 Nevala, N., Pehkonen, I., Koskela, I., Ruusuvoori, J., & Anttila, H. (2015). Workplace accommodation among
17 persons with disabilities: A systematic review of its effectiveness and barriers or facilitators. *Journal*
18 *of occupational rehabilitation*, 25(2), 432-448. doi: 10.1007/s10926-014-9548-z
19
20
- 21
22 Ng Fat, L. N., Scholes, S., Boniface, S., Mindell, J., & Stewart-Brown, S. (2017). Evaluating and establishing
23 national norms for mental wellbeing using the short Warwick–Edinburgh mental well-being scale
24 (SWEMWBS): findings from the health survey for England. *Quality of Life Research*, 26(5), 1129-
25 1144. doi: 10.1007/s11136-016-1454-8
26
27
- 28
29
30 NHS Staff Survey Coordination Centre. (2018). 2018 Results.
31
32 <https://www.nhsstaffsurveys.com/Page/1064/Latest-Results/2018-Results/> [Accessed 24/10/2019].
33
34
- 35 Office for National Statistics. (2109). Gender pay gap in the UK: 2019. London: OfNS
36
- 37 Ruotsalainen J.H., Verbeek J.H., Mariné A., & Serra C. (2015). Preventing occupational stress in healthcare
38 workers. *Cochrane Database of Systematic Reviews* (4). doi: 10.1002/14651858.CD002892.pub5
39
40
- 41 Rupert, P. A., & Kent, J. S. (2007). Gender and work setting differences in career-sustaining behaviors and
42 burnout among professional psychologists. *Professional Psychology: Research and Practice*, 38(1),
43 88- 96. doi: 10.1037/0735-7028.38.1.88
44
45
- 46
47 Rupert, P. A., & Morgan, D. J. (2005). Work setting and burnout among professional psychologists.
48 *Professional Psychology: Research and Practice*, 36(5), 544–550. doi: 10.1037/0735-7028.36.5.544
49
50
- 51
52 Rupert, P. A., Stevanovic, P., & Hunley, H. A. (2009). Work-family conflict and burnout among practicing
53 psychologists. *Professional Psychology: Research and Practice*, 40(1), 54. doi: 10.1037/a0012538
54
55
- 56 Shah, N., Cader, M., Andrews, W. P., Wijesekera, D., & Stewart-Brown, S. L. (2018). Responsiveness of the
57 Short Warwick Edinburgh Mental Well-Being Scale (SWEMWBS): evaluation a clinical
58 sample. *Health and quality of life outcomes*, 16(1), 239. doi: 10.1186/s12955-018-1060-2
59
60

- 1 Stevenson, J. & Rao, M. (2014). Explaining levels of wellbeing in Black and Minority Ethnic populations in
2 England. London: Institute for Health and Development.
3
4
- 5 Stewart-Brown, S., Tennant, A., Tennant, R., Platt, S., Parkinson, J., & Weich, S. (2009). Internal construct
6 validity of the Warwick-Edinburgh Mental Well-being Scale (WEMWBS): A Rasch analysis using data
7 from the Scottish Health Education Population Survey. *Health and Quality of Life Outcomes*, 7(1), 1–
8 8. doi: 10.1186/1477-7525-7-15.
9
10
- 11 Summers, E. M. A., Morris, R. C. & Bhutani, G. E. (2020). A measure to assess the workplace wellbeing of
12 psychological practitioners. *Clinical Psychology & Psychotherapy*, 27(1), 11-23.
13 doi.org/10.1002/cpp.2401
14
15
- 16 Tennant, R., Hiller, L., Fishwick, R., Platt, S., Joseph, S., Weich, S., ... & Stewart-Brown, S. (2007). The
17 Warwick-Edinburgh mental well-being scale (WEMWBS): development and UK validation. *Health
18 and Quality of Life Outcomes*, 5(1), 63-76. doi: 10.1186/1477-7525-5-63
19
20
21
22
23
24
- 25 World Health Organization. (2018). International classification of diseases for mortality and morbidity
26 statistics (11th Revision). Retrieved from <https://icd.who.int/browse11/l-m/en>
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Studies of burnout in psychological practitioners

Authors (year) <i>Country of study</i>	Sample size (% response rate)	Profession (%)	Years of experience	Burnout measure	Main Conclusion
Ackerley et al. (1988) <i>USA</i>	562 (35.4)	Licensed psychologists (100)	<i>M</i> = 18.8 years Range = 1-43	MBI (1986)	Each MBI subscale showed higher burnout than the norms for mental health workers. 'High burnout' threshold exceeded on EE & DP scales for over 1/3 of the sample
Berjot et al. (2017) <i>France</i>	664 (NR)	Professional psychologists (100)	<i>M</i> = 8.01	French- Canadian version MBI-HSS (Maslach et al., 1996) French translation (Dion & Tessier, 1994)	Over 60% over cut-off for burnout on EE & DP scales
Di Benedetto & Swadling (2014) <i>Australia</i>	167 (NR)	Registered psychologists (100)	NR	CBI (Kristensen et al., 2005)	14.4% above threshold for burnout on the overall score. Burnout negatively correlated with experience.
D'Souza et al. (2011) <i>Australia</i>	87 (NR)	Clinical psychologists (100)	1-5 years (39%) 6-10 years (29%) 11-15 years (14%) 16-20 years (6%) Over 20 years (12%)	CBI (Kristensen et al., 2005)	8% above threshold for burnout on overall score. Average of all three scales above burnout threshold Negative correlation with age.
Emery et al. (2009) <i>Australia</i>	190 (39)	Clinical psychologists (100)	< 5 years (31.1%) 5-9 years (24.7%) 10-14 years (13.7%) 15-19 years (10%) 20+ years (20.5%)	MBI-HSS (Maslach et al., 1996)	Only the PA scale average was above burnout threshold. No association with age.
Rupert & Kent (2007) <i>USA</i>	595 (49.6)	Clinical psychologists (83.2) Counselling psychologists (16.8)	<i>M</i> = 17.91 <i>SD</i> = 8.07	MBI-HSS (Maslach et al., 1996)	Overall mean was in average burnout range for EE & DP scales, low for PA scale. Negative correlation with age.
Rupert & Morgan (2005) <i>USA</i>	571 (47.6)	Clinical psychologists (80) Counselling psychologists (20)	<i>M</i> = 16.93 <i>SD</i> = 7.66	MBI-HSS (Maslach et al., 1996)	Average burnout on DP & EE scales, low on PA scale. Negative correlation with age.
Rupert et al. (2009) <i>USA</i>	487 (40.58)	Licensed psychologists (100)	<i>M</i> = 19.6 <i>SD</i> = 7.8	MBI-HSS (Maslach et al., 1996)	Average burnout on DP & EE scales, low on PA scale. Negative correlation with age.

Table 2 Demographic data by gender

	Male		Female	
	Count	%	Count	%
Age by gender**				
18 - 24	2	0.6%	27	2.0%
25 - 34	58	18.1%	401	30.1%
35 - 44	93	29.1%	377	28.3%
45 - 54	100	31.3%	354	26.5%
55 - 64	55	17.2%	145	10.9%
65 - 74	10	3.1%	19	1.4%
75 or older	0	0.0%	0	0.0%
Undisclosed	2	0.6%	11	0.8%
**Kruskal-Wallis Test, $p < .000$				
Contract type by gender				
Permanent/ Open ended	272	85.0%	1141	85.5%
Non-permanent/ Fixed-term/ Secondment	32	10.0%	154	11.5%
Self Employed/ Private Practice/ Independent	5	1.6%	14	1.0%
Other	11	3.4%	25	1.9%
Agenda for Change salary by gender*				
Band 3	2	0.7%	3	0.2%
Band 4	8	2.7%	49	3.9%
Band 5	23	7.7%	152	12.0%
Band 6	30	10.0%	157	12.4%
Band 7	92	30.8%	315	24.9%
Band 8a	54	18.1%	286	22.6%
Band 8b	27	9.0%	109	8.6%
Band 8c	35	11.7%	136	10.7%
Band 8d	18	6.0%	42	3.3%
Band 9	5	1.7%	6	0.5%
Do not wish to disclose	5	1.7%	12	0.9%
*Kruskal-Wallis Test, $p = .03$				
Years worked since qualified by gender**				
Up to 5 years	88	27.5%	515	38.6%
5 - 10 years	79	24.7%	308	23.1%
11 - 15 years	50	15.6%	198	14.8%
16 - 20 years	40	12.5%	127	9.5%
21 - 25 years	24	7.5%	82	6.1%
26 - 30 years	16	5.0%	56	4.2%
31 - 35 years	10	3.1%	32	2.4%
36 years or more	13	4.1%	16	1.2%
**Kruskal-Wallis Test, $p < .000$				

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 3 PPWWM means by group

	CATEGORY	N(%) [†]	PPWWM Mean(SD)	SWEMWBS Mean(SD)
PROFESSIONAL GROUP	Clinical Psychologist	814(49)	90.85(16.57)	22.08(3.36)
	Counselling Psychologist	39(2)	92.05(20.70)	22.97(4.34)
	High Intensity Therapist	152(9)	89.11(19.96)	21.69(4.77)
	Psychological Wellbeing Practitioner	226(13)	94.88(18.22)*	23.11(4.64)**
	Counsellor	95(6)	95.66(16.59)*	22.88(3.81)
	Cognitive Behaviour Therapist	148(9)	91.25(18.52)	22.07(4.07)
	Psychotherapist	42(3)	91.05(18.15)	22.58(4.71)
	Trainee Clinical Psychologist	28(2)	96.11(14.84)	20.54(2.86)**
	Nurse/Mental Health Nurse	20(1)	92.80(16.46)	22.13(4.39)
	Assistant Psychologist	24(1)	102.58(15.51)*	20.88(3.02)*
	Other	90(5)	91.29(17.34)	21.92(4.26)
Total	1678	91.88(17.55)***	22.21(3.92)***	
CONTRACT	Permanent/ Open ended	1434(85)	91.21(17.67)	22.19 (3.97)
	Non-permanent/ Fixed-term/ Secondment	189(11)	95.99(15.29)***	21.79(3.25)
	Self Employed/ Private Practice/ Independent	19(1)	103.84(17.92)**	25.66(3.90)**
	Other	36(2)	90.33(19.02)	23.48 (4.16)
	Total	1678	91.88(17.55)***	22.21(3.92)***
SECTOR	NHS	1471(88)	91.20(17.61)	22.10(3.97)
	Private organisation/ Independent	56(3)	97.20(16.64)*	23.09(3.10)
	Third sector/ Charitable organisation	72(4)	99.82(15.21)**	23.12(4.09)
	Equal NHS and non-NHS	22(1)	92.09(17.43)	22.71(2.95)
	Education	17(1)	97.00(16.64)	23.17(2.51)
	Other	40(2)	92.90(16.71)	22.81(3.31)
	Total	1678	91.88(17.55)***	22.21(3.92)
GENDER	Male	320(19)	92.10(17.83)	22.09(3.99)
	Female	1334(81)	91.86(17.35)	22.24(3.89)
	Total	1654	91.91(17.44)	22.21(3.91)
SEX ORIENTATION	Heterosexual	1412(89)	92.06(17.50)	22.29(3.91)
	Bisexual	81(5)	92.75(15.51)	21.07(3.41)
	Lesbian/Gay/Queer	93(6)	92.89(18.55)	22.29(4.79)
	Total	1586	92.14(17.46)	22.23(3.95)
ETHNICITY	Asian or Asian British (Bangladeshi/ Indian/ Pakistani/ Any other Asian)	66(4)	97.26(20.08)*	23.21(4.08)*
	Black or Black British (African/ Caribbean/ Any other Black)	40(2)	94.28(23.10)	24.01(6.02)
	Mixed (Any mixed ethnicity)	51(3)	87.14(18.71)	22.00(4.47)
	White (British/ Irish/ Any other White ethnicity)	1440(89)	91.79(17.03)	22.11(3.81)
	Other ethnic group (Chinese/ Any other)	25(2)	94.80(18.06)	23.16 (4.44)
	Total	1622	91.97(17.45)*	22.21(3.93)**
DI	Report Disability	279(17)	88.24(17.08)	21.23 (3.75)

1					
2		No Report of Disability	1353(83)	92.82(17.38)	22.41(3.91)
3	SABILI	TY			
4		Total	1632	92.04(17.41)***	22.21(3.91)***

† Based on the number answering the question

* $p < .05$; ** $p < .01$; *** $p < .001$: For subgroups the p-values refer to differences from the overall mean.

For Peer Review

Table 4 Significant correlations of key variables with PPWWM

Variable	Correlation [†]	N
Additional unpaid hours on average per week	<i>Tau-b</i> = -.189***	1656
How often in the past month have you wanted to leave the NHS	<i>Tau-b</i> = -.511***	1455
Age	<i>Tau-b</i> = -.043*	1678
Agenda for Change Band	<i>Tau-b</i> = -.036*	1678
Years served in current post	<i>Tau-b</i> = -.075***	1590
Number of years since qualifying	<i>Tau-b</i> = -.041*	1678
Harassment or bullying—Service Users/Public	<i>Tau-b</i> = -.147***	1678
Harassment or bullying—Managers	<i>Tau-b</i> = -.319***	1678
Harassment or bullying—Colleagues	<i>Tau-b</i> = -.165***	1678
Time in past week felt depressed	<i>Tau-b</i> = -.400***	1678
Time in the past week felt a failure	<i>Tau-b</i> = -.352***	1678


[†] All correlations used bootstrapping with 2000 iterations and were two-tailed.

* $p < .05$ *** $p < .0001$

Table 5 Factor Structure

Item	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	PPWWM Dimension / Construct
Factor 1: Professional and organisational satisfaction						
7. I feel I can balance less fulfilling aspects of my job with more enjoyable aspects	.43	.25	.38	.33	.12	Positive and negative job aspects/Work control and autonomy
8. I cannot see how the service/ organisation in which I work can ever be delivered effectively	.64	.06	.13	.13	.17	Organisational context/Organisational hopefulness-hopelessness
10. I am enabled to manage and organise my workload and diary	.40	.37	.18	.37	-.05	Positive and negative job aspects/Work control and autonomy
12. I feel confident the service/ organisation in which I work can adapt to meet future service demands	.66	.21	.16	.18	.16	Organisational context/Organisational hopefulness-hopelessness
18. I do not feel included in service/ organisational decisions that affect me	.59	.12	.21	-.08	.13	Organisational context/Organisational engagement
23. I am expected to reach unrealistic or unattainable targets	.64	.21	.06	.26	.14	Organisational context/Organisational targets
25. I feel service/ organisational targets are meaningful	.71	.09	.14	.14	.15	Organisational context/Organisational targets
Factor 2: Support and flexibility						
1. I do not feel there is always someone there for me when I need personal support	.15	.43	.24	.17	.15	Personal support-lack of support/Family support
4. Flexible working arrangements are supported in my service/organisation	.34	.37	.26	-.11	-.08	Positive and negative job aspects /Work-life balance
5. I feel supported by my line-manager to take positive risks without fear of reproach	.38	.60	.30	-.08	-.02	Personal support-lack of support/Line management
9. The clinical supervision I receive is containing and safe	.00	.78	.11	.27	.19	Personal support-lack of support/Clinical supervision
13. Clinical supervision meets my support needs	.08	.75	.10	.35	.22	Personal support-lack of support/Clinical supervision
20. My line-manager is approachable and responsive	.24	.62	.31	-.04	-.01	Personal support-lack of support/Management
21. My continuing professional development needs are supported	.50	.50	.15	-.03	.07	Positive and negative job aspects/Opportunities to learn
22. I am encouraged and supported to develop my skill-set and knowledge	.53	.53	.22	-.04	.09	Positive and negative job aspects/Opportunities to learn
Factor 3: Professional Role						
2. I feel I can seek support from my colleagues	-.01	.35	.68	.09	.11	Personal support-lack of support/Colleagues
3. I feel a sense of belonging to the service/ organisation in which I work	.40	.25	.59	-.02	.12	Organisational context/Organisational engagement
6. I work in an environment where my colleagues are caring and supportive towards each other	.06	.32	.70	.05	.12	Personal support-lack of support/Colleagues
11. I am clear of my role in relation to other professionals with whom I work	.32	.13	.47	.25	.01	Inter-professional Agents/Role clarity
14. My colleagues have realistic expectations of my professional role	.33	.23	.50	.31	.15	Inter-professional Agents/Role clarity
16. My colleagues value my professional contribution	.14	.12	.73	.19	.12	Positive and negative job aspects /Feeling valued
26 My specific skills as a psychological practitioner add value to the team/ service/ organisation	.25	.01	.55	.18	-.02	Positive and negative job aspects /Feeling valued
Factor 4: Physical Environment						
15. The physical environment and facilities in my workplace enable me to work efficiently and effectively	.30	.17	.19	.08	.81	Positive and negative job aspects /Physical environment and facilities
24. The physical environment and/ or facilities in my workplace adversely affect my workplace wellbeing	.26	.13	.11	.07	.85	Positive and negative job aspects /Physical environment and facilities
Factor 5: External Personal						
17. I have a good work/ life balance	.29	.15	.20	.65	.04	Positive and negative job aspects /Work-life balance
19. The personal support I receive from family and/or friends meets my needs	-.00	.03	.17	.71	.07	Personal support-lack of support/Family support

Figure 1. Distribution of the PPWWM



Note: Can be printed in grey-scale

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

For Peer Review