



Extensions of K.J. Roy's and R.E. Roadifer's Subjective Approach to Oil Resource Estimation

Baecher, G.B.

IIASA Working Paper



1975

Baecher, G.B. (1975) Extensions of K.J. Roy's and R.E. Roadifer's Subjective Approach to Oil Resource Estimation. IIASA Working Paper. WP-75-054 Copyright © 1975 by the author(s). <http://pure.iiasa.ac.at/374/>

Working Papers on work of the International Institute for Applied Systems Analysis receive only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute, its National Member Organizations, or other organizations supporting the work. All rights reserved. Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage. All copies must bear this notice and the full citation on the first page. For other purposes, to republish, to post on servers or to redistribute to lists, permission must be sought by contacting repository@iiasa.ac.at

EXTENSIONS OF K.J. ROY'S AND
R.E. ROADIFER'S SUBJECTIVE APPROACH
TO OIL RESOURCE ESTIMATION

Gregory B. Baecher

May 1975

WP-75-54

Working Papers are not intended for distribution outside of IIASA, and are solely for discussion and information purposes. The views expressed are those of the author, and do not necessarily reflect those of IIASA.

Messrs. Roy and Roadifer, in separate papers, have proposed similar methodologies for making regional resource estimates. The idea underlying their approach is quite interesting, and may have extensions incorporable in larger schemes of estimation.

Roy's and Roadifer's approach adopts a simple "structural" model of deposit size, and uses the subjective feelings of geologists to generate probability density functions (pdf's) of the model parameters. A pdf of regional resources is approximated by using Monte Carlo simulation.

Specifically, the model they adopt is of the form

$$P = A \times T \times PR \times RF \times c_0 ,$$

in which

P = reservoir or deposit potential ,

A = reservoir or deposit area ,

T = formation thickness ,

PR = proportion of the formation which is resource ,

RF = recovery factor ,

c_0 = constant .

We call this a "structural" model here because it is based on a physical or geological conception of resources, though a simple one. Examples of other structural models are those dealing with mineralization, sedimentation, migration and accumulation, etc. from first principles of physics and chemistry.

A second type of models are those of more purely empirical

nature: regression, factor analysis, pattern recognition, etc. These models are not based on fundamental concepts of geological processes, but on correlations and simple relationships extracted from past data. Perhaps the best known use of statistical empirical models in economic geology is that of Harris (1965).

The procedure of Roy and Roadifer can be extended to use statistical models as well as structural models. The advantage of this extension is that statistical models (if carefully developed) may be better estimators of resource potential than either simple structural models or purely subjective estimation (e.g., Harris, 1973). Further, statistical models make use of a larger set of geological variables than simple structural models, and can be applied to other types of resources for which simple structural models are not available. In either case, probabilistic estimates of regional resources made prior to extensive exploration should be used as a priori probabilities rather than final estimates, and updated by subsequent exploration data using the normal Bayesian procedure.

A proposed extension of Roy's and Roadifer's model is the following. Let some regression analysis have been performed on previous acquired data from geologically similar and extensively explored regions. Let the result of these analyses be expressed as a linear function of the common form

$$R = \underline{\beta} \underline{x} + \underline{e} \quad , \quad (1)$$

in which \underline{x} is a vector of relevant geological variables,* $\underline{\beta}$ is a vector of regression coefficients, and \underline{e} is a random error term with zero-mean and variance σ^2 . If this analysis has been undertaken within a Bayesian framework, the terms $\underline{\beta}$ and σ^2 will have some joint pdf, $f(\underline{\beta}, \sigma^2)$, which describes their uncertainty.

A priori, values of \underline{x} for the region in question are uncertain. However, these values may be estimated subjectively by several geologists familiar with the region, and may be coalesced by the method of Morris (1974). This leads to a joint subjective probability distribution of the geological variables, $f(x_1, \dots, x_n)$; or if the variables are assumed mutually independent, to a set of single variable pdf's, $f(x_1), \dots, f(x_n)$. Both Roy and Roadifer describe this process.

Inserting these pdf's of \underline{x} in equation 1, and using an error-in-the-variables approach to regression and prediction, a probability distribution for R, $f(R)$ can be generated which is a function of the uncertainty in \underline{x} and in $(\underline{\beta}, \sigma^2)$. The mathematics of this approach are currently

* Some of these might be time of formation, environment of deposition or mineralization, subsequent tectonic activity, etc.

available in the literature of econometrics and statistics, and if certain well known families of distributions are chosen to model uncertainty in the variables \underline{x} and \underline{e} , closed form solutions are available (Zellner, 1971). This would make reversion to Monte Carlo methods unnecessary.

Subsequent exploration data could be included in the analysis using $f(R)$ as a prior distribution, and updating as described in Kaufman (1974), or in other papers dealing with Bayesian sampling theory.

This total procedure might lead to more comprehensive regional estimates than any component of it in isolation, and may provide better *a priori* estimates than either simple structural models or purely subjective estimation.

Gregory B. Baecher
Laxenburg
19.V.75

REFERENCES

- Harris, D.P. (1973). "A subjective probability appraisal of metal endowment in Northern Sonora, Mexico." Econ. Geol., 68: 222-242.
- _____ (1965). An Application of Multivariate Statistical Analysis to Mineral Exploration. PhD thesis, Penn. State Univ.
- Kaufman, G.M. (1974). "Statistical methods for predicting the number and size distribution of undiscovered hydrocarbon deposits." in, Crandall, K.H., and J.W. Harbaugh, eds, Methods of Estimating the Volume of Undiscovered Oil and Gas Resources. Stanford Univ.
- Morris, P. (1974). "Decision analysis expert use." Mgt. Sci., 20: 1233-41.
- Roadifer, R.E. (1975). "A probability approach to estimate volumes of undiscovered oil and gas." IIASA Workshop on Energy Resources, preprint.
- Roy, K.J. (1975). "Hydrocarbon assessment using subjective probability and Monte Carlo methods." IIASA Workshop on Energy Resources, preprint.