



Criteria for Proportional Representation

Balinski, M.L. and Young, H.P.

**IIASA Research Report
December 1976**



Balinski, M.L. and Young, H.P. (1976) Criteria for Proportional Representation. IIASA Research Report. Copyright © December 1976 by the author(s). <http://pure.iiasa.ac.at/525/> All rights reserved. Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage. All copies must bear this notice and the full citation on the first page. For other purposes, to republish, to post on servers or to redistribute to lists, permission must be sought by contacting repository@iiasa.ac.at

CRITERIA FOR PROPORTIONAL REPRESENTATION

M. L. BALINSKI*
H. P. YOUNG*
DECEMBER 1976

* Graduate School and University Center, City University of New York, and
International Institute for Applied Systems Analysis, Laxenburg, Austria.

This work was supported in part by the National Science Foundation under
Contract MPS 75-07414 with the Graduate School of the City University of
New York.

Research Reports provide the formal record of research conducted by the
International Institute for Applied Systems Analysis. They are carefully
reviewed before publication and represent, in the Institute's best judgment,
competent scientific work. Views or opinions expressed herein, however,
do not necessarily reflect those of the National Member Organizations support-
ing the Institute or of the Institute itself.

**International Institute for Applied Systems Analysis
2361 Laxenburg, Austria**

Preface

The problem of how to make a “*fair division*” of resources among competing interests arises in many areas of application at IIASA. One of the tasks in the System and Decision Sciences area is the systematic investigation of different criteria of fairness and the formulation of allocation procedures based thereon.

A particular problem of fair division having wide application in governmental decision-making is the *apportionment* problem. An application has recently arisen in the debate over how many seats in the European Parliament to allocate to the different member countries. Discussions swirled around particular numbers, over which agreement was difficult to achieve. A systematic approach that seeks to establish bases for agreement on the criteria or “*principles*” for fair division should stand a better chance of acceptance in that it represents a scientific or system-analytic approach to the problem.

Abstract

Methods to allocate seats in proportional representation systems are investigated in terms of several underlying common-sense properties. In particular, the idea of stability is introduced, and the method of Jefferson (or d'Hondt) is characterized.

Criteria for Proportional Representation

INTRODUCTION

There exist wholesale numbers of possible election procedures. A basic characterization has been made of these into "plurality systems" and "proportional representation systems."

In a *plurality system* an elector usually casts one vote for the candidate or the (party) list of candidates of his or her choice in some election district, and the candidate or list receiving a majority or plurality is elected. Such systems are based on a notion of geographical representation. Mid-nineteenth century Europe saw an increasing dissatisfaction with plurality systems as unfair to minorities, for small political parties were effectively barred from having any representation whenever their adherents were distributed throughout many single-member election districts.

This led to the idea of *proportional representation* which, in its pure form, has electors cast one vote for a party or party list in a multi-member district and then, by some rule, metes numbers of seats "proportionally" among the parties according to their respective vote totals. Of course, variants of both types of system exist, as do complex mixtures of both.

This paper is focused on the pure form of the *proportional representation problem*: voters cast a single vote for a party in a multi-member district and the question is to determine the just number of representatives due each party. Exact proportionality cannot, in general, be achieved since representation must be integral. Some "rounding" must take place. Consider, for example, the problem of Table 1 with party vote totals as given and 36 seats to be allocated. Six different possible solutions are advanced...which *should* be chosen?

In fact the identical problem arises under plurality systems, but in a different guise. For, usually, a nation is divided into states or provinces and each single-member election district is wholly contained in one such subdivision. How many election districts or representatives should one geographical region be allocated? This problem is known as the *apportionment problem*. Of course, geographical apportionment can arise in proportional representation systems too. The apportionment problem in the United States arises from interpreting the somewhat vague Constitutional mandate "...Representatives...shall be apportioned among the several states...according to their respective members." Senator Daniel Webster of Massachusetts caught the spirit of the intended solution to the apportionment problem in his definition: "To

apportion is to distribute by right measure, to set off in just parts, to assign in due and proper proportion" [17, p. 107]. But, appearances to the contrary, an operational definition is not easily forthcoming and United States history is rich with controversies over proposed solutions and methods. It suffices to consult Table 1 to see why.

Table 1

Party	Votes Received	Exact Proportionality	Possible Allocations					
			SD	H HM	EP	W	Q	J
A	27,744	9.988	10	10	10	10	10	11
B	25,178	9.064	9	9	9	9	10	9
C	19,947	7.181	7	7	7	8	7	7
D	14,614	5.261	5	5	6	5	5	5
E	9,225	3.321	3	4	3	3	3	3
F	3,292	1.185	2	1	1	1	1	1
	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>
	100,000	36.000	36	36	36	36	36	36

Furthermore, what seems right, just and proper in dealing with apportionment in plurality systems may or may not seem so in dealing with allocation in proportional election systems. Different sets of properties of possible methods for solving the underlying problem are suggested by the practical situation in which the mathematical allocation problem is embedded.

FORMULATION

Let $\underline{p} = (p_1, \dots, p_s)$ be the (positive, integer) vote totals of s parties (or the number of citizens deserving representation in s regions), where each $p_i > 0$, and $h \geq 0$ is the number of seats to be allocated (or apportioned). The problem is to find an *allocation for h*: an s -tuple of non-negative integers (a_1, \dots, a_s) whose sum is h . A *solution* of the problem is a function \underline{f} which to any population vector \underline{p} and all h associates a unique apportionment for h , $a_i = f_i(\underline{p}, h) \geq 0$ where $\sum_i a_i = h$.

If \underline{f} is a solution, then \underline{f}^h will represent the function \underline{f} restricted to the domain (\underline{p}, h') where $0 \leq h' \leq h$. \underline{f}^h is a solution up to h , and \underline{f} is an extension of \underline{f}^h .

A specific allocation method may give several solutions, since "ties" may occur when using it, as for example when two parties have identical vote totals but must share an odd number of seats. For this reason it is useful to define an allocation method \underline{M} as a non-empty set of solutions. Thus, in particular, a solution up to some given h may have several different extensions in a method \underline{M} . In the sequel only symmetric methods of allocation are considered, that is, methods in which the ordering of the parties in the list (p_1, \dots, p_s) is immaterial. This is a clearly essential property.

The choice of an allocation, or of an allocation method, cannot and should not revolve about the rival numerical results of one solution over another. Rather, the issue is to decide upon a rule that is "fair," that is, whose qualitative properties satisfy criteria or "principles" acceptable to both citizen and politician. It is as Representative E.W. Gibson said on the floor of the U.S. Congress in 1929, "The apportionment of Representatives to the population is a mathematical problem. Then why not use a method that will stand the test...?" [5].

STABILITY

Three principal methods seem to have been considered for proportional representation systems: Sainte-Lagüe's [15], d'Hondt's [8,9], and Hamilton's [7], usually known as "la répartition au plus fort reste." In the apportionment literature Sainte-Lagüe's and d'Hondt's have found their places (see [10]), under other names, in a class of five so-called "modern" [4,13] methods which, from about 1920 through 1974, were the only ones collectively considered for apportioning the United States House of Representatives.

The five "modern" methods were first grouped by E.V. Huntington in 1921 [11] via an approach to allocation (or apportionment) based on pairwise comparisons of "inequality in representation." Given vote totals $\underline{p} = (p_1, \dots, p_s)$ and an allocation $\underline{a} = (a_1, \dots, a_s)$ for h consider the numbers p_i/a_i and a_i/p_i . These represent the number of votes per representative of party i and the number of representatives per vote of party i . If $p_i/a_i > p_j/a_j$, or $a_i/p_i < a_j/p_j$, or $a_j > a_i(p_j/p_i)$, or $\dots, (p_i/a_i)(a_j/p_j) > 1$, then party j is better off than party i . Given a particular measure or test of inequality between a pair of parties such as $|p_i/a_i - p_j/a_j|$ or $|a_i/p_i - a_j/p_j|$ it is natural to ask whether the amount of inequality can be reduced by a transfer of one seat from the better-off party to the less-well-off party. Given a measure or test T an allocation is

said to be *in equilibrium* if no transfer reduces the value of T for any pair of parties. Of course, certain conceivable measures T may not (and do not) admit equilibrium solutions for all vote-total distributions, but Huntington showed [10,11] that five measures T do.

Huntington's approach can be developed in several ways. One is via the tests of inequality in representation mentioned above. Another (see [13,10]) is by letting $d(a)$, a *divisor criterion*, be any real-valued monotone-increasing function on the non-negative integers such that $d(0) \geq 0$. Given a divisor criterion $d(a)$, a *divisor method* M of allocation is the set of solutions obtained recursively as follows:

$$(i) \quad f_i(\underline{p}, 0) = 0 \quad , \quad 1 \leq i \leq s \quad .$$

(ii) If $a_i = f_i(\underline{p}, h)$ is an M -allocation for h and k is some one party for which

$$p_k/d(a_k) \geq p_i/d(a_i) \quad \text{for } 1 \leq i \leq s \quad ,$$

then

$$f_k(\underline{p}, h+1) = a_k + 1 \quad , \quad f_i(\underline{p}, h+1) = a_i \quad \text{for } i \neq k \quad .$$

The meaning of such an allocation method is that the numbers $p_i/d(a_i)$ represent some measure of the "priority" of a party to receive one more seat, the next seat being given to the most deserving party. The divisor function $d(a)$ thus represents some sort of "weighting" of the current number of seats the party has.

The five "modern" methods proposed by Huntington are divisor methods, and their various names, associated tests of inequality and divisor criteria are given in Table 2 below. Huntington himself argued strongly and effectively for the method he called *Equal Proportions*. It is in equilibrium for the measure of inequality which is the relative difference* between the pair of numbers p_i/a_i and p_j/a_j , and also between a_i/p_i and a_j/p_j . This method is now the law for apportioning the U.S. House of Representatives [14].

*The relative difference between x and y is $|x - y|/\min(x, y)$.

Table 2. Huntington's five methods.

Method	In Equilibrium for Test T (where $p_i/a_i \geq p_j/a_j$)	Divisor Criterion $d(a)$
Smallest Divisors (SD)	$T_1: a_j - a_i(p_j/p_i)$	a
Harmonic Mean (HM)	$T_2: p_i/a_i - p_j/a_j$	$2a(a+1)/(2a+1)$
Equal Proportions (EP)	$T_3: (p_i a_j / p_j a_i) - 1$	$(a(a+1))^{1/2}$
Webster (W) or Major Fractions or Sainte-Lagüe Formula	$T_4: a_j/p_j - a_i/p_i$	$(a+1/2)$
Jefferson (J) or Greatest Divisors or d'Hondt or plus forte moyenne	$T_5: a_j(p_i/p_j) - a_i$	$a + 1$

The method of equal proportions (EP) necessarily first gives to each party one seat. As an example of a divisor method in use, Table 3 below gives EP allocations to parties for the example of Table 1 for a house size h ranging from 6 to 16 seats.

D'Hondt's method [8,9], or "la répartition à la plus forte moyenne," also called in the apportionment literature "the method of greatest divisors," was in fact first proposed by Thomas Jefferson [12] in 1792 and has therefore been called *Jefferson's method J* [3]. Sainte-Lagüe's method [15], also called in the apportionment literature "the method of major fractions," was in fact first suggested in embryonic form by Daniel Webster [17] in 1832 and has therefore been called the *Webster method W* [3]. A particular variant of \bar{W} is the modified method of odd numbers (see, e.g., [6]) used in some Scandinavian countries. It is defined by: $d(0) = 7/10$ (instead of $d(0) = 1/2$), and otherwise the divisors are identical with those of \bar{W} , $d(a) = a + 1/2$.

It is an interesting historical note that Sainte-Lagüe [15] came upon the Webster method quite independently via the idea of minimizing a total measure of the inequality of an allocation. He proposed that an allocation should minimize

$$\sum_i p_i \left(\frac{h}{p} - \frac{a_i}{p_i} \right)^2 ,$$

since in a perfect allocation $h/p = a_i/p_i$ for all i . The Webster method provides solutions which do this. In the same paper Sainte-Laguë suggests in words (though not in symbols) that one could be interested in minimizing

$$\sum_i a_i \left(\frac{p}{h} - \frac{p_i}{a_i} \right)^2 ,$$

but that "one is led to a more complex rule." In fact, this gives precisely the method of equal proportions.

Table 3. Sample EP allocations.

Party	A	B	C	D	E	F
Vote Total	27,744	25,178	19,947	14,614	9,225	3,292
House size						
6	1	1	1	1	1	1
7	2	1	1	1	1	1
8	2	2	1	1	1	1
9	2	2	2	1	1	1
10	3	2	2	1	1	1
11	3	2	2	2	1	1
12	3	3	2	2	1	1
13	3	3	3	2	1	1
14	4	3	3	2	1	1
15	4	4	3	2	1	1
16	4	4	3	2	2	1
⋮	⋮	⋮	⋮	⋮	⋮	⋮
36	10	9	7	6	3	1

It is useful to know that there is a "local" criterion by which to verify whether allocation belongs to a divisor method.

Lemma. $\bar{a} = (a_1, \dots, a_s)$ is an allocation for h of a divisor method with divisor criterion $d(a)$ if and only if

$$(1) \quad \min_j p_j/d(a_j-1) \geq \max_i p_i/d(a_i) \quad .$$

This is immediate by definition. \square

Parties in proportional representation systems are dynamic. They may group together for electoral purposes, but they also may splinter. It is likely that the behavior of parties will be influenced by the mathematical consequences of their coalescing as versus splintering. Specifically, it is pertinent to ask how the number of seats allocated by a method \underline{M} to the joint vote total of two parties coalesced into one compares with the seats allocated by \underline{M} to the two parties separately. Consider a symmetric method \underline{M} and a problem with vote totals p in which some party has p^* votes and another \bar{p} votes; and consider the result of coalescing the star- and the bar-party into one party with $p^* + \bar{p}$ votes, keeping all other party vote totals the same. Let $f \in \underline{M}$ be a solution in which a^* and \bar{a} are the number of seats allocated to the star- and the bar-party, respectively, in an allocation for h . Then \underline{M} is said to be *stable* if there exists an allocation for h in which the number b of seats allocated to the coalesced party satisfies $a^* + \bar{a} - 1 \leq b \leq a^* + \bar{a} + 1$.

Theorem 1. A divisor method \underline{M} with divisor criterion $d(a)$ satisfying

$$(2) \quad d(a_1+a_2) \leq d(a_1) + d(a_2) \leq d(a_1+a_2+1)$$

is stable.

Proof: Suppose that an allocation for h of a problem specified by the vote totals $p = (p_1, \dots, p_s)$ is $\bar{a} = (a_1, \dots, a_s)$ and that parties 1 and 2 form a coalition having vote total $p_1 + p_2$. Then we show that if $(a_1+a_2, a_3, \dots, a_s)$ is not an allocation for h then there exists one that gives the coalition either $a_1 + a_2 - 1$ or $a_1 + a_2 + 1$ seats.

If $a_1 + a_2$ is not a solution, then by the Lemma

$$\min\{(p_1+p_2)/d(a_1+a_2-1), \min_{i \neq 1,2} p_i/d(a_i-1)\} \neq \max\{(p_1+p_2)/d(a_1+a_2), \max_{j \neq 1,2} p_j/d(a_j)\} \quad .$$

Thus since a satisfies (1), either

$$(3) \quad (p_1+p_2)/d(a_1+a_2-1) < \max_{j \neq 1,2} p_j/d(a_j) = p_k/d(a_k) = \lambda \quad (k \neq 1,2)$$

or

$$(4) \quad (p_1+p_2)/d(a_1+a_2) > \min_{i \neq 1,2} p_i/d(a_i-1) = p_\ell/d(a_\ell-1) = \delta \quad (\ell \neq 1,2) .$$

Before proceeding, notice that if $s_i, v_i \geq 0$ ($i=1,2$) then

$$\max \{s_1/v_1, s_2/v_2\} \geq (s_1+s_2)/(v_1+v_2) \geq \min \{s_1/v_1, s_2/v_2\} .$$

This inequality is used repeatedly.

Consider the first case (3). From (2) we have

$$\begin{aligned} (p_1+p_2)/d(a_1+a_2-2) &\geq (p_1+p_2)/(d(a_1-1)+d(a_2-1)) \geq \min\{p_1/d(a_1-1), p_2/d(a_2-1)\} \\ &\geq \max_i p_i/d(a_i) \geq \lambda \end{aligned}$$

and find

$$\begin{aligned} \min\{(p_1+p_2)/d(a_1+a_2-2), p_k/d(a_k), \min_{i \neq 1,2,k} p_i/d(a_i-1)\} &\geq \lambda \\ &\geq \max\{(p_1+p_2)/d(a_1+a_2-1), p_k/d(a_k+1), \max_{i \neq 1,2,k} p_i/d(a_i)\} , \end{aligned}$$

showing that $(a_1+a_2-1, a_3, \dots, a_k+1, \dots, a_s)$ is an allocation for h .

On the other hand, consider (4). Since by (2)

$$\begin{aligned} (p_1+p_2)/d(a_1+a_2+1) &\leq (p_1+p_2)/(d(a_1)+d(a_2)) \leq \max\{p_1/d(a_1), p_2/d(a_2)\} \\ &\leq \min_j p_j/d(a_j-1) \leq \delta , \end{aligned}$$

we find

$$\min\{(p_1+p_2)/d(a_1+a_2), p_\ell/d(a_\ell-2) \min_{i \neq 1,2,\ell} p_i/d(a_i-1)\} \geq \delta$$

$$\geq \max\{(p_1+p_2)/d(a_1+a_2+1), p_\ell/d(a_\ell-1), \max_{j \neq 1,2,\ell} p_j/d(a_j)\} ,$$

showing that $(a_1+a_2+1, a_3, \dots, a_\ell-1, \dots, a_s)$ is an allocation for h . \square

Corollary. Hamilton's five divisor methods are stable.

The method that has been and is most often proposed for proportional representation happens to be the seemingly most natural one. Although known by several names, including "la répartition au plus fort reste" and "Vinton's method of 1850," it was apparently first proposed by Alexander Hamilton [7] in 1792 and has therefore been called the *Hamilton method* H [3]. We define the *exact quota* of party j to be $q_j = q_j(p, h) = p_j h / \sum_i p_i$. It is the exactly proportional number of seats deserved by party j and the number that one would wish to allocate to j were it integral. Otherwise, each party should, it seems, receive at least as many seats as $\lfloor q_j \rfloor$ (the largest integer less than or equal to q_j). This motivates the allocations of the *Hamilton method*: First, give to each party i $\lfloor q_i \rfloor$ seats; then, order the parties by their fractional remainders $d_j = q_j - \lfloor q_j \rfloor \geq 0$ in a priority list $d_{j_1} \geq \dots \geq d_{j_s}$. Second, give one additional seat to each of the first $h - \sum \lfloor q_j \rfloor$ parties on the list. If there are ties then there exist distinct arrangements of the priority list and, hence, possibly several solutions. In fact, this method, under the name of the *Vinton method of 1850*, was adopted as the solution to the apportionment problem for the U.S. House of Representatives for the censuses of 1850 through 1900.

Theorem 2. The Hamilton method is stable.

Proof: Consider any two parties, say $i = 1, 2$, and suppose that in a particular problem party i has an exact quota $q_i = n_i + r_i, n_i \geq 0$ integer and $0 \leq r_i < 1$, and let a_i be their allocations at h . Then for the problem in which parties 1 and 2 form a coalition, its exact quota for h is $q_1 + q_2 = n_1 + n_2 + r_1 + r_2$. Let b be the number of seats given the coalition by the Hamilton method.

We consider several cases. First, if $b = n_1 + n_2$ then $r_1 + r_2 < 1$, implying that the same total number of parties is rounded

up in both problems. If $r_1, r_2 > 0$ then it must be that $a_i = n_i$ ($i=1,2$). For otherwise one of the two parties would have a remainder r_i high enough in the list to warrant an extra seat while $r_1 + r_2 > r_1, r_2$ is not high enough, which cannot be. If $r_1 = 0$ then $a_1 = n_1$ and $a_2 = n_2$ or $a_2 = n_2 + 1$; in either case the criterion for stability is satisfied.

If $b = n_1 + n_2 + 1$, then since $a_i = n_i$ or $n_i + 1$ there is nothing more to show. If $b = n_1 + n_2 + 2$, then $r_1 + r_2 > 1$. Suppose stability is not satisfied, i.e., that $a_i = n_i$ ($i=1,2$). Then for some party $k \neq 1,2$, $r_1 + r_2 - 1 \geq r_k$ while $r_1 \leq r_k$ and $r_2 \leq r_k$. Thus $2r_k - 1 \geq r_k$ and $r_k \geq 1$, a contradiction. \square

MONOTONICITY AND CONSISTENCY

The Hamilton method came under sharp criticism and was subsequently abandoned by the U.S. Congress for, in 1881, it chanced to admit the not so congenial "Alabama paradox" [16]. Consider the vote totals of Table 1 and apply H to apportion 37 and 38 seats (Table 4): parties D and E both lose a seat as the total number of seats to be allocated increases! For the data of the 1880

Table 4.

Party	A	B	C	D	E	F	Totals
Vote Total	27,744	25,178	19,947	14,614	9,225	3,292	100,000
Exact quota 35	9.711	8.812	6.982	5.115	3.229	1.152	35
H allocation 35	10	9	7	5	3	1	35
Exact quota 36	9.998	9.064	7.181	5.261	3.321	1.185	36
H allocation 36	10	9	7	5	4	1	36
Exact quota 37	10.265	9.316	7.380	5.407	3.413	1.218	37
H allocation 37	10	9	7	6	4	1	37
Exact quota 38	10.543	9.568	7.580	5.533	3.506	1.251	38
H allocation 38	11	10	8	5	3	1	38
Exact quota 39	10.820	9.819	7.779	5.699	3.598	1.284	39
H allocation 39	11	10	8	6	3	1	39

census, Alabama suffered by dropping from 8 to 7 seats as h increased from 299 to 300. This phenomenon is not rare numerically. For example, the populations of the states used to apportion the U.S. Congress in 1901 gave rise to many similar occasions. Between $h = 381$ and $h = 391$, Maine changed between 2 and 3 seats six times and led a Representative of that State to exclaim "...it does seem as though mathematics and science has combined to make a shuttlecock* and battledore* of the State of Maine in connection with the scientific basis upon which this bill is presented" (8 January 1901).

An evident property of any allocation method is that the number of seats accorded to any party not *decrease* if the house size *increases*. A method \underline{M} is said to be *monotone* if for any \underline{M} -solution \underline{f} , and all \underline{p}, h

$$f_i(\underline{p}, h+1) \geq f_i(\underline{p}, h) \quad , \quad i = 1, \dots, s \quad .$$

\underline{H} fails to satisfy this test and hence was emphatically rejected.

Divisor methods are obviously monotone. Indeed, it was for this reason that they were formulated. Notice, moreover, that divisor methods have a certain inner regularity; namely, the decision as to which party of any pair most deserves the extra seat as the house size is increased by 1 depends only upon the populations and seats already allocated to those parties singly, and not on the vote vector \underline{p} or the vector of seats so far allocated \underline{a} .

More generally, consider a method \underline{M} and suppose that it has a solution \underline{f} allocating to a party with p^* votes a^* seats and to a party with \bar{p} votes \bar{a} seats in a house h , while \underline{f} allocates to the star-party $a^* + 1$ seats, and to the bar-party \bar{a} seats in a house $h + 1$. Then the star-party is said to have *weak priority* at that point, and this is written $(p^*, a^*) \preceq (\bar{p}, \bar{a})$. A natural criterion for any method is that the claims of any two parties for an extra seat should depend *only* on their respective vote totals and allocations. Specifically, if for some other allocation problem with votes \underline{p}' there are parties having votes p^* and \bar{p} that are allocated, by a solution of \underline{M} , a^* and \bar{a} seats respectively, and also $(\bar{p}, \bar{a}) \preceq (p^*, a^*)$, then the parties are said to be *tied*, and this is written $(p^*, a^*) \sim (\bar{p}, \bar{a})$. A method is said to be *consistent* if it treats tied parties equally, that is, if $(p^*, a^*) \sim (\bar{p}, \bar{a})$ implies \underline{f}^h has both an extension giving the star-party $a^* + 1$ seats at $h + 1$, and an extension giving the bar-party $\bar{a} + 1$ seats at $h + 1$. Any two parties will naturally compare their resultant numbers of seats: a change in priorities could not but be viewed as conflicting with common sense. Note that any consistent method is automatically symmetric.

*The projectile and racquet, respectively, in badminton.

Divisor methods are clearly consistent. The Hamilton method is not. For consider the H allocations of Table 4. Parties D and E receive 5 and 3 seats, respectively, for $h = 35$ and uniquely received 5 and 4 for $h = 36$. However, they receive 5 and 3, respectively, for $h = 38$, and uniquely 6 and 3 for $h = 39$. This provides one more argument against the Hamilton method.

Let $r(p,a)$ be any real-valued function of two real variables called a *rank-index* (possibly including $\pm \infty$ for certain values of p and a). Given a rank-index, a (generalized) *Huntington method* [3] of allocation M is the set of all solutions f obtained recursively as follows:

- (i) $f_i(p,0) = 0$, $1 \leq i \leq s$.
- (ii) if $a_i = f_i(p,h)$ is an M allocation for h and k is some one party for which

$$r(p_k, a_k) \geq r(p_i, a_i) \text{ for } 1 \leq i \leq s \text{ ,}$$

then

$$f_k(p, h+1) = a_k + 1 \text{ , } f_i(p, h+1) = a_i \text{ for } i \neq k \text{ .}$$

Huntington methods are a direct generalization of divisor methods. They are clearly monotone and consistent. Less obvious is

Theorem 3 [2]. *Any consistent, monotone method of allocation is a Huntington method.*

There do, nevertheless, exist rather dubious Huntington methods. For example, take $r(p,a) = a/p$. Then the first party to receive one seat must necessarily receive all. We will say that a method is *balanced* if two parties having identical vote totals are always allocated numbers of seats that differ by at most 1. Divisor methods are balanced; so is the Hamilton method. But Huntington methods in general are not. A method that is not balanced could not be countenanced.

COALITIONS AND SCHISMS

In the context of proportional representation it is important to ask, not only whether methods are stable, but also whether they tend to encourage parties to merge or to splinter. For *political* stability it would usually be considered desirable to have methods of allocation that encourage parties to coalesce by assuring that this would never cost seats, but could in fact result in an increase in the total number of seats allocated jointly to the parties.

To make these ideas precise, consider a symmetric stable method \underline{M} and a problem with vote totals p in which one party has p^* votes and another \bar{p} votes; and consider the result of coalescing the star- and the bar-party into one party with $p^* + \bar{p}$ votes, keeping all other party vote totals the same. Let $\underline{f} \in \underline{M}$ be a solution in which a^* and \bar{a} are the number of seats allocated to the star- and the bar-party, respectively, in an allocation for h . Then \underline{M} is said to *encourage coalitions* if there exists an allocation giving the coalesced party b seats with $b \geq a^* + \bar{a}$, and to *encourage schisms* if $b \leq a^* + \bar{a}$.

A method \underline{M} is said to be *unique* satisfying certain properties if any other set \underline{M}' of allocation solutions with the same properties is also a set of solutions by \underline{M} , that is, $\underline{M}' \subseteq \underline{M}$.

Theorem 4. The Jefferson method \underline{J} is the unique consistent, monotone and balanced method that encourages coalitions.

Viewed in this light the Jefferson method presents strong credentials for being adopted in a proportional representation system. Sainte-Laguë appears to have realized the tendency of \underline{J} to encourage coalitions, but he gave no proofs and his statement has the curiosity of referring to a comparison: "In comparing the two rules, one can show that the d'Hondt rule (\underline{J}) favors the grouping of parties which, by coalescing, may receive more seats; whereas the method of least squares (\underline{W}) favors neither groupings nor schisms," [15, p. 378].

Proof: Since \underline{J} is a divisor method, it is easily seen that it is consistent, monotone, and balanced. To show that \underline{J} encourages coalitions, let \underline{a} be a Jefferson allocation of h for given vote totals \underline{p} , and consider the vote totals $(p_1+p_2, p_3, \dots, p_s)$ in which parties 1 and 2 have formed a coalition. If $(a_1+a_2, a_3, \dots, a_s)$ is not a solution, then by the Lemma,

$$(5) \quad \min\{(p_1+p_2)/(a_1+a_2), \min_{i \neq 1,2} p_i/a_i\} \neq \max\{(p_1+p_2)/(a_1+a_2+1), \max_{j \neq 1,2} p_j/(a_j+1)\}$$

Since \underline{a} is a Jefferson allocation,

$$(p_1+p_2)/(a_1+a_2) \geq \min\{p_1/a_1, p_2/a_2\} \geq \max_{j \neq 1,2} p_j/(a_j+1) \quad ,$$

so the only reason (5) can hold is that

$$(6) \quad (p_1+p_2)/(a_1+a_2+1) > \min_{i \neq 1,2} p_i/a_i = p_\ell/a_\ell = \delta \quad (\ell \neq 1,2) \quad .$$

On the other hand,

$$(7) \quad \begin{aligned} (p_1+p_2)/(a_1+a_2+2) &\leq \max \{p_1/(a_1+1), p_2/(a_2+1)\} \\ &\leq \min_j p_j/a_j \leq \delta \quad , \end{aligned}$$

hence, combining (6) and (7),

$$\begin{aligned} \min \{ (p_1+p_2)/(a_1+a_2+1), p_\ell/(a_\ell-1), \min_{j \neq 1, 2, \ell} p_j/a_j \} &\geq \delta \\ &\geq \max \{ (p_1+p_2)/(a_1+a_2+2), p_\ell/a_\ell, \max_{j \neq 1, 2, \ell} p_j/(a_j+1) \} \quad , \end{aligned}$$

which shows (by the Lemma) that $(a_1+a_2+1, a_3, \dots, a_\ell-1, \dots, a_s)$ is a Jefferson allocation for h .

Conversely, let \underline{M} be any method having the stated properties. Then \underline{M} is a Huntington method, hence (by Theorem 3) has a rank-index $r(p, a)$. Since \underline{M} is balanced, $r(p, a) > r(p, a+1) > \dots > r(p, a+k)$ for any integer $k > 0$. It will be shown that $r(p, a)$ is equivalent to $p/(a+1)$.

As a first step we show that $r(p, a) = r(np, na+n-1)$ for all integers $n \geq 1$. Consider the vote vector $\underline{p} = (p_1, \dots, p_{n+1}) = (p, \dots, p)$ and house size $h = (n+1)a + n$ for any integer $a \geq 0$. Since \underline{M} is balanced we can assume an allocation for h $(a, a+1, \dots, a+1)$. Now consider the vote vector (p, np) and a corresponding two-party allocation for h , say (x, y) . By coalition encouragement there exists a solution with $y \geq na + n$ and hence $x \leq a$. There is a lowest $h' \leq h$ such that the second party (with vote total np) has y seats and the first some $x - k$ seats, $k \geq 0$. Thus $r(np, y-1) \geq r(p, x-k) \geq r(p, x)$ and so

$$(8) \quad r(p, a) \leq r(p, x) \leq r(np, y-1) \leq r(np, na+n-1) \quad .$$

Consider, instead, the vote vector $(p_1, \dots, p_{2n}) = (p, \dots, p)$ and house size $h = 2na + n$. \underline{M} balanced implies that it must have an allocation of form $(a+1, \dots, a+1, a, \dots, a)$ with $a_i = a + 1$ for $1 \leq i \leq n$ and $a_i = a$ for $n + 1 \leq i \leq 2n$. Now consider the vote totals $(p_1, \dots, p_n, p_{n+1}) = (p, \dots, p, np)$ and let (x_1, \dots, x_n, y) be an allocation of \underline{M} for $h = 2na + n$. Since \underline{M} is stable, $y \leq na + n - 1$,

and thus there is an i with $1 \leq i \leq n$ having $x_i \geq a + 1$. There is a lowest $h' \leq h$ such that party i has x_i seats and the $(n+1)$ st party (with vote total np) some $y - k$ seats, $k \geq 0$. Thus $r(p, x_i - 1) \geq r(np, y - k) \geq r(np, y)$ and so

$$(9) \quad r(p, a) \geq r(p, x_i - 1) \geq r(np, y) \geq r(np, na + n - 1)$$

which, with (8), shows that $r(p, a) = r(np, na + n - 1)$.

To complete the proof suppose $p/(a+1) = p'/(a'+1)$. Then

$$r(p, a) = r(p'p, p'a + p' - 1) = r(pp', pa' + p - 1) = r(p', a') \quad ,$$

implying that $r(p, a) = R(p/(a+1))$ for some function R . R is order-preserving, for suppose $p/(a+1) < p'/(a'+1)$. Then $pa' + p < p'a + p'$, and

$$\begin{aligned} R(p/(a+1)) &= R(pp'/(p'a + p')) = r(pp', p'a + p' - 1) \\ &< r(p'p, pa' + p - 1) = R(p'p/(pa' + p)) = R(p'/a' + 1) \quad , \end{aligned}$$

which completes the proof. \square

The method of smallest divisors is, in a certain sense, "symmetric" with \underline{J} . This suggests

Theorem 5. The method of smallest divisors \underline{SD} is the unique consistent, monotone and balanced method that encourages schisms.

Proof: We first show that $r(p, a) = r(np, na)$ for all integers $n \geq 1$. Consider the vote vector $(p_1, \dots, p_{n+1}) = (p, \dots, p)$ and house size $h = na + a + 1$ for any integer $a \geq 0$. \underline{M} balanced implies that an allocation at h has form $(a+1, a, \dots, a)$. Now consider (p, np) and any allocation of \underline{M} (x, y) . By schism encouragement there exists a solution with $y \leq na$ and hence $x \geq a + 1$. An argument identical to that given in Theorem 4 shows that for $k \geq 0$

$$(10) \quad r(p, a) \geq r(p, x - 1) \geq r(np, y - k) \geq r(np, y) \geq r(np, na) \quad .$$

Consider, instead, the vote vector $(p_1, \dots, p_{2n}) = (p, \dots, p)$ and house size $h = 2na + n$. \underline{M} balanced implies $(a, \dots, a, a+1, \dots, a+1)$ is an allocation of h . Now consider the vote vector $(p_1, \dots, p_n, p_{n+1}) = (p, \dots, p, np)$. Since \underline{M} is stable there exists an

allocation (x_1, \dots, x_n, y) with $y \geq na + 1$ and so there must be some i , $1 \leq i \leq n$, with $x_i \leq a$. So, we obtain as before

$$r(p, a) \leq r(p, x_i) \leq r(np, y-1) \leq r(np, na) \quad ,$$

which, with (10), shows that $r(p, a) = r(np, na)$. The proof that SD results from the stated properties is completed by showing that $r(p, a) = R(p/a)$ for some order-preserving function R , as in the proof of Theorem 4.

That SD is consistent, monotone and balanced is clear. To show that it encourages schisms requires an argument that precisely parallels the corresponding part of the proof of Theorem 4. \square

SATISFYING QUOTA

The ideal, exactly proportional number of seats "due" party j is the exact quota $q_j(p, h) = p_j h / \sum_i p_i$. Given \underline{p} and h , if $q_i = q_i(\underline{p}, h)$ is integer for all i then $a_i = q_i$ is a seemingly perfect solution. In general, of course, we cannot expect *any* of the exact quotas to be integer. On the other hand, it is natural to require that no party receive more seats than the result of "rounding" q_i up ($\lceil q_i \rceil$, the smallest integer greater than or equal to q_i) and no less than the result of "rounding" q_i down ($\lfloor q_i \rfloor$, the largest integer less than or equal to q_i). A method is said to *satisfy quota* if all of its allocations satisfy $\lfloor q_i \rfloor \leq a_i \leq \lceil q_i \rceil$. Allocations not satisfying quota seem to violate common sense, and have proven to be politically subject to attack [17].

Aside from the fairness question, a method satisfying quota clearly is balanced, and--while not necessarily stable in the strict sense defined above--is *almost stable* in the sense that if any two parties with a^* and \bar{a} seats coalesce, then the coalesced party receives b seats, where $a^* + \bar{a} - 2 \leq b \leq a^* + \bar{a} + 2$.

The Hamilton method (\underline{H}) clearly satisfies quota--indeed can be said to have been motivated by this desire. However, as we have seen, it violates two other basic principles. Aside from \underline{H} , what other methods satisfy quota? The emphasis historically, particularly since Huntington's work, has been on divisor methods. However, none of these can satisfy quota. In fact, a subsequent result (Theorem 7) implies the following.

Theorem 6. There exists no allocation method that is monotone, consistent and satisfies quota.

Since these three properties seem to be essential ones, an immediate question is: what can be weakened in order to admit the existence of some method?

In the presence of monotonicity and satisfying quota it is reasonable to consider a slight weakening of consistency. Consider a method M and let $f \in M$ with $f_j(p, h) = a_j$. Then party j is said to be *eligible at $h+1$* for its (a_j+1) st seat if $a_j + 1 \leq \{p_j(h+1) / \sum_i p_i\}$; equivalently, $a_j < q_j(p, h+1) = p_j(h+1) / \sum_i p_i$, that is if party j can be given an extra seat without exceeding upper quota. We say that a method M is *quota-consistent** if it is consistent as between eligible parties. More precisely, (p^*, a) has *weak eligible priority* over (\bar{p}, \bar{a}) written $(p^*, a^*) \succ^e (\bar{p}, \bar{a})$, if in some problem one party having p^* votes and a^* seats has weak priority over another party having \bar{p} votes and \bar{a} seats and both are *eligible* to receive another seat. If both $(p^*, a^*) \succ^e (\bar{p}, \bar{a})$ and $(\bar{p}, \bar{a}) \succ^e (p^*, a^*)$ we write $(p^*, a^*) \sim^e (\bar{p}, \bar{a})$. For M to be *quota-consistent* it is required that whenever $(p^*, a^*) \sim^e (\bar{p}, \bar{a})$ and both parties are eligible at $h+1$ but f gives the $(h+1)$ st seat to the p^* -party, then there is an extension of f^h giving the $(h+1)$ st instead to the \bar{p} -party.

The *quota method* Q of allocation is the set of all solutions f obtained recursively as follows:

- (i) $f_i(p, 0) = 0 \quad 1 \leq i \leq s$.
- (ii) Let $a_i = f_i(p, h)$ be an allocation for h of Q and $E(h+1)$ the set of parties eligible to receive an extra seat at $h + 1$. If $k \in E(h+1)$ is some one party for which

$$p_k / (a_k + 1) \geq p_i / (a_i + 1) \text{ for all } i \in E(h+1)$$

then

$$f_k(p, h+1) = a_k + 1 \quad , \quad f_i(p, h+1) = a_i \text{ for } i \neq k \quad .$$

* Called "consistent" in [1,3].

Theorem 7 [4]. Q is the unique allocation method that is house-monotone, quota-consistent and satisfies quota.

This theorem gives powerful reasons for acceptance of the quota method Q for both problems of apportionment in majority systems and problems of allocation in proportional representation systems. In practice one finds that Q has a tendency to produce solutions that round up the exact quotas of large parties more often than those of small parties. This seems reasonable for the application of Q to proportional representation systems in that it inferentially asks for a "large" vote before according any representation at all. Notice, however, that no large party is allowed more seats than its upper quota. This same property might appear to cause a difficulty for the application of Q to apportionment problems, since in this situation it is usually necessary for each geographical entity to receive some minimum number of seats (e.g., 1 or 2). For this case the method Q has an immediate generalization that allows the specification of any desired "unbiased" minimum representations for districts, and a uniqueness theorem similar to Theorem 7 obtains [3].

CONCLUSIONS

This paper has addressed the problem of the allocation of integral representation to parties having vote totals in a proportional representation system. The principal point is that methods of allocation should not be chosen by bickering over numbers, nor, indeed, through *ad hoc* claims of various mechanical procedures, but rather by analysis of the properties of methods. The issue is to decide upon a method whose qualitative properties are equitable for the situation at hand. For proportional representation systems this analysis commends one of two methods: the quota method Q or the Jefferson method J .

The Jefferson method claims recognition because it is monotone, consistent, balanced and encourages coalitions. Specifically, encouraging coalitions would seem to be precisely the type of stability desired for a body politic operating a proportional representation system. However, a major defect of J is that it fails the seemingly most common-sense test of satisfying quota. The Quota method merits recognition because it does satisfy quota, is consistent subject to that property, and is monotone and balanced. But it is only almost stable, and, in fact, does not necessarily encourage coalitions.

The choice between these methods of allocation should be made in terms of which *criteria* are viewed as most important for the situation in question.

References

- [1] Balinski, M.L., and H.P. Young, A New Method for Congressional Apportionment, *Proceedings of the National Academy of Sciences, U.S.A.*, 71 (1974), 4602-4606.
- [2] Balinski, M.L., and H.P. Young, On Huntington Methods of Apportionment, submitted for publication.
- [3] Balinski, M.L., and H.P. Young, The Quota Method of Apportionment, *American Mathematical Monthly*, 82 (1975), 701-730.
- [4] Bliss, G.A., E.W. Brown, L.P. Eisenhart, and R. Pearl, Report to the President of the National Academy of Sciences, 9 February 1929, in *Congressional Record*, 70th Congress, 2nd Session, 70 (1929), 4966-4967.
- [5] *Congressional Record*, 70th Congress, 2nd Session, 70 (1929), 1500.
- [6] Cotteret, J.M., and C. Emeri, *Les systèmes électoraux*, Presses Universitaires de France, Paris, 1970.
- [7] Hamilton, Alexander, *The Papers of Alexander Hamilton, Vol. XI (February 1792 - June 1792)*, Harold C. Syrett, ed., Columbia University Press, New York, 1966, 228-230.
- [8] d'Hondt, V., *La représentation proportionnelle des partis par un électeur*, Ghent, 1878.
- [9] d'Hondt, V., *Système pratique et raisonné de représentation proportionnelle*, Muquardt, Brussels, 1882.
- [10] Huntington, E.V., The Apportionment of Representatives in Congress, *Transactions of the American Mathematical Society*, 30 (1928), 85-110.
- [11] Huntington, E.V., The Mathematical Theory of the Apportionment of Representatives, *Proceedings of the National Academy of Sciences, U.S.A.*, 7 (1921), 123-127.
- [12] Jefferson, T., *The Works of Thomas Jefferson, Vol. VI*, Paul Leicester Ford, ed., G.P. Putnam and Sons, New York, 1904, 460-471.
- [13] Morse, M., J. von Neumann, and L.P. Eisenhart, Report to the President of the National Academy of Sciences, 28 May, 1948.

- [14] U.S. Public Law 291, H.R. 2665, 55 Stat. 761, 15 November, 1941.
- [15] Sainte-Lagüe, La représentation et la méthode des moindres carrés, *Comptes Rendus de l'Académie des Sciences*, 151 (1910), 377-378.
- [16] Seaton, C.W., Report to the Chairman of the Committee of the Census, 25 October 1881, in *Apportionment Among the Several States*, House of Representatives, 56th Congress, 2nd Session, Report No. 2130, 20 December 1900.
- [17] Webster, D., *The Writings and Speeches of Daniel Webster*, Vol. VI, National Edition, Little, Brown and Company, Boston, Massachusetts, 1903, 102-123.