



Differentiability of a Support Function of an E-Subgradient

Lemarechal, C. and Nurminski, E.A.

IIASA Working Paper

WP-80-101

June 1980



Lemarechal, C. and Nurminski, E.A. (1980) Differentiability of a Support Function of an E-Subgradient. IIASA Working Paper. WP-80-101 Copyright © 1980 by the author(s). <http://pure.iiasa.ac.at/1368/>

Working Papers on work of the International Institute for Applied Systems Analysis receive only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute, its National Member Organizations, or other organizations supporting the work. All rights reserved. Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage. All copies must bear this notice and the full citation on the first page. For other purposes, to republish, to post on servers or to redistribute to lists, permission must be sought by contacting repository@iiasa.ac.at

NOT FOR QUOTATION
WITHOUT PERMISSION
OF THE AUTHOR

DIFFERENTIABILITY OF A SUPPORT FUNCTION
OF AN E-SUBGRADIENT

Claude Lemarechal
Evgeni Nurminski

June 1980
WP-80-101

Working Papers are interim reports on work of the International Institute for Applied Systems Analysis and have received only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute or of its National Member Organizations.

INTERNATIONAL INSTITUTE FOR APPLIED SYSTEMS ANALYSIS
A-2361 Laxenburg, Austria

Differentiability of a support function of an ϵ -subgradient mapping

Claude Lemarechal

INRIA

Eugeni Nurminski

IIASA

ABSTRACT

Directional differentiability of a support function of an ϵ -subgradient set-valued mapping is proved and formula for a directional derivative is given.

The notion of an ϵ -subgradient proposed by R.T. Rockafellar (Rockafellar 1972a), appears to be rather useful for convex nondifferentiable optimization. This is mainly due to their practical advantages from a computational point of view, but it is also important that this mapping has richer analytical properties than the subgradient mapping traditionally studied in convex analysis.

During the last few years it was observed that this mapping has strong continuity properties (Nurminski (1978) a, Hiriart-Urruty 1979a), both in finite dimensional and abstract spaces. Here we state a certain kind of differentiability properties of the support function of ϵ -subgradient mapping which may lead to a definition of second order differentiability of convex functions. Throughout the paper we stay within the framework of convex analysis, so $f(x)$ will be a convex locally Lipschitzian function, $f^*(g)$ its conjugate and $\partial_\epsilon f(x)$ will be the set of ϵ -subgradients. For a number of technical reasons we need an additional assumption that the epigraph of function $f(x)$ does not contain nonvertical straight lines. It can be assured for instance by a coercivity assumption with respect to function $f(x)$.

Definition. The function $V_p(x)$ defined as

$$V_p(x) = \max_{g \in \partial_\epsilon f(x)} pg \quad (1)$$

is called the support function of an ϵ -subdifferential $\partial_\epsilon f(x)$.

The support function $V_p(x)$ has an equivalent representation:

$$V_p(x) = \inf_{t > 0} \frac{f(x + tp) - f(x) + \epsilon}{t} \quad (2)$$

which is easy to obtain from dual consideration of problem (1). Notice that due to coercivity problem (2) always has a bounded solution.

Duality can also be used for obtaining some auxiliary results.

Lemma 1. For fixed x and p let g_x and t_x be any solutions of the problems (1) and (2) respectively. Then

$$g_x \in \partial f(x + t_x p)$$

Proof. Notice that the result is obvious for $\epsilon = 0$. Also if $\epsilon > 0$ then $t_x > 0$. Consider the Lagrange function of problem (1):

$$L(g, t) = pg - \frac{1}{t} (f^*(g) - xg)$$

in which the terms not depending on g are omitted. It follows from the optimality of g_x and t_x that

$$0 \in \partial_g L(g_x, t_x)$$

or

$$0 \in p - \frac{1}{t_x} (\partial f^*(g_x) - x)$$

For g_x one has the inclusion

$$x + t_x p \in \partial f^*(g_x)$$

which is equivalent to

$$g_x \in \partial f(x + t_x p)$$

Q.E.D.

Lemma 2. Let $y = x + \tau d$. Then for any solution t_x of problem (2), any $g \in \partial f(x)$ and $g_y \in \partial_\epsilon f(y)$ such that

$$V_p(y) = pg_y$$

one has

$$V_p(y) - V_p(x) \leq \frac{\tau}{t_x} (g_y - g)d$$

Proof. Let

$$V_p(y) = \inf_{t_y} \frac{1}{t_y} (f(y + t_y p) - f(y) + \epsilon) = \frac{1}{t_y} (f(y + t_y p) - f(y) + \epsilon)$$

$$t_y > 0$$

$$V_p(x) = \frac{1}{t_x} (f(x + t_x p) - f(x) + \epsilon)$$

Then for any $g \in \partial f(x)$

$$t_y V_p(y) = f(y + t_y p) - f(y) + \epsilon \leq f(y + t_y p) - f(x) + \epsilon - g(y - x)$$

and for any $g_y \in \partial f(y + t_y p)$

$$t_x V_p(x) = f(x + t_x p) - f(x) + \epsilon \geq f(y + t_y p) - f(x) + \epsilon + g_y(x + t_x p - y - t_y p)$$

because of Lemma 1 $g_y \in \partial f(y + t_y p)$.

Subtracting these two inequalities we obtain now, taking $g_y \in \partial f(y + t_y p)$ such that

$$V_p(y) = pg_y,$$

$$t_y V_p(y) - t_x V_p(x) \leq g_y(y - x) + (t_y - t_x) V_p(y) = g(y - x)$$

and consequently

$$t_x (V_p(y) - V_p(x)) \leq \tau (g_y - g)d$$

Q.E.D.

These results make it possible to study the differentiability properties of $V_p(x)$.

Let T_x be the set of solutions of problem (2) and G_x be the solution set of problem (1).

The following theorem holds:

Theorem. The support function $V_p(x)$ is directionally differentiable and its directional derivative is given by the formula:

$$V'_p(x, d) = \max_{g_x \in G_x} \min_{\substack{t \in T_x \\ g \in \partial f(x)}} \frac{g_x - g}{t} d$$

Proof. Let $y = x + \tau d$, $\tau > 0$. Using Lemma 2 twice one can obtain two-side bounds of the kind:

$$\frac{V_p(y) - V_p(x)}{\tau} \leq \min_{\substack{g \in \partial f(x) \\ t \in T_x}} \frac{g_y - g}{t} d$$

$$\frac{V_p(y) - V_p(x)}{\tau} \geq \max_{\substack{g \in \partial f(y) \\ t \in T_y}} \frac{g_x - g}{t} d$$

for any $g_x \in G_x$ and $g_y \in G_y$. Passing to the limit when τ goes to $+0$ one can assume that correspondent sequences of ϵ -subgradients and subgradients in upper and lower boundaries converge to some limits. From u.s.c of T_x , see for instance (Hogan 1973a). with related bibliography, it follows that the correspondent sequence of solutions $t^*, \epsilon T_y$ can be assumed convergent to some limit in T_x as well.

Then

$$\lim_{t \rightarrow +0} \frac{V_p(y) - V_p(x)}{\tau} \geq \lim_{t \rightarrow +0} \max_{\substack{g \in \partial f(x) \\ t \in T_y}} \frac{g_x - g}{t} d = \frac{g_x - \bar{g}}{\bar{t}_x} d \geq \min_{\substack{t \in T_x \\ g \in \partial f(x)}} \frac{g_x - g}{t} d$$

for arbitrary $g_x \in G_x$ and some $\bar{g} \in \partial f(x)$.

Also

$$\lim_{t \rightarrow +0} \frac{V_p(y) - V_p(x)}{\tau} \leq \lim_{t \rightarrow 0} \min_{\substack{g \in \partial f(x) \\ t \in T_x}} \frac{g_y - g}{t} d = \min_{\substack{t \in T_x \\ g \in \partial f(x)}} \frac{\bar{g}_x - g}{t} d$$

where $\bar{g}_x \in \partial_\epsilon f(x)$.

So

$$\min_{\substack{t \in T_x \\ g \in \partial f(x)}} \frac{g_x - g}{t} d \leq \min_{\substack{t \in T_x \\ g \in \partial f(x)}} \frac{\bar{g}_x - g}{t} d \leq \max_{g \in G_x} \min_{\substack{t \in T_x \\ g \in \partial f(x)}} \frac{g_x - g}{t} d$$

for arbitrary $g_x \in G_x$ and some $\bar{g}_x \in G_x$ which means that

$$\max_{g \in G_x} \min_{\substack{t \in T_x \\ g \in \partial f(x)}} \frac{g_x - g}{t} d \geq \min_{\substack{t \in T_x \\ g \in \partial f(x)}} \max_{g \in G_x} \frac{g_x - g}{t} d$$

So far as strict inequality is impossible in this case then

$$\lim_{t \rightarrow 0} \frac{V_p(y) - V_p(x)}{t} = \lim_{t \rightarrow 0} \frac{V_p(y) - V_p(x)}{t} = \max_{g_x \in G_x} \min_{\substack{g \in \partial f(x) \\ t_x \in T_x}} \frac{g_x - g}{t} d \quad (3)$$

Q.E.D.

A final remark should be made on the order of *min max* operations in expression (3). The proof of the theorem shows that it is irrelevant in which order these operations are performed. An additional argument for that is that the function $f(x, y, \theta) = \theta(x - y)b$ for $0 < \theta_0 \leq \theta \leq \theta_0$ has a saddle point in variables (y, θ) and x , where x and y are taken from some compact sets.

References

- Rockafellar 1972a. R.T. Rockafellar, *Convex Analysis*, Princeton University Press (1972).
- Nurminski (1978)a. E. Nurminski, "Continuity of E-subgradient mappings," *Kibernetika (Cybernetics)* 5, pp.148-149 (790-791) (1977 (1978)).
- Hiriart-Urruty 1979a. J.-B. Hiriart-Urruty, "Lipschitz r-continuity of the approximate subdifferential of a convex function," Technical report, Universite de Clermont-Ferrand II Complexe Scientifique des Cezeaux Departement de Mathematiques Appliquees Boite Postale n 45 (D).

Hogan1973a W.W. Hogan, "Point-to-set maps in mathematical programming," *SIAM Review*
15, pp.591-603 (1973).