



What We Have Also Learned: Adaptive Speciation is Theoretically Possible

Doebeli, M., Dieckmann, U., Metz, J.A.J. and Tautz,
D.

IIASA Interim Report
March 2005



Doebeli, M., Dieckmann, U., Metz, J.A.J. and Tautz, D. (2005) What We Have Also Learned: Adaptive Speciation is Theoretically Possible. IIASA Interim Report. Copyright © 2005 by the author(s). <http://pure.iiasa.ac.at/7817/>

Interim Report on work of the International Institute for Applied Systems Analysis receive only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute, its National Member Organizations, or other organizations supporting the work. All rights reserved. Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage. All copies must bear this notice and the full citation on the first page. For other purposes, to republish, to post on servers or to redistribute to lists, permission must be sought by contacting repository@iiasa.ac.at



International Institute for
Applied Systems Analysis
Schlossplatz 1
A-2361 Laxenburg, Austria

Tel: +43 2236 807 342
Fax: +43 2236 71313
E-mail: publications@iiasa.ac.at
Web: www.iiasa.ac.at

Interim Report

IR-05-018

What We Have Also Learned: Adaptive Speciation is Theoretically Plausible

Michael Doebeli (doebeli@zoology.ubc.ca)
Ulf Dieckmann (dieckmann@iiasa.ac.at)
Johan A.J. Metz (metz@rulsfb.leidenuniv.nl)
Diethard Tautz (tautz@uni-koeln.de)

Approved by

Leen Hordijk
Director, IIASA

March 2005



The Adaptive Dynamics Network at IIASA fosters the development of new mathematical and conceptual techniques for understanding the evolution of complex adaptive systems.

Focusing on these long-term implications of adaptive processes in systems of limited growth, the Adaptive Dynamics Network brings together scientists and institutions from around the world with IIASA acting as the central node.

Scientific progress within the network is collected in the IIASA Studies in Adaptive Dynamics series.

- No. 1 Metz JAJ, Geritz SAH, Meszéna G, Jacobs FJA, van Heerwaarden JS: *Adaptive Dynamics: A Geometrical Study of the Consequences of Nearly Faithful Reproduction*. IIASA Working Paper WP-95-099 (1995). van Strien SJ, Verduyn Lunel SM (eds): *Stochastic and Spatial Structures of Dynamical Systems*, Proceedings of the Royal Dutch Academy of Science (KNAW Verhandelingen), North Holland, Amsterdam, pp. 183-231 (1996).
- No. 2 Dieckmann U, Law R: *The Dynamical Theory of Coevolution: A Derivation from Stochastic Ecological Processes*. IIASA Working Paper WP-96-001 (1996). *Journal of Mathematical Biology* 34:579-612 (1996).
- No. 3 Dieckmann U, Marrow P, Law R: *Evolutionary Cycling of Predator-Prey Interactions: Population Dynamics and the Red Queen*. IIASA Preprint (1995). *Journal of Theoretical Biology* 176:91-102 (1995).
- No. 4 Marrow P, Dieckmann U, Law R: *Evolutionary Dynamics of Predator-Prey Systems: An Ecological Perspective*. IIASA Working Paper WP-96-002 (1996). *Journal of Mathematical Biology* 34:556-578 (1996).
- No. 5 Law R, Marrow P, Dieckmann U: *On Evolution under Asymmetric Competition*. IIASA Working Paper WP-96-003 (1996). *Evolutionary Ecology* 11:485-501 (1997).
- No. 6 Metz JAJ, Mylius SD, Dieckmann O: *When Does Evolution Optimize? On the Relation Between Types of Density Dependence and Evolutionarily Stable Life History Parameters*. IIASA Working Paper WP-96-004 (1996).
- No. 7 Ferrière R, Gatto M: *Lyapunov Exponents and the Mathematics of Invasion in Oscillatory or Chaotic Populations*. *Theoretical Population Biology* 48:126-171 (1995).
- No. 8 Ferrière R, Fox GA: *Chaos and Evolution*. IIASA Preprint (1996). *Trends in Ecology and Evolution* 10:480-485 (1995).
- No. 9 Ferrière R, Michod RE: *The Evolution of Cooperation in Spatially Heterogeneous Populations*. IIASA Working Paper WP-96-029 (1996). *The American Naturalist* 147:692-717 (1996).
- No. 10 van Dooren TJM, Metz JAJ: *Delayed Maturation in Temporally Structured Populations with Non-Equilibrium Dynamics*. IIASA Working Paper WP-96-070 (1996). *Journal of Evolutionary Biology* 11:41-62 (1998).
- No. 11 Geritz SAH, Metz JAJ, Kisdi É, Meszéna G: *The Dynamics of Adaptation and Evolutionary Branching*. IIASA Working Paper WP-96-077 (1996). *Physical Review Letters* 78:2024-2027 (1997).
- No. 12 Geritz SAH, Kisdi É, Meszéna G, Metz JAJ: *Evolutionary Singular Strategies and the Adaptive Growth and Branching of the Evolutionary Tree*. IIASA Working Paper WP-96-114 (1996). *Evolutionary Ecology* 12:35-57 (1998).
- No. 13 Heino M, Metz JAJ, Kaitala V: *Evolution of Mixed Maturation Strategies in Semelparous Life-Histories: The Crucial Role of Dimensionality of Feedback Environment*. IIASA Working Paper WP-96-126 (1996). *Philosophical Transactions of the Royal Society of London Series B* 352:1647-1655 (1997).
- No. 14 Dieckmann U: *Can Adaptive Dynamics Invade?* IIASA Working Paper WP-96-152 (1996). *Trends in Ecology and Evolution* 12:128-131 (1997).
- No. 15 Meszéna G, Czibula I, Geritz SAH: *Adaptive Dynamics in a 2-Patch Environment: A Simple Model for Allopatric and Parapatric Speciation*. IIASA Interim Report IR-97-001 (1997). *Journal of Biological Systems* 5:265-284 (1997).
- No. 16 Heino M, Metz JAJ, Kaitala V: *The Enigma of Frequency-Dependent Selection*. IIASA Interim Report IR-97-061 (1997). *Trends in Ecology and Evolution* 13:367-370 (1998).
- No. 17 Heino M: *Management of Evolving Fish Stocks*. IIASA Interim Report IR-97-062 (1997). *Canadian Journal of Fisheries and Aquatic Sciences* 55:1971-1982 (1998).
- No. 18 Heino M: *Evolution of Mixed Reproductive Strategies in Simple Life-History Models*. IIASA Interim Report IR-97-063 (1997).
- No. 19 Geritz SAH, van der Meijden E, Metz JAJ: *Evolutionary Dynamics of Seed Size and Seedling Competitive Ability*. IIASA Interim Report IR-97-071 (1997). *Theoretical Population Biology* 55:324-343 (1999).
- No. 20 Galis F, Metz JAJ: *Why Are There So Many Cichlid Species? On the Interplay of Speciation and Adaptive Radiation*. IIASA Interim Report IR-97-072 (1997). *Trends in Ecology and Evolution* 13:1-2 (1998).

- No. 21 Boerlijst MC, Nowak MA, Sigmund K: *Equal Pay for all Prisoners/ The Logic of Contrition*. IIASA Interim Report IR-97-073 (1997). American Mathematical Society Monthly 104:303-307 (1997). Journal of Theoretical Biology 185:281-293 (1997).
- No. 22 Law R, Dieckmann U: *Symbiosis Without Mutualism and the Merger of Lineages in Evolution*. IIASA Interim Report IR-97-074 (1997). Proceedings of the Royal Society of London Series B 265:1245-1253 (1998).
- No. 23 Klinkhamer PGL, de Jong TJ, Metz JAJ: *Sex and Size in Cosexual Plants*. IIASA Interim Report IR-97-078 (1997). Trends in Ecology and Evolution 12:260-265 (1997).
- No. 24 Fontana W, Schuster P: *Shaping Space: The Possible and the Attainable in RNA Genotype-Phenotype Mapping*. IIASA Interim Report IR-98-004 (1998). Journal of Theoretical Biology 194:491-515 (1998).
- No. 25 Kisdi É, Geritz SAH: *Adaptive Dynamics in Allele Space: Evolution of Genetic Polymorphism by Small Mutations in a Heterogeneous Environment*. IIASA Interim Report IR-98-038 (1998). Evolution 53:993-1008 (1999).
- No. 26 Fontana W, Schuster P: *Continuity in Evolution: On the Nature of Transitions*. IIASA Interim Report IR-98-039 (1998). Science 280:1451-1455 (1998).
- No. 27 Nowak MA, Sigmund K: *Evolution of Indirect Reciprocity by Image Scoring/ The Dynamics of Indirect Reciprocity*. IIASA Interim Report IR-98-040 (1998). Nature 393:573-577 (1998). Journal of Theoretical Biology 194:561-574 (1998).
- No. 28 Kisdi É: *Evolutionary Branching Under Asymmetric Competition*. IIASA Interim Report IR-98-045 (1998). Journal of Theoretical Biology 197:149-162 (1999).
- No. 29 Berger U: *Best Response Adaptation for Role Games*. IIASA Interim Report IR-98-086 (1998).
- No. 30 van Dooren TJM: *The Evolutionary Ecology of Dominance-Recessivity*. IIASA Interim Report IR-98-096 (1998). Journal of Theoretical Biology 198:519-532 (1999).
- No. 31 Dieckmann U, O'Hara B, Weisser W: *The Evolutionary Ecology of Dispersal*. IIASA Interim Report IR-98-108 (1998). Trends in Ecology and Evolution 14:88-90 (1999).
- No. 32 Sigmund K: *Complex Adaptive Systems and the Evolution of Reciprocation*. IIASA Interim Report IR-98-100 (1998). Ecosystems 1:444-448 (1998).
- No. 33 Posch M, Pichler A, Sigmund K: *The Efficiency of Adapting Aspiration Levels*. IIASA Interim Report IR-98-103 (1998). Proceedings of the Royal Society London Series B 266:1427-1435 (1999).
- No. 34 Mathias A, Kisdi É: *Evolutionary Branching and Coexistence of Germination Strategies*. IIASA Interim Report IR-99-014 (1999).
- No. 35 Dieckmann U, Doebeli M: *On the Origin of Species by Sympatric Speciation*. IIASA Interim Report IR-99-013 (1999). Nature 400:354-357 (1999).
- No. 36 Metz JAJ, Gyllenberg M: *How Should We Define Fitness in Structured Metapopulation Models? Including an Application to the Calculation of Evolutionarily Stable Dispersal Strategies*. IIASA Interim Report IR-99-019 (1999). Proceedings of the Royal Society of London Series B 268:499-508 (2001).
- No. 37 Gyllenberg M, Metz JAJ: *On Fitness in Structured Metapopulations*. IIASA Interim Report IR-99-037 (1999). Journal of Mathematical Biology 43:545-560 (2001).
- No. 38 Meszéna G, Metz JAJ: *Species Diversity and Population Regulation: The Importance of Environmental Feedback Dimensionality*. IIASA Interim Report IR-99-045 (1999).
- No. 39 Kisdi É, Geritz SAH: *Evolutionary Branching and Sympatric Speciation in Diploid Populations*. IIASA Interim Report IR-99-048 (1999).
- No. 40 Ylikarjula J, Heino M, Dieckmann U: *Ecology and Adaptation of Stunted Growth in Fish*. IIASA Interim Report IR-99-050 (1999). Evolutionary Ecology 13:433-453 (1999).
- No. 41 Nowak MA, Sigmund K: *Games on Grids*. IIASA Interim Report IR-99-038 (1999). Dieckmann U, Law R, Metz JAJ (eds): The Geometry of Ecological Interactions: Simplifying Spatial Complexity, Cambridge University Press, Cambridge, UK, pp. 135-150 (2000).
- No. 42 Ferrière R, Michod RE: *Wave Patterns in Spatial Games and the Evolution of Cooperation*. IIASA Interim Report IR-99-041 (1999). Dieckmann U, Law R, Metz JAJ (eds): The Geometry of Ecological Interactions: Simplifying Spatial Complexity, Cambridge University Press, Cambridge, UK, pp. 318-332 (2000).
- No. 43 Kisdi É, Jacobs FJA, Geritz SAH: *Red Queen Evolution by Cycles of Evolutionary Branching and Extinction*. IIASA Interim Report IR-00-030 (2000). Selection 2:161-176 (2001).
- No. 44 Meszéna G, Kisdi É, Dieckmann U, Geritz SAH, Metz JAJ: *Evolutionary Optimisation Models and Matrix Games in the Unified Perspective of Adaptive Dynamics*. IIASA Interim Report IR-00-039 (2000). Selection 2:193-210 (2001).
- No. 45 Parvinen K, Dieckmann U, Gyllenberg M, Metz JAJ: *Evolution of Dispersal in Metapopulations with Local Density Dependence and Demographic Stochasticity*. IIASA Interim Report IR-00-035 (2000). Journal of Evolutionary Biology 16:143-153 (2003).
- No. 46 Doebeli M, Dieckmann U: *Evolutionary Branching and Sympatric Speciation Caused by Different Types of Ecological Interactions*. IIASA Interim Report IR-00-040 (2000). The American Naturalist 156:S77-S101 (2000).
- No. 47 Heino M, Hanski I: *Evolution of Migration Rate in a Spatially Realistic Metapopulation Model*. IIASA Interim Report IR-00-044 (2000). The American Naturalist 157:495-511 (2001).
- No. 48 Gyllenberg M, Parvinen K, Dieckmann U: *Evolutionary Suicide and Evolution of Dispersal in Structured Metapopulations*. IIASA Interim Report IR-00-056 (2000). Journal of Mathematical Biology 45:79-105 (2002).
- No. 49 van Dooren TJM: *The Evolutionary Dynamics of Direct Phenotypic Overdominance: Emergence Possible, Loss Probable*. IIASA Interim Report IR-00-048 (2000). Evolution 54: 1899-1914 (2000).
- No. 50 Nowak MA, Page KM, Sigmund K: *Fairness Versus Reason in the Ultimatum Game*. IIASA Interim Report IR-00-57 (2000). Science 289:1773-1775 (2000).
- No. 51 de Feo O, Ferrière R: *Bifurcation Analysis of Population Invasion: On-Off Intermittency and Basin Riddling*. IIASA Interim Report IR-00-074 (2000). International Journal of Bifurcation and Chaos 10:443-452 (2000).

- No. 52 Heino M, Laaka-Lindberg S: *Clonal Dynamics and Evolution of Dormancy in the Leafy Hepatic Lophozia Silvicola*. IIASA Interim Report IR-01-018 (2001). *Oikos* 94:525-532 (2001).
- No. 53 Sigmund K, Hauert C, Nowak MA: *Reward and Punishment in Minigames*. IIASA Interim Report IR-01-031 (2001). *Proceedings of the National Academy of Sciences of the USA* 98:10757-10762 (2001).
- No. 54 Hauert C, De Monte S, Sigmund K, Hofbauer J: *Oscillations in Optional Public Good Games*. IIASA Interim Report IR-01-036 (2001).
- No. 55 Ferrière R, Le Galliard J: *Invasion Fitness and Adaptive Dynamics in Spatial Population Models*. IIASA Interim Report IR-01-043 (2001). Clobert J, Dhondt A, Danchin E, Nichols J (eds): *Dispersal*, Oxford University Press, pp. 57-79 (2001).
- No. 56 de Mazancourt C, Loreau M, Dieckmann U: *Can the Evolution of Plant Defense Lead to Plant-Herbivore Mutualism*. IIASA Interim Report IR-01-053 (2001). *The American Naturalist* 158: 109-123 (2001).
- No. 57 Claessen D, Dieckmann U: *Ontogenetic Niche Shifts and Evolutionary Branching in Size-Structured Populations*. IIASA Interim Report IR-01-056 (2001). *Evolutionary Ecology Research* 4:189-217 (2002).
- No. 58 Brandt H: *Correlation Analysis of Fitness Landscapes*. IIASA Interim Report IR-01-058 (2001).
- No. 59 Dieckmann U: *Adaptive Dynamics of Pathogen-Host Interactions*. IIASA Interim Report IR-02-007 (2002). Dieckmann U, Metz JAJ, Sabelis MW, Sigmund K (eds): *Adaptive Dynamics of Infectious Diseases: In Pursuit of Virulence Management*, Cambridge University Press, Cambridge, UK, pp. 39-59 (2002).
- No. 60 Nowak MA, Sigmund K: *Super- and Coinfection: The Two Extremes*. IIASA Interim Report IR-02-008 (2002). Dieckmann U, Metz JAJ, Sabelis MW, Sigmund K (eds): *Adaptive Dynamics of Infectious Diseases: In Pursuit of Virulence Management*, Cambridge University Press, Cambridge, UK, pp. 124-137 (2002).
- No. 61 Sabelis MW, Metz JAJ: *Perspectives for Virulence Management: Relating Theory to Experiment*. IIASA Interim Report IR-02-009 (2002). Dieckmann U, Metz JAJ, Sabelis MW, Sigmund K (eds): *Adaptive Dynamics of Infectious Diseases: In Pursuit of Virulence Management*, Cambridge University Press, Cambridge, UK, pp. 379-398 (2002).
- No. 62 Cheptou P, Dieckmann U: *The Evolution of Self-Fertilization in Density-Regulated Populations*. IIASA Interim Report IR-02-024 (2002). *Proceedings of the Royal Society of London Series B* 269:1177-1186 (2002).
- No. 63 Bürger R: *Additive Genetic Variation Under Intraspecific Competition and Stabilizing Selection: A Two-Locus Study*. IIASA Interim Report IR-02-013 (2002). *Theoretical Population Biology* 61:197-213 (2002).
- No. 64 Hauert C, De Monte S, Hofbauer J, Sigmund K: *Volunteering as Red Queen Mechanism for Co-operation in Public Goods Games*. IIASA Interim Report IR-02-041 (2002). *Science* 296:1129-1132 (2002).
- No. 65 Dercole F, Ferrière R, Rinaldi S: *Ecological Bistability and Evolutionary Reversals under Asymmetrical Competition*. IIASA Interim Report IR-02-053 (2002). *Evolution* 56:1081-1090 (2002).
- No. 66 Dercole F, Rinaldi S: *Evolution of Cannibalistic Traits: Scenarios Derived from Adaptive Dynamics*. IIASA Interim Report IR-02-054 (2002). *Theoretical Population Biology* 62:365-374 (2002).
- No. 67 Bürger R, Gimelfarb A: *Fluctuating Environments and the Role of Mutation in Maintaining Quantitative Genetic Variation*. IIASA Interim Report IR-02-058 (2002). *Genetical Research* 80:31-46 (2002).
- No. 68 Bürger R: *On a Genetic Model of Intraspecific Competition and Stabilizing Selection*. IIASA Interim Report IR-02-062 (2002). *Amer. Natur.* 160:661-682 (2002).
- No. 69 Doebeli M, Dieckmann U: *Speciation Along Environmental Gradients*. IIASA Interim Report IR-02-079 (2002). *Nature* 421:259-264 (2003).
- No. 70 Dercole F, Irisson J, Rinaldi S: *Bifurcation Analysis of a Prey-Predator Coevolution Model*. IIASA Interim Report IR-02-078 (2002). *SIAM Journal on Applied Mathematics* 63:1378-1391 (2003).
- No. 71 Le Galliard J, Ferrière R, Dieckmann U: *The Adaptive Dynamics of Altruism in Spatially Heterogeneous Populations*. IIASA Interim Report IR-03-006 (2003). *Evolution* 57:1-17 (2003).
- No. 72 Taborsky B, Dieckmann U, Heino M: *Unexpected Discontinuities in Life-History Evolution under Size-Dependent Mortality*. IIASA Interim Report IR-03-004 (2003). *Proceedings of the Royal Society of London Series B* 270:713-721 (2003).
- No. 73 Gardmark A, Dieckmann U, Lundberg P: *Life-History Evolution in Harvested Populations: The Role of Natural Predation*. IIASA Interim Report IR-03-008 (2003). *Evolutionary Ecology Research* 5:239-257 (2003).
- No. 74 Mizera F, Meszéna G: *Spatial Niche Packing, Character Displacement and Adaptive Speciation Along an Environmental Gradient*. IIASA Interim Report IR-03-062 (2003). *Evolutionary Ecology Research* 5: 363-382 (2003).
- No. 75 Dercole F: *Remarks on Branching-Extinction Evolutionary Cycles*. IIASA Interim Report IR-03-075 (2003). *Journal of Mathematical Biology* 47: 569-580 (2003).
- No. 76 Hofbauer J, Sigmund K: *Evolutionary Game Dynamics*. IIASA Interim Report IR-03-078 (2003). *Bulletin of the American Mathematical Society* 40: 479-519 (2003).
- No. 77 Ernande B, Dieckmann U, Heino M: *Adaptive Changes in Harvested Populations: Plasticity and Evolution of Age and Size at Maturation*. IIASA Interim Report IR-03-058 (2003). *Proceedings of the Royal Society of London Series B-Biological Sciences*, 271: 415-423 (2004).
- No. 78 Hanski I, Heino M: *Metapopulation-Level Adaptation of Insect Host Plant Preference and Extinction-Colonization Dynamics in Heterogeneous Landscapes*. IIASA Interim Report IR-03-028 (2003). *Theoretical Population Biology* 63:309-338 (2003).
- No. 79 van Doorn G, Dieckmann U, Weissing FJ: *Sympatric Speciation by Sexual Selection: A Critical Re-Evaluation*. IIASA Interim Report IR-04-003 (2004). *American Naturalist* 163: 709-725 (2004).
- No. 80 Egas M, Dieckmann U, Sabelis MW: *Evolution Restricts the Coexistence of Specialists and Generalists - the Role of Trade-off Structure*. IIASA Interim Report IR-04-004 (2004). *American Naturalist* 163: 518-531 (2004).

- No. 81 Ernande B, Dieckmann U: *The Evolution of Phenotypic Plasticity in Spatially Structured Environments: Implications of Intraspecific Competition, Plasticity Costs, and Environmental Characteristics*. IIASA Interim Report IR-04-006 (2004). *Journal of Evolutionary Biology* 17 (3): 613-628 (2004).
- No. 82 Cressman R, Hofbauer J: *Measure Dynamics on a One-Dimensional Continuous Trait Space: Theoretical Foundations for Adaptive Dynamics*. IIASA Interim Report IR-04-016 (2004).
- No. 83 Cressman R: *Dynamic Stability of the Replicator Equation with Continuous Strategy Space*. IIASA Interim Report IR-04-017 (2004).
- No. 84 Ravigné V, Olivieri I, Dieckmann U: *Implications of Habitat Choice for Protected Polymorphisms*. IIASA Interim Report IR-04-005 (2004). *Evolutionary Ecology Research* 6: 125-145 (2004).
- No. 85 Nowak MA, Sigmund K: *Evolutionary Dynamics of Biological Games*. IIASA Interim Report IR-04-013 (2004). *Science* 303: 793-799 (2004).
- No. 86 Vukics A, Asbóth J, Meszéna G: *Speciation in Multidimensional Evolutionary Space*. IIASA Interim Report IR-04-028 (2004). *Physical Review E* 68 4 (2003).
- No. 87 de Mazancourt C, Dieckmann U: *Trade-off Geometries and Frequency-dependent Selection*. IIASA Interim Report IR-04-039 (2004).
- No. 88 Cadet CR, Metz JAJ, Klinkhamer PGL: *Size and the Not-So-Single Sex: disentangling the effects of size on sex allocation*. IIASA Interim Report IR-04-084 (2004). *American Naturalist*, 164: 779-792 (2004).
- No. 89 Rueffler C, van Dooren TJM, Metz JAJ: *Adaptive Walks on Changing Landscapes: Levins' Approach Extended*. IIASA Interim Report IR-04-083 (2004). *Theoretical Population Biology*, 65: 165-178 (2004).
- No. 90 de Mazancourt C, Loreau M, Dieckmann U: *Understanding Mutualism When There is Adaptation to the Partner*. IIASA Interim Report IR-05-016 (2005).
- No. 91 Dieckmann U, Doebeli M: *Pluralism in Evolutionary Theory*. IIASA Interim Report IR-05-017 (2005).
- No. 92 Doebeli M, Dieckmann U, Metz JAJ, Tautz D: *What We Have Also Learned*. IIASA Interim Report IR-05-018 (2005).

Issues of the IIASA Studies in Adaptive Dynamics series can be obtained at www.iiasa.ac.at/Research/ADN/Series.html or by writing to adn@iiasa.ac.at.

What we have also learned:

Adaptive speciation is theoretically plausible

Michael Doebeli¹, Ulf Dieckmann², Johan A.J. Metz^{3,2}, and Diethard Tautz⁴

¹ Departments of Zoology and Mathematics, University of British Columbia, 6270

University Boulevard, Vancouver, BC V6T 1Z4, Canada

Email: doebeli@zoology.ubc.ca, phone: +1-604-822-3326

² Adaptive Dynamics Network, International Institute for Applied Systems Analysis

A-2361 Laxenburg, Austria

Email: dieckmann@iiasa.ac.at; phone: +43-2236-807-386

³ Institute of Biology, Leiden University, P.O. Box 9516, NL-2300 RA Leiden, The

Netherlands

Email: metz@rulsfb.leidenuniv.nl; phone: +31-71-5274937

⁴ Genetics Institute, University of Köln, Weyertal 121, D-50931 Köln, Germany

Email: tautz@uni-koeln.de; phone: +49-221-470-2465

In a recent article in *Evolution* entitled “Models of speciation: what have we learned in 40 years?” Gavrilets (2003) aimed at reviewing the insights that evolutionary biologists have gleaned from mathematical models of speciation over the past decades. Despite this nominal ambition, there have been important developments in speciation research that were barely touched on in Gavrilets’ review.

Our aim here is not to point out factual errors in Gavrilets’ article, but to highlight crucial omissions. In particular, we focus on the innovations brought about by research into adaptive speciation, which in our opinion have received unduly short shrift in Gavrilets’ article. In general, the past years have seen a systematic shift in speciation research from the traditional emphasis on geographical patterns of speciation to a broader perspective of stressing the mechanisms and processes of evolutionary diversification (e.g., Schluter 2000, Kondrashov 2001, Mallet 2001, Via 2001). These processes include adaptive speciation, in which the splitting of lineages is an adaptive response to disruptive selection driven by biological interactions. Based on the modeling effort of a whole group of scientists it has by now become clear that adaptive speciation is a plausible evolutionary process in many different evolutionary scenarios (e.g., Metz et al. 1996, Doebeli 1996, Meszéna et al. 1997, Geritz et al. 1998, Kisdi 1999, Dieckmann and Doebeli 1999, Higashi et al. 1999, Kondrashov and Kondrashov 1999, Kisdi and Geritz 1999, Drossel and McKane 2000, Geritz and Kisdi 2000, Doebeli and Dieckmann 2000, Law et al. 2001, Kaneko and Yomo 2002, Mizera and Meszéna 2003, Doebeli and Dieckmann 2003, Van Doorn et al. 2004).

Adaptive speciation requires ecological contact between the diverging lineages and is therefore often equated with sympatric speciation, even though disruptive selection

can also be a potent driver of speciation in parapatry. The possibility of adaptive speciation has been dismissed by Mayr (1963) and Dobzhansky (1970) as a plausible alternative to speciation through isolation by distance. The question of whether speciation under conditions of ecological contact, without isolation by distance, is a theoretically plausible evolutionary process hinges upon two key factors: first, on the ecological conditions under which frequency-dependent interactions are likely to generate disruptive selection, and second, on the evolution of assortative mating mechanisms in populations experiencing disruptive selection.

Investigating, by means of models, the ecological conditions under which sympatric speciation can occur has a long tradition that started with Maynard Smith (1966). The bulk of such models rely on rather simple genetic and ecological assumptions, typically involving two discrete ecological character states corresponding to two discrete ecological niches, and one or two loci determining mate choice (see Kawecki 2004 for a review). The model by Udovic (1980) that Gavrilets discusses in his article is an example of this class of models, as are most models for sympatric host-race formation (e.g., Diehl and Bush 1989, but see Fry 2003). It is known that the conditions for the maintenance of disruptive selection are restrictive in such models (Kawecki 2004). Together with the fact that the ecological assumptions in these models often appear to be rather special anyway, this has contributed to the perception that the origin and maintenance of diversity due to frequency-dependent selection requires special circumstances (e.g., Kassen 2002).

Extending earlier approaches toward more realistic (and, at the same time, more general) ecological settings by introducing the notion of competitive speciation,

Rosenzweig (1978) provided a conceptual framework for thinking about how frequency-dependent selection on quantitative characters determining the utilization of continuously distributed resources (or niches) could lead to adaptive diversification. However, this promising line of research was rarely taken up in subsequent mathematical models of speciation, with Seger (1985) and Doebeli (1996) representing two of the few exceptions. A second line of research was opened by Christiansen (1991), Brown & Pavlovic (1992), and Abrams et al. (1993), who showed how the evolution of quantitative characters driven by frequency-dependent ecological interactions can converge on points in phenotype space at which selection turns disruptive. However, at the time these seemingly disparate examples were not yet recognized as special cases of a general principle. Moreover, these studies did not actually address the problem of speciation, restricting attention to the emergence of disruptive selection.

Even if a population does experience persistent disruptive selection, adaptive speciation in sexual populations requires the evolution of assortative mating mechanisms. Assortative mating can either be directly based on the ecological trait that is under disruptive selection, or it can be based on ecologically neutral marker traits, a distinction that corresponds to the 1-allele and 2-allele models of Felsenstein (1981) (for reviews see Kirkpatrick and Ravigné 2002, and Dieckmann and Doebeli 2004). When assortative mating is based on marker traits, a linkage disequilibrium between these marker traits and the ecological trait must develop for assortativeness to be able to latch onto the ecological trait. It has long been thought that this requirement significantly hinders adaptive speciation (Felsenstein 1981).

However, in recent years substantial progress has been made with regard to both understanding the ecological conditions of adaptive diversification, and the evolution of assortative mating mechanisms. In particular, we think that fundamental advances have been made with regard to the first aspect: the mathematical theory of adaptive dynamics (Metz et al. 1992, 1996, Dieckmann and Law 1996, Geritz et al. 1997, 1998) has provided a general framework for studying the emergence of disruptive selection induced by ecological interactions, which is embodied by the concept of evolutionary branching (Metz et al. 1996, Geritz et al. 1998). Evolutionary branching points are phenotypes characterized by a set of general and simple mathematical conditions that determine when directional selection can lead to disruptive selection and, further, to the emergence of protected dimorphisms. These conditions can be applied to any particular ecological scenario that may underlie the adaptive evolution of quantitative traits. Therefore, the concept of evolutionary branching serves as a potentially unifying principle for identifying the ecological conditions of adaptive diversification. Based on this principle, a multitude of theoretical studies in different evolutionary contexts have provided analytical results about the conditions under which adaptive splitting is likely to occur; see Kisdi and Gyllenberg (2004) for an overview of pertinent studies. Thus, adaptive dynamics theory allows us to discover the potential richness of adaptive speciation processes: based on the analytical conditions required for evolutionary branching it has become clear that ecological conditions for adaptive diversification are, as far as we can tell from theoretical studies, truly ubiquitous (see also Doebeli and Dieckmann 2000).

With regard to the second aspect of adaptive speciation, i.e., the evolution of assortative mating and reproductive isolation, two studies by Kondrashov and

Kondrashov (1999) and by Dieckmann and Doebeli (1999) have shown that in models with more realistic genetic assumptions than were used earlier on, adaptive speciation is a plausible process, even when only allowing for indirect assortative mating based on ecologically neutral marker traits. Thus, the conditions for the evolution of assortative mating under frequency-dependent disruptive selection are clearly less restrictive than earlier, simpler models had suggested. It also turns out that this conclusion is robust with regard to various changes in model assumptions, including costs of assortative mating (Bolnick 2004, Doebeli and Dieckmann 2004, Doebeli 2004).

The flurry of recent theoretical papers on the possibility of adaptive speciation reflects the fact that evolutionary biologists from all walks have started to realize that frequency-dependent selection can induce adaptive diversification, and that requirements for such processes are less restrictive than past dogma had us believe (e.g., Via 2001, Turelli et al. 2001). In his review, Gavrilets questions the value of the corresponding “dozens of new modeling papers” by suggesting that it is obvious that “selection promotes speciation”. However, only a short while ago the possibility of adaptive speciation seemed far from obvious to many evolutionary biologists. Traditionally, only two mechanisms were considered through which selection could facilitate speciation: first, local adaptation in geographically segregated populations might accelerate the build-up of reproductive incompatibilities due to pre- or postzygotic isolation mechanisms, and second, reinforcement upon secondary contact might enhance an already existing level of reproductive isolation (see, e.g., the review by Turelli et al. 2001). Overcoming this unnecessarily narrow perspective on the interplay between adaptation and speciation required exactly the flurry of papers that Gavrilets bemoans,

and the extent of this research activity is simply a consequence of speciation research being freed, after decades, from what one might portray as the ‘shackles of allopatry.’

In our opinion, the big news in recent speciation research is that many different ecological selection scenarios can easily give rise to selection pressures under which adaptive speciation is likely to occur. Understanding how these selective scenarios emerge from biological interactions is as important as understanding how the ensuing split into reproductively isolated subunits unfolds genetically. Ignoring this by focusing solely on traditional population genetic approaches and on traditional geographical classifications does not do justice to the exciting and dynamic state of the field. In particular, resurrecting an ecological perspective on speciation holds many promises for tying in speciation research with modern empirical and experimental approaches (e.g., Schluter 1994, Rainey and Travisano 1998, Schlieven et al. 2001, Friesen et al. 2004).

Contrary to what Gavrilets alleges in his review, the recent modeling efforts have indeed led to many analytical results. In fact, adaptive dynamics theory is exactly the kind of framework that yields analytical results similar to those presented by Gavrilets (2003) in his last example, and it is able to deliver such insights in a vast variety of different ecological and behavioral settings (see, e.g., Metz et al. 1996, Geritz et al. 1998, Kisdi and Geritz 1999, Doebeli and Dieckmann 2000). This is what one would want from a general theory. The bulk of analytical results obtained so far concern the ecological conditions for adaptive diversification, i.e., the existence or not of evolutionary branching points. Analytical results concerning the evolution of assortative mating mechanisms in multi-locus models for sexual populations are generally hard to come by. However, it should be pointed out that extensive numerical simulations can also yield complete

classifications of system behavior (e.g., Doebeli and Dieckmann 2003), and can lead to statements that are just as universal as those derived by purely analytical means. Overall, given the many analytical results about conditions for diversification obtained using adaptive dynamics theory, we cannot agree with Gavrilets' assessment that "What is missing in the theoretical speciation research are general and transparent analytical results comparable to those in other areas of theoretical population genetics and ecology." Besides the fact that we now have a comprehensive mathematical framework that explains why adaptive diversification should be a ubiquitous and robust process, general analytical results have been obtained both for ecological and for sexual selection (e.g., Doebeli and Dieckmann 2000, Van Doorn et al. 2004).

Many empiricists have welcomed these new theoretical developments. Results from adaptive dynamics theory have shed new light on existing empirical work (e.g., Schluter 1994, Schliewen et al. 1994, Rosenzweig et al. 1994 (see Doebeli 2002), Johanneson et al. 1995, Skúlason et al. 1995, Treves et al. 1998, Rainey and Travisano 1998, Schliewen et al. 2001, Jiggins and Mallet 2001, Jones et al. 2003), and have inspired new empirical work that tests the theoretical predictions, both by analyzing existing data (e.g., Bolnick et al. 2003) and by using evolutionary experiments (e.g., Bolnick 2001, Friesen et al. 2004).

In addition, having available a general theoretical framework allows us to compare adaptive speciation with other processes of evolutionary diversification, such as the evolution of sexual dimorphism (Bolnick and Doebeli 2003, Van Doorn et al., submitted) or the evolution of ecological niche widths (Egas et al., submitted, Ackermann and Doebeli, submitted). With time, these efforts are likely to yield a fairly complete

picture about the likelihood of adaptive speciation in various scenarios of ecological and sexual selection. In the end, theoretical advances have to be brought to fruition by modifying general theory so that it yields models that are applicable to particular situations. A multitude of models is needed to reflect the complexity of speciation, and specificity is not a problem if the models can be understood within a common conceptual framework. Excising such healthy pluralism from speciation research would seem unwise.

In this note, we did not endeavor to reflect all of theoretical speciation research. Instead we took a necessarily biased view by concentrating on the innovations brought about by research into adaptive speciation. It has become clear that the traditional geographical classification of speciation modes is no longer appropriate to capture the essential complexity of many speciation processes (e.g., Mizera and Meszéna 2003, Doebeli and Dieckmann 2003). By emphasizing adaptive *processes* rather than restricting attention to biogeographical *patterns* of diversification, theoretical and experimental speciation research have taken off again to new shores. These exciting developments were ignored in the review by Gavrilets. There are other omissions in Gavrilets' article, most notably perhaps the body of theory pertaining to the problem of reinforcement (e.g., Liou and Price 1994, Kirkpatrick and Servedio 1999, Servedio 2000). Reinforcement is of fundamental importance for many processes of ecological speciation as defined by Schluter (Schluter 2000; see also the Introduction in Dieckmann et al. 2004), and is related to the problem of the evolution of assortative mating mechanisms in processes of adaptive speciation. This further illustrates that Gavrilets' adherence to an old geographical classification of speciation that is fraught with problems, and the omission

of whole bodies of work that have reinvigorated speciation research in the last decade, led to an unproductive bias on the representation of the field. While there is nothing wrong with Gavrilets reviewing his own contributions to speciation theory, which are substantial, we feel that a broader representation of an exciting and reinvigorated field would have been appropriate.

References:

- Abrams, P. A., Matsuda, H. and Harada, Y. 1993. Evolutionarily unstable fitness maxima and stable fitness minima of continuous traits. *Evolutionary Ecology* **7**: 465-487.
- Ackermann, M. and Doebeli, M. 2004. Evolution of niche width and adaptive diversification. (submitted)
- Bolnick, D.I. 2001. Intraspecific competition favours niche width expansion in *Drosophila melanogaster*. *Nature* **410**: 463-466.
- Bolnick, D.I. 2004. Waiting for sympatric speciation. *Evolution* **58**: 895-899.
- Bolnick, D.I., and Doebeli, M. 2003. Sexual dimorphism and adaptive speciation: Two sides of the same ecological coin. *Evolution* **57**: 2433-2449.
- Bolnick, D.I. Svanback, R., Fordyce, J.A., Yang, L.H., Davis, J.M., Hulsey, C.D. and Forister, M.L. 2003. The ecology of individuals: Incidence and implications of individual specialization. *The American Naturalist* **161**: 1-28.
- Brown, J.S. and Pavlovic, N.B. 1992. Evolution in heterogeneous environments – effects of migration on habitat specialization. *Evolutionary Ecology* **6**: 360-382.
- Bush, G. L. 1994. Sympatric speciation in animals: New wine in old bottles. *Trends in Ecology and Evolution* **9**: 285-288.
- Christiansen, F. B. 1991. On conditions for evolutionary stability for a continuously varying trait. *Theoretical Population Biology* **7**: 13-38.
- Dieckmann, U. and Doebeli, M. 1999. On the origin of species by sympatric speciation. *Nature* **400**: 354-357.
- Dieckmann, U. and Doebeli, M. 2004. Adaptive dynamics of speciation: Sexual populations. pp. 76-111 in U. Dieckmann, M. Doebeli, J.A.J. Metz and D. Tautz, eds. *Adaptive Speciation*. Cambridge University Press, Cambridge, UK.
- Dieckmann, U. and Law, R. 1996. The dynamical theory of coevolution: A derivation from stochastic ecological processes. *Journal of Mathematical Biology* **43**:1308-1311.
- Dieckmann, U., Doebeli, M., Metz, J.A.J. and Tautz, D. (eds.) 2004. *Adaptive Speciation*. Cambridge University Press, Cambridge, UK.
- Diehl, S. R., and Bush, G. L. 1989. The role of habitat preference in adaptation and speciation. pp. 345-365 in D. Otte and J. Endler, eds. *Speciation and its consequences*. Sinauer Associates, Sunderland, MA.

- Dobzhansky, T., 1970 *Genetics of the Evolutionary Process*. Columbia University Press, New York.
- Doebeli, M. 1996. A quantitative genetic competition model for sympatric speciation. *Journal of Evolutionary Biology* **9**: 893-909.
- Doebeli, M. 2002. A model for the evolutionary dynamics of cross-feeding polymorphisms in microorganisms. *Population Ecology* **44**: 59-70.
- Doebeli, M. 2004. Adaptive speciation when assortative mating is based on female preference for male marker traits. (In preparation)
- Doebeli, M. and Dieckmann, U. 2000. Evolutionary branching and sympatric speciation caused by different types of ecological interactions. *The American Naturalist* **156**: S77-S101.
- Doebeli, M. and Dieckmann, U. 2003. Speciation along environmental gradients *Nature* **421**: 259-264.
- Doebeli, M. and Dieckmann, U. 2004. Adaptive dynamics as a mathematical tool for studying the ecology of speciation processes. *Journal of Evolutionary Biology* (submitted)
- Drossel, B. and McKane, A. 2000. Competitive speciation in quantitative genetic models. *Journal of Theoretical Biology* **204**: 467-478.
- Egas, M., Sabelis, M. W., and Dieckmann, U. Evolution of specialization and ecological character displacement of herbivores along a gradient of plant quality. (submitted)
- Eshel, I. 1983. Evolutionary and continuous stability. *Journal of Theoretical Biology* **103**: 99-111.
- Felsenstein, J. 1981. Skepticism towards Santa Rosalia, or why are there so few kinds of animals? *Evolution* **35**: 124-238.
- Friesen, M., Saxer, G., Travisano, M. and Doebeli, M. 2004. Experimental evidence for sympatric ecological diversification due to frequency-dependent competition in *Escherichia coli*. *Evolution* **58**: 245-260.
- Fry, J. D. 2003. Multilocus models of sympatric speciation: Bush versus Rice versus Felsenstein. *Evolution* **57**, 1735-1746.
- Gavrilets, S. 2003. Models of speciation: What have we learned in 40 years? *Evolution* **57**: 2197-2215.

- Geritz, S.A.H., Metz, J.A.J., Kisdi, É. and Meszéna, G. 1997. Dynamics of adaptation and evolutionary branching. *Physical Review Letters* **78**: 2024-2027.
- Geritz, S.A.H. and Kisdi, É. 2000. Adaptive dynamics in diploid sexual populations and the evolution of reproductive isolation. *Proceedings of the Royal Society of London B* **267**: 1671-1678.
- Geritz, S.A.H., Kisdi, É., Meszéna, G. and Metz, J.A.J. 1998. Evolutionarily singular strategies and the adaptive growth and branching of the evolutionary tree. *Evolutionary Ecology Research* **12**: 35-57.
- Higashi, M., Takimoto, G. and Yamamura, N. 1999. Sympatric speciation by sexual selection. *Nature* **402**: 523-526.
- Jiggins, C.D., Naisbit, R.E., Coe, R.L. and Mallet, J. 2001. Reproductive isolation caused by colour pattern mimicry. *Nature* **411**: 302-305.
- Johannesson, K., Rolán-Alvarez, E. and Ekendahl, A. 1995. Incipient reproductive isolation between two sympatric morphs of the intertidal snail *Littorina saxatilis*. *Evolution* **49**: 1180-1190.
- Jones, A.G., Moore, G.I., Kvarnemo, C., Walker, D. and Avise, J.C. 2003. Sympatric speciation as a consequence of male pregnancy in seahorses. *Proceedings of the National Academy of Sciences USA* **100**: 6598-6603.
- Kaneko, K. and Yomo, T. 2002. Symbiotic sympatric speciation through interaction-driven phenotype differentiation. *Evolutionary Ecology Research* **4**: 317-350.
- Kawecki, T. J. 2004. Genetic theories of sympatric speciation. pp. 36-53 in U. Dieckmann, M. Doebeli, J.A.J. Metz and D. Tautz, eds. *Adaptive Speciation*. Cambridge University Press, Cambridge, UK.
- Kirkpatrick, M. and Ravigné, V. 2002 Speciation by natural and sexual selection: models and experiments. *American Naturalist* **159**: S22-S35.
- Kisdi, É. 1999. Evolutionary branching under asymmetric competition. *Journal of Theoretical Biology* **197**: 149-162.
- Kisdi, É. and Geritz, S.A.H. 1999. Adaptive dynamics in allele space: Evolution of genetic polymorphism by small mutations in a heterogeneous environment. *Evolution* **53**: 993-1008.
- Kisdi, É. and Gyllenberg, M. 2004. Adaptive dynamics and the paradigm of diversity. *Journal of Evolutionary Biology* (submitted)

- Kondrashov, A.S. 1986. Multilocus model of sympatric speciation. III. Computer simulations. *Theoretical Population Biology* **29**: 1-15.
- Kondrashov, A.S. 2001. Speciation: Darwin revisited. *Trends in Ecology and Evolution* **16**: 412.
- Kondrashov, A.S. and Kondrashov, F.A. 1999. Interactions among quantitative traits in the course of sympatric speciation. *Nature* **400**: 351-354.
- Law, R., Bronstein, J.L. and Ferrière, R. 2001. On mutualists and exploiters: Plant-insect coevolution in pollinating seed-parasite systems. *Journal of Theoretical Biology* **212**: 373-389.
- Liou, L.W. and Price, T.D. 1994. Speciation by reinforcement of premating isolation. *Evolution* **48**: 1451-1459
- Mallet, J. 2001. The speciation revolution. *Journal of Evolutionary Biology* **14**: 887-888.
- Maynard Smith, J. 1996. Sympatric speciation. *The American Naturalist* **100**: 637-650.
- Mayr, E. 1963. *Animal Species and Evolution*. Cambridge, MA, USA: Harvard University Press.
- Meszéna, G., Czibula, I. and Geritz, S.A.H. 1997. Adaptive dynamics in a 2-patch environment: A toy model for allopatric and parapatric speciation. *Journal of Biological Systems* **5**: 265-284.
- Metz, J.A.J., Geritz, S.A.H., Meszéna, G., Jacobs, F.J.A. and van Heerwaarden, J.S. 1996. Adaptive dynamics: A geometrical study of the consequences of nearly faithful reproduction. pp. 183-231 in van Strien, S.J. and Verduyn Lunel, S.M., eds. *Stochastic and Spatial Structures of Dynamical Systems, Proceedings of the Royal Dutch Academy of Science (KNAW Verhandelingen)*. Dordrecht, Netherlands: North Holland.
- Mizera, F. and Meszéna, G. (2003). Spatial niche packing, character displacement and adaptive speciation along an environmental gradient. *Evolutionary Ecology Research* **5**: 363-382.
- Rainey, P.B. and Travisano, M. 1998. Adaptive radiation in a heterogeneous environment. *Nature* **394**: 69-72.
- Rosenzweig, M.L. 1978. Competitive speciation. *Biological Journal of the Linnean Society* **10**: 275-289.
- Rosenzweig, R.F., Sharp, R.R., Treves, D.S. and Adams, J. 1994. Microbial evolution in a simple unstructured environment: Genetic differentiation in *Escherichia coli*. *Genetics* **137**: 903-917.

- Schluter, D. 1994. Experimental evidence that competition promotes divergence in adaptive radiation. *Science* **266**: 798-800.
- Schluter, D. 2000. *The Ecology of Adaptive Radiation*. Oxford University Press, Oxford.
- Schliewen, U.K., Tautz, D. and Pääbo, S. 1994. Sympatric speciation suggested by monophyly of crater lake cichlids. *Nature* **368**: 629-632.
- Schliewen, U., Rassmann, K., Markmann, M., Markert, J., Kocher, T. and Tautz, D. 2001. Genetic and ecological divergence of a monophyletic cichlid species pair under fully sympatric conditions in Lake Ejagham, Cameroon. *Molecular Ecology* **10**: 1471-1488.
- Seger, J. 1985. Intraspecific resource competition as a cause of sympatric speciation. In *Evolution: Essays in Honour of John Maynard Smith*, eds. Greenwood, P.J., Harvey, P.H. and Slatkin, M., pp. 43-53, Cambridge University Press, Cambridge.
- Kirkpatrick, M. and Servedio, M.R. 1999. The reinforcement of mating preferences on an island. *Genetics* **151**: 865-884
- Servedio, M.R. 2000. Reinforcement and the genetics of nonrandom mating. *Evolution* **54**: 21-29.
- Skúlason, S., Snorrason, S.S. and Jonsson, B. 1999. Sympatric morphs, populations and speciation in freshwater fish with emphasis on arctic charr. pp. 70-92 in Magurran, A.E. and May, R.M., eds. *Evolution of Biological Diversity*. Oxford, UK: Oxford University Press.
- Taylor, P. Evolutionary stability in one-parameter models under weak selection. *Theoretical Population Biology* **36**: 125-143.
- Travisano, M. and Rainey, P.B. 2000. Studies of adaptive radiation using model microbial systems. *The American Naturalist* **156**: S35-S44.
- Treves, D.S., Manning, S. and Adams, J. 1998. Repeated evolution of an acetate-crossfeeding polymorphism in long-term populations of *Escherichia coli*. *Molecular Biology and Evolution* **15**: 789-797.
- Turelli, M., Barton, N.H. and Coyne, J.A. 2001. Theory and speciation. *Trends in Ecology and Evolution* **16**: 330-343.
- Van Dooren, T. J. M., Durinx, M. and Demon, I. Sexual dimorphism or evolutionary branching? (submitted)

Van Doorn, G.S., Dieckmann, U. and Weissing, F.J. 2004. Sympatric speciation by sexual selection: A critical re-evaluation. *The American Naturalist*, in press.

Van Doorn, G.S., Luttikhuisen, P.C. and Weissing, F.J. 2001. Sexual selection at the protein level drives the extraordinary divergence of sex related genes during sympatric speciation. *Proceedings of the Royal Society of London B* **268**: 2155-2161.

Via, S. 2001. Sympatric speciation in animals: the ugly duckling grows up. *Trends in Ecology and Evolution* 16:381-390.