International Institute for Applied Systems Analysis Www.iiasa.ac.at

Fuzzy Random Noncooperative Two-level Linear Programming through Absolute Deviation Minimization Using Possibility and Necessity

HAN

HE III

HETTER

QH1

Sakawa, M. and Kato, K.

IIASA Interim Report May 2009 Sakawa, M. and Kato, K. (2009) Fuzzy Random Noncooperative Two-level Linear Programming through Absolute Deviation Minimization Using Possibility and Necessity. IIASA Interim Report. IR-09-021 Copyright © 2009 by the author(s). http://pure.iiasa.ac.at/9128/

Interim Report on work of the International Institute for Applied Systems Analysis receive only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute, its National Member Organizations, or other organizations supporting the work. All rights reserved. Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage. All copies must bear this notice and the full citation on the first page. For other purposes, to republish, to post on servers or to redistribute to lists, permission must be sought by contacting repository@iiasa.ac.at



Interim Report IR-09-021

Fuzzy Random Noncooperative Two-level Linear Programming through Absolute Deviation Minimization Using Possibility and **Necessity**

Masatoshi Sakawa (sakawa@hiroshima-u.ac.jp) Kosuke Kato(kosuke-kato@hiroshima-u.ac.jp)

Approved by

Marek Makowski (marek@iiasa.ac.at) Leader, Integrated Modeling Environment Project

May 2009

Interim Reports on work of the International Institute for Applied Systems Analysis receive only limited review. Views or opinions expressed herein do not necessarily represent those of the Institute, its National Member Organizations, or other organizations supporting the work.

Foreword

In this paper, assuming noncooperative behavior of the decision makers, we consider solution methods for decision making problems in hierarchical organizations under fuzzy random environments. Taking into account vagueness of judgments of decision makers, fuzzy goals are introduced into the formulated fuzzy random noncooperative two-level linear programming problems. Considering the possibility and necessity measure that each objective function fulfills the corresponding fuzzy goal, we transform the fuzzy random two-level linear programming problems to minimize each objective function with fuzzy random variables into stochastic two-level programming problems to maximize the degree of possibility and necessity that each fuzzy goal is fulfilled. Through the use of absolute deviation minimization in stochastic programming, the transformed stochastic two-level programming problems can be reduced to deterministic two-level programming problems. It should be emphasized here that the absolute deviation minimization model is suitable for risk-averse decision makers and it is more tractable than the variance minimization model. For the transformed problems, extended concepts of Stackelberg solutions are introduced and computational methods are also presented. It is significant to note that the extended Stackelberg solutions can be obtained through the combined use of the variable transformation method and the Kth best algorithm for two-level linear programming problems. A numerical example is provided to illustrate the proposed methods.

Abstract

This paper considers fuzzy random two-level linear programming problems under noncooperative behavior of the decision makers. Having introduced fuzzy goals of decision makers together with the possibility and necessity measure, following absolute deviation minimization, fuzzy random two-level programming problems are transformed into deterministic ones. Extended Stackelberg solutions are introduced and computational methods are also presented.

Keywords: Two-level linear programming problem; fuzzy random variables; Stackelberg solutions; possibility; necessity; absolute deviation minimization.

Acknowledgments

Masatoshi Sakawa appreciates the hospitality and the working environment during his two-months Guest Scholar affiliation with the Integrated Modeling Project. The research presented in this paper was completed and the paper written during this time.

About the Authors

Masatoshi Sakawa joined the Integrated Modeling Environment in April 2009. His research and teaching activities are in the area of systems engineering, especially mathematical optimization, multiobjective decision making, fuzzy mathematical programming and game theory. In addition to over 300 articles in national and international journals, he is an author and coauthor of 5 books in English and 14 books in Japanese. At present Dr. Sakawa is a Professor at Hiroshima University, Japan and is working with the Department of Artificial Complex Systems Engineering. Dr. Sakawa received BEng, MEng, and DEng degrees in applied mathematics and physics at Kyoto University, in 1970, 1972, and 1975 respectively. From 1975 he was with Kobe University, where from 1981 he was an Associate Professor in the Department of Systems Engineering. From 1987 to 1990 he was Professor of the Department of Computer Science at Iwate University and from March to December 1991 he was an Honorary Visiting Professor at the University of Manchester Institute of Science and Technology (UMIST), Computation Department, sponsored by the Japan Society for the Promotion of Science (JSPS). He was also a Visiting Professor of the Institute of Economic Research, Kyoto University from April 1991 to March 1992. In 2002 Dr. Sakawa received the Georg Cantor Award of the International Society on Multiple Criteria Decision Making.

Kosuke Kato is an Associate Professor at Department of Artificial Complex Systems Engineering, Hiroshima University, Japan. He received B.E. and M.E. degrees in biophysical engineering from Osaka University, in 1991 and 1993, respectively. He received D.E. degree from Kyoto University in 1999. His current research interests are evolutionary computation, large-scale programming and multiobjective/multi-level programming under uncertain environments.

Contents

1	Intr	oduction	1			
2	Fuzzy random two-level linear programming problems					
	2.1	Fuzzy goals	4			
	2.2	Possibility and necessity	5			
3	Stackelberg solutions through absolute deviation minimization					
	3.1	Possibility case	7			
	3.2	Necessity case	13			
4	Nun	nerical example	16			
5	Con	clusions	18			

Fuzzy Random Noncooperative Two-level Linear Programming through Absolute Deviation Minimization Using Possibility and Necessity

Masatoshi Sakawa (sakawa@hiroshima-u.ac.jp) * ** Kosuke Kato(kosuke-kato@hiroshima-u.ac.jp)*

1 Introduction

In the real world, we often encounter situations where there are two or more decision makers in an organization with a hierarchical structure, and they make decisions in turn or at the same time so as to optimize their objective functions. Decision making problems in decentralized organizations are often modeled as Stackelberg games [46], and they are formulated as two-level mathematical programming problems [45, 44]. In the context of two-level programming, the decision maker at the upper level first specifies a strategy, and then the decision maker at the lower level specifies a strategy so as to optimize the objective with full knowledge of the action of the decision maker at the upper level. In conventional multi-level mathematical programming models employing the solution concept of Stackelberg equilibrium, it is assumed that there is no communication among decision makers, or they do not make any binding agreement even if there exists such communication. Computational methods for obtaining Stackelberg solutions to two-level linear programming problems are classified roughly into three categories: the vertex enumeration approach [5], the Kuhn-Tucker approach [3, 4, 5, 15], and the penalty function approach [51]. The subsequent works on two-level programming problems under noncooperative behavior of the decision makers have been appearing [34, 35, 14, 36, 9, 11] including some applications to aluminum production process [33], pollution control policy determination [2], tax credits determination for biofuel producers [10], pricing in competitive electricity markets [12], supply chain planning [39] and so forth.

However, to utilize two-level programming for resolution of conflict in decision making problems in real-world decentralized organizations, it is important to realize that simultaneous considerations of both fuzziness [41, 42, 43] and randomness [48, 6, 47] would be required. Fuzzy random variables, first introduced by Kwakernaak [25], have been developing [24, 37, 29], and an overview of the developments of fuzzy random variables was found in [13]. Studies on linear programming problems with fuzzy random variable coefficients, called fuzzy random linear programming problems, were initiated by Wang and Qiao [50], Qaio, Zhang and Wang [38] as seeking the probability distribution of the optimal solution and optimal value. Optimization models for fuzzy random

^{*}Graduate School of Engineering, Hiroshima University.

^{**}Corresponding author.

linear programming problems were first considered by Luhandjula et al. [30, 32] and further developed by Liu [27, 28] and Rommelfanger [40]. A brief survey of major fuzzy stochastic programming models was found in the paper by Luhandjula [31]. As we look at recent developments in the fields of fuzzy random programming, we can see continuing advances [16, 18, 20, 17, 19, 22, 40, 21, 1, 52].

Under these circumstances, in this paper, assuming noncooperative behavior of the decision makers, we consider solution methods for decision making problems in hierarchical organizations under fuzzy random environments. Taking into account vagueness of judgments of decision makers, fuzzy goals are introduced into the formulated noncooperative two-level linear programming problems involving fuzzy random variables. Considering the possibility and necessity measure that each objective function fulfills the corresponding fuzzy goal, we transform the fuzzy random two-level linear programming problems to minimize each objective function with fuzzy random variables into stochastic two-level programming problems to maximize the degree of possibility and necessity that each fuzzy goal is fulfilled. Through the use of absolute deviation minimization [23] in stochastic programming, the transformed stochastic two-level programming problems can be reduced to deterministic two-level programming problems. It is significant to note that the absolute deviation minimization model is suitable for risk-averse decision makers and it is more tractable than the variance minimization model. For the transformed problems, extended concepts of Stackelberg solutions are introduced and computational methods are also presented. It is shown that extended Stackelberg solutions can be obtained through the combined use of the variable transformation method by Charnes et al. [8] and the Kth best algorithm by Bialas et al. [5].

2 Fuzzy random two-level linear programming problems

Fuzzy random variables, first introduced by Kwakernaak [25], have been defined in various ways [25, 37, 24, 29]. For example, as a special case of fuzzy random variables given by Kwakernaak, Kruse and Meyer [24] defined a fuzzy random variable as follows.

Definition 1 (Fuzzy random variable) Let (Ω, B, P) be a probability space, $F(\mathcal{R})$ the set of fuzzy numbers with compact supports and X a measurable mapping $\Omega \to F(\mathcal{R})$. Then X is a fuzzy random variable if and only if given $\omega \in \Omega$, $X_{\alpha}(\omega)$ is a random interval for any $\alpha \in (0, 1]$, where $X_{\alpha}(\omega)$ is an α -level set of the fuzzy set $X(\omega)$.

Although there exist some minor differences in several definitions of fuzzy random variables, fuzzy random variables could be roughly understood to be a random variable whose observed values are fuzzy sets.

In this paper, we deal with fuzzy random noncooperative two-level linear programming problems formulated as:

$$\begin{array}{l} \underset{\text{DM1}}{\text{minimize}} \quad z_1(\boldsymbol{x}_1, \boldsymbol{x}_2) = \tilde{\boldsymbol{C}}_{11} \boldsymbol{x}_1 + \tilde{\boldsymbol{C}}_{12} \boldsymbol{x}_2 \\ \underset{\text{DM2}}{\text{minimize}} \quad z_2(\boldsymbol{x}_1, \boldsymbol{x}_2) = \tilde{\boldsymbol{C}}_{21} \boldsymbol{x}_1 + \tilde{\boldsymbol{C}}_{22} \boldsymbol{x}_2 \\ \text{subject to} \quad A_1 \boldsymbol{x}_1 + A_2 \boldsymbol{x}_2 \leq \boldsymbol{b} \\ \quad \boldsymbol{x}_1 \geq \boldsymbol{0} , \ \boldsymbol{x}_2 \geq \boldsymbol{0} \end{array} \right\}.$$

$$(1)$$

It is significant to note here that randomness and fuzziness of the coefficients are denoted by the "dash above" and "wave above" i.e., "-" and "~", respectively. In this formulation, \boldsymbol{x}_1 is an n_1 dimensional decision variable column vector for the decision maker at the upper level (DM1), \boldsymbol{x}_2 is an n_2 dimensional decision variable column vector for the decision for the decision maker at the lower level (DM2), $z_1(\boldsymbol{x}_1, \boldsymbol{x}_2)$ is the objective function for DM1 and $z_2(\boldsymbol{x}_1, \boldsymbol{x}_2)$ is the objective function for DM2. Elements \tilde{C}_{ljk} , $k = 1, 2, \ldots, n_j$ of coefficient vectors \tilde{C}_{lj} , l = 1, 2, j = 1, 2 are fuzzy random variables characterized by the membership function:

$$\mu_{\tilde{C}_{ljk}}(\tau) = \begin{cases} \max\left\{0, 1 - \frac{\bar{d}_{ljk} - \tau}{\beta_{ljk}}\right\} &, \text{ if } \tau \leq \bar{d}_{ljk} \\\\ \max\left\{0, 1 - \frac{\tau - \bar{d}_{ljk}}{\gamma_{ljk}}\right\} &, \text{ otherwise,} \end{cases}$$

where \bar{d}_{ljk} is a random variable that takes an observed value d_{ljks_l} under a scenario $s_l \in \{1, 2, \ldots, S_l\}$ whose probability is p_{ls_l} , and parameters β_{ljk} and γ_{ljk} , representing left and right spreads of $\mu_{\tilde{C}_{ljk}}(\cdot)$, are positive numbers. This definition of fuzzy random variables was first appeared in the literature by Katagiri et al. [20]. Figure 1 illustrates an example of the membership function of a fuzzy random variable \tilde{C}_{ljk} .



Figure 1: An example of a membership function of a fuzzy random variable.

Fuzzy random two-level linear programming problems formulated as (1) are often seen in actual decision making situations. For example, consider a supply chain planning [39] where the distribution center (DM1) and the production part (DM2) hope to minimize the distribution cost and the production cost respectively. Since coefficients of these objective functions are often affected by the economic conditions varying at random, they can be regarded as random variables. In addition, since observed values of them are often ambiguous and estimated by fuzzy numbers, they are expressed by fuzzy random variables. Hence, the supply chain planning problem can be formulated as a two-level linear programming problem involving fuzzy random variable coefficients.

Observing that each coefficient \bar{C}_{ljk} is a fuzzy random variable defined as a random variable whose observed values are L-R fuzzy numbers, each objective function $\tilde{C}_l x = \tilde{C}_{l1} x_1 + \tilde{C}_{l2} x_2$ is also a fuzzy random variable whose observed values are fuzzy numbers

characterized by the membership function

$$\mu_{\tilde{\boldsymbol{C}}_{l}\boldsymbol{x}}(\upsilon) = \begin{cases} \max\left\{0, 1 - \frac{\bar{\boldsymbol{d}}_{l}\boldsymbol{x} - \upsilon}{\boldsymbol{\beta}_{l}\boldsymbol{x}}\right\} &, \text{ if } \upsilon \leq \bar{\boldsymbol{d}}_{l}\boldsymbol{x} \\\\ \max\left\{0, 1 - \frac{\upsilon - \bar{\boldsymbol{d}}_{l}\boldsymbol{x}}{\boldsymbol{\gamma}_{l}\boldsymbol{x}}\right\} &, \text{ otherwise.} \end{cases}$$

An example of a membership function of an objective function for DMl is shown in Figure 2.



Figure 2: An example of a membership function of an objective function for DMl.

It should be emphasized here that problem (1) is not a well-defined problem due to both fuzziness and randomness of the coefficients, and it cannot be minimized in the sense of deterministic two-level linear programming. Therefore, it is necessary to interpret the problem from some point of view and to transform the problem into the deterministic equivalent one. Realizing this difficulty, in this paper, we assume that decision makers prefer to maximize the degree of possibility or necessity that objective function values satisfy fuzzy goals.

2.1 Fuzzy goals

Considering vague natures of decision makers' judgments, it is natural to assume that decision makers may have vague or fuzzy goals for each of the objective functions. In a minimization problem, a goal stated by decision makers may be to achieve "substantially less than or equal to some value." This type of statement can be quantified by eliciting a corresponding membership function. In this paper, in view of the linearity of the formulated problems, the fuzzy goals \tilde{G}_l such as " $z_l(\boldsymbol{x}_1, \boldsymbol{x}_2)$ should be substantially less than or equal to a certain value" are assumed to be quantified by the linear membership functions:

$$\mu_{\tilde{G}_{l}}(y) = \begin{cases} 1 & , & \text{if } y \leq z_{l}^{1} \\ \frac{y - z_{l}^{0}}{z_{l}^{1} - z_{l}^{0}} & , & \text{if } z_{l}^{1} < y \leq z_{l}^{0} \\ 0 & , & \text{if } y > z_{l}^{0}. \end{cases}$$
(2)

Figure 3 illustrates a possible shape of the membership function for the fuzzy goal.



Figure 3: An example of a membership function $\mu_{\tilde{G}_l}(\cdot)$ of a fuzzy goal \tilde{G}_l .

2.2 Possibility and necessity

Having determined the fuzzy goals of the decision makers, if we regard $\mu_{\tilde{C}_l \boldsymbol{x}}(\cdot)$ as a possibility distribution on the basis of the concept of possibility measure, the degree of possibility $\Pi_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l)$ that the fuzzy goal \tilde{G}_l is fulfilled under the possibility distribution $\mu_{\tilde{C}_l \boldsymbol{x}}(\cdot)$ is given by:

$$\Pi_{\tilde{\boldsymbol{C}}_{l}\boldsymbol{x}}(\tilde{G}_{l}) = \sup_{y} \min\left\{\mu_{\tilde{\boldsymbol{C}}_{l}\boldsymbol{x}}(y), \ \mu_{\tilde{G}_{l}}(y)\right\}, \ l = 1, 2.$$
(3)

Figure 4 illustrates the degree of possibility that the fuzzy goal \tilde{G}_l is fulfilled under the possibility distribution $\mu_{\tilde{G}_l x}(\cdot)$.



Figure 4: The degree of possibility $\Pi_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l)$

Using the degree of possibility, problem (1) to minimize each objective function $\tilde{C}_l x$ can be transformed into the following stochastic two-level programming problem to maximize the degree of possibility for each objective function $\prod_{\tilde{C}_l x} (\tilde{G}_l)$:

$$\begin{array}{c} \underset{DM1}{\text{maximize }} \Pi_{\tilde{\tilde{C}}_{1}\boldsymbol{x}}(\tilde{G}_{1}) \\ \underset{DM2}{\text{maximize }} \Pi_{\tilde{\tilde{C}}_{2}\boldsymbol{x}}(\tilde{G}_{2}) \\ \text{subject to } A_{1}\boldsymbol{x}_{1} + A_{2}\boldsymbol{x}_{2} \leq \boldsymbol{b} \\ \boldsymbol{x}_{1} \geq \boldsymbol{0}, \ \boldsymbol{x}_{2} \geq \boldsymbol{0} \end{array} \right\}.$$

$$(4)$$

On the other hand, if decision makers are more risk-averse or wish to avoid a risk, decision making using possibility may be inappropriate since the obtained solution becomes too optimistic. In such a situation, decision making using necessity seems to be suitable for pessimistic decision makers. To be more specific, the degree of necessity $N_{\tilde{C}_l x}(\tilde{G}_l)$ that the fuzzy goal \tilde{G}_l is fulfilled under the possibility distribution $\mu_{\tilde{C}_l x}(\cdot)$ is given by:

$$N_{\tilde{\boldsymbol{C}}_{l}\boldsymbol{x}}(\tilde{G}_{l}) = \inf_{y} \max\left\{1 - \mu_{\tilde{\boldsymbol{C}}_{l}\boldsymbol{x}}(y), \ \mu_{\tilde{G}_{l}}(y)\right\}, \ l = 1, 2.$$

$$(5)$$

Figure 5 illustrates the degree of necessity that the fuzzy goal \tilde{G}_l is fulfilled under the possibility distribution $N_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l)$.



Figure 5: The degree of necessity $N_{\tilde{\bar{C}}_l x}(\tilde{G}_l)$

Quite similar to the possibility case, using the degree of necessity, problem (1) can be transformed into the following stochastic two-level programming problem to maximize the degree of necessity for each objective function $N_{\tilde{C}_l x}(\tilde{G}_l)$:

$$\begin{array}{c} \underset{\text{DM1}}{\text{maximize }} N_{\tilde{\tilde{C}}_{1}\boldsymbol{x}}(\tilde{G}_{1}) \\ \underset{\text{DM2}}{\text{maximize }} N_{\tilde{\tilde{C}}_{2}\boldsymbol{x}}(\tilde{G}_{2}) \\ \text{subject to } A_{1}\boldsymbol{x}_{1} + A_{2}\boldsymbol{x}_{2} \leq \boldsymbol{b} \\ \boldsymbol{x}_{1} \geq \boldsymbol{0}, \ \boldsymbol{x}_{2} \geq \boldsymbol{0} \end{array} \right\} .$$

$$(6)$$

For each of the objective functions in (4) and (6), if we set

$$z_l^0 = \max_{s_l \in \{1, 2, \dots, S_l\}} \max_{\boldsymbol{x} \in X} \sum_{j=1}^2 \sum_{k=1}^{n_j} d_{ljks_l} x_{jk}$$
(7)

$$z_l^1 = \min_{s_l \in \{1, 2, \dots, S_l\}} \min_{\boldsymbol{x} \in X} \sum_{j=1}^2 \sum_{k=1}^{n_j} d_{ljks_l} x_{jk},$$
(8)

the degree of possibility (3) and the degree of necessity (5) can be rewritten as:

$$\Pi_{\tilde{\boldsymbol{C}}_{l}\boldsymbol{x}}(\tilde{G}_{l}) = \frac{\sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \beta_{ljk} - \bar{d}_{ljk} \right\} x_{jk} + z_{l}^{0}}{\sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \beta_{ljk} x_{jk} - z_{l}^{1} + z_{l}^{0}}$$
(9)

$$N_{\tilde{C}_{l}\boldsymbol{x}}(\tilde{G}_{l}) = \frac{-\sum_{j=1}^{2}\sum_{k=1}^{n_{j}} \bar{d}_{ljk}x_{jk} + z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}} \gamma_{ljk}x_{jk} - z_{l}^{1} + z_{l}^{0}}$$
(10)

where denominators in (9) and (10) are assumed to be positive since each of β_{ljk} , γ_{ljk} and x_{jk} , $l = 1, 2, j = 1, 2, k = 1, 2, ..., n_j$ is nonnegative and $z_l^1 < z_l^0$.

In this way, it follows that both of the problems (4) and (6) are stochastic two-level programming problems whose objective functions $\Pi_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l)$ and $N_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l)$ are linear fractional and vary randomly depending on random variables \bar{d}_{ljk} .

3 Stackelberg solutions through absolute deviation minimization

In this section, assuming the decision makers are risk-averse, we reduce the transformed stochastic two-level programming problems (4) and (6) to deterministic two-level programming problems through the absolute deviation minimization model [23]. It is significant to note that the absolute deviation minimization model is suitable for risk-averse decision makers and more tractable than the variance minimization model.

3.1 Possibility case

Following the absolute deviation minimization model, the maximization of $\Pi_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l)$ is replaced with the minimization of its absolute deviation $\operatorname{E}\left[\left|\Pi_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l) - \operatorname{E}\left[\Pi_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l)\right]\right|\right]$ as follows:

$$\begin{array}{l} \underset{\text{DM1}}{\text{minimize }} Z_{1}^{\Pi,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) = \mathbb{E}\left[\left|\Pi_{\tilde{\boldsymbol{C}}_{1}\boldsymbol{x}}(\tilde{G}_{1}) - \mathbb{E}\left[\Pi_{\tilde{\boldsymbol{C}}_{1}\boldsymbol{x}}(\tilde{G}_{1})\right]\right|\right] \\ \underset{\text{DM2}}{\text{minimize }} Z_{2}^{\Pi,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) = \mathbb{E}\left[\left|\Pi_{\tilde{\boldsymbol{C}}_{2}\boldsymbol{x}}(\tilde{G}_{2}) - \mathbb{E}\left[\Pi_{\tilde{\boldsymbol{C}}_{2}\boldsymbol{x}}(\tilde{G}_{2})\right]\right|\right] \\ \underset{\text{subject to }}{\text{subject to }} A_{1}\boldsymbol{x}_{1} + A_{2}\boldsymbol{x}_{2} \leq \boldsymbol{b} \\ \boldsymbol{x}_{1} \geq \boldsymbol{0}, \ \boldsymbol{x}_{2} \geq \boldsymbol{0} \end{array}\right\}. \tag{11}$$

Since each \bar{d}_{ljk} is a random variable that takes an observed value d_{ljks_l} under a scenario $s_l \in \{1, 2, ..., S_l\}$ whose probability is p_{s_l} , $\mathbf{E}\left[\left|\Pi_{\tilde{\boldsymbol{C}}_l\boldsymbol{x}}(\tilde{G}_l) - \mathbf{E}\left[\Pi_{\tilde{\boldsymbol{C}}_l\boldsymbol{x}}(\tilde{G}_l)\right]\right|\right]$ in (11) are rewritten as:

$$\begin{split} & \mathbf{E}\left[\left|\Pi_{\tilde{\mathbf{C}}_{l}\boldsymbol{x}}(\tilde{G}_{l})-\mathbf{E}\left[\Pi_{\tilde{\mathbf{C}}_{l}\boldsymbol{x}}(\tilde{G}_{l})\right]\right|\right] \\ & = \mathbf{E}\left[\left|\frac{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\left\{\beta_{ljk}-\bar{d}_{ljk}\right\}x_{jk}+z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\beta_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}-\mathbf{E}\left[\frac{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\left\{\beta_{ljk}-\bar{d}_{ljk}\right\}x_{jk}+z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\beta_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}\right]\right]\right] \\ & = \sum_{s_{l}=1}^{S_{l}}p_{ls_{l}}\left|\frac{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\left\{\beta_{ljk}-d_{ljks_{l}}\right\}x_{jk}+z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\beta_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}-\sum_{s_{l}=1}^{S_{l}}p_{ls_{l}}\frac{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\left\{\beta_{ljk}-d_{ljks_{l}}\right\}x_{jk}+z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\beta_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}\right| \end{split}$$

$$= \sum_{s_l=1}^{S_l} p_{ls_l} \left| \frac{\sum_{j=1}^{2} \sum_{k=1}^{n_j} \left\{ \sum_{s_l=1}^{S_l} p_{ls_l} d_{ljk} - d_{ljks_l} \right\} x_{jk} + z_l^0}{\sum_{j=1}^{2} \sum_{k=1}^{n_j} \beta_{ljk} x_{jk} - z_l^1 + z_l^0} \right|.$$

If we introduce the auxiliary variables

$$\begin{aligned} r_{ls_{l}}^{+} &= \frac{1}{2} \left[\left| \sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{ljk} - d_{ljks_{l}} \right\} x_{jk} + z_{l}^{0} \right| \\ &+ \left(\sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{ljk} - d_{ljks_{l}} \right\} x_{jk} + z_{l}^{0} \right) \right] \\ r_{ls_{l}}^{-} &= \frac{1}{2} \left[\left| \sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{ljk} - d_{ljks_{l}} \right\} x_{jk} + z_{l}^{0} \right| \\ &- \left(\sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{ljk} - d_{ljks_{l}} \right\} x_{jk} + z_{l}^{0} \right| \end{aligned}$$

 $r^+_{ls_l} \geq 0, \, r^-_{ls_l} \geq 0$ and the following relations hold:

$$\begin{aligned} r_{ls_{l}}^{+} + r_{ls_{l}}^{-} &= \left| \sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{ljk} - d_{ljks_{l}} \right\} x_{jk} + z_{l}^{0} \right| \\ r_{ls_{l}}^{+} - r_{ls_{l}}^{-} &= \sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{ljk} - d_{ljks_{l}} \right\} x_{jk} + z_{l}^{0} \\ r_{ls_{l}}^{+} \cdot r_{ls_{l}}^{-} &= 0. \end{aligned}$$

In this way, through absolute deviation minimization using possibility, (11) can be reduced to the following deterministic two-level programming problem:

$$\begin{array}{l} \text{minimize} \ Z_{1}^{\Pi,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) = \frac{\sum\limits_{j=1}^{S_{1}} p_{1s_{1}}\left(r_{1s_{1}}^{+} + r_{1s_{1}}^{-}\right)}{\sum\limits_{j=1}^{2} \sum\limits_{k=1}^{n_{j}} \beta_{1jk}x_{jk} - z_{1}^{1} + z_{1}^{0}} \\ \\ \text{minimize} \ Z_{2}^{\Pi,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) = \frac{\sum\limits_{j=1}^{S_{2}} p_{2s_{2}}\left(r_{2s_{2}}^{+} + r_{2s_{2}}^{-}\right)}{\sum\limits_{j=1}^{2} \sum\limits_{k=1}^{n_{j}} \beta_{2jk}x_{jk} - z_{1}^{1} + z_{2}^{0}} \\ \\ \text{subject to} \ A_{1}\boldsymbol{x}_{1} + A_{2}\boldsymbol{x}_{2} \le \boldsymbol{b} \\ \\ \\ \sum\limits_{j=1}^{2} \sum\limits_{k=1}^{n_{j}} \left\{\sum\limits_{s_{l}=1}^{S_{l}} p_{ls_{l}}d_{ljk} - d_{ljks_{l}}\right\} x_{jk} - r_{ls_{l}}^{+} + r_{ls_{l}}^{-} = -z_{l}^{0}, \end{array} \right\} .$$
(12)

$$\begin{array}{l} l = 1, 2, \ s_l = 1, 2, \dots, S_l \\ r_{ls_l}^+ \cdot r_{ls_l}^- = 0, \ l = 1, 2, \ s_l = 1, 2, \dots, S_l \\ \boldsymbol{x}_1 \ge \boldsymbol{0}, \ \boldsymbol{x}_2 \ge \boldsymbol{0}, \ r_{ls_l}^+ \ge 0, \ r_{ls_l}^- \ge 0, \ l = 1, 2, \ s_l = 1, 2, \dots, S_l \end{array}$$

It should be noted here that for noncooperative two-level programming problems, DM1 first specifies a decision and then DM2 determines a decision so as to optimize the objective function of self with full knowledge of the decision of DM1. According to this rule, DM1 also makes a decision so as to optimize the objective function of self. The solution defined as the procedure is called a Stackelberg solution.

Realizing that (12) is a deterministic two-level programming problem, we are now ready to introduce the extended concepts of Stackelberg solution for the original fuzzy random two-level linear programming problem (1).

Definition 2 (AD-P-Stackelberg solution) A feasible solution $(\boldsymbol{x}_1^*, \boldsymbol{x}_2^*) \in X$ is called an AD-P-Stackelberg solution, meaning a Stackelberg solution through absolute deviation minimization using possibility, if $(\boldsymbol{x}_1^*, \boldsymbol{x}_2^*)$ is an optimal solution to the following two-level linear fractional programming problem:

$$\begin{array}{c} \underset{\boldsymbol{x}_{1}}{\min initial} Z_{1}^{\Pi,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) \\ \text{where } \boldsymbol{x}_{2} \text{ solves} \\ \min initial} Z_{2}^{\Pi,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) \\ \text{subject to } A_{1}\boldsymbol{x}_{1} + A_{2}\boldsymbol{x}_{2} \leq \boldsymbol{b} \\ \sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}}d_{ljk} - d_{ljks_{l}} \right\} x_{jk} - r_{ls_{l}}^{+} + r_{ls_{l}}^{-} = -z_{l}^{0}, \\ l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l} \\ r_{ls_{l}}^{+} \cdot r_{ls_{l}}^{-} = 0, \ l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l} \\ \boldsymbol{x}_{1} \geq \boldsymbol{0}, \ \boldsymbol{x}_{2} \geq \boldsymbol{0}, \ r_{ls_{l}}^{+} \geq 0, \ r_{ls_{l}}^{-} \geq 0, \ l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l} \end{array} \right\} .$$
(13)

Observing that (13) is a two-level linear fractional programming problem when complementary conditions are relaxed, it is now appropriate to consider some effective computational methods for obtaining AD-P-Stackelberg solutions.

Following the definition of Stackelberg solutions, for any feasible decision \hat{x}_1 given by DM1, DM2 is assumed to select a decision $x_2(\hat{x}_1)$ which is an optimal solution to the following problem:

$$\begin{array}{l} \text{minimize} \quad \frac{\sum\limits_{s_{2}=1}^{S_{2}} p_{2s_{2}} \left(r_{2s_{2}}^{+} + r_{2s_{2}}^{-} \right) }{\sum\limits_{k=1}^{n_{2}} \beta_{22k} x_{2k} - z_{2}^{1} + z_{2}^{0} + \sum\limits_{k=1}^{n_{1}} \beta_{21k} \hat{x}_{1k} } \\ \text{subject to} \quad A_{2} \boldsymbol{x}_{2} \leq \boldsymbol{b} - A_{1} \hat{\boldsymbol{x}}_{1} \\ \sum\limits_{k=1}^{n_{2}} \left(\sum\limits_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{l2k} - d_{l2ks_{l}} \right) x_{2k} - r_{ls_{l}}^{+} + r_{ls_{l}}^{-} \\ = -z_{l}^{0} - \sum\limits_{k=1}^{n_{1}} \left(\sum\limits_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{l1k} - d_{l1ks_{l}} \right) \hat{x}_{1k}, \ l = 1, 2, \ s_{l} = 1, 2, \dots, S_{l} \\ r_{ls_{l}}^{+} \cdot r_{ls_{l}}^{-} = 0, \ l = 1, 2, \ s_{l} = 1, 2, \dots, S_{l} \\ \boldsymbol{x}_{2} \geq \boldsymbol{0}, \ r_{ls_{l}}^{+} \geq 0, \ r_{ls_{l}}^{-} \geq 0, \ l = 1, 2, \ s_{l} = 1, 2, \dots, S_{l} \end{array} \right)$$

The optimal solution $x_2(\hat{x}_1)$ to (14) is called a rational reaction for \hat{x}_1 . Let us denote the set of rational reactions for \hat{x}_1 by $RR(\hat{x}_1)$. Then, DM1 should select a solution (x_1, x_2)

to optimize $Z_1^{\Pi,AD}(\boldsymbol{x}_1,\boldsymbol{x}_2)$ from among the inducible region $IR = \{(\boldsymbol{x}_1,\boldsymbol{x}_2) \mid (\boldsymbol{x}_1,\boldsymbol{x}_2) \in$ $X, x_2 \in RR(x_1)$. To be more explicit, DM1 selects an optimal solution to the following problem:

minimize
$$Z_{1}^{\Pi,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2})$$

subject to $A_{1}\boldsymbol{x}_{1} + A_{2}\boldsymbol{x}_{2} \leq \boldsymbol{b}$
 $\sum_{j=1}^{2}\sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}}d_{ljk} - d_{ljks_{l}} \right\} x_{jk} - r_{ls_{l}}^{+} + r_{ls_{l}}^{-} = -z_{l}^{0},$
 $l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l}$
 $r_{ls_{l}}^{+} \cdot r_{ls_{l}}^{-} = 0, \ l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l}$
 $\boldsymbol{x}_{1} \geq \boldsymbol{0}, \ \boldsymbol{x}_{2} \in RR(\boldsymbol{x}_{1}), \ r_{ls_{l}}^{+} \geq 0, \ r_{ls_{l}}^{-} \geq 0, \ l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l}$
(15)

It should be emphasized here that the optimal solution to (15) is an AD-P-Stackelberg solution.

For two-level linear fractional programming problems with linear fractional objective functions and linear constraints, it is shown that Stackelberg solutions exist at some extreme point of the feasible region [7]. Although (13) is not a two-level linear fractional programming problem, if complementary conditions $r_{ls_l}^+ \cdot r_{ls_l}^- = 0, l = 1, 2,$ $s_l = 1, 2, \ldots, S_l$ in (13) are relaxed, it should be emphasized here that the resulting relaxed problem becomes a two-level linear fractional programming one. Hence, from the property of Stackelberg solutions to two-level linear fractional programming problems, we can omit complementary conditions from (13) since these conditions automatically hold at any extreme point of the feasible region. In this way, we can consider the following relaxed problem:

minimize
$$Z_1^{\Pi,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2) = \frac{\sum\limits_{s_1=1}^{S_1} p_{1s_1} \left(r_{1s_1}^+ + r_{1s_1}^- \right)}{\sum\limits_{j=1}^{2} \sum\limits_{k=1}^{n_j} \beta_{1jk} x_{jk} - z_1^1 + z_1^0}$$

where \boldsymbol{x}_2 solves

where \boldsymbol{x}_2 solves

minimize
$$Z_{2}^{\Pi,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) = \frac{\sum_{s_{2}=1}^{S_{2}} p_{2s_{2}}\left(r_{2s_{2}}^{+} + r_{2s_{2}}^{-}\right)}{\sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \beta_{2jk} x_{jk} - z_{2}^{1} + z_{2}^{0}}$$

subject to $A_{1}\boldsymbol{x}_{1} + A_{2}\boldsymbol{x}_{2} \leq \boldsymbol{b}$
 $\sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{\sum_{s_{l}=1}^{S_{l}} p_{ls_{l}}d_{ljk} - d_{ljks_{l}}\right\} x_{jk} - r_{ls_{l}}^{+} + r_{ls_{l}}^{-} = -z_{l}^{0},$
 $l = 1, 2, s_{l} = 1, 2, \dots, S_{l}$

$$(16)$$

$$l = 1, 2, \ s_l = 1, 2, \ldots, S_l$$

 $oldsymbol{x}_1 \ge oldsymbol{0}, \ oldsymbol{x}_2 \ge oldsymbol{0}, \ r_{ls_l}^+ \ge 0, \ r_{ls_l}^- \ge 0, \ l = 1, 2, \ s_l = 1, 2, \ldots, S_l$

Observing that (16) is a two-level linear fractional programming problem, we can construct the following computational method for obtaining AD-P-Stackelberg solutions through the combined use of the variable transformation method by Charnes and Cooper [8] and the Kth best algorithm for two-level linear programming problems by Bialas et al. [5].

The computational method for obtaining AD-P-Stackelberg solutions

Step 1: Let i := 1. Removing the objective function of DM2 from (16), solve the following problem:

minimize
$$Z_{1}^{\Pi,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) = \frac{\sum_{s_{1}=1}^{S_{1}} p_{1s_{1}}\left(r_{1s_{1}}^{+} + r_{1s_{1}}^{-}\right)}{\sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \beta_{1jk}x_{jk} - z_{1}^{1} + z_{1}^{0}}$$
subject to $A_{1}\boldsymbol{x}_{1} + A_{2}\boldsymbol{x}_{2} \leq \boldsymbol{b}$
 $\sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{\sum_{s_{l}=1}^{S_{l}} p_{ls_{l}}d_{ljk} - d_{ljks_{l}}\right\} x_{jk} - r_{ls_{l}}^{+} + r_{ls_{l}}^{-} = -z_{l}^{0},$
 $l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l}$
 $\boldsymbol{x}_{1} \geq \boldsymbol{0}, \ \boldsymbol{x}_{2} \geq \boldsymbol{0}, \ r_{ls_{l}}^{+} \geq 0, \ r_{ls_{l}}^{-} \geq 0, \ l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l}$
(17)

Observing that (17) is a linear fractional programming problem and the denominator of the objective function is positive as discussed in (9), it can be transformed into an equivalent linear programming problem by the variable transformation method by Charnes and Cooper [8]. To be more specific, introducing the variable transformation

$$t = \frac{1}{\sum_{j=1}^{2} \sum_{k=1}^{n_j} \beta_{1jk} x_{jk} - z_1^1 + z_1^0}$$

and letting $\boldsymbol{y}_1 = t \cdot \boldsymbol{x}_1$, $\boldsymbol{y}_2 = t \cdot \boldsymbol{x}_2$, $\boldsymbol{q}^+ = t \cdot \boldsymbol{r}^+$, $\boldsymbol{q}^- = t \cdot \boldsymbol{r}^-$, (17) is transformed into the following linear programming problem:

$$\begin{array}{l} \text{minimize} \quad \sum_{s_{1}=1}^{S_{1}} p_{1s_{1}} \left(q_{1s_{1}}^{+} + q_{1s_{1}}^{-} \right) \\ \text{subject to} \quad A_{1} \boldsymbol{y}_{1} + A_{2} \boldsymbol{y}_{2} - \boldsymbol{b}t \leq \boldsymbol{0} \\ \sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{ljk} - d_{ljks_{l}} \right\} y_{jk} - q_{ls_{l}}^{+} + q_{ls_{l}}^{-} + z_{l}^{0} t = 0, \\ \quad l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l} \\ \sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \beta_{1jk} y_{jk} - (z_{1}^{1} - z_{1}^{0}) t = 1 \\ \boldsymbol{y}_{1} \geq \boldsymbol{0}, \ \boldsymbol{y}_{2} \geq \boldsymbol{0}, \ \boldsymbol{q}^{+} \geq \boldsymbol{0}, \ \boldsymbol{q}^{-} \geq \boldsymbol{0}, \ t \geq 0 \end{array} \right\} .$$
(18)

Observing that (18) is a linear programming problem, we can obtain an optimal solution by the simplex method. Using the optimal solution to (18) denoted by $(\boldsymbol{y}_{1[1]}^T, \boldsymbol{y}_{2[1]}^T, (\boldsymbol{q}_{1]}^+)^T, (\boldsymbol{q}_{1]}^-)^T, t_{1]})^T$, we can obtain

$$(oldsymbol{x}_{1[1]}^T,oldsymbol{x}_{2[1]}^T)^T := (oldsymbol{y}_{1[1]}^T/t_{[1]},oldsymbol{y}_{2[1]}^T/t_{[1]})^T$$

which is an extreme point of the feasible region of (17) as shown in [49]. Let W be a set of feasible extreme points to be searched and U a set of feasible extreme points that had been searched. Let $W := \{(\boldsymbol{x}_{1[1]}^T, \boldsymbol{x}_{2[1]}^T)^T\}$ and $U := \emptyset$. Go to step 2.

Step 2: In order to check whether the present extreme point $(\boldsymbol{x}_{1[i]}^T, \boldsymbol{x}_{2[i]}^T)^T$ exists in the inducible region *IR*, i.e., $\boldsymbol{x}_{2[i]}$ is a rational reaction for $\boldsymbol{x}_{1[i]}$ or not, we solve the following problem:

Observing that (19) is also a linear fractional programming problem and the denominator of the objective function is positive, we can apply the variable transformation method by Charnes and Cooper [8] to (19). Namely, introducing the variable transformation

$$u = \frac{1}{\sum_{k=1}^{n_2} \beta_{22k} x_{2k} - z_2^1 + z_2^0 + \sum_{k=1}^{n_1} \beta_{21k} x_{1k[i]}}$$

and letting $w_2 := u \cdot x_2$, $o^+ := u \cdot r^+$, $o^- := u \cdot r^-$, (19) is transformed into the following linear programming problem:

minimize
$$\sum_{s_{2}=1}^{S_{2}} p_{2s_{2}} \left(o_{2s_{2}}^{+} + o_{2s_{2}}^{-} \right)$$

subject to $A_{2} w_{2} - (b - A_{1} x_{1[i]}) u \leq \mathbf{0}$

$$\sum_{k=1}^{n_{2}} \left(\sum_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{l2k} - d_{l2ks_{l}} \right) w_{2k} - o_{ls_{l}}^{+} + o_{ls_{l}}^{-}$$

$$+ \left(z_{l}^{0} + \sum_{k=1}^{n_{1}} \left(\sum_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{l1k} - d_{l1ks_{l}} \right) x_{1k[i]} \right) u = 0,$$

$$l = 1, 2, \ s_{l} = 1, 2, \dots, S_{l}$$

$$\sum_{k=1}^{n_{2}} \beta_{22k} w_{2k} - \left(z_{2}^{1} - z_{2}^{0} - \sum_{k=1}^{n_{1}} \beta_{21k} x_{1k[i]} \right) u = 1$$

$$w_{2} \geq \mathbf{0}, \ \mathbf{o}^{+} \geq \mathbf{0}, \ \mathbf{o}^{-} \geq \mathbf{0}, \ u \geq 0$$

$$(20)$$

Observing that (20) is a linear programming problem, we can obtain an optimal solution $(\boldsymbol{w}_{2[i]}^T, (\boldsymbol{o}_{[i]}^+)^T, (\boldsymbol{o}_{[i]}^-)^T, u_{[i]})^T$ by the simplex method. If $\boldsymbol{w}_{2[i]}/u_{[i]}$ is equal to $\boldsymbol{x}_{2[i]}$, then the current extreme point $(\boldsymbol{x}_{1[i]}^T, \boldsymbol{x}_{2[i]}^T)^T$ exists in *IR*, i.e., it is an AD-P-Stackelberg solution and the algorithm is terminated. Otherwise, go to step 3.

Step 3: Let $W_{[i]}$ be a set of feasible extreme points which is adjacent to $(\boldsymbol{x}_{1[i]}^T, \boldsymbol{x}_{2[i]}^T)^T$ and satisfies $Z_1^{\Pi,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2) \geq Z_1^{\Pi,AD}(\boldsymbol{x}_{1[i]}, \boldsymbol{x}_{2[i]})$. Let $U := U \cup \{(\boldsymbol{x}_{1[i]}^T, \boldsymbol{x}_{2[i]}^T)^T\}$ and $W := (W \cup W_{[i]}) \setminus U$, and go to step 4. **Step 4:** Let i := i + 1. Choose an extreme point $(\boldsymbol{x}_{1[i]}^T, \boldsymbol{x}_{2[i]}^T)^T$ such that

$$Z_1^{\Pi,AD}(m{x}_{1[i]},m{x}_{2[i]}) = \min_{(m{x}_1^T,m{x}_2^T)^T\in W} Z_1^{\Pi,AD}(m{x}_1,m{x}_2),$$

and return to step 2.

It should be noted here that the proposed computational method uses nothing but the variable transformation method, the simplex method and the pivot operation for obtaining an AD-P-Stackelberg solution.

3.2 Necessity case

Quite similar to the possibility case, following the absolute deviation minimization model, the maximization of $N_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l)$ in (6) is replaced with the minimization of its absolute deviation $\mathbf{E}\left[\left|N_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l) - \mathbf{E}\left[N_{\tilde{C}_l \boldsymbol{x}}(\tilde{G}_l)\right]\right|\right]$ as follows:

$$\begin{array}{l} \underset{\text{DM1}}{\text{minimize }} Z_1^{N,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2) = \mathbb{E}\left[\left| N_{\tilde{\boldsymbol{C}}_1 \boldsymbol{x}}(\tilde{G}_1) - \mathbb{E}\left[N_{\tilde{\boldsymbol{C}}_1 \boldsymbol{x}}(\tilde{G}_1) \right] \right| \right] \\ \underset{\text{DM2}}{\text{minimize }} Z_2^{N,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2) = \mathbb{E}\left[\left| N_{\tilde{\boldsymbol{C}}_2 \boldsymbol{x}}(\tilde{G}_2) - \mathbb{E}\left[N_{\tilde{\boldsymbol{C}}_2 \boldsymbol{x}}(\tilde{G}_2) \right] \right| \right] \\ \text{subject to } A_1 \boldsymbol{x}_1 + A_2 \boldsymbol{x}_2 \leq \boldsymbol{b} \\ \boldsymbol{x}_1 \geq \boldsymbol{0}, \ \boldsymbol{x}_2 \geq \boldsymbol{0} \end{array} \right\},$$
(21)

where $\mathbf{E}\left[\left|N_{\tilde{\tilde{C}}_{l}\boldsymbol{x}}(\tilde{G}_{l})-\mathbf{E}\left[N_{\tilde{\tilde{C}}_{l}\boldsymbol{x}}(\tilde{G}_{l})\right]\right|\right]$ is rewritten as:

$$\begin{split} & \mathbf{E}\left[\left|N_{\tilde{\mathcal{C}}_{l}\boldsymbol{x}}(\tilde{G}_{l})-\mathbf{E}\left[N_{\tilde{\mathcal{C}}_{l}\boldsymbol{x}}(\tilde{G}_{l})\right]\right|\right] \\ &= \mathbf{E}\left[\left|\frac{-\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\bar{d}_{ljk}x_{jk}+z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\gamma_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}-\mathbf{E}\left[\frac{-\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\bar{d}_{ljk}x_{jk}+z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\gamma_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}\right]\right|\right] \\ &= \sum_{s_{l}=1}^{S_{l}}p_{ls_{l}}\left|\frac{-\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}d_{ljks_{l}}x_{jk}+z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\gamma_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}-\sum_{s_{l}=1}^{S_{l}}p_{ls_{l}}\frac{-\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}d_{ljks_{l}}x_{jk}+z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\gamma_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}-\sum_{s_{l}=1}^{S_{l}}p_{ls_{l}}\frac{-\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\gamma_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}{\sum_{j=1}^{2}\sum_{k=1}^{n_{j}}\gamma_{ljk}x_{jk}-z_{l}^{1}+z_{l}^{0}}\right|. \end{split}$$

Introducing the auxiliary variables $r_{ls_l}^+$ and $r_{ls_l}^-$ as defined in the possibility case, (21) can be reduced to the following deterministic two-level programming problem:

minimize
$$Z_1^{N,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2) = \frac{\sum_{s_1=1}^{S_1} p_{1s_1} \left(r_{1s_1}^+ + r_{1s_1}^- \right)}{\sum_{j=1}^{2} \sum_{k=1}^{n_j} \gamma_{1jk} x_{jk} - z_1^1 + z_1^0}$$

minimize $Z_2^{N,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2) = \frac{\sum_{s_2=1}^{S_2} p_{2s_2} \left(r_{2s_2}^+ + r_{2s_2}^- \right)}{\sum_{j=1}^{2} \sum_{k=1}^{n_j} \gamma_{2jk} x_{jk} - z_2^1 + z_2^0}$
while to $A, \boldsymbol{x}_1 + A, \boldsymbol{x}_2 \leq \boldsymbol{b}$. (22)

subject to
$$A_1 \boldsymbol{x}_1 + A_2 \boldsymbol{x}_2 \leq \boldsymbol{b}$$

 $\sum_{j=1}^{2} \sum_{k=1}^{n_j} \left\{ \sum_{s_l=1}^{S_l} p_{ls_l} d_{ljk} - d_{ljks_l} \right\} x_{jk} - r_{ls_l}^+ + r_{ls_l}^- = -z_l^0,$
 $l = 1, 2, \ s_l = 1, 2, \dots, S_l$
 $r_{ls_l}^+ \cdot r_{ls_l}^- = 0, \ l = 1, 2, \ s_l = 1, 2, \dots, S_l$
 $\boldsymbol{x}_1 \geq \boldsymbol{0}, \ \boldsymbol{x}_2 \geq \boldsymbol{0}, \ r_{ls_l}^+ \geq 0, \ r_{ls_l}^- \geq 0, \ l = 1, 2, \ s_l = 1, 2, \dots, S_l$

Observing that (22) is a deterministic two-level programming problem, we can introduce the extended concepts of Stackelberg solution for the original fuzzy random twolevel linear programming problem (1).

Definition 3 (AD-N-Stackelberg solution) A feasible solution $(\mathbf{x}_1^*, \mathbf{x}_2^*) \in X$ is called an AD-N-Stackelberg solution, meaning a Stackelberg solution through absolute deviation minimization using necessity, if $(\mathbf{x}_1^*, \mathbf{x}_2^*)$ is an optimal solution to the following two-level linear fractional programming problem:

$$\begin{array}{c} \underset{\mathbf{x}_{1}}{\min initial} \mathbb{Z}_{1}^{N,AD}(\mathbf{x}_{1},\mathbf{x}_{2}) \\ \text{where } \mathbf{x}_{2} \text{ solves} \\ \min initial} \mathbb{Z}_{2}^{N,AD}(\mathbf{x}_{1},\mathbf{x}_{2}) \\ \text{subject to } A_{1}\mathbf{x}_{1} + A_{2}\mathbf{x}_{2} \leq \mathbf{b} \\ \sum_{j=1}^{2} \sum_{k=1}^{n_{j}} \left\{ \sum_{s_{l}=1}^{S_{l}} p_{ls_{l}}d_{ljk} - d_{ljks_{l}} \right\} \mathbb{x}_{jk} - r_{ls_{l}}^{+} + r_{ls_{l}}^{-} = -z_{l}^{0}, \\ \mathbb{1} = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l} \\ \mathbb{1} = 1, 2, \ldots, S_{l} \\ \mathbb$$

Similarly to the possibility case, for any feasible decision \hat{x}_1 given by DM1, the ratio-

nal reaction $x_2(\hat{x}_1)$ of DM2 can be obtained by solving the following problem:

$$\begin{array}{l} \text{minimize} \quad \frac{\sum\limits_{s_{2}=1}^{S_{2}} p_{2s_{2}} \left(r_{2s_{2}}^{+} + r_{2s_{2}}^{-} \right) }{\sum\limits_{k=1}^{n_{2}} \gamma_{22k} x_{2k} - z_{2}^{1} + z_{2}^{0} + \sum\limits_{k=1}^{n_{1}} \gamma_{21k} \hat{x}_{1k} } \\ \text{subject to} \quad A_{2} \boldsymbol{x}_{2} \leq \boldsymbol{b} - A_{1} \hat{\boldsymbol{x}}_{1} \\ \sum\limits_{k=1}^{n_{2}} \left(\sum\limits_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{l2k} - d_{l2ks_{l}} \right) x_{2k} - r_{ls_{l}}^{+} + r_{ls_{l}}^{-} \\ = -z_{l}^{0} - \sum\limits_{k=1}^{n_{1}} \left(\sum\limits_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{l1k} - d_{l1ks_{l}} \right) \hat{x}_{1k}, \ l = 1, 2, \ s_{l} = 1, 2, \dots, S_{l} \\ r_{ls_{l}}^{+} \cdot r_{ls_{l}}^{-} = 0, \ l = 1, 2, \ s_{l} = 1, 2, \dots, S_{l} \\ \boldsymbol{x}_{2} \geq \boldsymbol{0}, \ r_{ls_{l}}^{+} \geq 0, \ r_{ls_{l}}^{-} \geq 0, \ l = 1, 2, \ s_{l} = 1, 2, \dots, S_{l} \end{array} \right)$$

and DM1 should select an optimal solution $(\boldsymbol{x}_1, \boldsymbol{x}_2)$ to the following problem:

minimize
$$Z_1^{N,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2)$$

subject to $A_1 \boldsymbol{x}_1 + A_2 \boldsymbol{x}_2 \leq \boldsymbol{b}$
 $\sum_{j=1}^2 \sum_{k=1}^{n_j} \left\{ \sum_{s_l=1}^{S_l} p_{ls_l} d_{ljk} - d_{ljks_l} \right\} x_{jk} - r_{ls_l}^+ + r_{ls_l}^- = -z_l^0,$
 $l = 1, 2, \ s_l = 1, 2, \dots, S_l$
 $r_{ls_l}^+ \cdot r_{ls_l}^- = 0, \ l = 1, 2, \ s_l = 1, 2, \dots, S_l$
 $\boldsymbol{x}_1 \geq \boldsymbol{0}, \ \boldsymbol{x}_2 \in RR(\boldsymbol{x}_1), \ r_{ls_l}^+ \geq 0, \ r_{ls_l}^- \geq 0, \ l = 1, 2, \ s_l = 1, 2, \dots, S_l$
(25)

Observe that the optimal solution to (25) is an AD-N-Stackelberg solution.

As discussed in the possibility case, we can consider the following relaxed problem:

$$\begin{array}{l} \text{minimize} \ Z_{1}^{N,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) = \frac{\sum\limits_{j=1}^{S_{1}} p_{1s_{1}} \left(r_{1s_{1}}^{+} + r_{1s_{1}}^{-}\right)}{\sum\limits_{j=1}^{2} \sum\limits_{k=1}^{n_{j}} \gamma_{1jk} x_{jk} - z_{1}^{1} + z_{1}^{0}} \\ \text{where} \ \boldsymbol{x}_{2} \text{ solves} \\ \text{minimize} \ Z_{2}^{N,AD}(\boldsymbol{x}_{1},\boldsymbol{x}_{2}) = \frac{\sum\limits_{s_{2}=1}^{S_{2}} p_{2s_{2}} \left(r_{2s_{2}}^{+} + r_{2s_{2}}^{-}\right)}{\sum\limits_{j=1}^{2} \sum\limits_{k=1}^{n_{j}} \gamma_{2jk} x_{jk} - z_{1}^{1} + z_{2}^{0}} \\ \text{subject to} \ A_{1}\boldsymbol{x}_{1} + A_{2}\boldsymbol{x}_{2} \le \boldsymbol{b} \\ \sum_{j=1}^{2} \sum\limits_{k=1}^{n_{j}} \left\{\sum\limits_{s_{l}=1}^{S_{l}} p_{ls_{l}} d_{ljk} - d_{ljks_{l}}\right\} x_{jk} - r_{ls_{l}}^{+} + r_{ls_{l}}^{-} = -z_{l}^{0}, \\ \ l = 1, 2, \ s_{l} = 1, 2, \ \ldots, S_{l} \\ \boldsymbol{x}_{1} \ge \boldsymbol{0}, \ \boldsymbol{x}_{2} \ge \boldsymbol{0}, \ r_{ls_{l}}^{+} \ge 0, \ r_{ls_{l}}^{-} \ge 0, \ l = 1, 2, \ s_{l} = 1, 2, \ldots, S_{l} \end{array} \right\}$$

The computational method for obtaining AD-N-Stackelberg solutions follows very much in the same fashion as that for obtaining AD-P-Stackelberg solutions and will, therefore, be omitted.

4 Numerical example

In order to demonstrate the feasibility and efficiency of the proposed computational methods, consider the following two-level linear programming problem involving fuzzy random variable coefficients:

$$\begin{array}{ll} \underset{\text{DM1}}{\text{minimize}} & z_1(\boldsymbol{x}_1, \boldsymbol{x}_2) = \tilde{\boldsymbol{C}}_{11} \boldsymbol{x}_1 + \tilde{\boldsymbol{C}}_{12} \boldsymbol{x}_2 \\ \underset{\text{DM2}}{\text{minimize}} & z_2(\boldsymbol{x}_1, \boldsymbol{x}_2) = \tilde{\boldsymbol{C}}_{21} \boldsymbol{x}_1 + \tilde{\boldsymbol{C}}_{22} \boldsymbol{x}_2 \\ \text{subject to} & \boldsymbol{a}_{11} \boldsymbol{x}_1 + \boldsymbol{a}_{12} \boldsymbol{x}_2 \leq b_1 \\ & \boldsymbol{a}_{21} \boldsymbol{x}_1 + \boldsymbol{a}_{22} \boldsymbol{x}_2 \leq b_2 \\ & \boldsymbol{a}_{31} \boldsymbol{x}_1 + \boldsymbol{a}_{32} \boldsymbol{x}_2 \leq b_3 \\ & \boldsymbol{a}_{41} \boldsymbol{x}_1 + \boldsymbol{a}_{42} \boldsymbol{x}_2 \leq b_4 \\ & \boldsymbol{x}_1 = (x_{11}, x_{12}, x_{13})^T \geq \boldsymbol{0} \\ & \boldsymbol{x}_2 = (x_{21}, x_{22}, x_{23})^T \geq \boldsymbol{0} \end{array} \right)$$

where \tilde{C}_{lj} , l = 1, 2, j = 1, 2 are vectors whose elements \tilde{C}_{ljk} , $k = 1, 2, ..., n_j$ are fuzzy random variables.

Values of coefficients in constraints, values of d_{ljks_l} for each s_l , β_{ljk} and γ_{ljk} , l = 1, 2 are shown in Tables 1, 2 and 3, respectively.

	x_{11}	x_{12}	x_{13}	x_{21}	x_{22}	x_{23}	b
$oldsymbol{a}_1$	2	3	1	2	3	3	65
$oldsymbol{a}_2$	4	4	2	3	2	1	80
$oldsymbol{a}_3$	2	4	3	3	2	2	105
$oldsymbol{a}_4$	-3	-2	-2	-4	-1	-4	-70

Table 1: Values of coefficients in constraints

Table 2: Values of d_{1jks_1} for each $s_1 \in \{1, 2, 3\}$, β_{1jk} and γ_{1jk} .

	x_{11}	x_{12}	x_{13}	x_{21}	x_{22}	x_{23}
Scenario $s_1 = 1 \ (p_{11} = 0.25)$	2.3	-1.0	1.3	-1.3	-1.8	2.0
Scenario $s_1 = 2 (p_{12} = 0.40)$	2.0	-1.3	2.0	1.1	-2.1	2.4
Scenario $s_1 = 3 \ (p_{13} = 0.35)$	1.9	-2.4	2.7	-1.5	-1.2	3.8
β_{1jk}	0.8	1.2	0.7	0.9	1.3	0.6
γ_{1jk}	0.8	1.1	0.5	0.6	0.9	1.0

Calculating z_l^0 and z_l^1 from (7) and (8) yields $z_1^0 = 121$, $z_1^1 = -48$, $z_2^0 = 76$ and $z_2^1 = -81$. Considering these values, membership functions of fuzzy goals for objective functions are determined as shown in Figure 6.

For illustrative purposes, we first derive an AD-P-Stackelberg solution to (27). For this numerical example, in step 1, after transforming (17) into (18) by the variable transformation method, (18) is solved by the simplex method. For the obtained value of

	x_{11}	x_{12}	x_{13}	x_{21}	x_{22}	x_{23}
Scenario $s_2 = 1 \ (p_{21} = 0.45)$	3.0	1.7	-1.6	-1.4	-1.6	1.7
Scenario $s_2 = 2 \ (p_{22} = 0.15)$	1.7	1.3	-2.3	-0.8	-1.9	2.6
Scenario $s_2 = 3 \ (p_{23} = 0.40)$	2.3	0.9	-1.0	-2.0	-1.2	3.5
eta_{2jk}	0.7	1.2	0.8	0.5	0.9	1.1
γ_{2ik}	0.7	0.9	0.6	1.0	0.8	0.9

Table 3: Values of d_{2jks_2} for each $s_2 \in \{1, 2, 3\}$, β_{2jk} and γ_{2jk} .



Figure 6: Membership functions $\mu_{\tilde{G}_l}(\cdot)$, l = 1, 2.

 $(\boldsymbol{x}_{1[1]}^T, \boldsymbol{x}_{2[1]}^T)^T = (14.58, 0.00, 0.83, 3.88, 9.08, 0.00)^T$, let $W := \{(\boldsymbol{x}_{1[1]}^T, \boldsymbol{x}_{2[1]}^T)^T\}$, $U := \emptyset$. In step 2, after transforming (19) into (20) by the variable transformation method, we solve (20) by the simplex method in order to obtain the rational reaction for $\boldsymbol{x}_{1[1]}$. Since the optimal solution to (20) $\tilde{\boldsymbol{w}}_{2[1]}/\boldsymbol{u}_{[1]} = (3.23, 11.70, 0.00)^T$ is not equal to $\boldsymbol{x}_{2[1]} = (3.88, 9.08, 0.00)^T$, the current extreme point $(\boldsymbol{x}_{1[1]}^T, \boldsymbol{x}_{2[1]}^T)^T$ is not an AD-P-Stackelberg solution. In step 3, we enumerate feasible extreme points $(\boldsymbol{x}_1^T, \boldsymbol{x}_2^T)^T$ which are adjacent to $(\boldsymbol{x}_{1[1]}^T, \boldsymbol{x}_{2[1]}^T)^T$ and satisfy $Z_1^{\Pi,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2) \ge Z_1^{\Pi,AD}(\boldsymbol{x}_{1[1]}, \boldsymbol{x}_{2[1]})$, and make $W_{[1]}$. Then, let $U := U \cup (\boldsymbol{x}_{1[1]}^T, \boldsymbol{x}_{2[1]}^T)^T$ in W whose $Z_1^{\Pi,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2)$ is the least and let it be the next extreme point $(\boldsymbol{x}_1^T, \boldsymbol{x}_2^T)^T$ in W whose $Z_1^{\Pi,AD}(\boldsymbol{x}_1, \boldsymbol{x}_2)$ is the least and let it be the next extreme point $(\boldsymbol{x}_{1[i+1]}^T, \boldsymbol{x}_{2[i+1]}^T)^T$. Then, let i := i + 1 and return to step 2. By repeating the procedures, we can obtain an AD-P-Stackelberg solution

$$(\boldsymbol{x}_{1,ADP}^T, \boldsymbol{x}_{2,ADP}^T)^T = (0.00, 12.67, 10.30, 1.36, 0.00, 4.66)^T$$

where

$$Z_1^{\Pi,AD}(\boldsymbol{x}_{1,ADP}, \boldsymbol{x}_{2,ADP}) = 0.0163, \quad Z_2^{\Pi,AD}(\boldsymbol{x}_{1,ADP}, \boldsymbol{x}_{2,ADP}) = 0.0745$$

On the other hand, using the computational method quite similar to that for obtaining an AD-P-Stackelberg solution, we can obtain an AD-N-Stackelberg solution

$$(\boldsymbol{x}_{1,ADN}^T, \boldsymbol{x}_{2,ADN}^T)^T = (4.92, 11.34, 0.47, 3.05, 0.00, 4.85)^T$$

where

$$Z_1^{N,AD}(\boldsymbol{x}_{1,ADN}, \boldsymbol{x}_{2,ADN}) = 0.0287, \quad Z_2^{N,AD}(\boldsymbol{x}_{1,ADN}, \boldsymbol{x}_{2,ADN}) = 0.0338.$$

5 Conclusions

In this paper, assuming noncooperative behavior of the decision makers, computational methods for obtaining Stackelberg solutions to two-level linear programming problems involving fuzzy random variable coefficients have been presented. Considering vague natures of decision makers' judgments, fuzzy goals were introduced into the formulated fuzzy random noncooperative two-level linear programming problems. On the basis of the possibility and necessity measure that each objective function fulfills the corresponding fuzzy goal, the fuzzy random two-level linear programming problems to minimize each objective function with fuzzy random variables were transformed into stochastic twolevel programming problems to maximize the degree of possibility and necessity that each fuzzy goal is fulfilled. Through the use of absolute deviation minimization in stochastic programming, the transformed stochastic two-level programming problems were reduced to deterministic two-level programming problems. For the transformed problems, AD-P- and AD-N-Stackelberg solutions were introduced and computational methods were also presented. It is significant to note here that AD-P- and AD-N-Stackelberg solutions can be obtained through the combined use of the variable transformation method and the Kth best algorithm for two-level linear programming problems. To illustrate the proposed computational methods, a numerical example for obtaining AD-P- and AD-N-Stackelberg solutions was provided. Extensions to other stochastic programming models will be considered elsewhere. Further considerations from the view point of fuzzy random cooperative two-level programming will be required in the near future.

References

- [1] Ammar EE. On solutions of fuzzy random multiobjective quadratic programming with applications in portfolio problem. Information Sciences 2008; 178; 468-484.
- [2] Amouzegar MA, Moshirvaziri K. Determining optimal pollution control policies: an application of bilevel programming. European Journal of Operational Research 1999; 119; 100-120.
- [3] Bard JF, Falk JE. An explicit solution to the multi-level programming problem. Computers and Operations Research 1982; 9; 77-100. (1982)
- [4] Bard JF, Moore JT. A branch and bound algorithm for the bilevel programming problem. SIAM Journal on Scientific and Statistical Computing 1990; 11; 281-292.
- [5] Bialas WF, Karwan MH. Two-level linear programming. Management Science 1984; 30; 1004-1020.
- [6] Birge JR, Louveaux F. Introduction to Stochastic Programming. Springer: London; 1997.
- [7] Calvete HI, Gale C. A note on bilevel linear fractional programming problem. European Journal of Operational Research 2004; 152; 296-299.
- [8] Charnes A, Cooper WW. Programming with linear fractional functionals. Naval Research Logistic Quarterly 1962; 9; 181-186.

- [9] Colson B, Marcotte P, Savard G. A trust-region method for nonlinear bilevel programming: algorithm and computational experience. Computational Optimization and Applications 2005; 30; 211-227.
- [10] Dempe S, Bard JF. Bundle trust-region algorithm for bilinear bilevel programming. Journal of Optimization Theory and Applications 2001; 110; 265-288.
- [11] Faisca NP, Dua V, Rustem B, Saraiva PM, Pistikopoulos EN. Parametric global optimisation for bilevel programming. Journal of Global Optimization 2007; 38; 609-623.
- [12] Fampa M, Barroso LA, Candal D, Simonetti L. Bilevel optimization applied to strategic pricing in competitive electricity markets. Computational Optimization and Applications 2008; 39; 121-142.
- [13] Gil MA, Lopez-Diaz M, Ralescu DA. Overview on the development of fuzzy random variables. Fuzzy Sets and Systems 2006; 157; 2546-2557.
- [14] GÜMÜS ZH, Floudas CA. Global optimization of nonlinear bilevel programming problems. Journal of Global Optimization 2001; 20; 1-31.
- [15] Hansen P, Jaumard B, Savard G. New branch-and-bound rules for liner bilevel programming. SIAM Journal of Scientific and Statistical Computing 1992; 13; 1194-1217.
- [16] Katagiri H, Ishii H, Sakawa M. On fuzzy random linear knapsack problems. Central European Journal of Operations Research 2004; 12; 59-70.
- [17] Katagiri H, Mermri EB, Sakawa M, Kato K, Nishizaki I. A possibilistic and stochastic programming approach to fuzzy random MST problems. IEICE Transaction on Information and Systems 2005; E88-D; 1912-1919.
- [18] Katagiri H, Sakawa M, Ishii H. Fuzzy random bottleneck spanning tree problems using possibility and necessity measures. European Journal of Operational Research 2004; 152; 88-95.
- [19] Katagiri H, Sakawa M, Ishii H. A study on fuzzy random portfolio selection problems using possibility and necessity measures. Scientiae Mathematicae Japonicae 2005; 61; 361-369.
- [20] Katagiri H, Sakawa M, Kato K, Nishizaki I. A fuzzy random multiobjective 0-1 programming based on the expectation optimization model using possibility and necessity measures. Mathematical and Computer Modelling 2004; 40; 411-421.
- [21] Katagiri H, Sakawa M, Kato K, Nishizaki I. Interactive multiobjective fuzzy random linear programming: maximization of possibility and probability. European Journal of Operational Research 2008; 188; 530-539.
- [22] Katagiri H, Sakawa M, Nishizaki I. Interactive decision making using possibility and necessity measures for a fuzzy random multiobjective 0-1 programming problem. Cybernetics and Systems 2006; 37; 59-74.

- [23] Konno H, Shirakawa H, Yamazaki H. A mean-absolute deviation-skewness portfolio optimization model. Annals of Operations Research 1993; 45; 205-220.
- [24] Kruse R, Meyer KD. Statistics with Vague Data. Dordrechut-Holland: D. Riedel Publishing Company; 1987.
- [25] Kwakernaak H. Fuzzy random variables I. definitions and theorems. Information Sciences 1978; 15; 1-29.
- [26] Liang R, Gao J, Iwamura K. Fuzzy random dependent-chance bilevel programming with applications. In: Liu D, Fei S, Hou Z, Zhang H, Sun C (Eds), Advances in Neural Networks - ISNN 2007, Springer-Verlag; Heidelberg; 2007. p. 257–266.
- [27] Liu B. Fuzzy random chance-constrained programming. IEEE Transaction on Fuzzy Systems 2001; 9; 713-720.
- [28] Liu B. Fuzzy random dependent-chance programming. IEEE Transaction on Fuzzy Systems 2001; 9; 721-726.
- [29] Liu YK, Liu B. Fuzzy random variables: a scalar expected value operator. Fuzzy Optimization and Decision Making 2003; 2; 143-160.
- [30] Luhandjula MK. Fuzziness and randomness in an optimization framework. Fuzzy Sets and Systems 1996; 77; 291-297.
- [31] Luhandjula MK. Fuzzy stochastic linear programming: survey and future research directions. European Journal of Operational Research 2006; 174; 1353-1367.
- [32] Luhandjula MK, Gupta M.M. On fuzzy stochastic optimization. Fuzzy Sets and Systems 1996; 81; 47-55.
- [33] Nicholls MG. The applications of non-linear bi-level programming to the aluminium industry. Journal of Global Optimization 1996; 8; 245-261.
- [34] Nishizaki I, Sakawa M. Stackelberg solutions to multiobjective two-level linear programming problems. Journal of Optimization Theory and Applications 1999; 103; 161-182.
- [35] Nishizaki I, Sakawa M. Computational methods through genetic algorithms for obtaining Stackelberg solutions to two-level mixed zero-one programming problems. Cybernetics and Systems: An International Journal 2000; 31; 203-221.
- [36] Nishizaki I, Sakawa M, Katagiri H. Stackelberg solutions to multiobjective two-level linear programming problems with random variable coefficients. Central European Journal of Operations Research 2003; 11; 281-296.
- [37] Puri ML, Ralescu DA. Fuzzy random variables. Journal of Mathematical Analysis and Applications 1986; 114; 409-422.
- [38] Qaio Z, Zhang Y, Wang G-Y. On fuzzy random linear programming. Fuzzy Sets and Systems 1994; 65; 31-49.

- [39] Roghanian E, Sadjadi SJ, Aryanezhad MB. A probabilistic bi-level linear multiobjective programming problem to supply chain planning. Applied Mathematics and Computation 2007; 188; 786-800.
- [40] Rommelfanger H. A general concept for solving linear multicriteria programming problems with crisp, fuzzy or stochastic values. Fuzzy Sets and Systems 2007; 156; 1892-1904.
- [41] Sakawa M. Fuzzy Sets and Interactive Multiobjective Optimization. Plenum Press: New York; 1993.
- [42] Sakawa M. Large Scale Interactive Fuzzy Multiobjective Programming. Physica-Verlag: Heidelberg; 2000.
- [43] Sakawa M. Genetic Algorithms and Fuzzy Multiobjective Optimization. Kluwer Academic Publishers: Boston; 2001.
- [44] M. Sakawa, I. Nishizaki, Cooperative and Noncooperative Multi-Level Programming, Springer, Norwell (in press).
- [45] Shimizu K, Ishizuka Y, Bard JF. Nondifferentiable and Two-Level Mathematical Programming. Kluwer Academic Publishers: Boston; 1997.
- [46] Simaan M, Cruz Jr. JB. On the Stackelberg strategy in nonzero-sum games. Journal of Optimization Theory and Applications 1973; 11; 533-555.
- [47] Slowinski R, Teghem J (Eds). Stochastic Versus Fuzzy Approaches to Multiobjective Mathematical Programming under Uncertainty. Kluwer Academic Publishers: Dordrecht/Boston/London; 1990.
- [48] Stancu-Minasian IM. Stochastic Programming with Multiple Objective Functions. D. Reidel Publishing Company: Dordrecht; 1984.
- [49] Stancu-Minasian IM. Fractional Programming. Kluwer Academic Publishers: Dordrecht/Boston/London; 1992.
- [50] Wang GY, Qiao Z. Linear programming with fuzzy random variable coefficients. Fuzzy Sets and Systems 1993; 57; 295-311.
- [51] White DJ, Anandalingam G. A penalty function approach for solving bi-level linear programs. Journal of Global Optimization 1993; 3; 397-419.
- [52] Xu J, Liu Y. Multi-objective decision making model under fuzzy random environment and its application to inventory problems. Information Sciences 2008; 178; 2899-2914.