

PathTracer: High-sensitivity detection of differential pathway activity in tumours

Ståle Nygård^{1,2,*}, Ole Christian Lingjærde^{1,3,4,*}, Carlos Caldas⁵, Eivind Hovig^{1,6}, Anne-Lise Børresen-Dale³, Aslaug Helland^{3,7,8}, and Vilde D. Haakensen^{3,7,†}

¹Centre for Bioinformatics, Department of Informatics, University of Oslo, Oslo, Norway

²Bioinformatics core facility, Institute for Cancer Research, Oslo University Hospital, Oslo, Norway

³Department of Cancer Genetics, Institute for Cancer Research, Oslo University Hospital, Oslo, Norway

⁴KG Jebsen Centre for B-cell malignancies, Institute for Clinical Medicine, University of Oslo, Norway

⁵Cancer Research UK, Cambridge Research Institute, Li Ka Shing Centre, University of Cambridge, Cambridge, UK

⁶Department of Tumor Biology, Institute for Cancer Research, Oslo University Hospital, Oslo, Norway


⁷Department of Oncology, Oslo University Hospital, Oslo, Norway

⁸Institute of Clinical Medicine, University of Oslo, Oslo, Norway

*These authors contributed equally to this study

†Corresponding author vilde.haakensen@gmail.com

Histogram of PDS vs PTS correlation



Histogram of correlation between pathifier deregulation scores (PDS) and PathTracer scores (PTS) based on the metabric breast cancer dataset.