



Zhang, W. H., Carter, B. M., Day, G. J., Govan, N., Jackson, C., & Perriman, A. W. (2019). Sequential Electrostatic Assembly of a Polymer Surfactant Corona Increases Activity of the Phosphotriesterase arPTE. *Bioconjugate Chemistry*, 30, 2771-2776. <https://doi.org/10.1021/acs.bioconjchem.9b00664>

Peer reviewed version

Link to published version (if available):
[10.1021/acs.bioconjchem.9b00664](https://doi.org/10.1021/acs.bioconjchem.9b00664)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via American Chemical Society at <https://pubs.acs.org/doi/10.1021/acs.bioconjchem.9b00664>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Sequential electrostatic assembly of a polymer surfactant corona increases activity of the phosphotriesterase arPTE.

William H. Zhang¹, Benjamin M. Carter¹, Graham Day¹, Norman Govan², Colin Jackson^{3*}, Adam Perriman^{1*}

¹University of Bristol, School of Cellular and Molecular Medicine. BS8 1TD, United Kingdom.

²Defence Science and Technology Laboratory. SP4 0JQ, United Kingdom.

³Australian National University, Research School of Chemistry. ACT 2601, Australia.

Abstract

We present a new methodology for the generation of discrete molecularly-dispersed enzyme–polymer-surfactant bioconjugates. Significantly, we demonstrate that a more-than three-fold increase in the catalytic efficiency of the diffusion-limited phosphotriesterase arPTE can be achieved through sequential electrostatic addition of cationic and anionic polymer surfactants, respectively. Here, the polymer surfactants assemble on the surface of the enzyme via ion exchange to yield a compact corona. The observed rate enhancement is consistent with a mechanism whereby the polymer-surfactant corona gives rise to a decrease in the dielectric constant in the vicinity of the active site of the enzyme, accelerating the rate-determining product diffusion step. The facile methodology has significant potential for increasing the efficiency of enzymes and could therefore have a substantial positive impact for industrial enzymology.

Main text

Phosphotriesterases (PTEs, also known as organophosphate hydrolases) have evolved to rapidly detoxify organophosphates (OPs). This has attracted significant interest as a potential means to counteract poisoning from OP-based chemical warfare agents and pesticides.(1, 2) These highly toxic compounds are still used as a tool of conflict and covert action, with recent examples including sarin attacks in Syria and the use of a Novichock agent in the United Kingdom.(3) The effectiveness of PTEs are directly dependent on the rate at which they can detoxify these compounds, as well as the lifetime of the enzyme.(4) The real-world performance of enzymes can often be improved through covalent modifications such as PEGylation,(5, 6) or immobilization and entrapment of enzymes at surfaces.(7-9) Indeed, enzyme immobilization can offer great improvements to thermostability, enable enzymes to be integrated into large-scale high-throughput production systems, and crucially, allow these enzymes to be recycled multiple times.(10-12) However, these approaches often impart detrimental effects such as a decrease in enzyme activity, which can preclude an enzyme from practical use.(10, 11, 13, 14) Here, this decrease in activity can result from surface-induced denaturation, obstruction of the active site, or reduction in the rate of substrate/product diffusion.

We have previously developed a methodology to modify protein surfaces *via* electrostatic assembly of a polymer surfactant corona.(15-18) This process involves chemical cationization *via* nucleophilic addition of diamines to the solvent-exposed acidic residues to generate super-cationic proteins, followed by electrostatic grafting of anionic polymer surfactant molecules to the surface. Once formed, these complexes were ionically stable, as verified through analytical ultracentrifugation in the presence of high salt,(17, 19) and similarly, the presence of fatty

substrates such as lipids does not dissociate the enzyme-surfactant complex.(15) Significantly, the resulting hybrid constructs were used to produce a portfolio of new biomaterials that exhibited a wide range of interesting properties, including biologically-active solvent-free biofluids with extreme thermostability,(15, 17, 18) hierarchically self-assembled membranes films with recyclable catalytic activities,(20, 21) and artificial cell membrane-binding proteins with oxygen delivery and extracellular matrix forming capacities.(22, 23) However, as commonly reported for other covalent approaches to protein modification, the chemical cationization required to supercharge the proteins resulted in a significant loss in protein structure and enzymatic activity.(15, 22, 24) Accordingly, we describe a non-covalent approach to generate molecularly dispersed protein–polymer surfactant bioconjugates (**Figure 1**). This new methodology involves sequential electrostatic grafting of PEG-based cationic and anionic surfactants *via* ion exchange to the naturally-occurring charged residues on the surface of the enzyme. This gives rise to monodispersed enzyme–polymer surfactant complexes (denoted [Enzyme][S⁺][S⁻]) that are analogous to previously described bioconjugates,(15, 22) but can be readily produced without the need for covalent modification. Significantly, for the *Agrobacterium radiobacter* phosphotriesterase (arPTE),(25) we observe a more-than three-fold increase in the catalytic efficiency by optimising the composition of the polymer surfactant corona.




Figure 1. Schematic depicting the conjugation process, involving the sequential electrostatic addition of polymer-surfactants. First, cationic surfactants (Ethoquad, red) are complexed to an unmodified protein (arPTE shown above) through electrostatic forces to the solvent exposed anionic amino acids (red sites on the protein). This is then followed by the addition of an anionic surfactant (oxidized IGEPAL, blue), which are likewise complexed to the protein through electrostatic interactions with the solvent exposed cationic amino acids (blue sites on the protein). The result is a surfactant corona that encapsulates the protein.

Kinetic assays of paraoxon hydrolysis using [arPTE][S⁺][S⁻] were performed over a range of anionic polymer surfactant molecular weights, with higher molecular weight surfactants tending towards higher activities (**Figure S1**). Moreover, no activity enhancement was observed after the addition of only the cationic surfactant Ethoquad (**Figure S1**). Reversing the order of addition of the surfactants (*i.e.*, [S⁻][S⁺]) for some surfactants resulted in enzyme aggregation and precipitation, and accordingly, was not pursued further. Previous computational models reveal the formation of micelle-like structures, which limits the interactions between the charged heads groups of the different surfactants and thus facilitates the formation of the corona despite the potential for surfactant-surfactant ionic interactions.⁽²⁶⁾ The greatest enhancement to catalytic activity occurred after the sequential addition of Ethoquad followed by oxidized IGEPAL-890 (**Figures 2 and S2**), which resulted in a two-fold increase in the substrate turnover rate (k_{cat}) when compared to unmodified arPTE ($267 \pm 12 \text{ s}^{-1}$ to $575 \pm 23 \text{ s}^{-1}$). Moreover, this was accompanied by an improvement in the substrate binding affinity (K_{m} ; 56 ± 10 to $36 \pm 6 \text{ }\mu\text{M}$), which translates to a more-than three-fold increase in the specificity constant (catalytic efficiency, $k_{\text{cat}}/K_{\text{m}}$; 4.7×10^6 to $1.6 \times 10^7 \text{ M}^{-1} \text{ S}^{-1}$). Accordingly, this protocol was used for all subsequent experiments, and the resulting bioconjugate is hereafter denoted by [arPTE][S⁺][S⁻].





Figure 2. a) The Michaelis–Menten curve of the enzyme arPTE (blue) compared and [arPTE][S⁺][S⁻] (red) demonstrating a large increase to catalytic rate at room temperature. Assays were performed under the conditions described in the supplementary methods. b) The corresponding k_{cat} and K_{m} of arPTE and [arPTE][S⁺][S⁻] derived from the Michaelis–Menten curve shown in in a).



To verify polymer-surfactant binding to the surface of arPTE, synchrotron radiation small angle X-ray scattering (SR-SAXS) experiments were performed to evaluate the dimensions of the unmodified and surfactant complexed enzyme ([arPTE][S⁺][S⁻]). A 12.6 Å increase in the radial average (r_{ave}) was calculated from the pair-distance distribution function ($P(r)$, **Figure 3A**) of [arPTE][S⁺][S⁻] relative to arPTE, which is comparable to computational models based on the analogous chemically-cationized single-surfactant systems.(26) Additionally, there was a small change in the Porod exponent between arPTE and [arPTE][S⁺][S⁻] (3.8 to 3.5, **Table S1**), which is to be expected due to the addition of flexible surfactant chains. Bead models produced from the SR-SAXS analyses revealed that the size and the shape of the unmodified arPTE is commensurate with published crystal structures of the dimeric enzyme (PDB ID: 2D2J), and that [arPTE][S⁺][S⁻] has an increased size corresponding to the addition of a compact polymer surfactant corona to the dimer (**Figures 3B and S3A-B**). This size increase supports an electrostatically-driven assembly process to form an encapsulating polymer surfactant layer, resulting in discrete, monodispersed surfactant-enzyme dimer complexes.(27)




Figure 3. a) The pair-distance distribution function ($P(r)$) of arPTE (blue) and [arPTE][S⁺][S⁻] (red) obtained through SR-SAXS. The surfactant conjugation process increases the radial average (r_{ave}) of the enzyme from 36.2 Å to 48.8 Å. b) The crystal structure of arPTE (PDB ID: 2D2J, purple) superimposed on the bead-model obtained from SR-SAXS analysis of arPTE (green) and [arPTE][S⁺][S⁻] (blue). The models confirm that the oligomeric state of the enzyme is still maintained after conjugation with the surfactant corona, and that the surfactant creates a consistent shell around the enzyme.

Circular dichroism (CD) measurements revealed superimposable spectra for arPTE and [arPTE][S⁺][S⁻] (**Figure S4A**), which indicated that surfactant complexation did not affect the secondary structure. This suggests that the rate enhancement did not occur *via* fortuitous stabilization of a structural conformation, although conformational changes beyond the resolution of CD cannot be discounted.

The observed catalytic efficiency of many enzymes are dependent on other extraneous contributions, such as surface-induced deactivation.(28) This effect can be significant when

working with enzymes at very low concentrations, as is necessary to collect the linear initial rate for highly efficient enzymes such as arPTE. Experimentally, these adsorption effects can be prevented with the addition of bovine serum albumin (BSA) to block non-specific protein binding. (29) Accordingly, both arPTE and [arPTE][S⁺][S⁻] were assayed against paraoxon with and without the presence of 0.1% w/v BSA. The addition BSA did not initially result in a change in the catalytic activity in either system (assay periods < 10 minutes) (**Figure 4A**). However, assays performed over a four-hour period showed a gradual decay in the activity of arPTE, which did not occur for [arPTE][S⁺][S⁻], [arPTE][S⁺][S⁻] with 0.1% w/v BSA, and arPTE with 0.1% w/v BSA (**Figure 4B**). To investigate if the loss in activity of the native enzyme was due to slow unfolding (rather than surface-induced deactivation), a time-dependent circular dichroism thermostability study of arPTE and [arPTE][S⁺][S⁻] was performed (**Figures S4B-C**). While a high-temperature CD gradient did show that [arPTE][S⁺][S⁻] had a slight enhancement to overall thermostability, an isothermal time-course experiment at 40 °C showed that no slow denaturation of either [arPTE][S⁺][S⁻] or arPTE occurs even at temperatures elevated slightly above ambient conditions. Accordingly, this data supports a scenario where either the surfactant corona or BSA stabilised the enzyme, and where the presence of BSA did not enhance the activity of the enzyme, but rather, prevented the loss of activity over time. Moreover, this loss of activity would only be significant for assays conducted over long time periods, and therefore could only represent a small fraction of the improvement in the catalytic efficiency observed for [arPTE][S⁺][S⁻].




Figure 4. a) The normalized catalytic rate (k_{cat}) of arPTE and [arPTE][S⁺][S⁻] at room temperature with and without the presence of 0.1% w/v BSA, showing that BSA itself does not appear to directly influence catalytic rate. b) Activity of a stock of 4 nM enzyme over time at room temperature, showing that hydrophobic denaturation can significantly affect the observed rate of arPTE (blue) over extended periods of time. arPTE with BSA (red), [arPTE][S⁺][S⁻] (green) and [arPTE][S⁺][S⁻] with BSA (purple) do not exhibit this loss in activity. All assays were performed as described in the supplementary methods in a buffer of 30 mM HEPES, 100 μ M cobalt chloride at pH 8.0.

Previous studies of PTEs have established that the rate limiting step of these high turnover rate enzymes is the diffusion of the reaction product away from the active site.(30) It is therefore possible that the formation of a compact amphiphilic polymer-surfactant corona may accelerate the diffusion step *via* product partitioning through a reduction in the local dielectric constant at the active site. This mechanism is supported by the conserved tertiary and secondary structure reported above (verified through SAXS and CD respectively), and the positive correlation between the molecular weight of the surfactant and the enzymatic activity. This is not unprecedented, as an increase in activity has previously been noted post-modification of PTEs,(31) with an in-depth study of PTE nanoparticle complexes also showing an increase in activity.(32, 33) The authors attributed this to an increase in the diffusion of the product away from the active site, promoted by a change in the immediate microenvironment around the enzyme. Using the same theoretical model as Breger *et al.* that considers changes in substrate and product on and off rates,(32) we show that our experimentally observed data is consistent with an increase to the product release/diffusion step, rather than a substrate binding step (**Figure S5**). Finally, we observe that compared to paraoxon, the magnitude of the rate enhancement is larger for coumaphos (**Figures S6A-B**), a substrate with a more hydrophobic product, further supporting the theory that diffusion of hydrophobic products is better enabled by the amphiphilic corona. Enzyme enhancement through substrate/product partitioning has been previously described, but through enzyme immobilization,(34) thus the retention of enzyme mobility is an important outcome. However, as alterations to the surface of arPTE can have impact on activity,(30, 35) we cannot discount the possibility that there is some fortuitous enhancement of conformational mobility, resulting in an increase in the rate of product release. An additional consideration is a change in the oligomeric equilibrium between the monomer and dimer influencing the catalytic rate, however as we see a difference in the rate enhancement between different substrates, this is unlikely.

To further probe the effect of polymer surfactant complexation on enzyme structure and function, the dual surfactant complexation method was applied to three other PTEs: pdPTE-C23, α -Esterase7 (α Est7) and DFPase (**Figure S7A-C**).(36-38) Phosphotriesterase-C23 (pdPTE-C23) is a PTE mutant from *Pseudomonas diminuta* designed for greater catalytic efficiency against the G and V series nerve agents.(36) α Est7 is a carboxylesterase that has evolved independently to break down a number of organophosphate pesticides and is structurally distinct from the PTE enzyme superfamily.(37) Lastly, DFPase, found in *Loligo vulgaris*, is capable of hydrolysing diisopropylfluorophosphate (DFP), and like α Est7, is structurally distinct.(38) All three enzymes successfully formed enzymatically active polymer surfactant-complexes through the dual-surfactant method. As with [arPTE][S⁺][S⁻], SR-SAXS confirmed the formation of discrete enzyme-polymer surfactant complexes with compact coronas (**Figures S7D-F**). Interestingly, polymer-surfactant complexation resulted in a decrease in k_{cat} and an increase in K_m for pdPTE-C23 and α Est7. As α Est7 has evolved to hydrolyse fatty ester substrates akin to the hydrophobic tails of the surfactant, this is likely a consequence of steric inhibition of the active site.(15, 24) In a similar fashion, pdPTE-C23 has an engineered mutation of a phenylalanine residue to a glutamate (F132E) at the active site,(36) which would promote inhibition *via* electrostatic binding of S⁺. To confirm that the observed inhibition was not due to denaturation and was a result of active site inhibition, we also assayed the activity after treating pdPTE-C23 with uncharged surfactants. As expected, at high surfactant concentrations (1% w/v of assay) pdPTE-C23 showed inhibition comparable to [pdPTE-C23][S⁻] (**Figure S8**). However after dialysis, activity was restored to pdPTE-C23 treated with uncharged surfactants, but not for [pdPTE-C23][S⁻]. This demonstrates that the decreased rate is through active site inhibition by the surfactant, and also verifies that the surfactant corona complex is maintained even at dilute concentrations. Finally, for DFPase we observed no change in affinity or rate between the unmodified enzyme and the bioconjugate

[DFPase][S⁺][S⁻]. As the rate limiting step of DFPase is the dissociation of fluorine from the phosphorous during the catalytic cycle,(38) the lack of rate enhancement (or decrease) observed here supports our proposed mechanism of improved product off rate.

In conclusion, we have demonstrated that our electrostatic dual surfactant conjugation methodology can be readily applied to generate aqueous molecular dispersion of active enzyme-polymer surfactant complexes without the need for covalent modifications. Moreover, we show that the approach can be tuned to enhance the catalytic efficiency of arPTE by more than three-fold, and we reconcile this improvement with an acceleration in the product diffusion rate *via* hydrophobic partitioning. We also show that the surface-bound polymer surfactant corona protects arPTE from surface-induced deactivation. Although we could readily apply our methodology to other PTEs to give catalytically active complexes, it is clear that rate enhancement is dependent on parameters such as catalytic mechanism, surface charge, hydrophobicity and active site geometry. In principle, the use of amphiphilic coronas is compatible with other beneficial enzyme modifications such as immobilization. The covalent linkage of the surfactant or the enzyme to a surface may impart properties such as improved selectivity,(34) and may enhance the enzyme beyond what is achievable through simple immobilization alone, but at the cost of enzyme mobility. With the correct selection of polymer-surfactants and rational enzyme design, it should be possible to improve the performance of industrially-relevant enzymes under both aqueous and non-aqueous conditions.(39, 40)

Supporting Information

The Supporting Information is available free of charge on the

ACS Publications website at DOI: XXX.

Contains experimental methods such as protein expression, synthesis and complexation, and assay procedures, as well as supplementary figures and tables. (link)

Author information

Corresponding author: *chawp@bristol.ac.uk; *cjackson@rsc.anu.edu.au

Acknowledgements

This research was funded by the EPSRC (EP/N026586/1) that was awarded in collaboration with the Defence Science and Technology Laboratory (DSTL).

We are grateful for the time allotment from Diamond, which allowed SR-SAXS to be performed at the Diamond Light Source I22 beamline, and SEC-SR-SAXS to be performed using the Diamond B21 mail-in service. We would like to thank Jayne Ede and Ian Shortman from the Defence Science and Technology Laboratory (DSTL) for intellectual input and discussion. We would also like to thank Eleanor Campbell and Galen Corey (From the Australian National University) for supplying the arPTE and α Est7 plasmids respectively.

References

- (1) Tsai, P. C., Fox, N., Bigley, A. N., Harvey, S. P., Barondeau, D. P., and Raushel, F. M. (2012) Enzymes for the homeland defense: optimizing phosphotriesterase for the hydrolysis of organophosphate nerve agents. *Biochemistry* 51, 6463-75.
- (2) Jacquet, P., Daude, D., Bzdrenga, J., Masson, P., Elias, M., and Chabriere, E. (2016) Current and emerging strategies for organophosphate decontamination: special focus on hyperstable enzymes. *Environ. Sci. Pollut. Res. Int.* 23, 8200-18.

- (3) Vale, J. A., Marrs, T. O., and Maynard, R. C. (2018) Novichok: a murderous nerve agent attack in the UK. *Clin. Toxicol. (Phila)*. 56, 1093-1097.
- (4) Gupta, R. D., Goldsmith, M., Ashani, Y., Simo, Y., Mullokandov, G., Bar, H., Ben-David, M., Leader, H., Margalit, R., Silman, I., *et al.* (2011) Directed evolution of hydrolases for prevention of G-type nerve agent intoxication. *Nat. Chem. Biol.* 7, 120-5.
- (5) Swierczewska, M., Lee, K. C., and Lee, S. (2015) What is the future of PEGylated therapies? *Expert Opin. Emerg. Drugs* 20, 531-6.
- (6) Dozier, J. K., and Distefano, M. D. (2015) Site-Specific PEGylation of Therapeutic Proteins. *Int. J. Mol. Sci.* 16, 25831-64.
- (7) Rabuka, D. (2010) Chemoenzymatic methods for site-specific protein modification. *Curr. Opin. Chem. Biol.* 14, 790-6.
- (8) Zhang, Y. F., Ge, J., and Liu, Z. (2015) Enhanced Activity of Immobilized or Chemically Modified Enzymes. *ACS Catalysis* 5, 4503-4513.
- (9) Denard, C. A., Ren, H., and Zhao, H. (2015) Improving and repurposing biocatalysts via directed evolution. *Curr. Opin. Chem. Biol.* 25, 55-64.
- (10) DiCosimo, R., McAuliffe, J., Poulouse, A. J., and Bohlmann, G. (2013) Industrial use of immobilized enzymes. *Chem. Soc. Rev.* 42, 6437-74.
- (11) Sheldon, R. A., and Brady, D. (2018) The limits to biocatalysis: pushing the envelope. *Chem. Commun. (Camb.)* 54, 6088-6104.
- (12) Sheldon, R. A., and Woodley, J. M. (2018) Role of Biocatalysis in Sustainable Chemistry. *Chem. Rev.* 118, 801-838.
- (13) Lawrence, P. B., and Price, J. L. (2016) How PEGylation influences protein conformational stability. *Curr. Opin. Chem. Biol.* 34, 88-94.
- (14) Keefe, A. J., and Jiang, S. (2011) Poly(zwitterionic)protein conjugates offer increased stability without sacrificing binding affinity or bioactivity. *Nat Chem* 4, 59-63.
- (15) Brogan, A. P. S., Sharma, K. P., Perriman, A. W., and Mann, S. (2014) Enzyme activity in liquid lipase melts as a step towards solvent-free biology at 150 °C. *Nat. Commun.* 5, 5058.
- (16) Gallat, F. X., Brogan, A. P., Fichou, Y., McGrath, N., Moulin, M., Hartlein, M., Combet, J., Wuttke, J., Mann, S., Zaccai, G., *et al.* (2012) A polymer surfactant corona dynamically replaces water in solvent-free protein liquids and ensures macromolecular flexibility and activity. *J. Am. Chem. Soc.* 134, 13168-71.
- (17) Perriman, A. W., Colfen, H., Hughes, R. W., Barrie, C. L., and Mann, S. (2009) Solvent-free protein liquids and liquid crystals. *Angew. Chem. Int. Ed. Engl.* 48, 6242-6.
- (18) Perriman, A. W., Brogan, A. P., Colfen, H., Tsoureas, N., Owen, G. R., and Mann, S. (2010) Reversible dioxygen binding in solvent-free liquid myoglobin. *Nat Chem* 2, 622-6.
- (19) Xiao, W. J., Green, T. I. P., Liang, X. W., Delint, R. C., Perry, G., Roberts, M. S., Le Vay, K., Back, C. R., Ascione, R., Wang, H. L., *et al.* (2019) Designer artificial membrane binding proteins to direct stem cells to the myocardium. *Chem. Sci.* 10, 7610-7618.
- (20) Sharma, K. P., Collins, A. M., Perriman, A. W., and Mann, S. (2013) Enzymatically active self-standing protein-polymer surfactant films prepared by hierarchical self-assembly. *Adv. Mater.* 25, 2005-10.
- (21) Farrugia, T., Perriman, A. W., Sharma, K. P., and Mann, S. (2017) Multi-enzyme cascade reactions using protein-polymer surfactant self-standing films. *Chem. Commun. (Camb.)* 53, 2094-2097.
- (22) Armstrong, J. P. K., Shakur, R., Horne, J. P., Dickinson, S. C., Armstrong, C. T., Lau, K., Kadiwala, J., Lowe, R., Seddon, A., Mann, S., *et al.* (2015) Artificial membrane-binding proteins stimulate oxygenation of stem cells during engineering of large cartilage tissue. *Nat. Commun.* 6, 7405.
- (23) Deller, R. C., Richardson, T., Richardson, R., Bevan, L., Zampetakis, I., Scarpa, F., and Perriman, A. W. (2019) Artificial cell membrane binding thrombin constructs drive in situ fibrin hydrogel formation. *Nat. Commun.* 10, 1887.

- (24) Perez, B., Coletta, A., Pedersen, J. N., Petersen, S. V., Periolo, X., Pedersen, J. S., Sessions, R. B., Guo, Z., Perriman, A., and Schiott, B. (2018) Insight into the molecular mechanism behind PEG-mediated stabilization of biofluid lipases. *Sci. Rep.* **8**, 12293.
- (25) Ely, F., Hadler, K. S., Gahan, L. R., Guddat, L. W., Ollis, D. L., and Schenk, G. (2010) The organophosphate-degrading enzyme from *Agrobacterium radiobacter* displays mechanistic flexibility for catalysis. *Biochem. J.* **432**, 565-73.
- (26) Brogan, A. P., Sessions, R. B., Perriman, A. W., and Mann, S. (2014) Molecular dynamics simulations reveal a dielectric-responsive coronal structure in protein-polymer surfactant hybrid nanoconstructs. *J. Am. Chem. Soc.* **136**, 16824-31.
- (27) Zhang, Y., Patil, A. J., Perriman, A. W., and Mann, S. (2013) Enhanced catalytic activity in organic solvents using molecularly dispersed haemoglobin-polymer surfactant constructs. *Chem. Commun. (Camb.)* **49**, 9561-3.
- (28) Brode, P. F., Erwin, C. R., Rauch, D. S., Lucas, D. S., and Rubingh, D. N. (1994) Enzyme Behavior at Surfaces - Site-Specific Variants of Subtilisin Bpn' with Enhanced Surface Stability. *J. Biol. Chem.* **269**, 23538-23543.
- (29) Goebel-Stengel, M., Stengel, A., Tache, Y., and Reeve, J. R., Jr. (2011) The importance of using the optimal plasticware and glassware in studies involving peptides. *Anal. Biochem.* **414**, 38-46.
- (30) Jackson, C. J., Foo, J. L., Tokuriki, N., Afriat, L., Carr, P. D., Kim, H. K., Schenk, G., Tawfik, D. S., and Ollis, D. L. (2009) Conformational sampling, catalysis, and evolution of the bacterial phosphotriesterase. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 21631-6.
- (31) Kim, M., Gkikas, M., Huang, A., Kang, J. W., Suthiwangcharoen, N., Nagarajan, R., and Olsen, B. D. (2014) Enhanced activity and stability of organophosphorus hydrolase via interaction with an amphiphilic polymer. *Chem. Commun. (Camb.)* **50**, 5345-8.
- (32) Breger, J. C., Ancona, M. G., Walper, S. A., Oh, E., Susumu, K., Stewart, M. H., Deschamps, J. R., and Medintz, I. L. (2015) Understanding How Nanoparticle Attachment Enhances Phosphotriesterase Kinetic Efficiency. *ACS Nano* **9**, 8491-503.
- (33) Hondred, J. A., Breger, J. C., Garland, N. T., Oh, E., Susumu, K., Walper, S. A., Medintz, I. L., and Claussen, J. C. (2017) Enhanced enzymatic activity from phosphotriesterase trimer gold nanoparticle bioconjugates for pesticide detection. *Analyst* **142**, 3261-3271.
- (34) Rodrigues, R. C., Ortiz, C., Berenguer-Murcia, A., Torres, R., and Fernandez-Lafuente, R. (2013) Modifying enzyme activity and selectivity by immobilization. *Chem. Soc. Rev.* **42**, 6290-307.
- (35) Campbell, E., Kaltenbach, M., Correy, G. J., Carr, P. D., Porebski, B. T., Livingstone, E. K., Afriat-Jurnou, L., Buckle, A. M., Weik, M., Hoffelder, F., *et al.* (2016) The role of protein dynamics in the evolution of new enzyme function. *Nat. Chem. Biol.* **12**, 944-950.
- (36) Cherny, I., Greisen, P., Jr., Ashani, Y., Khare, S. D., Oberdorfer, G., Leader, H., Baker, D., and Tawfik, D. S. (2013) Engineering V-type nerve agents detoxifying enzymes using computationally focused libraries. *ACS Chem. Biol.* **8**, 2394-403.
- (37) Jackson, C. J., Liu, J. W., Carr, P. D., Younus, F., Coppin, C., Meirelles, T., Lethier, M., Pandey, G., Ollis, D. L., Russell, R. J., *et al.* (2013) Structure and function of an insect alpha-carboxylesterase (alphaEsterase7) associated with insecticide resistance. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 10177-82.
- (38) Wymore, T., Field, M. J., Langan, P., Smith, J. C., and Parks, J. M. (2014) Hydrolysis of DFP and the nerve agent (S)-sarin by DFPase proceeds along two different reaction pathways: implications for engineering bioscavengers. *J. Phys. Chem. B* **118**, 4479-89.
- (39) Klibanov, A. M. (2001) Improving enzymes by using them in organic solvents. *Nature* **409**, 241-6.
- (40) Ke, T., and Klibanov, A. M. (1998) On enzymatic activity in organic solvents as a function of enzyme history. *Biotechnol. Bioeng.* **57**, 746-750.

