

## Research Article

# Potential Antioxidant Anthraquinones Isolated from *Rheum emodi* Showing Nematicidal Activity against *Meloidogyne incognita*

Brijesh Tripathi,<sup>1,2</sup> Rohit Bhatia,<sup>3,4</sup> Alka Pandey,<sup>4</sup> Jitender Gaur,<sup>5</sup> Gautam Chawala,<sup>6</sup> Suresh Walia,<sup>1</sup> Eun Ha Choi,<sup>7</sup> and Pankaj Attri<sup>7</sup>

<sup>1</sup> Division of Agricultural Chemicals, Indian Agricultural Research Institute, New Delhi 110012, India

<sup>2</sup> Department of Science and Technology, New Delhi 110016, India

<sup>3</sup> Department of Chemistry, Institute of Home Economics, Hauz Khas, New Delhi 110016, India

<sup>4</sup> Department of Chemistry, University of Delhi, New Delhi 110 007, India

<sup>5</sup> Council of Scientific and Industrial Research-National Physical Laboratory, Dr. K. S. Krishnan Marg, New Delhi 110012, India

<sup>6</sup> Division of Nematology, Indian Agricultural Research Institute, New Delhi 110012, India

<sup>7</sup> Plasma Bioscience Research Center and Department of Electrical and Biological Physics, Kwangwoon University, Seoul 139-701, Republic of Korea

Correspondence should be addressed to Rohit Bhatia; rohit\_bhatia10@yahoo.co.in and Pankaj Attri; chem.pankaj@gmail.com

Received 16 August 2014; Accepted 28 October 2014; Published 25 November 2014

Academic Editor: Iciar Astiasaran

Copyright © 2014 Brijesh Tripathi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Antioxidant and nematicidal properties were evaluated for *R. emodi* extractives which are extracted by standardizing and adopting accelerated solvent extraction (ASE) method along with traditional Soxhlet extraction. The extracted material was separated using flash chromatography and the separation conditions and solvents were standardized for the extracted plant constituents. The purity was detected by using analytical reverse phase high pressure liquid chromatography (HPLC). LC-MS/MS detection in the direct infusion mode of the isolated, purified products afforded four anthraquinones, characterized by their infrared spectra (IR) and <sup>1</sup>H spectra as chrysophanol, physcion, emodin, and aloe-emodin. Five anthraquinone glucoside derivatives and piceatannol-3-O-β-d-glucopyranoside have also been detected from the extracted product. During *in vitro* evaluation the antioxidant potential of methanolic crude extract (CE1) was the highest, followed by ethyl acetate crude extract (CE2) and chloroform extract (CE3) in DPPH radical scavenging activity. The CE1 also demonstrated outstanding nematicidal activity as compared with other extracts, pure anthraquinones, and even positive control azadirachtin. The study conclusively demonstrated the antioxidant potential of *R. emodi* extracts and also its ability in extenuating the *Meloidogyne incognita* (root-knot nematode). The bioassay results can be extrapolated to actual field condition and clinical studies.

## 1. Introduction

*Rheum emodi* (family: Polygonaceae), commonly known as Indian or Himalayan rhubarb, is a perennial herb and a potential source of bioactive molecules against pests and diseases. It is distributed in the temperate and subtropical alpine region of the Himalayas. Ethnomedical uses of *Rheum emodi* have been recorded from India, China, Nepal, and Pakistan for 57 different types of ailments [1, 2]. *Rheum emodi* contains different secondary metabolites that are categorized as

anthraquinones, stilbenes, anthrones, chromones, oxantrone ethers and esters, flavonoids, carbohydrates, lignans, phenols, and sterols. The major constituents of this herb are hydroxyanthraquinones and their glycosides which are responsible for various physiological activities. Some of the bioactive compounds identified in these plants extracts include rhein, emodin, aloe-emodin, chrysophanol, physcion, and their glycosides [3]. Other complex compounds have also been identified as torachryson-8-O-β-d-glucopyranoside, sulphated emodin glucoside, and acetylated chrysophanol glucoside

[4]. Revandchinone-1, 3, and 4 exhibited significant antifungal activity against *Aspergillus niger* and *Rhizopus oryzae* and antibacterial activity against gram positive (*Bacillus subtilis*, *Bacillus sphaericus*, and *Staphylococcus aureus*) and gram negative (*Klebsiella aerogenes*, *Chromobacterium violaceum*, and *Pseudomonas aeruginosa*) bacteria [5]. Most of the compounds have been evaluated for their medicinal activities. Hydroxyl anthraquinones and their glycosides have been found effective against animal pathogens, namely, *Candida albicans*, *Aspergillus fumigatus*, *Cryptococcus neoformans*, and *Trichophyton mentagrophytes* [6, 7]. The petroleum ether and chloroform extracts of the rhizomes have been reported to exhibit moderate antifungal and antibacterial activities; the benzene extract of *R. emodi* inhibited the growth of *H. pylori* [8].

However, only few reports are available on the agricultural pest control use of *R. emodi* extracts including nematocidal activity which is also far and fewer. So, the present study is undertaken to explore the possibility of using *R. emodi* extracts and isolated bioactive molecules in nematode control and antioxidant potential evaluation by DPPH radical scavenging activity. In the literature several methods have been reported about efficient extractions and HPLC analysis of *R. emodi* [9]. But, we have standardized accelerated solvent extraction (ASE) method for efficient extraction of bioactive constituents of *R. emodi* and compared it with Soxhlet extraction method. This was followed by separation on flash chromatography and purity check by analytical reverse phase high pressure liquid chromatography (HPLC). LC-MS/MS method in ESI mode has been employed in the direct infusion mode for the detection and confirmation of the structure of isolated, purified products which were further characterized by their spectroscopic details using infrared (IR) and  $^1\text{H}$  NMR spectroscopy.

## 2. Materials and Methods

**2.1. Chemicals and Plant Materials.** Laboratory and analytical grade chemicals, reagents, and solvents were locally procured. Quercetin, gallic acid, and 1,1-diphenyl-2-picrylhydrazyl (DPPH) were procured from Merck India Ltd. Authentic samples of the rhizomes of *Rheum emodi* were procured, Delhi [10]. The nematode *Meloidogyne incognita* (root-knot nematode) cultures were obtained, cultured, and identified from the farms of the Division of Nematology, IARI, New Delhi.

### 2.2. Extraction and Isolation

**2.2.1. Accelerated Solvent Extraction.** The extraction was carried out using a Dionex ASE 300 (Sunnyvale, CA, USA) accelerated solvent extractor. A 50 g portion of the root powder of *R. emodi* was packed tightly in a 100 mL stainless steel vessel and extracted with methanol at different temperatures (40, 50, and 60°C). The extractions were performed under pressure at 10.34 Mpa, with 5 min equilibration, 5 min static time, 60% purge volume, and a 60 sec purge for a total of three cycles. About 80–100 mL of solvent was collected in each extraction. The solvent was removed under reduced pressure

at 40°C in a rotary evaporator (Heidolph, Germany) to obtain dark yellow coloured viscous residue. All the operations were repeated thrice and the extracts were kept at –20°C until analysis.

**2.2.2. Soxhlet Extraction.** *R. emodi* rhizome powder (150 gm) was Soxhlet extracted with methanol (500 mL) for 8 hours. It was concentrated under reduced pressure and then partitioned with chloroform and ethyl acetate. The chloroform extract gave positive test for anthraquinones and showed four bands on the TLC plate. Four bands were scrapped and eluted with methanol. The purity of individual bands was checked by HPLC.

### 2.3. Chromatography and Spectroscopy

**2.3.1. Thin Layer Chromatography (TLC).** Activated silica gel coated plates (6 cm × 20 cm) of 0.25 mm thickness were used. Preparative TLC was performed on silica gel coated glass plates (20 cm × 20 cm) of 0.25 mm thickness. The preparative plates were developed in different solvent combinations like (i) petroleum ether : ethyl acetate : formic acid (2 : 8 : 0.1) and (ii) hexane : chloroform : acetic acid (9 : 1 : 0.1). Plates were air-dried and visualized with iodine vapor. The individual bands were scraped and extracted with methanol and the solvent evaporated under vacuo to obtain the compounds.

**2.3.2. Flash Chromatography.** Methanol fraction of *R. emodi* was also fractioned using flash chromatography (Biotage, Isolera<sup>R</sup>). The elution was done automatically by programming the instrument with hexane-chloroform ( $\text{CHCl}_3$ ) [(3:1), (2:1), (1:1), (1:2) v/v],  $\text{CHCl}_3$  and ethyl acetate (EtOAc), and  $\text{CHCl}_3$ -methanol (MeOH) [(9:1), (8:2), (7:3) v/v]. The fractions of same  $R_f$  (TLC) were combined and concentrated in a rotary evaporator at 40°C to obtain sufficient quantity of pure anthraquinones like emodin, aloe-emodin, physcion, chrysophanol, and so forth.

**2.3.3. High Performance Liquid Chromatography (HPLC).** Analytical reverse phase HPLC (WATERS e2695) fitted with quaternary gradient pump, autosampler, controller Empower 2 software, waters 2998 PDA detector, and waters spherisorb<sup>R</sup> 5  $\mu\text{m}$  ODS2 (4.6 × 250 mm) analytical column was used for all the analysis. Anthraquinones were separated under gradient elution at a flow rate of 1 mL min<sup>-1</sup> using a mobile phase of ACN : methanol (95 : 5) and water (0.2%  $\text{CH}_3\text{COOH}$  v/v). A 20  $\mu\text{L}$  volume of sample was injected each time and peaks were detected at  $\lambda_{\text{max}}$  of 290. The retention time ( $R_t$ ) for each compound was recorded. Water used in HPLC analysis was obtained from Millipore water purifier system with resistivity of 18.2 M $\Omega$ cm.

**2.3.4. Fourier Transform Infrared Spectroscopy (FT-IR).** FT-IR was performed by Bruker Alpha making KBr palates, using Hydraulic press and the data processed using Opus\_65 software.

**2.3.5. Nuclear Magnetic Resonance Spectroscopy.** The  $^1\text{H}$  spectra of the compounds were recorded on Bruker (400 MHz)




FIGURE 1: ESI-MS total ion chromatogram of crude *R. emodi* MeOH fraction.

instrument. Samples were dissolved in deuterated chloroform ( $\text{CDCl}_3$ ), dimethyl sulfoxide (DMSO), or carbon tetrachloride ( $\text{CCl}_4$ ) as per the requirement. Tetramethylsilane (TMS) was used as an internal standard. Chemical shifts were recorded in  $\delta$  (ppm) relative to TMS.

**2.3.6. Electrospray Mass Spectroscopy (ESI-MS).** Liquid chromatography coupled with mass spectrometry (LC-ESI-MS) is a powerful technique to analyze complex botanical extracts [11–15]. It facilitates rapid and accurate identification of chemical compounds, especially when a pure standard is not available. The technique has been extensively used in the past for the analysis of phenolic compounds [16]. The most active extract of *R. emodi* was analyzed for its active constituents and their structures confirmed by electrospray mass spectrometry (ESI-MS) in the direct infusion mode.

Mass spectrometry was carried out using Thermo Electron spectrometer (Thermo Electron Corporation, USA). Detection of mass was done by electron-spray ionization (ESI) source with Finnigan LCQ tune plus program fitted with MAX-detector. Xcalibur software was used for the purpose of identification, quantification, and fragmentation of required masses. The MS parameters optimized in direct infusion mode were spray voltage 3.5 to 5 kV, sheath gas flow rate  $10 \text{ mL min}^{-1}$ , auxiliary gas flow rate  $5 \text{ mL min}^{-1}$ , spray current 0.5, capillary temperature  $225^\circ\text{C}$ , capillary voltage 20–35 V, and tube lens offset 40–65. The mass spectra were recorded in negative ion mode (Figure 1).

**2.4. Characterization of *R. emodi* Anthraquinones.** The preparative thin layer chromatography of the chloroform extracts afforded four compounds: chrysophanol, physcion, emodin, and aloe-emodin. The bands on the TLC were scrapped and eluted with methanol. The purity of the




FIGURE 2: HPLC of isolated pure anthraquinones.

individual bands was checked by TLC followed by their analysis by HPLC (Figure 2).

The spectral characteristics of pure anthraquinones are as follows.

**Chrysophanol.** The first compound was crystallized from chloroform-acetone as yellow powder [m.p.  $194^\circ\text{C}$ , Rf 0.9 in hexane + ethyl acetate + formic acid (2 + 8 + 0.1 by volume)].  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 2.45 (3-H, s, H-3), 7.15 (1H, d, H-2), 7.59 (1H, d, H-4), 7.30 (1H, dd, H-7), 7.86 (1H, dd, H-5), 7.69 (1H, d, H-6), 12.03 (1H, s, OH-1), 12.14 (1H, s, OH-8); UV ( $\lambda_{\text{max}}$  MeOH): 228, 256, 279, 287, 429 nm; IR ( $\nu_{\text{MeOH}}$ ): 3470, 3080, 2940, 1680, 1630, 1590, 1460, 1470, 1260, 1210  $\text{cm}^{-1}$ ; ESI-MS:  $m/z$  253.3  $[\text{M} - \text{H}]^-$ , 225.3  $[\text{M} - 28]^-$ , 209.1  $[\text{M} - 28 - 15]^-$ .

**Physcion.** The second compound was crystallized from chloroform to give yellow needles [m.p.  $204^\circ\text{C}$ , Rf 0.8 in hexane + ethyl acetate + formic acid (2 + 8 + 0.1 by volume)].  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 12.13 (1H, s, OH-1), 12.33 (1H, s, OH-8), 6.69 (1H,

*d*, H-2), 7.09 (1H, *d*, H-7), 7.37 (1H, *s*, H-4), 7.63 (1H, *d*, H-5), 2.46 (3H, H-3), 3.94 (3-H, H-6); **IR** ( $\nu$  MeOH): 1675, 1630, 1612, 1570, 1380  $\text{cm}^{-1}$ ; ESI-MS:  $m/z$  283.3  $[\text{M} - \text{H}]^-$ , 269.3  $[\text{M} - 15]^-$ , 253.3  $[\text{M} - 31]^-$  and  $m/z$  235.2  $[\text{M} - 31-18]^-$ .

**Emodin**. The third compound was crystallized from diethyl ether + ethyl acetate (80 + 20) to give pinkish powder [m.p. 258°C, Rf 0.40 in hexane + ethyl acetate + formic acid (2 + 8 + 0.1 by volume)]. <sup>1</sup>H-NMR ( $\text{CDCl}_3$ )  $\delta$ : 12.30 (1H, *s*, OH-1), 12.80 (1H, *s*, OH-8), 6.69 (1H, *d*, H-2), 7.09 (1H, *d*, H-7), 7.37 (1H, *s*, H-4), 7.63 (1H, *d*, H-5), 2.46 (3H, H-3), 5.65 (1H, H-6); **IR** ( $\nu$  MeOH): 3390, 1675, 1631, 2950  $\text{cm}^{-1}$ ; ESI-MS:  $m/z$  269.2  $[\text{M} - \text{H}]^-$ , 241.3  $[\text{M} - 28]^-$  and  $m/z$  225.2  $[\text{M} - 28-17]^-$ .

**Aloe-Emodin**. The fourth compound was crystallized from chloroform: acetone (3 + 1) as yellow needles [m.p. 220°C, Rf 0.30 in hexane + ethyl acetate + formic acid (2 + 8 + 0.1 by volume)]. <sup>1</sup>H-NMR ( $\text{CDCl}_3$ )  $\delta$ : 12.08 (1H, *s*, OH-1), 12.10 (1H, *s*, OH-8), 7.32 (1H, *d*, H-2), 7.35 (1H, *dd*, H-7), 7.69 (1H, *d*, H-4), 7.80 (1H, *d*, H-6), 7.85 (1H, *dd*, H-5), 4.35 (2H, H-3), 5.35 (1H, OH-3); ESI-MS:  $m/z$  269.3  $[\text{M} - \text{H}]^-$ ,  $m/z$  241.2  $[\text{M} - 28]^-$ ; **IR** ( $\nu$  MeOH): 3300, 3050, 2890, 1650, 1610, 1590, 1250  $\text{cm}^{-1}$ .

**2.5. Extraction of Nematodes from Plant Root.** *M. incognita* infected plant material was collected from the glass house of Division of Nematology, I.A.R.I., New Delhi. The infected roots of brinjal (cv. Pusa Purple Long) were washed thoroughly with tap water to remove soil adhering to the roots. *Meloidogyne* egg mass was collected from the brinjal under stereoscopic microscope. The egg masses were kept on double layered tissue paper supported by wire mesh and kept 2-3 days in incubator at 25–30°C and 70% RH for hatching. The number of freshly hatched second stage ( $J_2$ s) juveniles was counted per 1 mL aliquot of distilled water.

Fresh nematode suspension (1 mL) was taken in a square counting dish (marked with 24 equal squares) under stereoscopic binocular microscope. The total number of nematodes in one mL was determined by counting the root-knot juveniles in each square. This process was repeated for five times and the average was taken. Suspension of the  $J_2$ s was diluted further to get approximately 25  $J_2$ s per mL.

## 2.6. Test Procedure

**2.6.1. Antioxidant Activity.** The antioxidant activity of test extracts such as CE1, CE2, CE3, and standard compounds gallic acid and quercetin was assessed by radical scavenging effect of 1,1-diphenyl-2-picrylhydrazyl (DPPH) [17–19]. The diluted working solution of the test extracts and the standards (1–100 mg/L) were prepared in methanol. 1 mL of DPPH (0.002%) solution in methanol was mixed separately with 1 mL of the sample solution and the standard solution. Different concentrations (50, 100, 500, and 1000 mg/L) of test extracts and the standards were pipetted to the test tubes and volume adjusted to 3 mL with methanol. 1 mL of alcoholic DPPH (0.002%) solution was added to each sample and the samples were vortexed and then incubated in dark at room temperature for 30 min. Methanol (1 mL)

with DPPH solution (0.002%, 1 mL) was used as blank. The spectrophotometric measurements at 517 nm against blank samples were made with a pair of matched quartz cuvettes using Analytik Jena UV-Vis spectrophotometer (SPCORD 200). The optical density was recorded and radical scavenging activity was expressed as percentage inhibition of DPPH radical and was calculated by the following equation:

$$\begin{aligned} \text{Inhibition (\%)} &= \left[ \text{Absorbance of control} - \left( \frac{\text{Absorbance of sample}}{\text{Absorbance of control}} \right) \right] \\ &\times 100. \end{aligned} \quad (1)$$

Extract concentration providing 50% inhibition ( $\text{IC}_{50}$ ) was calculated from the graph plotted between inhibition percentages against extract concentration with the help of statistical package (GW BASIC).

**2.6.2. Nematicidal Activity.** Root-knot nematode suspension (2 mL) containing 50 juveniles was taken in petri plates; an equal volume of the test solutions was added separately to obtain the desired test concentrations of 200, 100, 50, and 25  $\text{mgL}^{-1}$ , respectively. Each treatment was replicated thrice separately for 24, 48, and 72 hours. For each treatment a control was taken by adding 2 mL of nematode suspension with equal volume of ethanol-Tween 80 emulsified water at 25–30°C. After 24 h of exposure, the petri plates for each treatment were observed under stereoscopic binocular microscope for determining mortality. A revival test was performed for each treatment by decanting the test solvents and adding distilled water to the petri plates. After 48 h of exposure readings were taken again for each treatment. The revived juveniles were counted and deducted from the number of immobile juveniles obtained in the previous reading. Similar process was followed after 72 hr exposure time. The nematodes were considered dead if appeared motionless, when probed with a fine needle. The percent mortality in treatment as well as in control for each compound was calculated and the corrected percent mortality was calculated by using the following Abbot's formula (1925) [20]:

$$\text{Corrected Mortality (\%)} = \left[ \frac{(T - C)}{(100 - C)} \right] \times 100, \quad (2)$$

where  $T$  = total mortality in treatment and  $C$  = total mortality in control.

The mortality data were subjected to Probit analysis.  $\text{LC}_{50}$  value ( $\text{mgL}^{-1}$ ) was calculated from the concentration ( $\text{mgL}^{-1}$ ) and the corresponding % corrected mortality data with the help of statistical package (GW BASIC).

## 3. Results and Discussion

**3.1. Extraction and Characterization of Anthraquinones.** The mass spectra of four hydroxyanthraquinones identified in




FIGURE 3: ESI-MS of rhabarb anthraquinones.

rhubarb are shown in Figure 3. ESI-MS of anthraquinones provides more structural information than the APCI-MS [21, 22]. In the ESI-MS spectrum of chrysophanol, a small molecular ion peak at  $m/z$  253 [ $M - H$ ]<sup>-</sup> and a stable large fragment ion at  $m/z$  225 [ $M - 2H$ ]<sup>-</sup> ion were observed as a result of deprotonation of one or two protons from phenolic-OH groups. A fragment ion at  $m/z$  225 originated as a result of the direct loss of CO from [ $M - H$ ]<sup>-</sup>. The  $m/z$  225 ion was stable and did not result in further fragmentation. The CO elimination may preferably originate from C-10, since the carbonyl group at C-9 has intramolecular hydrogen bonding with the  $\alpha$ -hydroxyl groups at C-1 and C-8 and is therefore difficult to be cleaved. Emodin and aloe-emodin are isomers and can be differentiated by their ESI-MS spectra. Both the compounds exhibit deprotonated molecular ion peak at  $m/z$  269 [ $M - H$ ]<sup>-</sup>. The fragmentation of emodin was initiated by the elimination of CO to produce  $m/z$  241, followed by loss of one hydroxyl group to give fragment ion at  $m/z$  225. The stable  $m/z$  269.3 [ $M - H$ ]<sup>-</sup> ion of aloe-emodin does not lead to formation of any peak at 225 rather, its fragmentation formed a peak at  $m/z$  253.3 [ $M - OH$ ]<sup>-</sup>. Physcion showed a prominent molecular ion peak at  $m/z$  283 [ $M - H$ ]<sup>-</sup>. It contained a methyl group that easily eliminates 15 mass units to give a significant ion at  $m/z$  269.2 followed by elimination of methoxy group  $m/z$  253.3 [ $M - OCH_3$ ]<sup>-</sup>. This ion could

further lose one molecule of H<sub>2</sub>O to produce fragment ion at  $m/z$  235.2.

**Chrysophanol.** The first compound was crystallized from chloroform-acetone as yellow leaflets, m. p. 194°C. Rf is 0.9 (hexane:ethyl acetate:formic acid; (2:8:0.1)). The IR (KBr) $\gamma_{\max}$  of the purified metabolite showed a strong hydroxyl group absorption band at 3470 cm<sup>-1</sup>, together with two carbonyl group absorption bands at 1680 and 1633 cm<sup>-1</sup>. It further shows absorption maxima at UV ( $\lambda^{\text{MeOH}}_{\max}$ ): 228, 256, 279, 287, and 429 nm. Additionally, its <sup>1</sup>H NMR spectrum showed two sharp singlets at  $\delta$  (ppm) 12.03 (OH-1) and 12.14 (OH-8) characteristic of two chelated hydroxyl groups and signals for five aromatic protons, 7.15 (1H, *d*, H-2), 7.59 (1H, *d*, H-4), 7.30 (1H, *dd*, H-7), 7.86 (1H, *dd*, H-5), and 7.69 (1H, *d*, H-6) and for the aromatic methyl group of an anthraquinone 2.45 (*s*, H-3). On the basis of its spectroscopic data, and by comparing it with those reported in the literature, the molecule was identified as chrysophanol (1,8-dihydroxy-3-methylantracene-9,10-dione).

**Physcion.** The second compound was crystallized from chloroform to give yellow needles, m.p. 204°C. Rf is 0.8 in hexane:ethyl acetate:formic acid (2:8:0.1) and 0.85 in benzene:chloroform:acetic acid (9:1:0.1). The IR (KBr)

TABLE 1: IC<sub>50</sub> (mg/L) and DPPH scavenging activity (%) of *Rheum emodi* (rhubarb) extracts.

Compound	% DPPH scavenging activity				$\chi^2_{\text{obs}}$ , (3d.f, 95%)	Regression equation	Fiducial limit	IC <sub>50</sub> (mg/L)
	5 mg/L	10 mg/L	50 mg/L	100 mg/L				
CE1	20.12	38.75	70.12	87.95	1.14	$Y = 3.18 + 1.44X$	14.69–22.73	18.28
CE2	21.56	32.71	68.52	88.33	5.94	$Y = 3.09 + 1.48X$	15.66–23.97	19.37
CE3	19.80	35.47	68.41	85.59	0.971	$Y = 3.15 + 1.41X$	16.25–25.28	20.27
Quercetin hydrate	43.38	67.17	87.78	91.41	5.59	$Y = 4.32 + 0.98X$	3.12–7.14	4.95
Gallic acid	31.79	44.64	85.06	92.43	2.96	$Y = 3.59 + 1.34X$	8.73–14.54	11.26

Results shown are means of three experiments.

$\gamma_{\text{max}}$  of the purified metabolite showed a strong carbonyl group absorption bands at 1675 and 1630  $\text{cm}^{-1}$ . It further shows absorption maxima at UV ( $\lambda^{\text{MeOH max}}$ ): 255, 250, and 430 nm. Additionally, its  $^1\text{H}$  NMR spectrum showed two sharp singlets at  $\delta$  12.13 and 12.33 and signals for five aromatic protons [ $\delta$  2.46 (CH<sub>3</sub>), 6.69(1H), 7.09 (1H), 7.37 (1H), 7.63 (1H)] and for the aromatic methoxy group of an anthraquinone  $\delta$  3.94 (OCH<sub>3</sub>). On the basis of its spectroscopic data, and by comparing it with those reported in the literature, the molecule was identified as physcion (1,8-dihydroxy-6-methyl-3-methoxyanthracene-9,10-dione).

*Emodin*. The third compound was crystallized from hexane: ethyl acetate (80 : 20) to give pinkish leaflets, m.p. 258°C. The R<sub>f</sub> is 0.40 (hexane : ethyl acetate : formic acid; 2 : 8 : 0.1) and 0.45. (hexane : chloroform : acetic acid (9 : 1 : 0.1)). Emodin showed absorption maxima at 437, 299, and 260 nm. IR of emodin showed frequency at 3390  $\text{cm}^{-1}$  assigned to hydroxyl groups. Characteristic bands for anthraquinones at 1675  $\text{cm}^{-1}$  and 1631  $\text{cm}^{-1}$  were obtained which is assigned to the free carbonyl group and the conjugated carbonyl group stretching frequencies, respectively. The observed frequency at 1475  $\text{cm}^{-1}$  is assigned to a skeletal ring stretching frequency. Its  $^1\text{H}$  NMR spectrum showed two sharp singlet at  $\delta$  12.30 and  $\delta$  12.80 and signals for four aromatic protons [ $\delta$  6.69 (1H), 7.09 (1H), 7.37 (1H), 7.63 (1H)] and for the aromatic methyl group of an anthraquinone at  $\delta$  2.46 (CH<sub>3</sub>). On the basis of its spectroscopic data, molecule was identified as emodin (1, 8-dihydroxy-6-methyl-3-hydroxyanthracene-9,10-dione).

*Aloe-Emodin*. The fourth compound was crystallized from chloroform: acetone (3:1) as yellow needles, m.p. 220°C. The R<sub>f</sub> is 0.30 (hexane: ethyl acetate: formic acid; 2 : 8 : 0.1) and 0.32 (hexane: chloroform: acetic acid (9 : 1 : 0.1)). Aloe-emodin showed absorption maxima at 225, 224, 276, 287, and 457 nm, respectively. IR of aloe-emodin shows frequency at 3390  $\text{cm}^{-1}$  assigned to hydroxyl groups. Characteristic bands for anthraquinones are 1675  $\text{cm}^{-1}$  and 1631  $\text{cm}^{-1}$  assigned to the free carbonyl group and the conjugated carbonyl group stretching frequencies, respectively. The observed frequency at 1475  $\text{cm}^{-1}$  is assigned to a skeletal ring stretching frequency. Its  $^1\text{H}$  NMR spectrum showed two sharp singlets at  $\delta$  12.08 and 12.10 and signals for five aromatic protons: [ $\delta$ , 7.32 (1H, 5-H), 7.35 (1H, 4-H), 7.69 (1H, 6-H), 7.80 (1H, 7-H), 7.85 (1H, 2-H)] and for the characteristics aromatic methylene group at  $\delta$  4.85 (CH<sub>2</sub>) and for aromatic hydroxyl at  $\delta$  5.35(OH) of

an anthraquinone. On the basis of its spectroscopic data, the molecule was identified as aloe-emodin (1, 8-dihydroxy-3-hydroxymethylanthracene-9,10-dione).

Anthraquinone glucosides also produced [M – H]<sup>–</sup> ions in the negative ESI source (Figure 4) like their free anthraquinones. The MS/MS fragmentation were predominated by the elimination of glucosyl residue to give [Aglycone – H]<sup>–</sup> that is, [A – H]<sup>–</sup> ions as the base peak. The aglycones were further identified based on the product ion spectra of [A – H]<sup>–</sup> ions, and by comparing to the MS fragmentation behaviour of free anthraquinones described earlier. Based on their deprotonated ion peaks in ESI-MS (negative ion mode), five anthraquinone glucoside have been identified as emodin 8-O-(6'-O-malonyl)-glucoside [M – H, 518], emodin 1-O-glucoside [M – H, 432], emodin 8-O-glucoside [M – H, 432], rhein 8-O-glucoside [M – H, 445], and chrysophanol 8-O-glucoside [M – H, 415]. Another glucoside, namely, piceatannol-3-O-beta-d-glucopyranoside [M – H, 406], has also been detected in the *R. emodi* methanol extract concentrate (Figure 4). Our results were in conformity with those reported by earlier authors [23].

**3.2. Antioxidant Activity.** The extracts of *R. emodi* were analyzed for antioxidant activity by using DPPH free radical scavenging activity. The anthraquinone derivatives, such as aloe-emodin, emodin, rhein, chrysophanol, and physcion, are reported to possess antiangiogenic activity, by preventing blood vessel formation in zebra-fish embryos [24]. The anticancer effect of aloe-emodin has been established in two human cancer cell lines, Hep G2 and Hep 3B. Aloe-emodin inhibited cell proliferation and induced apoptosis in both examined cell lines by different antiproliferative mechanisms [25]. In our study all the extracts showed a concentration dependent scavenging of DPPH radicals (Figure 5). CE1 with (IC<sub>50</sub> 18.28 mg/mL) was most active, followed by ethyl CE2 (IC<sub>50</sub> 19.37 mg/mL) and CE3 (IC<sub>50</sub> 20.27 mg/mL) (Table 1). All the extracts were comparatively less active than both quercetin hydrate (QH) (4.95 mg/L) and gallic acid (GA) (11.26 mg/L) in quenching DPPH radical than standard antioxidants. The extracts contained a high number of phenolic compounds, which were found to have significant positive correlation with free radicals (DPPH and OH) scavenging efficacies. The results were in agreement with those of [26] who found the methanolic extract to be a more active radical scavenger than aqueous extract.

The HPLC analysis had shown the methanolic extract of *R. emodi* rhizomes to be rich in the major anthraquinones




FIGURE 4: ESI-MS/MS of rhabarb anthraquinone glucosides.

and their glucosides, namely, emodin, aloe-emodin, chrysophanol, physcion, emodin glucoside, chrysophanol glucoside, and revandchinone, whereas ethyl acetate and chloroform extract was found to be low in aloe-emodin and other glucosides. The lower  $IC_{50}$  value for methanolic extract in free radical scavenging activity of DPPH is attributed to the fact that the extract is rich in bioactive constituents including anthraquinones. However, our results are in contrast to the study [4] where the compounds like marsupsin and

maesopsin obtained from the rhizome/root extracts of *R. emodi* are found to possess antioxidant activity, whereas chrysophanol, physcion, and emodin as well as their 8-O-glucosides were found to be inactive.

**3.3. Nematicidal Activity.** After 72 hr of exposure and observation time CE1 ( $ED_{50}$  54.84 mg/L) was most active followed by CE2 ( $ED_{50}$  98.10 mg/L) and CE3 (135.23 mg/L). Among the pure anthraquinones chrysophanol ( $ED_{50}$  102.59 mg/L)

TABLE 2: Antinemic activity ( $ED_{50}$ ) of isolated anthraquinones and solvent extractives of *Rheum emodi* against  $j_2s$  of *Meloidogyne incognita* after 72 hrs.

Compounds	$\chi^2_{obs}$ (3d.f, 95%)	Regression equation	$ED_{50}$ (mg/L)	Fiducial limit
CE1	1.91	$Y = 2.74 + 1.30X$	54.84	44.51–67.57
CE2	6.73	$Y = 2.73 + 1.13X$	98.10	74.34–129.46
CE3	6.47	$Y = 2.26 + 1.27X$	135.23	101.27–180.58
Physcion	2.50	$Y = 2.33 + 1.32X$	102.61	82.80–136.25
Aloe-emodin	7.39	$Y = 2.25 + 1.27X$	148.50	109.17–202.01
Chrysophanol	2.15	$Y = 2.03 + 1.42X$	102.59	81.50–129.15
Emodin	2.62	$Y = 2.03 + 1.38X$	139.95	106.64–183.6
Aza (20%)	0.42	$Y = 3.42 + 0.769X$	112.70	73.14–173.67




FIGURE 5: Free radical scavenging activity of the extracts of *Rheum emodi* and standard antioxidants.

and physcion ( $ED_{50}$  102.61 mg/L) were equally active followed by emodin (139.95 mg/L) and aloe-emodin (148.50 mg/L). The CE1 has also outperformed the positive control (Aza 20%) (112.70 mg/L) (Table 2). The results revealed that crude methanolic extract (CE1) was rich in anthraquinones, glucosides, and other phenolics contained in *R. emodi* which were very effective in controlling *M. incognita* menace. The pure compounds were, however, comparatively less lethal than crude fractions. The observations substantiate the fact that bioactive constituents collectively showed symbiotic effect as the positive control azadirachtin (20%) was less active than the crude extract (CE1). The revelation opens a new horizon that the CE1 could be formulated as potent nematocidal and the same can be evaluated in the field conditions.

#### 4. Conclusion

The present study conclusively demonstrated that the anthraquinones and other phenolics of *R. emodi* especially

in the methanolic extract (CE1) have great potential and can be suitably utilized into environmentally benign pest control strategies. The potential antioxidant activity of the *R. emodi* constituents as revealed by the DPPH assay strengthens the importance of anthraquinones in ethno-medical use as well as in contemporary medicine. Further LC-MS analysis revealed that the activity of CE1 was mainly due to the synergistic combinations of its anthraquinone and their glucosides along with other phenolics. The constituents can suitably become one of the lead molecules to be further exploited in various spheres of pest and disease control mechanisms after suitable clinical testing.

#### Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

#### Acknowledgments

The authors express their gratitude to the Head of Division of Agricultural Chemicals, Nematology, and NAIP (ICAR) for providing necessary facilities and Director of IARI and Dean P. G. School, IARI, New Delhi, for financial assistance Senior Research Fellow (SRF) to Brijesh Tripathi. This work has been supported by a National Research Foundation of Korea (NRF) grant funded by the Ministry of Science, ICT and Future Planning (MSIP) (NRF-2010-0027963 and 20120009639). This work has also been partially supported by Kwangwoon University, 2014.

#### References

- [1] M. B. Rokaya, Z. Münzbergová, B. Timsina, and K. R. Bhattarai, "Rheum australe D. Don: a review of its botany, ethnobotany, phytochemistry and pharmacology," *Journal of Ethnopharmacology*, vol. 141, no. 3, pp. 761–774, 2012.
- [2] B. P. Nautiyal, V. Prakash, U. C. Maithani, R. S. Chauhan, H. Purohit, and M. C. Nautiyal, "Germinability, productivity and economic viability of *Rheum emodi* Wall. ex Meissn. cultivated at lower altitude," *Current Science*, vol. 84, no. 2, pp. 143–148, 2002.



- [3] B. A. Zargar, M. H. Masoodi, B. Ahmed, and S. A. Ganie, "Phytoconstituents and therapeutic uses of *Rheum emodi* wall. ex Meissn," *Food Chemistry*, vol. 128, no. 3, pp. 585–589, 2011.
- [4] L. Krenn, A. Presser, R. Pradhan et al., "Sulfemodin 8-O- $\beta$ -D-glucoside, a new sulfated anthraquinone glycoside, and antioxidant phenolic compounds from *Rheum emodi*," *Journal of Natural Products*, vol. 66, no. 8, pp. 1107–1109, 2003.
- [5] L. Krenn, R. Pradhan, A. Presser, G. Reznicek, and B. Kopp, "Anthrone C-glucosides from *Rheum emodi*," *Chemical and Pharmaceutical Bulletin*, vol. 52, no. 4, pp. 391–393, 2004.
- [6] K. S. Babu, P. V. Srinivas, B. Praveen, K. H. Kishore, U. S. Murty, and J. M. Rao, "Antimicrobial constituents from the rhizomes of *Rheum emodi*," *Phytochemistry*, vol. 62, no. 2, pp. 203–207, 2003.
- [7] S. K. Agarwal, S. S. Singh, S. Verma, and S. Kumar, "Antifungal activity of anthraquinone derivatives from *Rheum emodi*," *Journal of Ethnopharmacology*, vol. 72, no. 1-2, pp. 43–46, 2000.
- [8] M. Ibrahim, A. A. Khan, S. K. Tiwari, M. A. Habeeb, M. N. Khaja, and C. M. Habibullah, "Antimicrobial activity of *Sapindus mukorossi* and *Rheum emodi* extracts against *H pylori*: in vitro and in vivo studies," *World Journal of Gastroenterology*, vol. 12, no. 44, pp. 7136–7142, 2006.
- [9] P. P. Rai, T. D. Turner, and S. A. Matlin, "High-pressure liquid chromatography of naturally occurring anthraquinones," *Journal of Chromatography A*, vol. 110, no. 2, pp. 401–402, 1975.
- [10] M. M. A. Alam, K. Javed, and M. A. Jafri, "Effect of *Rheum emodi* (Revand Hindi) on renal functions in rats," *Journal of Ethnopharmacology*, vol. 96, no. 1-2, pp. 121–125, 2005.
- [11] Z. Cai, F. S. C. Lee, X. R. Wang, and W. J. Yu, "A capsule review of recent studies on the application of mass spectrometry in the analysis of Chinese medicinal herbs," *Journal of Mass Spectrometry*, vol. 37, no. 10, pp. 1013–1024, 2002.
- [12] X.-G. He, "On-line identification of phytochemical constituents in botanical extracts by combined high-performance liquid chromatographic-diode array detection-mass spectrometric techniques," *Journal of Chromatography A*, vol. 880, no. 1-2, pp. 203–232, 2000.
- [13] U. Justesen, "Collision-induced fragmentation of deprotonated methoxylated flavonoids, obtained by electrospray ionization mass spectrometry," *Journal of Mass Spectrometry*, vol. 36, no. 2, pp. 169–178, 2001.
- [14] N. Fabre, I. Rustan, E. De Hoffmann, and J. Quetin-Leclercq, "Determination of flavone, flavonol, and flavanone aglycones by negative ion liquid chromatography electrospray ion trap mass spectrometry," *Journal of the American Society for Mass Spectrometry*, vol. 12, no. 6, pp. 707–715, 2001.
- [15] M. Ye, D. Guo, G. Ye, and C. Huang, "Analysis of homoisoflavonoids in *Ophiopogon japonicus* by HPLC-DAD-ESI-MSn," *Journal of the American Society for Mass Spectrometry*, vol. 16, no. 2, pp. 234–243, 2005.
- [16] M. Ye, Y. Yan, and D.-A. Guo, "Characterization of phenolic compounds in the Chinese herbal drug Tu-Si-Zi by liquid chromatography coupled to electrospray ionization mass spectrometry," *Rapid Communications in Mass Spectrometry*, vol. 19, no. 11, pp. 1469–1484, 2005.
- [17] M. S. Blois, "Antioxidant determinations by the use of a stable free radical," *Nature*, vol. 181, no. 4617, pp. 1199–1200, 1958.
- [18] İ. Gülçin, I. G. Şat, Ş. Beydemir, M. Elmastaş, and Ö. I. Küfrevioğlu, "Comparison of antioxidant activity of clove (*Eugenia caryophyllata* Thunb) buds and lavender (*Lavandula stoechas* L.)," *Food Chemistry*, vol. 87, no. 3, pp. 393–400, 2004.
- [19] W. Brand-Williams, M. E. Cuvelier, and C. Berset, "Use of a free radical method to evaluate antioxidant activity," *LWT—Food Science and Technology*, vol. 28, no. 1, pp. 25–30, 1995.
- [20] W. S. Abbott, "A method of computing the effectiveness of an insecticide," *Journal of Economic Entomology*, vol. 18, no. 2, pp. 265–267, 1925.
- [21] D. J. Zeeb, B. C. Nelson, K. Albert, and J. J. Dallunge, "Separation and identification of twelve catechins in tea using liquid chromatography/atmospheric pressure chemical ionization-mass spectrometry," *Analytical Chemistry*, vol. 72, no. 20, pp. 5020–5026, 2000.
- [22] W. Li, C.-L. Chan, and H.-W. Lueng, "Liquid chromatography-atmospheric pressure chemical ionization mass spectrometry as a tool for the characterization of anthraquinone derivatives from Chinese herbal medicine," *Journal of Pharmacy and Pharmacology*, vol. 52, no. 6, pp. 723–729, 2000.
- [23] M. Ye, J. Han, H. Chen, J. Zheng, and D. Guo, "Analysis of phenolic compounds in rhubarbs using liquid chromatography coupled with electrospray ionization mass spectrometry," *Journal of the American Society for Mass Spectrometry*, vol. 18, no. 1, pp. 82–91, 2007.
- [24] Z.-H. He, M.-F. He, S.-C. Ma, and P. P.-H. But, "Anti-angiogenic effects of rhubarb and its anthraquinone derivatives," *Journal of Ethnopharmacology*, vol. 121, no. 2, pp. 313–317, 2009.
- [25] P.-L. Kuo, T.-C. Lin, and C.-C. Lin, "The antiproliferative activity of aloe-emodin is through p53-dependent and p21-dependent apoptotic pathway in human hepatoma cell lines," *Life Sciences*, vol. 71, no. 16, pp. 1879–1892, 2002.
- [26] R. Ashok Kumar, V. Rajkumar, and G. Guha, "Antioxidant and anti-cancer potentials of rheum emodi rhizome extracts," *Evidence-Based Complementary and Alternative Medicine*, vol. 2011, Article ID 697986, 9 pages, 2011.



**Hindawi**

Submit your manuscripts at  
<http://www.hindawi.com>

