



Citation for published version:

Xiao, Q, Chaput, J-P, Olds, T, Fogelholm, M, Hu, G, Lambert, EV, Maher, C, Maia, J, Onywera, V, Sarmiento, OL, Standage, M, Tremblay, MS, Tudor-Locke, C & Katzmarzyk, PT 2020, 'Sleep characteristics and health-related quality of life in 9- to 11-year-old children from 12 countries', *Sleep Health*, vol. 6, no. 1, pp. 4-14. <https://doi.org/10.1016/j.sleh.2019.09.006>

DOI:

[10.1016/j.sleh.2019.09.006](https://doi.org/10.1016/j.sleh.2019.09.006)

Publication date:

2020

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

CC BY-NC-ND

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Sleep characteristics and health-related quality of life in 9-11 year old children from 12 countries**

2 Qian Xiao¹, Jean-Philippe Chaput², Timothy Olds³, Mikael Fogelholm⁴, Gang Hu⁵, Estelle V. Lambert⁶,
3 Carol Maher³, Jose Maia⁷, Vincent Onywera⁸, Olga L. Sarmiento⁹, Martyn Standage¹⁰, Mark S. Tremblay²,
4 Catrine Tudor-Locke¹¹ and Peter T. Katzmarzyk⁵ for the ISCOLE Research Group

5

6 ¹ University of Iowa, Iowa city, USA

7 ² Children's Hospital of Eastern Ontario Research Institute, Ottawa, Canada

8 ³ University of South Australia, Adelaide, Australia

9 ⁴ University of Helsinki, Helsinki, Finland

10 ⁵ Pennington Biomedical Research Center, Baton Rouge, Louisiana, USA

11 ⁶ University of Cape Town, Cape Town, South Africa

12 ⁷ University of Porto, Porto, Portugal

13 ⁸ Kenyatta University, Nairobi, Kenya

14 ⁹ Universidad de los Andes, Bogota, Colombia

15 ¹⁰ University of Bath, Bath, United Kingdom

16 ¹¹ University of Massachusetts Amherst, Amherst, USA

17

18 Corresponding Author: Qian Xiao, Department of Health and Human Physiology and Department of
19 Epidemiology, University of Iowa, Iowa city, USA (email: qian-xiao@uiowa.edu ; phone: 319-335-9348).

20 **Abstract**

21 **Introduction**

22 Previous studies have linked short sleep duration, poor sleep quality, and late sleep timing with lower
23 health-related quality of life (HRQoL) in children. However, almost all studies relied solely on self-
24 reported sleep information and most were conducted in high income countries. To address these gaps,
25 we studied both device-measured and self-reported sleep characteristics in relation to HRQoL in a
26 sample of children from 12 countries that vary widely in terms of economic and human development.

27 **Methods**

28 The study sample included 6,626 children aged 9-11 years from Australia, Brazil, Canada, China,
29 Colombia, Finland, India, Kenya, Portugal, South Africa, the United Kingdom, and the United States.
30 Waist-worn actigraphy was used to measure total sleep time, bedtime, wake-up time, and sleep
31 efficiency on both weekdays and weekends. Children also reported ratings of sleep quantity and quality.
32 HRQoL was measured by the KIDSCREEN-10 survey. Multilevel regression models were used to
33 determine the relationships between sleep characteristics and HRQoL.

34 **Results**

35 Results showed considerable variation in sleep characteristics, particularly duration and timing, across
36 study sites. Overall, we found no association between device-measured total sleep time, sleep timing or
37 sleep efficiency and HRQoL. In contrast, self-reported ratings of poor sleep quantity and quality were
38 associated with HRQoL.

39 **Conclusions**

40 Self-reported, rather than device-based, measures of sleep are related to HRQoL in children. The
41 discrepancy related to sleep assessment methods highlights the importance of considering both device-
42 measured and self-reported measures of sleep in understanding its health effects.

43 **Key Words**

44 Sleep duration, total sleep time, sleep efficiency, sleep timing, health related quality of life

45 **Introduction**

46 The American Academy of Sleep Medicine (AASM) guidelines recognize sleep as a multidimensional
47 behavior and define healthy sleep as “adequate duration, appropriate timing, good quality, regularity,
48 and the absence of sleep disturbances or disorders.”¹ For children and adolescents, healthy sleep is
49 essential for healthy living, and insufficient sleep has been linked with a wide range of physical, mental
50 and social problems,² including obesity,³ hypertension,^{4,5} reduced insulin sensitivity and type 2
51 diabetes,^{6,7} depression,⁸ anxiety,⁹ impaired emotional regulation,^{10,11} sub-optimal cognitive and
52 academic performance¹²⁻¹⁴ and poor relationships with peers and family.¹⁵ The AASM recommends 9-12
53 hours of sleep on a regular basis for children aged 6-12 years to achieve optimal health.¹

54 Health-related quality of life (HRQoL) captures all three main domains of health (physical, mental, and
55 social) and is an important and widely used indicator of overall health in adults and children.¹⁶ Multiple
56 studies have examined the relationship between sleep and HRQoL in children and adolescents, yet
57 mixed findings have been reported. For example, several studies found that self-reported short sleep
58 duration, late sleep timing, and sleep disturbances were associated with poor physical, social, and
59 emotional functioning as well as poor overall HRQoL, while long sleep duration and better sleep quality
60 were related to reduced health complaints and better psychological health.^{15,17-24} However, other
61 studies have reported no association between self-reported sleep characteristics and HRQoL, or weak
62 and inconsistent findings.²⁵⁻²⁷ It is worth noting that almost all studies have been conducted in high
63 income countries, and little is known about sleep and HRQoL in children living in low-to-middle income
64 countries.

65 A major limitation of the current literature on sleep and HRQoL is that all the previous studies have
66 relied on self-reported sleep information. Previous validation studies have found substantial
67 discrepancies between self-reported and device-measured sleep in both adults and children.²⁸⁻³⁰

68 Moreover, these studies have reported that the size of such discrepancies varies by health status and
69 sleep characteristics, and people with poor health and sleep deficiencies tend to have larger differences
70 between device-measured and self-reported sleep measurements. Therefore, there is a need to better
71 understand the association between device-measured sleep characteristics in relation to HRQoL in
72 children.

73 To address the aforementioned limitations in the literature, we studied device-measured sleep
74 characteristics and HRQoL in an international sample of children, and compared results with those using
75 self-reported measures in the same study. We conducted country-specific analyses to examine and
76 compare the associations across 12 different nations with a wide range of geographic, economic and
77 socio-cultural variability. We hypothesized that short sleep, low sleep efficiency, later sleep timing and
78 larger day-to-day variabilities in sleep characteristics are associated with poor HRQoL in children.

79

80 **Methods**

81 *Study Population*

82 The International Study of Childhood Obesity, Lifestyle and the Environment (ISCOLE) (ClinicalTrials.gov:
83 NCT01722500) is an international cross-sectional study of children (9-11 years) from study sites in 12
84 countries (Australia, Brazil, Canada, China, Colombia, England, Finland, India, Kenya, Portugal, South
85 Africa, and USA). Details of ISCOLE have been previously published.³¹ The countries were selected to
86 maximize geographical, socio-economic and cultural variability: these countries differed widely in
87 several socio-economic indicators: Six countries (Australia, Canada, Finland, Portugal, England and
88 United States) are classified as high income according to World Bank Classification, five (Brazil,
89 Colombia, China, India and South Africa) classified as upper-middle income, and one (Kenya) classified as
90 low income.³² These countries also range from very high (0.93, Australia) to low (0.51, Kenya) in their

91 Human Development Index, a composite score based on life expectancy, gross national income, literacy
92 and school participation.³³ Details of economic indicators for all twelve countries have been published
93 before.³¹ In each country, children were recruited from one or more school districts that located
94 reasonably close to the local study center. Sampling included students from urban and suburban areas.
95 Rural areas were excluded to maximize comparisons across ISCOLE sites because in some countries,
96 study centers had limited access to rural populations. Each site aimed to recruit a sex-balanced sample
97 of 500 children. Technicians were trained to administer the questionnaire in a standardized fashion in
98 order to minimize bias, and provisions were made to administer the questionnaire via an interview for
99 participants with low levels of literacy. Questionnaires were translated to local languages and children
100 as young as 9 years showed sufficient comprehension of the study questionnaires. Of the total 7,372
101 children in the final ISCOLE sample, we excluded children who were missing any sleep variables
102 (N=1,081) or HRQoL information (N=25). The analytic sample for this study therefore includes 6,266
103 children. The Pennington Biomedical Research Center Institutional Review Board approved the ISCOLE
104 protocol and ethical review boards at each site approved local protocols. Written informed consent
105 from parents and child assent were obtained as required by local review boards.

106 *Health-Related Quality of Life*

107 Children in the study completed the KIDSCREEN-10 survey, which provides a global measure of HRQoL in
108 children.³⁴ The KIDSCREEN-10 is the brief form of KIDSCREEN-54, an instrument that was designed to
109 measure HRQoL among children aged 8-18 years and validated in numerous low-to-middle-income
110 countries.^{35,36} The KIDSCREEN-10 included 10 questions on children's physical activity, energy and
111 fitness, mood and emotions, social and leisure participation, social and family relationships, cognitive
112 capacity, and school experience. Responses to each question are recorded on a 5-point scale and
113 participant's scores were reversed when appropriate to ensure that higher scores indicate better

114 HRQoL. We summed each participant's score across questions to calculate Rasch person-parameters,
115 which were transformed into t-values with a mean of 50 and a standard deviation of approximately 10.³⁴

116 *Assessment of Sleep*

117 Sleep was measured by waist-worn accelerometers (Actigraph GT3X+, Actigraph LLC, Pensacola, FL,
118 USA). Actigraphy data collection was conducted during the school year. Data collection was purposefully
119 spread over this period to account for differences across seasons. Children were asked to wear the
120 device 24-hours per day for seven consecutive days, removing it only for water-related activities such as
121 bathing, showering and swimming. Details about the sleep variable assessments were reported
122 before.^{37,38} Briefly, for sleep variable calculation, we only included participants with at least three nights
123 (including one weekend night) of valid sleep (total sleep period time ≥ 160 min). Nocturnal total sleep
124 time was estimated from 1-min epochs using a validated, fully-automated algorithm for waist-worn
125 accelerometers, which captures total sleep time from sleep onset to the end of sleep and all episodes of
126 wakefulness after onset.³⁸

127 From the accelerometry data, we measured three primary sleep variables: 1) duration measured by total
128 sleep time; 2) timing measured by midpoint between sleep onset and offset; and 3) sleep efficiency
129 measured by total sleep time divided by total time spent in bed. We also included an additional sleep
130 timing variable, the onset of sleep time, as some previous studies indicated that bedtime was a
131 significant predictor of HRQoL.²² Finally, we also calculated mid-sleep time on free days corrected for
132 sleep debt on work days (MSFsc) as an indicator of chronotype.³⁹ For each sleep variable, we calculated
133 average values for weekday (Sunday through Thursday) and weekend (Friday and Saturday) nights
134 separately. We also derived two indicators of variability for each sleep characteristic: the overall
135 variability was measured by the standard deviation of the sleep characteristic for all valid nights; and the
136 weekday-to-weekend difference was defined as the difference between weekend and weekday values

137 (weekend mean minus weekday mean) for the sleep characteristic. We divided each sleep variable into
138 quartiles. The reference group was chosen to represent the group that was hypothesized to have the
139 highest HRQoL: the longest group (Q4) for total sleep time, the earliest group (Q1) for sleep onset and
140 sleep midpoint, the highest group for sleep efficiency (Q4), and the groups with the smallest absolute
141 value for overall and weekday-to-weekend difference (Q1 for overall variability of all three sleep
142 characteristics as well as weekday-to-weekend difference of midpoint, and Q2 for weekday-to-weekend
143 difference of total sleep time and efficiency). For total sleep time, we also created a 4-category variable
144 (<7 hours, 7-8 hours, 9-12 hours, and >12 hours) and used 9-12 hours as the reference group based on
145 the recommended amount of sleep for pediatric population. ¹

146 Self-reported sleep measures were obtained by asking children “during the past week, rate the quantity
147 of your sleep overall” or “during the past week, rate the quality of your sleep overall”. Children were
148 asked to choose from four answers: very good, fairly good, fairly bad, or very bad.

149 *Covariates*

150 A wide array of sociodemographic and lifestyle variables from child participants and parents were
151 collected as part of ISCOLE, including age and sex of child participants. Parents reported parental
152 education, and marital status. Parents at each study site also reported annual household income using a
153 monetary scale of 8 to 10 country-specific categories using currencies of each country. These categories
154 were grouped into four levels to facilitate multi-country analysis. The four levels were created to ensure
155 the most balanced distribution possible within each country. Child participants completed a Food
156 Frequency Questionnaire, which has demonstrated moderate reliability and low to moderate- validity.⁴⁰
157 Principal components analysis was used to identify two component scores that represent 1) a healthy
158 diet pattern (positive loadings for vegetables, fruits, whole grains, etc.) and 2) an unhealthy diet pattern
159 (positive loadings for fast food, soft drinks, sweets, etc.). Height and weight were measured objectively

160 using standard techniques.³¹ Body-mass index (BMI) was calculated as weight in kilograms divided by
161 height in meters squared and transformed to age- and sex-specific z scores using the World Health
162 Organization reference data.⁴¹

163 *Statistical Analysis*

164 To report the distribution of participant characteristics, we calculated means and standard deviations
165 for continuous variables and percentages for categorical variables. To examine the associations between
166 sleep variables and HRQoL, we used multi-level multiple linear regression (PROC MIXED) to calculate the
167 mean difference and 95% confidence intervals (CIs) for each quartile of the sleep variable, comparing to
168 the reference group. In the model we adjusted for age (continuous), sex (female, male), parental
169 education (less than high school, high school graduate or some college, completed college or
170 postgraduate degree), parental marital status (married, divorced or separated, never married,
171 widowed), household income (site-specific categories representing low, low-to-medium, medium-to-
172 high and high income levels), BMI z-score (continuous), healthy diet pattern score (continuous) and
173 unhealthy diet pattern score (continuous). Schools were included as random effects in all models, while
174 study sites were included as fixed effects in analysis that included the overall population. We also
175 considered the total number of valid days for sleep recording, as well as the number of valid weekdays
176 and weekend days as covariates, but adjusting for these variables only had a minimal impact on the
177 results, and we did not include them in the final model. Tests for linear trends were performed using the
178 median value for each quartile as a continuous variable. To control for false positives due to multiple
179 comparison, we used a Bonferroni correction of p-value ($p < 0.05/130 = 0.0004$) to determine statistical
180 significance.

181 **Results**

182 The distribution of participant characteristics by study site are presented in **Table 1**. The average age of
183 child participants was 10.4 years and ~55% were girls. Across 12 countries, we observed a relatively wide
184 distribution of BMI z scores (lowest in Kenya and highest in Portugal) and healthy (lowest in Brazil and
185 highest in Canada) and unhealthy diet patterns (lowest in Finland and highest in South Africa). Parental
186 education levels and marital status also varied substantially across sites.

187 The distribution of number of valid nights, HRQoL and sleep variables are presented in **Table 2**. Overall
188 the number of valid nights across different sites are similar. Finland had the lowest average number of
189 nights (4.79) while China had the highest (5.76). Most of the countries had more than 5.5 valid nights.
190 We observed considerable variation in HRQoL scores and total sleep time and sleep timing across
191 countries. Kenya had the lowest HRQoL score (47.1) and Portugal had the highest (53.0). Total sleep
192 time on weekdays varied between 8.12 hr (Portugal) to 9.53 hr (Australia) while weekend total sleep
193 time was between 8.55 hr (Finland) and 9.62 hr (Colombia). For overall variability in total sleep time,
194 China had the smallest standard deviation (0.98) while Colombia had the largest (1.52). For weekday-to-
195 weekend variability, the average total sleep time on weekends was generally longer than that on
196 weekdays, except for Australia and Canada, for which weekend sleep was ~0.3 hr shorter than weekday
197 sleep. We also observed a relatively wide range of midpoint of sleep on both weekdays (earliest: 1:46,
198 Kenya; latest: 3:08, Brazil) and weekends (earliest: 2:59, China and Portugal; latest: 4:07, Brazil). Overall
199 variability of sleep midpoint was between 0.64 hr (China) and 0.96 hr (Brazil). Weekend sleep midpoint
200 was later than weekday sleep midpoint for all countries, and the difference ranged from 0.70 (Canada)
201 to 1.14 hr (USA). In contrast, sleep efficiency was consistently high in all countries and on both weekdays
202 and weekends, ranging from 95% to 97% on average, and there was little overall or weekday to
203 weekend day variability in sleep efficiency. For self-reported sleep, only a small fraction of children
204 reported very bad or fairly bad sleep quality and quantity. China had the highest percent of children

205 reporting both bad sleep quality (15.3%) and quantity (14.3%), while Kenya (4.8%) and Portugal (3.2%)
206 had the lowest percent for reporting bad sleep quality and quantity, respectively.

207 Overall, we found little evidence supporting an association between HRQoL and total sleep time (**Table**
208 **3**), midpoint of sleep (**Table 4**) or sleep efficiency (**Table 5**). Most of the effect estimates were not
209 statistically significant and no consistent patterns emerged as the magnitude and direction of effect
210 estimates for all three sleep variables varied considerably among countries. The results were similar
211 before and after adjusting for covariates. For total sleep time and sleep efficiency, neither the average
212 values on weekdays or weekend days, nor the two measures of variability, were associated with HRQoL.
213 We also performed analysis using total sleep time as a four-category and the results were similarly null
214 (data not shown). For midpoint of sleep, the findings were also largely null. However, we found that the
215 average sleep midpoint on weekends was positively associated with HRQoL in England with a 5.49 point
216 increase in HRQoL score comparing the latest quartile to the earliest ($\beta_{Q4 \text{ vs } Q1}$ (95% CI), 5.39 (2.69, 8.09),
217 $p\text{-trend}=0.0002$). Results for timing of sleep onset were largely null (**Supplementary Table 1**). In
218 addition, we did not find an association between MSFsc and HRQoL (data not shown).

219 Finally, we examined subjective ratings of sleep quantity and quality in relation to HRQoL. First, we
220 found that the correlation between self-reported and actigraphy-measured sleep variables were
221 generally null or weak (Spearman correlation coefficient, -0.07-0.10, **Supplementary Table 2**). Next, In
222 contrast to the null findings for device-measured sleep variables, poorer ratings of both sleep quantity
223 and quality were associated with worse HRQoL in almost all countries (**Table 6**). In the full sample,
224 ratings of “very bad” or “fairly bad” for sleep quantity or quality were associated with a more than six
225 point reduction in HRQoL.

226 **Discussion**

227 In this international sample of 9-11 year-old children, we found that sleep characteristics, especially
228 device-measured duration and timing, vary substantially across different countries. Overall our findings
229 do not support a relationship between device-measured sleep characteristics and HRQoL, although
230 there appeared to be some evidence suggesting that device-measured sleep timing might be associated
231 with HRQoL in England. On the other hand, we found that children with poor self-reported ratings of
232 sleep quantity and quality on average reported lower HRQoL.

233 A growing body of literature has examined the relationship between multiple sleep characteristics and
234 HRQoL in children and adolescents. All of the previous studies used self-reported sleep measures and
235 most examined sleep duration. Several studies have reported a positive association between self-
236 reported sleep duration and HRQoL in children and adolescents in a wide range of countries, including
237 Israel,²⁰ US,¹⁷ Korea,¹⁸ Switzerland,¹⁹ and Spain.¹⁵ For example, in over 3,000 children (age 11-17 years)
238 from Houston, Texas, Roberts et al. found that self-reporting less than 6 hours of sleep on weekdays and
239 weekends was associated with a higher likelihood of reporting low life satisfaction, poor perceived
240 mental health, depressed mood, problems at school and lower grades, as well as drug use.¹⁷ In another
241 study that included a large sample of Korean adolescents (age 13-17 years), self-reported short sleep
242 duration (<7 hours) was associated with a higher probability of reporting depressive symptoms, suicidal
243 ideation and overweight and obesity, and lower probability of reporting “very healthy” for self-rated
244 health.¹⁸ In addition, a few studies also examined self-reported sleep characteristics related to sleep
245 quality, including daytime sleepiness and morning tiredness (nonrestorative sleep), and symptoms
246 related to insomnia, such as sleep latency and waking during the night.^{15,20,21} Overall their findings
247 suggested that self-reported poor sleep quality was related to multiple domains of health and well-
248 being, including perceived health status, life satisfaction, quality of relationships, and academic
249 performance.

250 A major gap in the literature was a lack of studies that examined device-measured sleep characteristics,
251 as all the aforementioned studies used survey questionnaires. In this current study, although we
252 confirmed a positive association between self-reported poor sleep quantity and quality and low HRQoL,
253 we found no association when evaluating device-measured sleep characteristics. In the adult
254 population, studies that have compared self-reported with device-measured sleep duration generally
255 found only a moderate correlation between the two ($r=0.45-0.47$).^{28,29} The studies generally found that
256 self-report comparatively overestimates sleep duration by ~1 hr, and the validity of self-reported sleep
257 was lower among people with poor health status, shorter sleep duration and higher device-measured
258 sleep variability. Although similar validation studies are limited in children, some evidence suggests that
259 there is also considerable discrepancy between self-reported and device-based sleep measurements in
260 children. For example, Alfano et al. reported no or weak correlations between self/parent-reported and
261 actigraphy-measured sleep characteristics.³⁰ Moreover, the study by Alfano et al. also found that the
262 differences between device-based and self-reported sleep were larger in children with anxiety disorders
263 when compared to healthy children. On one hand, these validation studies in children and adults
264 suggest that self-reported sleep patterns may not accurately reflect objective characteristics of the sleep
265 behavior, particularly in populations with poor sleep and health status, which may introduce both errors
266 and biases in studies focusing on the relationships between sleep and health outcomes. On the other
267 hand, it is worth noting that self-reported sleep can capture subjective perceptions about sleep and the
268 fact that it correlates better with health outcomes suggest it may better reflect self-appraisal of overall
269 health, which itself is an important domain of health and well-being. Therefore self-reported sleep
270 should not be dismissed as merely an inaccurate measure. Instead, we encourage future studies to
271 include both device-based and self-reported sleep measures as they may reveal how different aspects of
272 sleep behavior may be associated with health outcomes.

273 Besides sleep duration and quality, sleep timing has received increasing attention in the past decade and
274 has been recognized as an important aspect of sleep health. Particularly, late sleep timing has been
275 highlighted as a risk factor for multiple adverse health outcomes in children and adolescents, including
276 obesity and mental health.⁴² For school children, the schedule of school on weekdays has a substantial
277 impact on their sleep timing, particularly the timing for waking up. When school start time is early, those
278 children with a preference for a later sleep timing (or later chronotype) are more likely to suffer from
279 sleep deficiency and its related health consequences. Indeed, several studies reported a relationship
280 between a late sleep timing and poor HRQoL. For instance, Chen et al. reported that a late self-reported
281 bedtime was associated with poor feelings, poor daily activities, pain and an overall poor rating of health
282 in a group of Japanese children ages 12-13.²³ Similarly, in another study of Israeli adolescents who were
283 ~14 years old, Tizischinsky et al. found that a later self-reported chronotype was associated with lower
284 HRQoL.²⁰ Additionally, in a sample of nearly 5,000 Australian children with self-reported sleep timing
285 data, those with a bedtime later than the sample median reported poorer HRQoL when compared to
286 those with an earlier bedtime.²² However, overall our results do not support a relationship between
287 device-measured sleep timing and HRQoL, which again may be due to the differences between device-
288 based and self-reported sleep measurements. Interestingly we found that later device-measured sleep
289 timing on weekends was associated with a higher HRQoL in England. This particular result could be due
290 to chance alone, or it could be explained by unmeasured cultural or environmental factors unique to
291 England.

292 Our study has some important strengths. As discussed above, we used both device-measured and self-
293 reported sleep measurements, which could provide different perspectives about sleep and its
294 relationship with health outcomes. Moreover, we studied an international sample of children from 12
295 countries with vastly different cultural and economic conditions, which has expanded the previous
296 literature that focused on developed regions alone. However, our study also has several limitations. First

297 of all, we did not have information on sports training and schedules of other important extracurricular
298 activities, which may have a substantial impact on both sleep and HRQoL. Second, although children
299 were recruited from different countries, all of the recruitment centers were located in relatively large
300 cities and our sample was not representative of the population in each country. Specifically we lacked
301 information on children living in rural communities. Third, the distribution of device-measured sleep
302 efficiency was limited in range and almost all children had high sleep efficiency, probably due to the
303 tendency of waist-worn actigraphy to overestimate sleep duration (by approximately 1 hour) compared
304 to wrist-worn protocols.⁴³ Fourth, children's sleep can be impacted by school schedules and a wide
305 range of weekday and weekend activities, such as sports and other extracurricular lessons, church
306 attendance and community activities, all of which may vary across different schools and societies.
307 Unfortunately we do not have detailed information on these activities, and could not control for their
308 influence on sleep and HRQoL in our analysis. Fifth, seasonal and geographic variations in the time of
309 sunrise and sunset are also known to have an impact on sleep, but we did not collect data on sunrise
310 and sunset time during the period of actigraphy assessment and weren't able to control for their
311 potential impact on our results. Moreover, we only recorded sleep for seven days, which may not
312 represent the habitual sleep patterns of children. Also, self-reported sleep information was only
313 obtained at one time point and may not capture children's long-term perceptions about sleep. Finally,
314 most of the children included in our study were relatively healthy and on average they reported high
315 HRQoL. Therefore we were not able to examine the relationship between sleep and HRQoL in children
316 with less healthy status.

317 In conclusion, while lower self-reported sleep quantity and quality were associated with poor HRQoL,
318 device-measured sleep characteristics were not. These findings highlight the importance of using both
319 device-based and subjective measures of sleep to fully understand its relationship with health
320 outcomes, particularly self-reported outcomes such as HRQoL.

321 **Acknowledgements**

322 We wish to thank the ISCOLE External Advisory Board and the ISCOLE participants and their families who
323 made this study possible. The ISCOLE Research Group includes: **Coordinating Center, Pennington**
324 **Biomedical Research Center:** Peter T. Katzmarzyk, PhD (Co-PI), Timothy S. Church, MD, PhD (Co-PI),
325 Denise G. Lambert, RN (Project Manager), Tiago Barreira, PhD, Stephanie Broyles, PhD, Ben Butitta, BS,
326 Catherine Champagne, PhD, RD, Shannon Cocreham, MBA, Kara D. Denstel, MPH, Katy Drazba, MPH,
327 Deirdre Harrington, PhD, William Johnson, PhD, Dione Milauskas, MS, Emily Mire, MS, Allison Tohme,
328 MPH, Ruben Rodarte MS, MBA; **Data Management Center, Wake Forest University:** Bobby Amoroso,
329 BS, John Luopa, BS, Rebecca Neiberg, MS, Scott Rushing, BS; **Australia, University of South Australia:**
330 Timothy Olds, PhD (Site Co-PI), Carol Maher, PhD (Site Co-PI), Lucy Lewis, PhD, Katia Ferrar, B Physio
331 (Hon), Effie Georgiadis, BPsych, Rebecca Stanley, BAppSc (OT) Hon; **Brazil, Centro de Estudos do**
332 **Laboratório de Aptidão Física de São Caetano do Sul (CELAFISCS):** Victor Keihan Rodrigues Matsudo,
333 MD, PhD (Site PI), Sandra Matsudo, MD, PhD, Timoteo Araujo, MSc, Luis Carlos de Oliveira, MSc, Luis
334 Fabiano, BSc, Diogo Bezerra, BSc, Gerson Ferrari, MSc; **Canada, Children’s Hospital of Eastern Ontario**
335 **Research Institute:** Mark S. Tremblay, PhD (Site Co-PI), Jean-Philippe Chaput, PhD (Site Co-PI), Priscilla
336 Bélanger, MA, Mike Borghese, MSc, Charles Boyer, MA, Allana LeBlanc, PhD, Claire Francis,
337 MSc, Geneviève Leduc, PhD; **China, Tianjin Women’s and Children’s Health Center:** Pei Zhao, MD (Site
338 Co-PI), Gang Hu, MD, PhD (Site Co-PI), Chengming Diao, MD, Wei Li, MD, Weiqin Li, MSc, Enqing Liu, MD,
339 Gongshu Liu, MD, Hongyan Liu, MD, Jian Ma, MD, Yijuan Qiao, MD, Huiguang Tian, PhD, Yue Wang, MD,
340 Tao Zhang, MSc, Fuxia Zhang, MD; **Colombia, Universidad de los Andes:** Olga Sarmiento, MD, PhD (Site
341 PI), Julio Acosta, Yalta Alvira, BS, Maria Paula Diaz, Rocio Gamez, BS, Maria Paula Garcia, Luis Guillermo
342 Gómez, Lisseth Gonzalez, Silvia Gonzalez, RD, Carlos Grijalba, MD, Leidys Gutierrez, David Leal, Nicolas
343 Lemus, Etelvina Mahecha, BS, Maria Paula Mahecha, Rosalba Mahecha, BS, Andrea Ramirez, MD, Paola
344 Rios, MD, Andres Suarez, Camilo Triana; **Finland, University of Helsinki:** Mikael Fogelholm, ScD (Site-PI),
345 Elli Jalo, BS, Jemina Kivelä, Sari Räsänen, BS, Sanna Roito, BS, Taru Saloheimo, MS, Leena Arjanne; **India,**
346 **St. Johns Research Institute:** Anura Kurpad, MD, PhD (Site Co-PI), Rebecca Kuriyan, PhD (Site Co-PI),
347 Deepa P. Lokesh, BSc, Michelle Stephanie D’Almeida, BSc, Annie Mattilda R, MSc, Lygia Correa, BSc, Vijay
348 Dakshina Murthy, BSc; **Kenya, Kenyatta University:** Vincent Onywera, PhD (Site Co-PI), Mark S.
349 Tremblay, PhD (Site Co-PI), Lucy-Joy Wachira, PhD, Stella Muthuri, PhD; **Portugal, University of Porto:**
350 Jose Maia, PhD (Site PI), Alessandra da Silva Borges, BA, Sofia Oliveira Sá Cachada, MSc, Raquel Nichele
351 de Chaves, MSc, Thayse Natacha Queiroz Ferreira Gomes, PhD, MSc, Sara Isabel Sampaio Pereira, BA,
352 Daniel Monteiro de Vilhena e Santos, PhD, Fernanda Karina dos Santos, MSc, Pedro Gil Rodrigues da
353 Silva, BA, Michele Caroline de Souza, MSc; **South Africa, University of Cape Town:** Vicki Lambert, PhD
354 (Site PI), Matthew April, BSc (Hons), Monika Uys, BSc (Hons), Nirmala Naidoo, MSc, Nandi Synyanya,
355 Madelaine Carstens, BSc(Hons); **United Kingdom, University of Bath:** Martyn Standage, PhD (Site PI),
356 Sean Cumming, PhD, Clemens Drenowatz, PhD, Lydia Emm, MSc, Fiona Gillison, PhD, Julia Zakrzewski,
357 PhD; **United States, Pennington Biomedical Research Center:** Catrine Tudor-Locke, PhD (Site-PI), Ashley
358 Braud, Sheletta Donatto, MS, LDN, RD, Corbin Lemon, BS, Ana Jackson, BA, Ashunti Pearson, MS, Gina
359 Pennington, BS, LDN, RD, Daniel Ragus, BS, Ryan Roubion, John Schuna, Jr., PhD; Derek Wiltz. **The**
360 **ISCOLE External Advisory Board includes** Alan Batterham, PhD, Teesside University, Jacqueline Kerr,
361 PhD, University of California, San Diego; Michael Pratt, MD, Centers for Disease Control and Prevention,
362 Angelo Pietrobelli, MD, Verona University Medical School.

363

364 **Financial support**

365 This work was supported by The Coca-Cola Company. With the exception of requiring that the study be
366 global in nature, the funder had no role in the design and conduct of the study; collection, management,
367 analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and
368 decision to submit the manuscript for publication.

369 **References**

- 370 1. Paruthi S, Brooks LJ, D'Ambrosio C, et al. Recommended Amount of Sleep for Pediatric
371 Populations: A Consensus Statement of the American Academy of Sleep Medicine. *J Clin Sleep*
372 *Med.* 2016;12(6):785-786.
- 373 2. Chaput JP, Gray CE, Poitras VJ, et al. Systematic review of the relationships between sleep
374 duration and health indicators in school-aged children and youth. *Appl Physiol Nutr Metab.*
375 2016;41(6 Suppl 3):S266-282.
- 376 3. Fatima Y, Doi SA, Mamun AA. Longitudinal impact of sleep on overweight and obesity in children
377 and adolescents: a systematic review and bias-adjusted meta-analysis. *Obes Rev.*
378 2015;16(2):137-149.
- 379 4. Peach H, Gaultney JF, Reeve CL. Sleep characteristics, body mass index, and risk for hypertension
380 in young adolescents. *J Youth Adolesc.* 2015;44(2):271-284.
- 381 5. Paciencia I, Araujo J, Ramos E. Sleep duration and blood pressure: a longitudinal analysis from
382 early to late adolescence. *J Sleep Res.* 2016;25(6):702-708.
- 383 6. Rudnicka AR, Nightingale CM, Donin AS, et al. Sleep Duration and Risk of Type 2 Diabetes.
384 *Pediatrics.* 2017;140(3).
- 385 7. De Bernardi Rodrigues AM, da Silva Cde C, Vasques AC, et al. Association of Sleep Deprivation
386 With Reduction in Insulin Sensitivity as Assessed by the Hyperglycemic Clamp Technique in
387 Adolescents. *JAMA Pediatr.* 2016;170(5):487-494.
- 388 8. Roberts RE, Duong HT. The prospective association between sleep deprivation and depression
389 among adolescents. *Sleep.* 2014;37(2):239-244.
- 390 9. Ojio Y, Nishida A, Shimodera S, Togo F, Sasaki T. Sleep Duration Associated with the Lowest Risk
391 of Depression/Anxiety in Adolescents. *Sleep.* 2016;39(8):1555-1562.
- 392 10. Gruber R, Cassoff J, Frenette S, Wiebe S, Carrier J. Impact of sleep extension and restriction on
393 children's emotional lability and impulsivity. *Pediatrics.* 2012;130(5):e1155-1161.
- 394 11. Vriend JL, Davidson FD, Corkum PV, Rusak B, Chambers CT, McLaughlin EN. Manipulating sleep
395 duration alters emotional functioning and cognitive performance in children. *J Pediatr Psychol.*
396 2013;38(10):1058-1069.
- 397 12. Touchette E, Petit D, Seguin JR, Boivin M, Tremblay RE, Montplaisir JY. Associations between
398 sleep duration patterns and behavioral/cognitive functioning at school entry. *Sleep.*
399 2007;30(9):1213-1219.
- 400 13. Stea TH, Knutsen T, Torstveit MK. Association between short time in bed, health-risk behaviors
401 and poor academic achievement among Norwegian adolescents. *Sleep Med.* 2014;15(6):666-
402 671.
- 403 14. Astill RG, Van der Heijden KB, Van Ijzendoorn MH, Van Someren EJ. Sleep, cognition, and
404 behavioral problems in school-age children: a century of research meta-analyzed. *Psychol Bull.*
405 2012;138(6):1109-1138.
- 406 15. Segura-Jimenez V, Carbonell-Baeza A, Keating XD, Ruiz JR, Castro-Pinero J. Association of sleep
407 patterns with psychological positive health and health complaints in children and adolescents.
408 *Qual Life Res.* 2015;24(4):885-895.
- 409 16. Ravens-Sieberer U, Gosch A, Rajmil L, et al. KIDSCREEN-52 quality-of-life measure for children
410 and adolescents. *Expert Rev Pharmacoecon Outcomes Res.* 2005;5(3):353-364.
- 411 17. Roberts RE, Roberts CR, Duong HT. Sleepless in adolescence: prospective data on sleep
412 deprivation, health and functioning. *J Adolesc.* 2009;32(5):1045-1057.

- 413 18. Do YK, Shin E, Bautista MA, Foo K. The associations between self-reported sleep duration and
414 adolescent health outcomes: what is the role of time spent on Internet use? *Sleep Med.*
415 2013;14(2):195-200.
- 416 19. Perkinson-Gloor N, Lemola S, Grob A. Sleep duration, positive attitude toward life, and academic
417 achievement: the role of daytime tiredness, behavioral persistence, and school start times. *J*
418 *Adolesc.* 2013;36(2):311-318.
- 419 20. Tzischinsky O, Shochat T. Eveningness, sleep patterns, daytime functioning, and quality of life in
420 Israeli adolescents. *Chronobiol Int.* 2011;28(4):338-343.
- 421 21. Magee CA, Robinson L, Keane C. Sleep quality subtypes predict health-related quality of life in
422 children. *Sleep Med.* 2017;35:67-73.
- 423 22. Quach J, Price AMH, Bittman M, Hiscock H. Sleep timing and child and parent outcomes in
424 Australian 4-9-year-olds: a cross-sectional and longitudinal study. *Sleep Med.* 2016;22:39-46.
- 425 23. Chen X, Sekine M, Hamanishi S, et al. Lifestyles and health-related quality of life in Japanese
426 school children: a cross-sectional study. *Prev Med.* 2005;40(6):668-678.
- 427 24. Sampasa-Kanyinga H, Standage M, Tremblay MS, et al. Associations between meeting
428 combinations of 24-h movement guidelines and health-related quality of life in children from 12
429 countries. *Public Health.* 2017;153:16-24.
- 430 25. Price AMH, Quach J, Wake M, Bittman M, Hiscock H. Cross-sectional sleep thresholds for
431 optimal health and well-being in Australian 4-9-year-olds. *Sleep Med.* 2016;22:83-90.
- 432 26. Jalali-Farahani S, Amiri P, Chin YS. Are physical activity, sedentary behaviors and sleep duration
433 associated with body mass index-for-age and health-related quality of life among high school
434 boys and girls? *Health Qual Life Outcomes.* 2016;14:30.
- 435 27. Tsiros MD, Samaras MG, Coates AM, Olds T. Use-of-time and health-related quality of life in 10-
436 to 13-year-old children: not all screen time or physical activity minutes are the same. *Qual Life*
437 *Res.* 2017;26(11):3119-3129.
- 438 28. Lauderdale DS, Knutson KL, Yan LL, Liu K, Rathouz PJ. Self-reported and measured sleep
439 duration: how similar are they? *Epidemiology.* 2008;19(6):838-845.
- 440 29. Jackson CL, Patel SR, Jackson WB, 2nd, Lutsey PL, Redline S. Agreement between self-reported
441 and objectively measured sleep duration among white, black, Hispanic, and Chinese adults in
442 the United States: Multi-Ethnic Study of Atherosclerosis. *Sleep.* 2018;41(6).
- 443 30. Alfano CA, Patriquin MA, De Los Reyes A. Subjective - Objective Sleep Comparisons and
444 Discrepancies Among Clinically-Anxious and Healthy Children. *J Abnorm Child Psychol.*
445 2015;43(7):1343-1353.
- 446 31. Katzmarzyk PT, Barreira TV, Broyles ST, et al. The International Study of Childhood Obesity,
447 Lifestyle and the Environment (ISCOLE): design and methods. *BMC Public Health.* 2013;13:900.
- 448 32. Bank W. World development indicators 2012. *Washington, DC: Development Data Group, The*
449 *World Bank.* 2012.
- 450 33. UNDP. Human Development Report 2011, Sustainability and Equity: A Better Future for All.
451 UNDP New York; 2011.
- 452 34. Ravens-Sieberer U, Europe KG. *The Kidscreen questionnaires: quality of life questionnaires for*
453 *children and adolescents; handbook.* Pabst Science Publ.; 2006.
- 454 35. Parizi AS, Garmaroudi G, Fazel M, et al. Psychometric properties of KIDSCREEN health-related
455 quality of life questionnaire in Iranian adolescents. *Quality of Life Research.* 2014;23(7):2133-
456 2138.
- 457 36. Masquillier C, Wouters E, Loos J, Nöstlinger C. Measuring health-related quality of life of HIV-
458 positive adolescents in resource-constrained settings. *PloS one.* 2012;7(7):e40628.

- 459 37. Tudor-Locke C, Mire EF, Barreira TV, et al. Nocturnal sleep-related variables from 24-h free-living
460 waist-worn accelerometry: International Study of Childhood Obesity, Lifestyle and the
461 Environment. *Int J Obes Suppl.* 2015;5(Suppl 2):S47-52.
- 462 38. Barreira TV, Schuna JM, Jr., Mire EF, et al. Identifying children's nocturnal sleep using 24-h waist
463 accelerometry. *Med Sci Sports Exerc.* 2015;47(5):937-943.
- 464 39. O'Connor A, Wellenius G. Rural-urban disparities in the prevalence of diabetes and coronary
465 heart disease. *Public Health.* 2012;126(10):813-820.
- 466 40. Saloheimo T, González S, Erkkola M, et al. The reliability and validity of a short food frequency
467 questionnaire among 9–11-year olds: a multinational study on three middle-income and high-
468 income countries. *International journal of obesity supplements.* 2015;5(S2):S22.
- 469 41. Onis Md, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO
470 growth reference for school-aged children and adolescents. *Bulletin of the World health
471 Organization.* 2007;85:660-667.
- 472 42. Dolsen MR, Wyatt JK, Harvey AG. Sleep, Circadian Rhythms, and Risk Across Health Domains in
473 Adolescents With an Evening Circadian Preference. *J Clin Child Adolesc Psychol.* 2018:1-11.
- 474 43. Hjorth MF, Chaput J-P, Damsgaard CT, et al. Measure of sleep and physical activity by a single
475 accelerometer: can a waist-worn Actigraph adequately measure sleep in children? *Sleep and
476 Biological Rhythms.* 2012;10(4):328-335.
- 477

Table 1 Study characteristics by study site in 6,266 children in ISCOLE (2011-2013).

	No.	Female (%)	Age, yr mean (SD)	zBMI ^a mean (SD)	Healthy diet score mean (SD)	Unhealthy diet score mean (SD)	Highest Parental Education (%) ^b			Parental Marital Status (%) ^b			
							Less than high school	High school or some college	College or higher	Married	Divorced or separated	Never Married	Widowed
Overall	6266	54.8	10.4 (0.6)	0.46 (1.26)	0.00 (1.00)	-0.05 (0.95)	19.0	40.5	36.3	64.0	14.0	16.0	1.7
Australia (Adelaide)	464	53.7	10.7 (0.4)	0.58 (1.12)	0.24 (0.96)	-0.30 (0.73)	11.2	45.7	40.5	72.0	19.6	4.5	0.7
Brazil (Sao Paulo)	479	51.4	10.5 (0.5)	0.84 (1.41)	-0.43 (1.05)	0.10 (0.90)	21.7	50.3	20.9	52.2	15.7	21.9	2.9
Canada (Ottawa)	513	59.1	10.5 (0.4)	0.41 (1.21)	0.47 (0.98)	-0.49 (0.57)	2.0	26.1	70.8	75.8	13.8	8.8	0.8
China (Tianjin)	470	48.7	9.9 (0.5)	0.73 (1.54)	0.07 (0.90)	-0.26 (0.93)	34.5	44.5	20.6	93.0	6.2	0.0	0.6
Colombia (Bogota)	843	50.7	10.5 (0.6)	0.21 (1.04)	-0.45 (0.74)	-0.08 (0.55)	30.5	51.3	18.2	24.9	24.4	47.6	3.0
England (Bath and North East Somerset)	424	56.8	10.9 (0.5)	0.45 (1.09)	0.03 (0.92)	-0.15 (0.76)	2.8	46.0	42.5	59.4	18.9	13.0	0.2
Finland (Helsinki, Espoo, and Vantaa)	464	54.7	10.5 (0.4)	0.26 (1.04)	-0.16 (0.86)	-0.55 (0.44)	2.6	51.1	40.5	61.4	17.5	14.7	0.4
India (Bangalore)	548	54.4	10.4 (0.5)	0.23 (1.36)	-0.08 (0.89)	-0.10 (0.83)	5.1	21.2	72.6	95.6	1.1	0.4	1.8
Kenya (Nairobi)	469	54.8	10.2 (0.7)	-0.03 (1.21)	0.27 (0.99)	0.12 (1.01)	14.1	46.3	39.5	81.7	6.8	6.8	4.1
Portugal (Porto)	651	56.8	10.4 (0.3)	0.85 (1.15)	0.22 (1.04)	-0.35 (0.66)	41.2	30.0	19.1	69.1	14.3	7.1	0.5
South Africa (Cape Town)	461	61.2	10.3 (0.7)	0.27 (1.29)	0.23 (1.08)	1.14 (1.20)	41.7	32.3	12.4	52.5	8.7	20.4	3.9
USA (Baton Rouge)	480	58.1	9.9 (0.6)	0.74 (1.29)	-0.14 (1.14)	0.63 (1.40)	6.0	41.7	50.8	53.5	15.4	27.7	1.0

^a Body mass index z-score (World Health Organization)

^b Percentages do not add up to 100% due to missingness

Abbreviations: ISCOLE, International Study of Childhood Obesity, Lifestyle and the Environment; USA, United States of America

Table 2 Distribution of HRQoL and objective and subjective sleep variables by study site in ISCOLE

	Australia	Brazil	Canada	China	Colombia	England	Finland	India	Kenya	Portugal	South Africa	USA
HRQoL (mean (SD))	49.86 (8.63)	47.23 (7.72)	51.18 (9.2)	51.2 (11.48)	49.9 (8.11)	50.07 (8.78)	52.62 (8.71)	48.14 (9.26)	47.07 (9.86)	53.04 (10.54)	49.34 (10.88)	50.81 (10.37)
Actigraph sleep variables, (mean (SD))												
<i>No. of valid nights (mean (SD))</i>	5.67 (0.67)	5.67 (0.62)	5.73 (0.59)	5.76 (0.59)	5.69 (0.66)	5.50 (0.80)	4.79 (0.59)	5.44 (0.72)	5.17 (0.93)	5.67 (0.68)	5.57 (0.76)	5.28 (0.84)
<i>Weekday</i>												
Total sleep time, h	9.53 (0.74)	8.34 (1.04)	9.18 (0.92)	8.57 (0.69)	8.38 (0.99)	9.44 (0.79)	8.41 (1.02)	8.35 (0.84)	8.30 (0.98)	8.12 (0.95)	9.06 (0.87)	8.79 (1.05)
Sleep onset, HH:MM	21:19 (0:46)	23:59 (1:10)	21:03 (0:50)	21:01 (0:41)	21:11 (0:50)	21:07 (0:43)	22:25 (0:56)	22:28 (0:47)	21:19 (0:44)	23:53 (0:55)	21:20 (0:44)	21:04 (0:53)
Sleep offset, HH:MM	7:08 (0:38)	7:17 (1:23)	7:03 (0:44)	6:28 (0:24)	6:07 (0:49)	7:14 (0:35)	6:55 (0:39)	6:48 (0:40)	5:54 (0:44)	7:10 (0:42)	6:38 (0:42)	6:38 (0:45)
Midpoint of sleep, HH:MM	2:23 (0:35)	3:08 (1:10)	2:28 (0:38)	2:01 (0:26)	1:56 (0:40)	2:31 (0:31)	2:43 (0:38)	2:38 (0:36)	1:46 (0:33)	3:07 (0:40)	2:07 (0:34)	2:15 (0:39)
Sleep efficiency	0.95 (0.01)	0.96 (0.01)	0.96 (0.01)	0.97 (0.01)	0.96 (0.01)	0.96 (0.01)	0.97 (0.01)	0.97 (0.01)	0.96 (0.02)	0.97 (0.01)	0.96 (0.02)	0.96 (0.01)
<i>Weekend</i>												
Total sleep time, h	9.23 (1.35)	9.00 (1.42)	8.88 (1.24)	9.22 (1.11)	9.62 (1.27)	9.61 (1.34)	8.55 (1.45)	9.13 (1.19)	9.30 (1.37)	8.58 (1.39)	9.43 (1.36)	9.04 (1.37)
Sleep onset, HH:MM	22:22 (1:13)	23:20 (1:23)	22:12 (1:10)	22:34 (1:01)	22:44 (1:13)	22:23 (1:07)	23:44 (1:07)	22:09 (1:01)	21:01 (1:05)	23:13 (1:10)	22:40 (1:07)	22:04 (1:18)
Sleep offset, HH:MM	7:46 (1:10)	8:37 (1:22)	7:35 (1:09)	7:35 (1:02)	7:49 (1:14)	8:08 (1:09)	7:46 (1:10)	7:55 (1:08)	7:12 (1:10)	8:18 (1:18)	7:42 (1:09)	7:55 (1:16)
Midpoint of sleep, HH:MM	3:10 (1:00)	4:07 (1:11)	3:10 (0:59)	2:59 (0:52)	3:01 (1:03)	3:21 (0:55)	3:29 (0:54)	3:21 (0:54)	2:34 (0:54)	4:01 (1:02)	2:59 (0:55)	3:24 (1:06)
Sleep efficiency	0.96 (0.02)	0.96 (0.02)	0.96 (0.02)	0.96 (0.02)	0.96 (0.02)	0.96 (0.02)	0.97 (0.01)	0.97 (0.01)	0.96 (0.02)	0.97 (0.01)	0.96 (0.02)	0.96 (0.02)
<i>Overall variability^a</i>												
Total sleep time, h	1.04 (0.60)	1.50 (0.70)	1.11 (0.58)	0.98 (0.53)	1.52 (0.73)	1.06 (0.56)	1.24 (0.70)	1.13 (0.61)	1.17 (0.69)	1.23 (0.59)	1.21 (0.64)	1.26 (0.73)
Sleep onset, h	0.88 (0.48)	1.04 (0.54)	0.92 (0.52)	0.76 (0.41)	0.90 (0.46)	0.81 (0.44)	0.97 (0.66)	0.75 (0.46)	0.73 (0.45)	0.98 (0.53)	0.92 (0.5)	1.05 (0.56)
Sleep offset, h	0.82 (0.49)	1.34 (0.62)	0.85 (0.47)	0.81 (0.48)	1.43 (0.70)	0.88 (0.50)	0.97 (0.49)	0.97 (0.53)	1.01 (0.67)	1.09 (0.49)	0.97 (0.57)	1.08 (0.63)
Midpoint of sleep, h	0.69 (0.36)	0.96 (0.41)	0.71 (0.37)	0.64 (0.33)	0.94 (0.44)	0.69 (0.35)	0.79 (0.40)	0.68 (0.36)	0.69 (0.40)	0.85 (0.39)	0.76 (0.38)	0.88 (0.43)
Sleep efficiency	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.01 (0)	0.01 (0.01)	0.01 (0.01)	0.01 (0)	0.01 (0.01)	0.01 (0.01)
<i>Weekday-to-weekend difference^b</i>												
Total sleep time, h	-0.30 (1.41)	0.66 (1.75)	-0.31 (1.27)	0.66 (1.15)	1.24 (1.49)	0.17 (1.37)	0.14 (1.53)	0.78 (1.29)	1.00 (1.42)	0.46 (1.48)	0.37 (1.57)	0.24 (1.54)
Sleep onset, h	0.94 (1.08)	0.65 (1.12)	0.84 (1.05)	0.45 (0.89)	0.44 (1.09)	0.74 (0.96)	0.68 (1.16)	0.31 (0.90)	0.29 (1.02)	0.67 (1.14)	0.67 (1.08)	1.01 (1.18)
Sleep offset, h	0.65 (1.09)	1.33 (1.50)	0.55 (1.00)	1.11 (1.00)	1.69 (1.28)	0.92 (1.10)	0.85 (1.14)	1.11 (1.08)	1.30 (1.20)	1.14 (1.26)	1.06 (1.17)	1.27 (1.24)
Midpoint of sleep, h	0.80 (0.83)	0.99 (1.00)	0.70 (0.82)	0.79 (0.75)	1.07 (0.93)	0.83 (0.78)	0.77 (0.86)	0.71 (0.76)	0.80 (0.86)	0.91 (0.96)	0.87 (0.82)	1.14 (0.94)
Sleep efficiency	0 (0.01)	0 (0.02)	0 (0.01)	-0.01 (0.02)	0 (0.02)	0 (0.01)	0 (0.01)	0 (0.01)	0 (0.02)	0 (0.01)	0 (0.02)	0 (0.02)
Self-reported sleep variables, N (%)												
Sleep quantity, fairly bad or very bad	49 (8.2)	26 (5.4)	45 (8.8)	67 (14.3)	60 (7.1)	49 (11.6)	40 (8.6)	74 (13.5)	40 (8.5)	31 (4.8)	54 (11.7)	61 (12.7)
Sleep quality, fairly bad or very bad	39 (8.4)	22 (4.6)	42 (8.2)	72 (15.3)	35 (4.2)	64 (15.1)	33 (7.1)	44 (8.0)	15 (3.2)	24 (3.7)	33 (7.2)	59 (12.3)

^a Measured as standard deviation of all valid nights

^b Measured as the average of sleep variables for weekends minus that for weekdays

Abbreviations: HRQoL, health-related quality of life; ISCOLE, International Study of Childhood Obesity, Lifestyle and the Environment SD, standard deviation; USA, United States of America

Table 3 Multivariable adjusted^a associations between HRQoL and objectively-measured total sleep time in ISCOLE

	Quartiles of Sleep Variable				<i>p</i> -for-trend	1 h increase
	Q 1	Q 2	Q 3	Q 4		
	Weekday total sleep time, h					
Median and IQR	7.51 (7.08, 7.80)	8.38 (8.21, 8.53)	8.99 (8.83, 9.16)	9.78 (9.53, 10.14)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	-1.04 (-6.63, 4.55)	-2.47 (-5.16, 0.21)	-0.92 (-2.67, 0.82)	ref	0.08	0.56 (-0.53, 1.65)
Brazil	-2.00 (-4.03, 0.04)	-2.05 (-4.19, 0.10)	-3.06 (-5.34, -0.78)	ref	0.25	0.65 (-0.02, 1.32)
Canada	-0.50 (-3.20, 2.20)	-0.03 (-2.30, 2.23)	-0.38 (-2.30, 1.54)	ref	0.85	0.05 (-0.81, 0.92)
China	-3.38 (-7.23, 0.47)	-0.42 (-3.89, 3.04)	-3.14 (-6.69, 0.41)	ref	0.61	0.52 (-0.95, 1.99)
Colombia	-0.38 (-2.01, 1.25)	-0.27 (-1.95, 1.41)	0.53 (-1.26, 2.32)	ref	0.38	0.19 (-0.36, 0.73)
England	0.30 (-5.14, 5.73)	0.81 (-1.78, 3.41)	0.46 (-1.42, 2.35)	ref	0.54	-0.39 (-1.45, 0.66)
Finland	-0.28 (-2.75, 2.19)	-1.29 (-3.83, 1.24)	-1.13 (-3.63, 1.36)	ref	0.97	0.08 (-0.71, 0.87)
India	-0.20 (-2.79, 2.40)	-1.14 (-3.70, 1.42)	-0.37 (-3.05, 2.31)	ref	0.93	-0.27 (-1.19, 0.65)
Kenya	-0.29 (-3.00, 2.41)	-0.61 (-3.37, 2.15)	-0.42 (-3.41, 2.57)	ref	0.89	0.05 (-0.81, 0.91)
Portugal	0.80 (-2.20, 3.81)	0.66 (-2.47, 3.79)	1.56 (-1.68, 4.81)	ref	0.95	0.07 (-0.76, 0.90)
South Africa	2.90 (-0.52, 6.32)	0.04 (-2.57, 2.65)	-0.76 (-3.07, 1.54)	ref	0.24	-0.51 (-1.62, 0.60)
USA	1.34 (-1.45, 4.12)	1.04 (-1.57, 3.66)	0.74 (-1.72, 3.21)	ref	0.32	-0.53 (-1.43, 0.37)
Overall	-0.40 (-1.09, 0.30)	-0.40 (-1.07, 0.28)	-0.38 (-1.03, 0.28)	ref	0.30	0.09 (-0.15, 0.33)
	Weekend total sleep time, h					
Median and IQR	7.68 (7.00, 8.08)	8.82 (8.60, 9.01)	9.57 (9.37, 9.77)	10.50 (10.20, 11.03)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	-0.25 (-2.45, 1.95)	-1.28 (-3.49, 0.93)	-0.44 (-2.51, 1.64)	ref	0.63	0.00 (-0.57, 0.57)
Brazil	-0.60 (-2.60, 1.41)	0.22 (-1.86, 2.30)	-0.67 (-2.72, 1.38)	ref	0.76	0.12 (-0.37, 0.61)
Canada	0.42 (-2.11, 2.95)	1.56 (-0.95, 4.07)	-0.78 (-3.24, 1.68)	ref	0.31	-0.46 (-1.13, 0.20)
China	-0.19 (-3.39, 3.01)	-0.51 (-3.33, 2.31)	0.26 (-2.57, 3.09)	ref	0.74	-0.05 (-0.96, 0.85)
Colombia	-0.61 (-2.27, 1.05)	0.75 (-0.74, 2.24)	-0.63 (-1.97, 0.70)	ref	0.94	0.01 (-0.41, 0.43)
England	-2.35 (-4.82, 0.12)	-0.92 (-3.21, 1.37)	1.10 (-0.98, 3.19)	ref	0.05	0.61 (-0.02, 1.23)
Finland	-2.19 (-4.68, 0.30)	-0.08 (-2.73, 2.58)	-0.51 (-3.19, 2.18)	ref	0.05	0.25 (-0.31, 0.80)
India	0.26 (-1.94, 2.46)	1.05 (-1.00, 3.10)	-1.19 (-3.34, 0.97)	ref	0.33	0.04 (-0.57, 0.66)
Kenya	-1.30 (-3.79, 1.18)	-0.51 (-2.81, 1.79)	-1.39 (-3.66, 0.87)	ref	0.45	0.26 (-0.36, 0.88)
Portugal	-0.13 (-2.61, 2.36)	-0.76 (-3.42, 1.89)	1.12 (-1.62, 3.85)	ref	0.52	0.24 (-0.34, 0.82)
South Africa	-0.32 (-3.12, 2.48)	-0.44 (-3.02, 2.14)	0.39 (-2.22, 3.00)	ref	0.71	0.08 (-0.66, 0.81)
USA	0.86 (-1.85, 3.57)	-0.18 (-2.81, 2.45)	-0.39 (-3.20, 2.43)	ref	0.50	-0.33 (-1.01, 0.36)

Overall		-0.20 (-0.87, 0.47)	0.07 (-0.58, 0.72)	-0.14 (-0.78, 0.51)	ref	0.72	-0.01 (-0.18, 0.17)
Overall variability total sleep time, ^b h							
Median and IQR		0.56 (0.44, 0.65)	0.91 (0.82, 0.99)	1.29 (1.18, 1.42)	2.00 (1.74, 2.43)		
Adjusted mean difference and 95% CI in HRQoL							
Australia	ref	1.43 (-0.48, 3.34)	0.12 (-2.12, 2.36)	0.77 (-1.50, 3.05)	0.65	-0.23 (-1.54, 1.07)	
Brazil	ref	0.07 (-2.66, 2.80)	-0.40 (-3.02, 2.23)	-1.04 (-3.63, 1.54)	0.22	-0.80 (-1.78, 0.18)	
Canada	ref	0.63 (-1.41, 2.68)	2.23 (0.11, 4.36)	0.94 (-1.45, 3.33)	0.15	0.94 (-0.43, 2.30)	
China	ref	0.22 (-2.23, 2.67)	0.06 (-2.77, 2.88)	1.11 (-2.33, 4.55)	0.63	0.09 (-1.80, 1.97)	
Colombia	ref	0.09 (-1.87, 2.04)	-0.57 (-2.44, 1.29)	0.50 (-1.26, 2.25)	0.50	0.22 (-0.51, 0.95)	
England	ref	-1.12 (-3.23, 1.00)	-2.99 (-5.12, -0.86)	0.57 (-1.92, 3.05)	0.39	-0.18 (-1.63, 1.27)	
Finland	ref	-2.10 (-4.44, 0.23)	-1.68 (-3.90, 0.54)	-1.41 (-3.55, 0.74)	0.26	-0.42 (-1.55, 0.71)	
India	ref	-0.13 (-2.12, 1.86)	-0.26 (-2.33, 1.80)	1.25 (-0.89, 3.40)	0.34	0.85 (-0.37, 2.08)	
Kenya	ref	-0.27 (-2.54, 2.00)	0.25 (-2.10, 2.60)	0.35 (-2.06, 2.76)	0.70	-0.13 (-1.35, 1.09)	
Portugal	ref	-1.31 (-3.71, 1.09)	0.27 (-1.97, 2.50)	-1.02 (-3.38, 1.34)	0.75	-0.62 (-1.96, 0.72)	
South Africa	ref	-1.52 (-4.24, 1.20)	1.53 (-1.13, 4.20)	-0.86 (-3.66, 1.94)	0.83	0.22 (-1.26, 1.70)	
USA	ref	2.40 (-0.18, 4.97)	1.17 (-1.51, 3.84)	-0.06 (-2.59, 2.46)	0.70	-0.74 (-2.05, 0.56)	
Overall	ref	-0.11 (-0.75, 0.54)	0.13 (-0.52, 0.78)	0 (-0.66, 0.65)	0.84	-0.11 (-0.46, 0.24)	
Weekday-to-weekend difference of total sleep time, ^c h							
Median and IQR		-1.18 (-1.76, -0.78)	0.03 (-0.20, 0.24)	0.89 (0.67, 1.13)	2.17 (1.72, 2.76)		
Adjusted mean difference and 95% CI in HRQoL							
Australia		-0.65 (-2.42, 1.12)	ref	0.30 (-2.06, 2.65)	-1.89 (-4.86, 1.09)	0.95	-0.14 (-0.68, 0.40)
Brazil		-0.69 (-2.75, 1.37)	ref	-2.20 (-4.29, -0.11)	-0.81 (-2.80, 1.18)	0.52	-0.13 (-0.54, 0.27)
Canada		-0.01 (-1.84, 1.81)	ref	-1.16 (-3.47, 1.16)	-0.38 (-3.50, 2.73)	0.46	-0.41 (-1.03, 0.21)
China		-1.21 (-4.64, 2.21)	ref	-1.76 (-4.31, 0.79)	-0.94 (-3.77, 1.88)	0.65	-0.21 (-1.08, 0.66)
Colombia		-0.81 (-2.87, 1.25)	ref	-0.95 (-2.61, 0.71)	-0.58 (-2.13, 0.96)	0.87	-0.08 (-0.44, 0.28)
England		-2.22 (-4.28, -0.16)	ref	1.42 (-0.84, 3.68)	-1.02 (-3.58, 1.53)	0.06	0.68 (0.09, 1.27)
Finland		0.55 (-1.50, 2.60)	ref	-0.21 (-2.45, 2.03)	1.16 (-1.18, 3.49)	0.77	0.18 (-0.33, 0.70)
India		2.13 (-0.23, 4.50)	ref	0.59 (-1.38, 2.55)	1.53 (-0.46, 3.52)	0.89	0.14 (-0.44, 0.72)
Kenya		1.84 (-1.17, 4.84)	ref	1.67 (-0.84, 4.17)	2.03 (-0.41, 4.47)	0.41	0.22 (-0.38, 0.82)
Portugal		0.44 (-1.77, 2.65)	ref	0.45 (-1.83, 2.73)	1.77 (-0.49, 4.03)	0.23	0.18 (-0.36, 0.72)
South Africa		-2.90 (-5.55, -0.26)	ref	-1.91 (-4.63, 0.81)	-0.82 (-3.63, 1.99)	0.28	0.22 (-0.40, 0.83)
USA		-0.86 (-3.29, 1.58)	ref	-0.03 (-2.69, 2.64)	-1.00 (-3.72, 1.73)	0.98	-0.02 (-0.62, 0.59)
Overall		-0.25 (-0.90, 0.39)	ref	-0.40 (-1.04, 0.25)	-0.15 (-0.81, 0.51)	0.94	-0.03 (-0.19, 0.12)

^a Models were adjusted for age, sex, parental education, parental marital status, household income, healthy diet score, unhealthy diet score and body-mass index. School was included as a random effect.

^b Measured as standard deviation of all valid nights.

^c Measured as the weekend sleep duration – weekday sleep duration.

Abbreviation: CI, confidence interval; HRQoL, health-related quality of life; IQR, interquartile range; ISCOLE, International Study of Childhood Obesity, Lifestyle and the Environment USA, United States of America.

1 **Table 4** Multivariable adjusted^a associations between HRQoL and objectively-measured sleep midpoint in ISCOLE.

	Quartiles of Sleep Variable				<i>p</i> -for-trend	1 h increase
	Q 1	Q 2	Q 3	Q 4		
Weekday sleep midpoint, HH:MM						
Median and IQR	1:35 (1:20, 1:46)	2:09 (2:02, 2:15)	2:36 (2:29, 2:43)	3:16 (3:03, 3:38)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	ref	-2.07 (-4.23, 0.10)	-1.01 (-3.14, 1.12)	-1.10 (-3.50, 1.30)	0.54	-0.58 (-1.92, 0.77)
Brazil	ref	-0.74 (-3.47, 1.99)	-0.31 (-2.80, 2.18)	0.74 (-1.50, 2.97)	0.24	0.22 (-0.39, 0.82)
Canada	ref	0.72 (-1.56, 3.00)	-0.02 (-2.32, 2.28)	-2.08 (-4.41, 0.25)	0.04	-1.45 (-2.68, -0.22)
China	ref	0.38 (-2.19, 2.95)	-0.73 (-3.66, 2.20)	-0.86 (-5.73, 4.00)	0.54	-1.06 (-3.53, 1.40)
Colombia	ref	0.42 (-0.93, 1.77)	-0.30 (-1.81, 1.21)	-0.44 (-2.46, 1.57)	0.65	-0.02 (-0.84, 0.80)
England	ref	2.75 (-0.34, 5.84)	3.25 (0.23, 6.28)	4.34 (1.13, 7.56)	0.01	2.52 (0.92, 4.13)
Finland	ref	-0.48 (-3.81, 2.86)	-0.32 (-3.56, 2.92)	-1.45 (-4.62, 1.72)	0.25	-0.59 (-1.83, 0.65)
India	ref	-0.51 (-3.41, 2.39)	-0.38 (-3.17, 2.41)	-1.79 (-4.62, 1.04)	0.14	-0.71 (-1.97, 0.55)
Kenya	ref	2.16 (0.19, 4.14)	-2.06 (-5.25, 1.14)	-2.24 (-6.69, 2.21)	0.55	-0.92 (-2.46, 0.62)
Portugal	ref	-3.07 (-9.58, 3.44)	0.52 (-5.48, 6.51)	1.21 (-4.68, 7.10)	0.03	1.48 (0.29, 2.67)
South Africa	ref	0.20 (-2.16, 2.56)	0.36 (-2.40, 3.13)	0.73 (-3.13, 4.60)	0.69	0.31 (-1.51, 2.13)
USA	ref	-1.88 (-4.24, 0.48)	-0.70 (-3.31, 1.92)	-3.25 (-6.19, -0.31)	0.08	-1.68 (-3.16, -0.20)
Overall	ref	0.22 (-0.45, 0.88)	0.41 (-0.28, 1.09)	0.25 (-0.48, 0.98)	0.42	0.11 (-0.23, 0.45)
Weekend sleep midpoint, HH:MM						
Median and IQR	2:08 (1:48, 2:21)	2:54 (2:43, 3:03)	3:33 (3:23, 3:45)	4:34 (4:13, 5:03)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	ref	-0.63 (-2.68, 1.41)	-1.68 (-3.83, 0.47)	-1.65 (-3.87, 0.57)	0.09	-0.70 (-1.48, .007)
Brazil	ref	-0.20 (-3.18, 2.79)	0.54 (-2.30, 3.38)	0.50 (-2.08, 3.07)	0.53	-0.05 (-0.65, 0.55)
Canada	ref	-1.25 (-3.35, 0.85)	-1.64 (-3.84, 0.56)	-1.56 (-3.92, 0.80)	0.15	-0.73 (-1.55, 0.09)
China	ref	-0.06 (-2.60, 2.48)	3.00 (0.16, 5.84)	0.05 (-3.38, 3.49)	0.27	0.08 (-1.12, 1.28)
Colombia	ref	-0.95 (-2.34, 0.45)	0.99 (-0.43, 2.4)	0.45 (-1.17, 2.07)	0.22	0.17 (-0.35, 0.68)
England	ref	3.13 (0.69, 5.56)	3.68 (1.18, 6.18)	5.39 (2.69, 8.09)	0.0002	1.62 (0.71, 2.54)
Finland	ref	1.13 (-1.49, 3.75)	0.01 (-2.44, 2.46)	-0.42 (-3.00, 2.15)	0.41	-0.45 (-1.35, 0.46)
India	ref	0.13 (-2.10, 2.35)	0.63 (-1.51, 2.77)	0.49 (-1.84, 2.82)	0.57	0.04 (-0.80, 0.87)
Kenya	ref	0.38 (-1.69, 2.46)	-0.51 (-3.03, 2.00)	-1.66 (-4.92, 1.60)	0.40	-0.16 (-1.09, 0.77)
Portugal	ref	2.52 (-1.21, 6.26)	2.09 (-1.50, 5.67)	3.00 (-0.43, 6.43)	0.15	0.91 (0.13, 1.69)
South Africa	ref	-0.69 (-3.05, 1.68)	0.84 (-1.81, 3.49)	-1.51 (-4.55, 1.54)	0.69	-0.49 (-1.54, 0.56)

USA	ref	-0.36 (-3.07, 2.34)	1.37 (-1.22, 3.97)	-0.26 (-3.02, 2.49)	0.81	0.09 (-0.78, 0.95)
Overall	ref	0.10 (-0.55, 0.75)	0.80 (0.14, 1.46)	0.42 (-0.26, 1.10)	0.07	0.13 (-0.10, 0.35)
Overall variability of sleep midpoint, h^b						
Median and IQR		0.36 (0.28, 0.42)	0.60 (0.54, 0.65)	0.84 (0.77, 0.93)		1.27 (1.13, 1.48)
Adjusted mean difference and 95% CI in HRQoL						
Australia	ref	2.01 (0.06, 3.95)	-1.70 (-3.84, 0.44)	-0.80 (-3.04, 1.45)	0.12	-1.66 (-3.77, 0.45)
Brazil	ref	-2.10 (-4.79, 0.60)	-1.67 (-4.17, 0.83)	-1.93 (-4.37, 0.51)	0.30	-1.57 (-3.28, 0.13)
Canada	ref	-0.25 (-2.27, 1.77)	0.92 (-1.32, 3.16)	-0.09 (-2.52, 2.34)	0.75	-0.003 (-2.20, 2.19)
China	ref	-0.34 (-2.78, 2.10)	4.08 (1.27, 6.90)	-0.02 (-3.21, 3.18)	0.17	1.37 (-1.69, 4.42)
Colombia	ref	-1.08 (-2.94, 0.78)	-0.79 (-2.59, 1.01)	-0.39 (-2.08, 1.30)	0.94	-0.09 (-1.32, 1.13)
England	ref	0.17 (-1.94, 2.28)	1.62 (-0.64, 3.87)	1.62 (-1.01, 4.25)	0.11	2.67 (0.25, 5.10)
Finland	ref	1.94 (-0.31, 4.20)	3.25 (1.05, 5.44)	0.66 (-1.60, 2.92)	0.32	0.12 (-1.89, 2.14)
India	ref	-0.06 (-1.97, 1.85)	0.37 (-1.56, 2.31)	0.92 (-1.40, 3.25)	0.43	0.90 (-1.15, 2.94)
Kenya	ref	0.99 (-1.15, 3.12)	-1.54 (-3.87, 0.78)	-1.11 (-3.51, 1.28)	0.17	-1.33 (-3.41, 0.75)
Portugal	ref	0.20 (-2.34, 2.74)	-0.20 (-2.60, 2.20)	0.83 (-1.59, 3.25)	0.54	0.21 (-1.86, 2.28)
South Africa	ref	-0.36 (-3.01, 2.28)	0.22 (-2.42, 2.86)	-0.68 (-3.40, 2.05)	0.76	-0.58 (-3.09, 1.93)
USA	ref	1.58 (-1.41, 4.56)	0.91 (-1.81, 3.64)	0.61 (-2.08, 3.29)	0.89	0.63 (-1.50, 2.76)
Overall	ref	0.37 (-0.27, 1.01)	0.52 (-0.13, 1.17)	0.17 (-0.50, 0.83)	0.55	-0.06 (-0.64, 0.52)
Weekday-to-weekend difference of sleep midpoint, h^c						
Median and IQR		-0.03 (-0.35, 0.17)	0.57 (0.44, 0.69)	1.06 (0.93, 1.22)		1.59 (1.87, 2.31)
Adjusted mean difference and 95% CI in HRQoL						
Australia	ref	-0.61 (-2.68, 1.46)	0.07 (-2.10, 2.23)	-2.09 (-4.27, 0.10)	0.12	-0.70 (-1.62, 0.20)
Brazil	ref	0 (-2.17, 2.16)	0.79 (-1.19, 2.77)	-0.5 (-2.39, 1.40)	0.72	-0.37 (-1.07, 0.34)
Canada	ref	-0.91 (-2.91, 1.08)	-0.78 (-2.97, 1.41)	-0.93 (-3.30, 1.44)	0.42	-0.13 (-1.13, 0.86)
China	ref	-0.65 (-3.38, 2.09)	0.53 (-2.30, 3.36)	2.26 (-0.80, 5.33)	0.12	0.40 (-0.96, 1.76)
Colombia	ref	-0.49 (-2.19, 1.21)	0.56 (-1.05, 2.18)	0.50 (-1.06, 2.05)	0.30	0.22 (-0.36, 0.81)
England	ref	2.67 (0.40, 4.94)	2.58 (0.25, 4.91)	2.66 (0.21, 5.12)	0.05	1.05 (-0.02, 2.11)
Finland	ref	0.27 (-1.94, 2.48)	1.05 (-1.20, 3.31)	0.21 (-2.11, 2.53)	0.68	-0.15 (-1.10, 0.80)
India	ref	0.39 (-1.54, 2.32)	0.37 (-1.68, 2.42)	0.59 (-1.75, 2.92)	0.63	0.48 (-0.50, 1.46)
Kenya	ref	3.18 (0.91, 5.45)	1.32 (-0.93, 3.57)	0.47 (-2.05, 3.00)	0.93	0.21 (-0.78, 1.20)
Portugal	ref	-1.25 (-3.59, 1.08)	-0.82 (-3.11, 1.46)	-0.39 (-2.58, 1.80)	0.83	0.32 (-0.52, 1.17)
South Africa	ref	0.30 (-2.38, 2.97)	-0.54 (-3.23, 2.16)	-0.98 (-3.76, 1.81)	0.40	-0.73 (-1.90, 0.43)
USA	ref	0.44 (-2.51, 3.38)	1.73 (-1.20, 4.65)	1.87 (-0.83, 4.57)	0.12	0.83 (-0.14, 1.81)
Overall	ref	0.12 (-0.52, 0.76)	0.43 (-0.22, 1.08)	0.20 (-0.45, 0.86)	0.38	0.11 (-0.16, 0.37)

2 ^a Models were adjusted for age, sex, parental education, parental marital status, household income, healthy diet score, unhealthy diet score
 3 and body-mass index. School was included as a random effect.

4 ^b Measured as standard deviation of all valid nights.

5 ^c Measured as weekend sleep midpoint – weekday sleep midpoints.

6 Abbreviation: CI, confidence interval; HRQoL, health-related quality of life; IQR, interquartile range; ISCOLE, International Study of Childhood
 7 Obesity, Lifestyle and the Environment USA, United States of America.

8 **Table 5** Multivariable adjusted ^a associations between HRQoL and objectively-measured sleep efficiency in ISCOLE.

	Quartiles of Sleep Variable				<i>p</i> -for-trend	1% increase
	Q 1	Q 2	Q 3	Q 4		
	Weekday sleep efficiency, %					
Median and IQR	94.5 (93.8, 95.1)	95.9 (95.6, 96.1)	96.8 (96.6, 97.0)	97.8 (97.5, 98.2)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	1.11 (-1.81, 4.03)	0.50 (-2.60, 3.59)	1.82 (-1.37, 5.02)	ref	0.88	-0.17 (-0.71, 0.37)
Brazil	0.41 (-2.04, 2.85)	-0.08 (-2.59, 2.43)	0.34 (-2.38, 3.06)	ref	0.78	-0.04 (-0.52, 0.44)
Canada	1.88 (-0.41, 4.17)	-0.03 (-2.28, 2.21)	-1.28 (-3.59, 1.03)	ref	0.05	-0.74 (-1.32, -0.16)
China	2.85 (-0.30, 5.99)	-0.68 (-3.56, 2.20)	0.79 (-1.73, 3.32)	ref	0.24	-0.48 (-1.30, 0.34)
Colombia	0.21 (-1.46, 1.88)	-0.75 (-2.41, 0.92)	-0.75 (-2.46, 0.97)	ref	0.58	-0.08 (-0.49, 0.32)
England	-1.65 (-4.39, 1.09)	-0.52 (-3.32, 2.29)	-1.07 (-3.96, 1.82)	ref	0.30	0.15 (-0.40, 0.71)
Finland	0.97 (-1.79, 3.73)	-0.19 (-2.32, 1.95)	-1.46 (-3.40, 0.49)	ref	0.67	-0.05 (-0.74, 0.65)
India	0.72 (-1.53, 2.96)	0.91 (-1.03, 2.85)	0.63 (-1.27, 2.54)	ref	0.39	-0.42 (-0.98, 0.14)
Kenya	0.77 (-1.71, 3.25)	-0.05 (-2.55, 2.44)	0.67 (-1.82, 3.16)	ref	0.69	-0.18 (-0.75, 0.36)
Portugal	-2.89 (-6.87, 1.09)	0.08 (-2.22, 2.37)	0.86 (-0.97, 2.68)	ref	0.57	0.14 (-0.66, 0.94)
South Africa	-1.77 (-4.47, 0.93)	-0.84 (-3.50, 1.83)	0.04 (-2.70, 2.78)	ref	0.15	0.46 (-0.17, 1.09)
USA	-0.17 (-2.83, 2.49)	-0.80 (-3.44, 1.83)	1.44 (-1.22, 4.09)	ref	0.46	0.09 (-0.59, 0.77)
Overall	0.07 (-0.61, 0.75)	-0.50 (-1.16, 0.16)	0.06 (-0.58, 0.71)	ref	0.74	-0.03 (-0.19, 0.14)
	Weekend sleep efficiency, %					
Median and IQR	94.3 (93.5, 94.8)	95.8 (95.5, 96.1)	96.8 (96.6, 97.1)	97.9 (97.6, 98.4)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	1.19 (-1.28, 3.65)	1.41 (-1.14, 3.97)	0.24 (-2.40, 2.88)	ref	0.24	-0.17 (-0.64, 0.29)
Brazil	0.08 (-2.2, 2.36)	-0.18 (-2.53, 2.18)	1.02 (-1.49, 3.52)	ref	0.66	0.09 (-0.35, 0.52)
Canada	2.37 (0.20, 4.54)	1.59 (-0.64, 3.82)	-0.77 (-2.91, 1.36)	ref	0.008	-0.72 (-1.22, 0.22)
China	0.26 (-2.67, 3.19)	0.27 (-2.61, 3.15)	0.65 (-2.19, 3.50)	ref	0.94	-0.13 (-0.72, 0.46)
Colombia	0.57 (-1.02, 2.17)	-0.48 (-2.06, 1.11)	0.70 (-0.98, 2.38)	ref	0.82	-0.06 (-0.39, 0.27)

England	-0.67 (-3.14, 1.8)	-0.43 (-2.91, 2.06)	-1.04 (-3.67, 1.58)	ref	0.79	0.03 (-0.44, 0.49)
Finland	-0.60 (-3.48, 2.29)	-0.42 (-2.62, 1.78)	-0.21 (-2.1, 1.69)	ref	0.62	0.07 (-0.54, 0.68)
India	2.60 (0.47, 4.73)	-0.55 (-2.56, 1.46)	-0.94 (-2.85, 0.96)	ref	0.05	-0.69 (-1.18, -0.21)
Kenya	0.77 (-1.71, 3.26)	-0.56 (-3.10, 1.97)	1.44 (-1.30, 4.17)	ref	0.95	-0.25 (-0.74, 0.23)
Portugal	-2.17 (-5.60, 1.25)	0.06 (-2.21, 2.34)	0.85 (-1.01, 2.70)	ref	0.53	0.15 (-0.53, 0.84)
South Africa	-0.10 (-2.73, 2.54)	0.02 (-2.71, 2.74)	1.28 (-1.60, 4.15)	ref	0.72	0.20 (-0.34, 0.73)
USA	-0.30 (-3.05, 2.46)	-1.37 (-4.05, 1.30)	-1.14 (-3.79, 1.50)	ref	0.85	0.08 (-0.49, 0.66)
Overall	0.15 (-0.52, 0.81)	-0.23 (-0.88, 0.42)	0.12 (-0.52, 0.77)	ref	0.93	-0.05 (-0.19, 0.09)

Overall variability of sleep efficiency, ^b %

Median and IQR	0.66 (0.52, 0.75)	1.03 (0.94, 1.10)	2.38 (1.29, 1.49)	2.00 (1.77, 2.33)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	ref	-1.85 (-4.25, 0.55)	-0.59 (-2.87, 1.7)	-1.13 (-3.38, 1.13)	0.63	-0.13 (-1.43, 1.18)
Brazil	ref	-0.23 (-2.42, 1.96)	0.84 (-1.31, 2.98)	-0.18 (-2.25, 1.88)	0.97	-0.37 (-1.43, 0.69)
Canada	ref	-1.93 (-4.03, 0.17)	-1.90 (-4.06, 0.26)	0.78 (-1.59, 3.14)	0.72	0.79 (-0.55, 2.12)
China	ref	-0.64 (-3.77, 2.49)	-2.84 (-5.98, 0.31)	0.04 (-3.06, 3.14)	0.89	0.29 (-1.21, 1.80)
Colombia	ref	0.06 (-1.73, 1.85)	0.19 (-1.49, 1.88)	0.33 (-1.31, 1.97)	0.65	0.05 (-0.77, 0.86)
England	ref	-0.93 (-3.43, 1.56)	-1.01 (-3.49, 1.47)	-1.94 (-4.42, 0.54)	0.14	-0.72 (-1.97, 0.54)
Finland	ref	0.94 (-1.06, 2.93)	-1.60 (-3.70, 0.51)	0.22 (-2.48, 2.93)	0.49	-0.51 (-2.12, 1.10)
India	ref	-0.47 (-2.52, 1.58)	-1.26 (-3.32, 0.81)	0.81 (-1.32, 2.94)	0.68	1.48 (0.17, 2.78)
Kenya	ref	-1.83 (-4.44, 0.79)	-1.49 (-3.97, 0.98)	-0.38 (-2.79, 2.04)	0.98	0.46 (-0.67, 1.59)
Portugal	ref	0.25 (-1.63, 2.12)	0.72 (-1.67, 3.11)	-2.25 (-5.94, 1.44)	0.75	-0.40 (-2.42, 1.62)
South Africa	ref	-0.12 (-3.05, 2.80)	-1.60 (-4.51, 1.32)	-1.80 (-4.77, 1.16)	0.12	-0.57 (-2.02, 0.89)
USA	ref	0.41 (-2.27, 3.09)	-1.17 (-3.86, 1.51)	-1.78 (-4.34, 0.78)	0.09	-0.77 (-2.20, 0.67)
Overall	ref	-0.57 (-1.21, 0.08)	-1.14 (-1.79, -0.48)	-0.69 (-1.35, -0.03)	0.02	-0.18 (-0.55, 0.18)

Weekday-to-weekend difference of sleep efficiency, ^c %

Median and IQR	-1.56 (-2.24, -1.17)	-0.41 (-0.61, -0.20)	0.38 (0.19, 0.58)	1.45 (1.08, 2.01)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	-0.50 (-2.86, 1.85)	ref	0.45 (-1.84, 2.75)	-0.17 (-2.25, 1.91)	0.97	-0.06 (-0.59, 0.48)
Brazil	1.13 (-0.87, 3.13)	ref	1.02 (-0.90, 2.95)	0.89 (-1.00, 2.78)	0.40	0.13 (-0.33, 0.59)
Canada	-0.41 (-2.86, 2.03)	ref	-1.01 (-3.33, 1.32)	-0.99 (-3.40, 1.41)	0.35	-0.25 (-0.84, 0.35)
China	0.61 (-1.96, 3.19)	ref	-0.49 (-3.25, 2.28)	1.22 (-1.74, 4.18)	0.64	0.14 (-0.50, 0.79)
Colombia	-0.02 (-1.51, 1.48)	ref	0.23 (-1.32, 1.79)	-0.18 (-1.63, 1.26)	0.88	-0.01 (-0.35, 0.35)
England	1.27 (-1.13, 3.67)	ref	-0.79 (-3.15, 1.58)	0.23 (-2.04, 2.51)	0.74	-0.12 (-0.68, 0.44)
Finland	-1.04 (-3.39, 1.31)	ref	-0.37 (-2.75, 2)	0.55 (-1.80, 2.89)	0.43	0.11 (-0.53, 0.76)
India	-0.56 (-2.54, 1.43)	ref	-0.49 (-2.61, 1.62)	-1.66 (-3.78, 0.45)	0.15	-0.53 (-1.12, 0.05)

Kenya	-1.57 (-3.90, 0.76)	ref	-0.99 (-3.31, 1.33)	0.21 (-2.10, 2.52)	0.86	-0.13 (-0.64, 0.37)
Portugal	-0.84 (-3.22, 1.55)	ref	-1.07 (-3.43, 1.28)	0.81 (-1.95, 3.56)	0.70	0.09 (-0.78, 0.96)
South Africa	0.07 (-2.54, 2.68)	ref	-0.08 (-2.7, 2.55)	-0.85 (-3.54, 1.84)	0.56	-0.17 (-0.76, 0.43)
USA	0.96 (-1.69, 3.62)	ref	1.98 (-0.65, 4.61)	0.34 (-2.19, 2.86)	0.71	0.02 (-0.58, 0.62)
Overall	0.15 (-0.49, 0.79)	ref	-0.05 (-0.7, 0.6)	0.09 (-0.56, 0.73)	0.95	-0.04 (-0.20, 0.11)

9 ^a Models were adjusted for age, sex, parental education, parental marital status, household income, healthy diet score, unhealthy index score
10 and body-mass index. School was included as a random effect.

11 ^b Measured as standard deviation of all valid nights.

12 ^c Measured as the difference between weekday and weekend means.

13 Abbreviation: CI, confidence interval; HRQoL, health-related quality of life; IQR, interquartile range; ISCOLE, International Study of Childhood
14 Obesity, Lifestyle and the Environment; USA, United States of America.

15 **Table 6** Multivariable adjusted ^a associations between HRQoL and subjectively-measured sleep quantity and quality in ISCOLE

	Sleep Rating				<i>P-for-trend</i>
	Very good	Fairly good	Fairly bad	Very bad	
Sleep Quantity					
N (%)	2858 (45.6)	2812 (44.9)	437 (7.0)	159 (2.5)	
Adjusted mean difference and 95% CI in HRQoL					
Australia	ref	-3.58 (-5.27, -1.88)	-6.09 (-8.94, -3.24)	2.14 (-5.09, 9.36)	<.0001
Brazil	ref	-1.83 (-3.26, -0.39)	-6.17 (-10.16, -2.19)	-7.65 (-12.19, -3.10)	<.0001
Canada	ref	-5.47 (-7.11, -3.84)	-10.08 (-13.15, -7.02)	-9.70 (-15.68, -3.72)	<.0001
China	ref	-6.18 (-8.25, -4.11)	-10.28 (-13.47, -7.09)	-13.40 (-19.03, -7.77)	<.0001
Colombia	ref	-2.16 (-3.30, -1.02)	-4.83 (-7.45, -2.21)	-4.77 (-8.02, -1.52)	<.0001
England	ref	-3.34 (-5.21, -1.48)	-7.43 (-10.44, -4.43)	-8.43 (-14.43, -2.43)	<.0001
Finland	ref	-5.13 (-6.75, -3.50)	-7.74 (-10.77, -4.70)	-9.06 (-17.34, -0.78)	<.0001
India	ref	1.29 (-0.29, 2.87)	-3.50 (-6.04, -0.97)	-2.68 (-6.84, 1.49)	0.08
Kenya	ref	-1.95 (-3.70, -0.20)	-6.64 (-10.31, -2.97)	-1.53 (-6.67, 3.60)	0.002
Portugal	ref	-2.18 (-3.82, -0.55)	-5.02 (-9.46, -0.58)	1.37 (-5.44, 8.18)	0.01
South Africa	ref	-1.94 (-4.08, 0.19)	-5.12 (-9.34, -0.89)	-4.44 (-8.38, -0.51)	0.002
USA	ref	-1.81 (-3.78, 0.17)	-4.58 (-7.84, -1.33)	-3.67 (-8.66, 1.33)	0.003
Overall	ref	-2.56 (-3.04, -2.08)	-6.21 (-7.12, -5.30)	-5.31 (-6.75, -3.88)	<.0001
Sleep Quality					
N (%)	3501 (55.9)	2283 (36.4)	338 (5.4)	144 (2.3)	
Adjusted mean difference and 95% CI in HRQoL					
Australia	ref	-2.64 (-4.30, -0.99)	-4.28 (-7.58, -0.99)	-0.95 (-6.25, 4.35)	0.004
Brazil	ref	-2.31 (-3.86, -0.77)	-3.14 (-6.81, 0.53)	-7.37 (-14.08, -0.66)	0.0002
Canada	ref	-5.02 (-6.61, -3.42)	-5.50 (-8.84, -2.16)	-9.93 (-15.24, -4.62)	<.0001

China	ref	-6.06 (-8.15, -3.98)	-9.92 (-13.18, -6.67)	-11.88 (-16.33, -7.42)	<.0001
Colombia	ref	-3.38 (-4.61, -2.14)	-5.89 (-9.41, -2.37)	-7.61 (-11.45, -3.76)	<.0001
England	ref	-3.74 (-5.50, -1.97)	-7.65 (-10.27, -5.03)	-11.36 (-16.39, -6.32)	<.0001
Finland	ref	-4.81 (-6.36, -3.26)	-7.78 (-10.96, -4.6)	-7.40 (-15.55, 0.75)	<.0001
India	ref	-3.00 (-4.52, -1.48)	-6.36 (-9.47, -3.26)	-5.70 (-10.91, -0.50)	<.0001
Kenya	ref	-2.23 (-4.28, -0.17)	-9.88 (-16.69, -3.08)	-3.22 (-9.59, 3.16)	0.004
Portugal	ref	-2.29 (-3.93, -0.65)	-7.67 (-12.55, -2.78)	0.78 (-7.49, 9.04)	0.001
South Africa	ref	-4.80 (-7.58, -2.02)	-5.26 (-9.87, -0.64)	-8.95 (-14.54, -3.36)	<.0001
USA	ref	-4.63 (-6.61, -2.66)	-5.33 (-8.93, -1.73)	-8.41 (-12.41, -4.40)	<.0001
Overall	ref	-3.43 (-3.92, -2.94)	-6.08 (-7.09, -5.07)	-7.34 (-8.83, -5.85)	<.0001

16 ^a Models were adjusted for age, sex, parental education, parental marital status, household income, healthy diet score, unhealthy diet score and
17 body-mass index. School was included as a random effect.
18 Abbreviation: CI, confidence interval; HRQoL, health-related quality of life; ISCOLE, International Study of Childhood Obesity, Lifestyle and the
19 Environment; USA, United States of America.
20

21 **Supplementary Table 1** Multivariable adjusted^a associations between HRQoL and timing of sleep onset in ISCOLE.

	Quartiles of Sleep Variable				<i>p-for-trend</i>	1 h increase
	Q 1	Q 2	Q 3	Q 4		
Weekday timing of sleep onset, HH:MM						
Median and IQR of sleep variable	21:01 (20:44, 21:13)	21:43 (21:34, 21:52)	22:08 (22:13, 22:30)	22:55 (23:00, 23:30)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	ref	-1.91 (-3.78, -0.03)	-0.79 (-2.95, 1.38)	-1.53 (-4.43, 1.36)	0.23	-0.64 (-1.70, 0.42)
Brazil	ref	-0.50 (-3.58, 2.57)	-0.24 (-3.05, 2.57)	0.06 (-2.49, 2.60)	0.74	-0.04 (-0.64, 0.55)
Canada	ref	-1.75 (-3.79, 0.28)	-1.46 (-3.61, 0.69)	-1.02 (-3.4, 1.36)	0.33	-0.86 (-1.80, 0.08)
China	ref	0.89 (-1.87, 3.64)	-0.09 (-3.00, 2.82)	-1.56 (-5.28, 2.17)	0.37	-0.72 (-2.25, 0.81)
Colombia	ref	0.92 (-0.45, 2.29)	0.08 (-1.35, 1.51)	-0.34 (-2.04, 1.36)	0.72	-0.15 (-0.80, 0.49)
England	ref	2.17 (0.09, 4.24)	1.36 (-0.88, 3.6)	3.81 (0.65, 6.98)	0.05	1.60 (0.42, 2.78)
Finland	ref	-0.69 (-3.85, 2.48)	-1.02 (-4.03, 1.98)	-1.65 (-4.61, 1.31)	0.21	-0.31 (-1.15, 0.52)
India	ref	-1.14 (-4.43, 2.14)	-0.46 (-3.66, 2.73)	-0.82 (-3.99, 2.35)	0.92	-0.28 (-1.25, 0.69)
Kenya	ref	-0.08 (-2.06, 1.91)	0.45 (-1.89, 2.79)	-1.82 (-4.99, 1.36)	0.59	-0.55 (-1.70, 0.61)
Portugal	ref	-0.83 (-6.87, 5.21)	4.73 (-0.89, 10.34)	3.41 (-2.07, 8.89)	0.08	0.73 (-0.14, 1.60)
South Africa	ref	-2.81 (-5.12, -0.49)	-0.2 (-2.77, 2.38)	2.69 (-1.29, 6.67)	0.53	0.54 (-2.72, 1.23)
USA	ref	0.17 (-2.2, 2.54)	1.01 (-1.56, 3.58)	-0.47 (-3.30, 2.36)	0.98	-0.43 (-1.49, 0.64)
Overall	ref	-0.32 (-0.97, 0.33)	0.32 (-0.35, 0.98)	0.09 (-0.63, 0.81)	0.43	0.01 (-0.25, 0.27)
Weekend timing of sleep onset, HH:MM						
IQR of sleep variable	21:20 (20:55, 21:36)	22:14 (22:03, 22:26)	23:00 (22:48, 23:13)	0:13 (23:48, 0:52)		
Adjusted mean difference and 95% CI in HRQoL						
Australia	ref	-1.24 (-3.3, 0.82)	-2.1 (-4.24, 0.04)	-1.3 (-3.61, 1.01)	0.15	-0.44 (-1.06, 0.17)
Brazil	ref	1.88 (-1.08, 4.85)	2.04 (-0.61, 4.68)	1.61 (-0.84, 4.05)	0.44	-0.10 (-0.61, 0.40)
Canada	ref	-1.00 (-3.15, 1.15)	-1.72 (-4.05, 0.61)	-0.99 (-3.31, 1.32)	0.33	-0.26 (-0.93, 0.42)
China	ref	-0.13 (-2.74, 2.47)	2.29 (-0.46, 5.03)	0.75 (-2.72, 4.22)	0.22	0.08 (-0.92, 1.09)
Colombia	ref	0.22 (-1.08, 1.53)	1.12 (-0.43, 2.66)	0.46 (-1.22, 2.15)	0.29	0.12 (-0.31, 0.56)
England	ref	2.07 (-0.16, 4.31)	1.63 (-0.70, 3.96)	1.74 (-0.87, 4.35)	0.23	0.65 (-0.11, 1.41)
Finland	ref	-0.15 (-3.13, 2.82)	0.89 (-1.95, 3.74)	-1.30 (-4.10, 1.51)	0.27	-0.47 (-1.18, 0.23)
India	ref	0.70 (-1.64, 3.05)	0.52 (-1.75, 2.80)	0.28 (-2.30, 2.86)	0.94	-0.02 (-0.75, 0.72)
Kenya	ref	-1.35 (-3.38, 0.68)	1.39 (-1.13, 3.90)	-0.18 (-3.44, 3.07)	0.76	-0.27 (-1.04, 0.49)
Portugal	ref	5.70 (0.95, 10.45)	7.60 (3.14, 12.05)	6.22 (1.86, 10.58)	0.19	0.51 (-0.17, 1.18)
South Africa	ref	-2.14 (-4.45, 0.17)	-0.86 (-3.55, 1.84)	-1.1 (-4.24, 2.03)	0.45	-0.38 (-1.24, 0.47)
USA	ref	-0.62 (-3.41, 2.18)	1.71 (-0.93, 4.36)	0.04 (-2.68, 2.77)	0.57	0.20 (-0.51, 0.91)
Overall	ref	0.06 (-0.59, 0.71)	0.98 (0.31, 1.64)	0.35 (-0.34, 1.04)	0.07	0.08 (-0.10, 0.27)
Overall variability of timing of sleep onset, h ^b						
IQR of sleep variable	0.41 (0.31, 0.47)	0.65 (0.59, 0.72)	0.94 (0.86, 1.05)	1.49 (1.30, 1.82)		

Adjusted mean difference and 95% CI in HRQoL

Australia	ref	-1.40 (-3.57, 0.77)	-0.83 (-3.01, 1.34)	-2.14 (-4.36, 0.08)	0.11	-1.24 (-2.80, 0.32)
Brazil	ref	0.42 (-1.93, 2.77)	1.98 (-0.32, 4.29)	0.37 (-1.85, 2.59)	0.69	-0.50 (-1.79, 0.79)
Canada	ref	-0.12 (-2.34, 2.10)	1.19 (-1.07, 3.45)	1.29 (-1.00, 3.58)	0.16	1.47 (-0.04, 2.97)
China	ref	0.73 (-1.87, 3.33)	0.86 (-1.85, 3.56)	1.82 (-1.27, 4.90)	0.26	0.62 (-1.79, 3.03)
Colombia	ref	-0.31 (-1.89, 1.26)	0.17 (-1.43, 1.77)	-0.22 (-1.88, 1.43)	0.99	-0.12 (-1.26, 1.01)
England	ref	0.30 (-1.90, 2.50)	-0.87 (-3.13, 1.40)	-0.63 (-2.98, 1.73)	0.43	-0.21 (-2.07, 1.64)
Finland	ref	0.51 (-1.92, 2.94)	1.02 (-1.26, 3.30)	0.41 (-1.73, 2.56)	0.64	-0.30 (-1.50, 0.89)
India	ref	-1.33 (-3.19, 0.53)	0.35 (-1.64, 2.34)	0.37 (-1.90, 2.63)	0.58	0.38 (-1.19, 1.94)
Kenya	ref	-0.16 (-2.29, 1.97)	0.38 (-1.99, 2.75)	-0.54 (-2.95, 1.88)	0.81	-0.19 (-2.02, 1.64)
Portugal	ref	-2.38 (-4.84, 0.08)	-2.05 (-4.44, 0.33)	-1.22 (-3.49, 1.05)	0.53	-0.52 (-2.01, 0.96)
South Africa	ref	-1.41 (-4.29, 1.47)	0.05 (-2.80, 2.90)	-0.14 (-3.07, 2.79)	0.70	0.38 (-1.47, 2.22)
USA	ref	-3.09 (-6.12, -0.05)	-1.12 (-3.9, 1.66)	-1.32 (-4.02, 1.38)	0.74	0.48 (-1.14, 2.10)
Overall	ref	-0.50 (-1.14, 0.14)	0.13 (-0.51, 0.78)	-0.06 (-0.72, 0.59)	0.67	0.05 (-0.39, 0.50)

Weekend-weekday variability of timing of sleep onset, h^c

IQR of sleep variable		-0.48 (-0.85, -0.23)	0.27 (0.13, 0.40)	0.85 (0.70, 1.03)	1.82 (1.46, 2.42)	
Adjusted mean difference and 95% CI in HRQoL						
Australia	ref	-0.12 (-2.68, 2.44)	0.65 (-1.43, 2.74)	-0.91 (-2.94, 1.12)	0.47	-0.28 (-0.97, 0.41)
Brazil	ref	1.12 (-0.92, 3.17)	0.85 (-1.14, 2.84)	0.29 (-1.63, 2.21)	0.60	-0.11 (-0.73, 0.50)
Canada	ref	1.13 (-1.31, 3.57)	-0.09 (-2.25, 2.07)	1.44 (-0.70, 3.58)	0.64	0.22 (-0.52, 0.97)
China	ref	-0.16 (-2.92, 2.59)	0.99 (-1.72, 3.69)	1.44 (-1.62, 4.5)	0.23	0.46 (-0.66, 1.59)
Colombia	ref	-0.15 (-1.58, 1.27)	1.06 (-0.47, 2.59)	0.73 (-0.88, 2.33)	0.12	0.24 (-0.24, 0.73)
England	ref	-0.93 (-3.43, 1.58)	0.86 (-1.41, 3.13)	-1.21 (-3.55, 1.13)	0.90	-0.01 (-0.87, 0.84)
Finland	ref	1.05 (-1.26, 3.36)	-0.81 (-3.01, 1.39)	1.29 (-0.88, 3.45)	0.87	-0.23 (-0.91, 0.45)
India	ref	-1.13 (-2.98, 0.72)	-0.82 (-2.83, 1.19)	-0.28 (-2.74, 2.18)	0.59	0.17 (-0.63, 0.97)
Kenya	ref	-0.47 (-2.57, 1.62)	0.31 (-2.27, 2.88)	-0.46 (-3.16, 2.24)	0.79	-0.04 (-0.87, 0.79)
Portugal	ref	-1.33 (-3.73, 1.06)	-2.60 (-4.96, -0.25)	-2.10 (-4.46, 0.25)	0.20	0.07 (-0.63, 0.76)
South Africa	ref	0.46 (-2.21, 3.12)	-0.5 (-3.22, 2.21)	-0.83 (-3.48, 1.81)	0.31	-0.63 (-1.51, 0.25)
USA	ref	-0.65 (-3.76, 2.47)	0.78 (-1.92, 3.48)	1.09 (-1.33, 3.51)	0.18	0.44 (-0.31, 1.20)
Overall	ref	-0.27 (-0.91, 0.37)	0.08 (-0.56, 0.72)	0.11 (-0.53, 0.76)	0.25	0.10 (-0.11, 0.31)

22 ^a Models were adjusted for age, sex, parental education, parental marital status, household income, healthy diet score, unhealthy index score
 23 and body-mass index. School was included as a random effect.

24 ^b Measured as standard deviation of all valid nights.

25 ^c Measured as weekend sleep midpoint – weekday sleep midpoints.

26 Abbreviation: CI, confidence interval; HRQoL, health-related quality of life; IQR, interquartile range; USA, United States of America.

27

28 **Supplementary Table 2** Distribution of objective sleep variables by subjective sleep categories in ISCOLE.

	Self-reported sleep quantity						Self-reported sleep quality					
	Very good	Fairly good	Fairly bad	Very bad	<i>p</i> -value ^a	Correlation Coefficient ^b	Very good	Fairly good	Fairly bad	Very bad	<i>p</i> -value ^a	Correlation Coefficient ^b
<i>Weekday</i>												
Total sleep time, h	8.63 (1.01)	8.70 (1.04)	8.65 (1.03)	8.54 (0.99)	0.01	0.03	8.61 (1.02)	8.73 (1.04)	8.72 (0.97)	8.79 (0.97)	<.0001	0.07
Sleep onset, HH:MM	22:04 (0:59)	22:07 (1:00)	22:04 (0:58)	22:04 (1:02)	0.20	0.02	22:05 (1:01)	22:07 (1:00)	22:10 (0:56)	23:53 (0:50)	0.02	0.01
Sleep offset, HH:MM	6:41 (0:55)	6:49 (0:52)	6:43 (0:46)	6:37 (0:52)	<.0001	0.05	6:41 (0:55)	6:50 (0:50)	6:52 (0:46)	6:40 (0:50)	<.0001	0.10
Midpoint of sleep, HH:MM	2:23 (0:49)	2:29 (0:46)	2:24 (0:43)	2:21 (0:49)	0.0001	0.04	2:23 (0:49)	2:24 (0:46)	2:32 (0:42)	2:17 (0:41)	<.0001	0.06
Sleep efficiency	0.96 (0.01)	0.96 (0.01)	0.96 (0.01)	0.96 (0.01)	0.73	0.00	0.96 (0.01)	0.96 (0.01)	0.96 (0.01)	0.96 (0.01)	0.67	0.01
<i>Weekend</i>												
Total sleep time, h	9.18 (1.39)	9.11 (1.36)	9.11 (1.29)	9.05 (1.16)	0.23	-0.03	9.19 (1.41)	9.07 (1.32)	9.1 (1.31)	9.18 (1.21)	0.02	-0.03
Sleep onset, HH:MM	22:41 (1:19)	22:47 (1:17)	22:43 (1:16)	22:36 (1:18)	0.006	0.03	22:41 (1:20)	22:49 (1:17)	22:49 (1:17)	22:32 (1:08)	0.0002	0.04
Sleep offset, HH:MM	7:51 (1:16)	7:54 (1:14)	7:50 (1:15)	7:39 (1:08)	0.07	-0.002	7:52 (1:17)	7:52 (1:12)	7:55 (1:13)	7:42 (1:04)	0.24	0.00
Midpoint of sleep, HH:MM	3:16 (1:05)	3:21 (1:04)	3:17 (1:05)	3:08 (1:04)	0.007	0.02	3:17 (1:06)	3:21 (1:03)	3:23 (1:04)	3:07 (0:55)	0.01	0.03
Sleep efficiency	0.96 (0.02)	0.96 (0.02)	0.96 (0.02)	0.96 (0.02)	0.74	0.01	0.96 (0.02)	0.96 (0.02)	0.96 (0.02)	0.96 (0.02)	0.08	0.02
<i>Overall variability^b</i>												
Total sleep time, h	1.25 (0.69)	1.21 (0.65)	1.18 (0.63)	1.19 (0.63)	0.09	-0.02	1.26 (0.69)	1.18 (0.65)	1.21 (0.62)	1.15 (0.6)	<.0001	-0.05
Sleep onset, h	0.91 (0.51)	0.91 (0.54)	0.89 (0.51)	0.92 (0.53)	0.83	-0.01	0.91 (0.51)	0.9 (0.54)	0.93 (0.52)	0.92 (0.51)	0.57	0.00
Sleep offset, h	1.08 (0.62)	1.01 (0.58)	1.02 (0.59)	1.02 (0.58)	0.0003	-0.05	1.08 (0.62)	1 (0.58)	0.98 (0.55)	0.93 (0.55)	<.0001	-0.07
Midpoint of sleep, h	0.8 (0.41)	0.77 (0.4)	0.77 (0.42)	0.8 (0.39)	0.05	-0.04	0.8 (0.41)	0.77 (0.4)	0.76 (0.39)	0.76 (0.38)	0.08	-0.03
Sleep efficiency	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.06	0.02	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.29	0.01
<i>Weekday-to-weekend difference^c</i>												
Total sleep time, h	0.54 (1.53)	0.41 (1.53)	0.47 (1.49)	0.51 (1.39)	0.01	-0.03	0.58 (1.56)	0.34 (1.46)	0.39 (1.52)	0.39 (1.4)	<.0001	-0.07
Sleep onset, h	0.61 (1.11)	0.67 (1.1)	0.65 (1.06)	0.53 (1.04)	0.13	0.02	0.6 (1.11)	0.69 (1.09)	0.66 (1.08)	0.65 (1.03)	0.002	0.04
Sleep offset, h	1.16 (1.24)	1.08 (1.21)	1.12 (1.26)	1.04 (1.16)	0.11	-0.03	1.18 (1.26)	1.04 (1.18)	1.04 (1.21)	1.03 (1.06)	<.0001	-0.06
Midpoint of sleep, h	0.88 (0.89)	0.87 (0.87)	0.89 (0.88)	0.79 (0.86)	0.60	-0.01	0.89 (0.89)	0.87 (0.87)	0.85 (0.86)	0.84 (0.78)	0.76	-0.01
Sleep efficiency	0 (0.01)	0 (0.01)	0 (0.01)	0 (0.02)	0.98	0.01	0 (0.01)	0 (0.01)	0 (0.01)	0 (0.01)	0.39	0.01

^a Derived from ANOVA test

^b Spearman correlation coefficient

^c Measured as standard deviation of all valid nights

^d Measured as the average of sleep variables for weekends minus that for weekdays

Abbreviations: HRQoL, health-related quality of life; ISCOLE, International Study of Childhood Obesity, Lifestyle and the Environment SD, standard deviation; USA, United States of America