

BRAIN COMMUNICATIONS

Prion protein codon 129 polymorphism in mild cognitive impairment and dementia: the Rotterdam Study

Hata Karamujić-Čomić,^{1,2} Shahzad Ahmad,¹ Thom S. Lysen,^{1,2} Alis Heshmatollah,^{1,2} Gennady V. Roshchupkin,^{1,3,4} Meike W. Vernooij,^{1,4} Annemieke J. M. Rozemuller,^{5,6} Mohammad Arfan Ikram,^{1,2} Najaf Amin¹ and Cornelia M. van Duijn^{1,2,7}

Creutzfeldt–Jakob disease is a rare, fatal, neurodegenerative disease caused by the accumulation of abnormally folded prion proteins. The common polymorphism at codon 129 (methionine/valine) in the prion protein (*PRNP*) gene is the most important determinant of genetic susceptibility. Homozygotes of either allele have a higher risk of sporadic Creutzfeldt–Jakob disease. Various studies suggest that this polymorphism is also involved in other forms of dementia. We studied the association between the codon 129 polymorphism of the *PRNP* gene and mild cognitive impairment in 3605 participants from the Rotterdam Study using logistic regression analyses. Subsequently, we studied the association between this polymorphism and incident dementia, including Alzheimer’s disease, in 11 070 participants using Cox proportional hazard models. Analyses were adjusted for age and sex. We found the prevalence of mild cognitive impairment to be higher for carriers of the methionine/methionine genotype (odds ratio, 1.40; 95% confidence interval, 1.11–1.78; $P=0.005$) as well as for carriers of the valine/valine genotype (odds ratio, 1.37; 95% confidence interval, 0.96–1.97; $P=0.08$). The codon 129 polymorphism was not associated with the risk of incident dementia or Alzheimer’s disease. In conclusion, we found a statistically significant higher prevalence of mild cognitive impairment in carriers of the methionine/methionine genotype in the codon 129 polymorphism of the *PRNP* gene within this population-based study. No associations were found between the codon 129 polymorphism and dementia or Alzheimer’s disease in the general population.

- 1 Department of Epidemiology, Erasmus MC, University Medical Center, Rotterdam, Netherlands
- 2 National Prion Disease Registry, Department of Epidemiology, Erasmus MC, University Medical Center, Rotterdam, Netherlands
- 3 Department of Medical Informatics, Erasmus MC, University Medical Center, Rotterdam, Netherlands
- 4 Department of Radiology and Nuclear Medicine, Erasmus MC, University Medical Center, Rotterdam, Netherlands
- 5 Department of Pathology, Amsterdam Neuroscience, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, Netherlands
- 6 Department of Pathology, UMC Utrecht, Utrecht, Netherlands
- 7 Clinical Trial Service Unit and Epidemiological Studies Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK

Correspondence to: Prof. Dr. Cornelia M. van Duijn, Clinical Trial Service Unit and Epidemiological Studies Unit, Nuffield Department of Population Health, University of Oxford, Old Road Campus, Oxford OX3 7LF, UK

E-mail: cornelia.vanduijn@ndph.ox.ac.uk


Keywords: prions; mild cognitive impairment; dementia

Received December 11, 2019. Revised February 12, 2020. Accepted February 17, 2020. Advance Access publication March 20, 2020

© The Author(s) (2020). Published by Oxford University Press on behalf of the Guarantors of Brain.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Graphical Abstract



Introduction

Creutzfeldt–Jakob disease is a rare, fatal, neurodegenerative prion disease caused by the accumulation of abnormally folded prion proteins. These proteins are the product of the prion protein (*PRNP*) gene. In prion diseases, the cellular prion protein (PrP^{C}) converts into a misfolded infectious state, the pathological prion protein (PrP^{Sc}) (Prusiner, 1982). Creutzfeldt–Jakob disease occurs in different forms: sporadic, genetic, iatrogenic, and variant. The sporadic form of Creutzfeldt–Jakob disease is the most common form. A major driver of the development and course of sporadic Creutzfeldt–Jakob disease is a genetic variant at codon 129 (methionine/valine) of the *PRNP* gene (Parchi *et al.*, 2012). Persons homozygous for either the methionine or valine allele are at increased risk of sporadic Creutzfeldt–Jakob disease (Alperovitch *et al.*, 1999; Parchi *et al.*, 2009). It has been speculated that homozygosity for either allele results in the synthesis of identical proteins, enabling the propagation and aggregation of *PRNP* in the brain, thus influencing both the risk and progression of sporadic Creutzfeldt–Jakob disease (Alperovitch *et al.*, 1999; Parchi *et al.*, 2009). Clinically, sporadic Creutzfeldt–Jakob disease is characterized by rapidly progressive dementia, and ultimately death on average 6 months after diagnosis. The variant form of Creutzfeldt–Jakob disease is the zoonotic form of the disease, which is transmitted through meat and other food products contaminated with material from cattle with bovine spongiform encephalopathy (Brandel *et al.*, 2009). Up to 2017, all patients diagnosed with variant Creutzfeldt–Jakob disease had the methionine/methionine (MM) genotype on *PRNP* codon 129. Recently, the first patient with the methionine/valine

(MV) genotype on *PRNP* codon 129 with the variant form was reported (Mok *et al.*, 2017).

While the dementia seen in Creutzfeldt–Jakob disease patients is likely a result of prion protein aggregation, various studies suggest that the *PRNP* gene is also involved in other forms of dementia, but results are inconsistent (Rujescu *et al.*, 2003; Del Bo *et al.*, 2006; Jeong *et al.*, 2007; Poleggi *et al.*, 2008; Choi *et al.*, 2010; Golanska *et al.*, 2013; He *et al.*, 2013; Zhang *et al.*, 2016). Furthermore, studies performed on early cognitive decline and early-onset Alzheimer’s disease also show inconsistency regarding which of the homozygous carriers have a higher risk (Croes *et al.*, 2003; Dermaut *et al.*, 2003). In contrast, a meta-analysis on the role of the M129V polymorphism in Alzheimer’s disease suggests that individuals with at least one valine allele have a lower risk of Alzheimer’s disease (He *et al.*, 2013), while the International Genetics of Alzheimer Disease Project did not find an association (Kunkle *et al.*, 2019).

In this study, we report on the association between the *PRNP* M129V polymorphism and mild cognitive impairment and dementia, including Alzheimer’s disease, within the population-based Rotterdam Study.

Materials and methods

Setting and study population

The Rotterdam Study is a prospective population-based middle-aged and elderly cohort that started in 1990 in the district of Ommoord, in Rotterdam, The Netherlands. The study includes 14 926 participants and has three

subcohorts (Ikram *et al.*, 2017). At start of the study, all inhabitants of the district of Ommoord who were aged 55 years and older were invited to participate. At baseline, in 1990–93, of the 10 215 invited inhabitants, 7983 agreed to participate in the baseline examination (response rate 78%). In 2000, the cohort was extended with 3011 participants (67% of invitees). This extension consisted of all persons living in the study district who had become 55 years and older or had moved into the study district. A second extension was initiated in 2006, in which 3932 participants (65% of invitees) who were 45 years and older were included. Study rounds consist of a home interview and visits with extensive investigations at the dedicated research centre. Rounds are repeated every 4–6 years. Between 2002 and 2005 extensive neuropsychological tests were implemented in the Rotterdam Study (de Bruijn *et al.*, 2014). These tests are a requirement for the assessment of mild cognitive impairment. Participants without contraindications are also invited to undergo a brain MRI scan, which has been implemented routinely since 2005, as discussed previously (Ikram *et al.*, 2015). Participants are continuously monitored for diseases and mortality through linkage of the medical records from the general practitioners and municipality records. The Rotterdam Study has been approved by the Medical Ethics Committee of the Erasmus MC and by the Ministry of Health, Welfare and Sport of The Netherlands. All participants provided written informed consent to participate in the study and to obtain information from their treating physicians.

Genotyping

A total of 11 496 participants who were genotyped passed genotyping quality control (92% of all subjects with genotyping) (Niemeijer *et al.*, 2015). Exclusion criteria were a call rate <98%, Hardy–Weinberg P -value <10⁻⁶, minor allele frequency <0.01%, excess autosomal heterozygosity >0.336, sex mismatch and outlying identity-by-state clustering estimates. Imputations were performed using the Haplotype Reference Consortium panel (Loh *et al.*, 2016). We selected one single-nucleotide polymorphism within the *PRNP* gene: M129V. Imputation quality for this single-nucleotide polymorphism was high ($r^2 = 0.97$). We rounded the imputed single-nucleotide polymorphism dosages to 0 (MM), 1 (MV) and 2 (valine/valine (VV)).

Assessment of mild cognitive impairment

We defined mild cognitive impairment using the following criteria: (i) presence of subjective cognitive complaints, (ii) presence of objective cognitive impairment and (iii) absence of dementia, as previously described (de Bruijn *et al.*, 2014). The first criterion, presence of subjective cognitive complaints, was evaluated by an interview,

which consisted three questions on memory complaints and three questions on impaired daily functioning. If the participant confirmed the presence of one of these six complaints or impairments, subjective cognitive complaints were seen as present. The second criterion, presence of objective cognitive impairment, was assessed using a cognitive test battery comprising letter digit test, Stroop test, Purdue Pegboard test, fluency task and 15-word verbal learning test based on Rey's recall of words. Participants were classified as objectively cognitive impaired if they scored <1.5 standard deviation of the age-adjusted and education-adjusted mean of the study population. This has previously been explained in more detail (de Bruijn *et al.*, 2014). For this study, we included participants with genetic data available and mild cognitive impairment assessment during the implementation period of extensive neuropsychological tests, as a large number of participants were then screened for mild cognitive impairment for the first time.

Assessment of dementia

Dementia ascertainment involved cognitive screening at the study research centre. We further assessed individuals with a Mini-Mental State Examination score of <26 or a Geriatric Mental State Schedule organic level of >0 (de Bruijn *et al.*, 2015), by administering the Cambridge Mental Disorders of the Elderly Examination by a research physician. We also interviewed spouses or informants. A consensus panel headed by a consultant neurologist established the final diagnosis according to the standard criteria. We studied the outcomes of all-cause dementia (DSM-III-R) and Alzheimer's disease (NINCDS–ADRDA). For the assessment of dementia, and type of dementia, the latest follow-up information with available data was used to determine the disease state. Follow-up for dementia was near complete until 1 January 2015. Within this period, participants were censored at the date of dementia diagnosis, death or loss to follow-up.

Brain MRI measurements

Brain MRI scanning was performed using a 1.5-T scanner (General Electric Healthcare, Milwaukee, WI, USA), and it included T₁-weighted sequence, proton-density-weighted sequence, fluid-attenuated inversion recovery-weighted sequence and T₂-weighted sequences (Ikram *et al.*, 2015). Details on methods for the measurement of brain volume and hippocampal volume have been described earlier (Ikram *et al.*, 2015).

Statistical analysis

Logistic regression analyses were used to study the association between the M129V polymorphism and mild cognitive impairment. For mild cognitive impairment as outcome of interest, analyses were conducted in all

participants who had mild cognitive impairment assessment at the implementation of neuropsychological tests. Subsequently, the association between the M129V polymorphism and incident dementia, including incident Alzheimer's disease, was studied using Cox proportional hazard models. For these two outcomes of interest, analyses were conducted in all participants who had dementia status and dementia type assessed. We ran the analyses unadjusted (Model 1) and also while adjusting for age and sex (Model 2). As secondary analyses, we ran linear regression analyses to study the association between the M129V polymorphism and MRI-derived brain volume and hippocampal volume, while adjusting for age, sex and intracranial volume.

Data availability statement

Because of restrictions based on privacy regulations and informed consent of the participants, data cannot be made freely available in a public repository. Data can be obtained upon reasonable request. Requests for the Rotterdam Study data should be directed towards the management team (secretariat.epi@erasmusmc.nl).


Results

Descriptive statistics

We examined the 11 496 participants in the Rotterdam Study who had genotyping data available from the M129V polymorphism from the *PRNP* gene (Fig. 1). Of these participants, 4991 (43.4%) had the MM genotype, 5209 (45.3%) had the MV genotype and the remaining 1295 (11.3%) had the VV genotype. The carriers of the MM genotype tended to be slightly older than the two other genotype groups. In the overall population, the distribution of this genotype was in Hardy–Weinberg equilibrium. The general characteristics of the study population are shown in Table 1.

PRNP M129V and the prevalence of mild cognitive impairment

We found that carriers of the MM genotype have a higher prevalence of mild cognitive impairment than those who are heterozygous (odds ratio, 1.40; 95% confidence interval, 1.11–1.78; $P=0.005$), as shown in Table 2. Carriers of the VV genotype showed a borderline significant higher prevalence of mild cognitive impairment than heterozygous carriers (odds ratio, 1.37; 95% confidence interval, 0.96–1.97; $P=0.08$). Combining homozygous carriers (MM and VV) resulted in an odds ratio of 1.40 (95% confidence interval, 1.11–1.75; $P=0.004$). The allele and genotype distribution of the M129V polymorphism was not in Hardy–Weinberg equilibrium in mild cognitive impairment cases. There was an



excess of homozygotes (both MM and VV carriers) in the mild cognitive impairment cases.

PRNP M129V and other outcomes

We did not find significant evidence for the association of *PRNP* M129V polymorphism with incident dementia, including Alzheimer's disease, as shown in Table 3. In dementia and Alzheimer's disease patients and controls, the M129V polymorphism was in Hardy–Weinberg equilibrium. We also did not find a significant association of the M129V polymorphism with brain volume and hippocampal volume.

Discussion

In this study, we investigated the association of the M129V polymorphism of the *PRNP* gene with mild cognitive impairment and dementia. We found that homozygous carriers had more often mild cognitive impairment than heterozygous carriers. We did not find an association between this polymorphism and dementia. We found a significantly higher prevalence of mild cognitive impairment in carriers of the MM genotype than in carriers of the MV genotype and a non-significant higher prevalence of mild cognitive impairment in carriers of the VV genotype than in carriers of the MV genotype. In contrast, our results do not indicate a role of the M129V polymorphism of the *PRNP* gene in dementia, including

Table 1 Characteristics of the study population

	Genotype at codon 129 of the prion protein gene			P
	MM N = 4991	MV N = 5209	VV N = 1296	
Age at baseline, median (range)	63.2 (45.6–99.1)	62.6 (45.5–99.2)	62.6 (45.7–99.2)	0.06
Sex, female, n (%)	2878 (58)	3068 (59)	726 (56)	0.13
APOEε4 carrier, n (%) ^a	1377 (29)	1451 (29)	369 (30)	0.82

^aFour percent of missings in each genotype group.

APOEε4, apolipoprotein E epsilon 4; n, number of people; N, number of people at risk; M, methionine; V, valine.

Table 2 Association of PRNP M129V polymorphism with mild cognitive impairment

Genotype	Cases, n (%)	Controls, n (%)	Model 1 OR (95% CI)	P	Model 2 OR (95% CI)	P
MM	182 (51)	1414 (44)	1.42 (1.13–1.80)	0.003	1.40 (1.11–1.78)	0.005
MV	133 (37)	1471 (45)	1.00 (reference)		1.00 (reference)	
VV	45 (13)	360 (11)	1.38 (0.97–1.98)	0.08	1.37 (0.96–1.97)	0.08

Hardy–Weinberg equilibrium P: overall: 0.95, cases: 0.0099, controls: 0.44.

CI, confidence interval; MM, methionine/methionine; MV, methionine/valine; n, number of people; OR, odds ratio; VV, valine/valine.

Table 3 Risk of incident dementia, including Alzheimer's disease, stratified on PRNP M129V polymorphism

Genotype	Cases, n (%)	Cohort at risk, n (%)	Model 1 HR (95% CI)	P	Model 2 HR (95% CI)	P
Dementia						
MM	602 (43)	4802 (43)	1.01 (0.90–1.13)	0.89	1.01 (0.90–1.13)	0.90
MV	631 (44)	5015 (45)	1.00 (reference)		1.00 (reference)	
VV	176 (13)	1253 (11)	1.12 (0.95–1.32)	0.18	1.16 (0.98–1.37)	0.09
Alzheimer's disease						
MM	480 (43)	4680 (44)	1.01 (0.89–1.15)	0.84	1.01 (0.87–1.15)	0.82
MV	500 (45)	4884 (45)	1.00 (reference)		1.00 (reference)	
VV	125 (11)	1202 (11)	1.02 (0.84–1.24)	0.87	1.06 (0.87–1.29)	0.56

Hardy–Weinberg equilibrium P for dementia: overall: 0.30, cases: 0.59, controls: 0.19. Hardy–Weinberg equilibrium P for Alzheimer's disease: overall: 0.18, cases: 0.76, controls: 0.19.

CI, confidence interval; HR, hazard ratio; MM, methionine/methionine; MV, methionine/valine; n, number of people; VV, valine/valine.

Alzheimer's disease, nor in brain volume and hippocampal volume. The mechanism underlying the association between the M129V polymorphism of the PRNP gene and neurodegeneration is thought to be caused by the effect the M129V polymorphism has on the aggregation of pathological prion proteins in the brain. The aggregation of pathological prion proteins is a biological process similar to the aggregation of tau protein and amyloid accumulation (Walker, 2018), which are key features in Alzheimer's disease (Bloom, 2014). While most neurodegenerative diseases are known to involve more than one misfolded or abnormally aggregated protein, a recent study demonstrated prion protein aggregation to be an independent pathogenic mechanism, with no cross-seeding between prion protein and misfolded amyloid beta (Rossi *et al.*, 2019), although prion protein is a stress protein that is elevated in (amyloid beta) plaques (Kellett and Hooper, 2009). This suggests that the neurodegenerative changes due to prion protein accumulation could occur regardless of other ongoing neurodegenerative processes.

This could be a possible explanation for our finding that the M129V polymorphism of the PRNP gene plays a role in mild cognitive impairment, and not in dementia. Our study showed that the M129V polymorphism of the PRNP gene may increase the prevalence of mild cognitive impairment. Mild cognitive impairment is often seen as pre-stage of Alzheimer's disease, related to apolipoprotein E. This study showed the heterogeneity of mild cognitive impairment as also other genes and other non-Alzheimer's disease dementing disorders are included in the mild cognitive impairment population.

Several limitations in this study should be addressed. Our analyses of the association of the M129V polymorphism with mild cognitive impairment have been performed cross-sectionally. We were not able to calculate the life-time risk of the M129V polymorphism on mild cognitive impairment in the participants from our population-based cohort study. Another potential limitation is the generalizability of our findings, as our study was performed in mainly Caucasians. Previous studies have

studied the effect of the M129V polymorphism of the *PRNP* gene in mild cognitive impairment and different types of dementia in Asians (Jeong *et al.*, 2007; Choi *et al.*, 2010; Zhang *et al.*, 2016). No association was found between the M129V polymorphism and mild cognitive impairment in a Korean population (Choi *et al.*, 2010). Our findings are therefore not generalizable to other ethnicities.

Our study also had several strengths. We used a population-based cohort study, and our study had a relatively large sample size. Although the sample size of the participants who underwent initial mild cognitive impairment assessment was smaller than the overall sample with available genotyping data and dementia assessment, we had sufficient power to demonstrate the association between the M129V polymorphism and mild cognitive impairment.

In conclusion, we found a statistically significant higher prevalence of mild cognitive impairment in carriers of the MM genotype in the M129V polymorphism of the *PRNP* gene in the Rotterdam Study, but no associations were found between this polymorphism and incidence of dementia, including Alzheimer's disease. Future studies should further elucidate the role of the M129V polymorphism of the *PRNP* gene in cognitive function, dementia and other neurodegenerative traits.

Acknowledgements

We are thankful to the study participants, the staff from the Rotterdam Study, the participating general practitioners and the participating pharmacists. We acknowledge F.J.A. van Rooij as data manager, B.C.T. Leening-Kieboom as study coordinator and J. Verkroost-van Heemst for her invaluable contribution to data collection. Furthermore, we acknowledge Prof. Dr. R.G. Will from the University of Edinburgh for sharing his invaluable expertise on prion diseases.

Funding

This work is funded by the European Union's Horizon 2020 research and innovation programme as part of the Common mechanisms and pathways in Stroke and Alzheimer's disease (CoSTREAM) project. The current study is supported by Memorabel supported by The Netherlands Organization for the Health Research and Development (ZonMw). The Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, Rotterdam, ZonMw, the Research Institute for Diseases in the Elderly (RIDE), the Ministry of Education, Culture and Science, the Ministry for Health, Welfare and Sports, the European Commission (DG XII) and the Municipality of Rotterdam.

Prof. Dr. Cornelia M. van Duijn is funded by the National Institute of Health (NIH), National Institute on Aging (NIA) with the following grants: Gut Liver Brain Biochemical Axis

in Alzheimer's Disease – R01 (Grant Project No. 1RF1AG058942-01) and Metabolomic Signatures for Disease Sub-classification and Target Prioritization in AMP-AD – U01 (Grant Project No. 5U01AG061359-02).

Competing interests

The authors report no competing interests.

References

- Alperovitch A, Zerr I, Pocchiari M, Mitrova E, de Pedro Cuesta J, Hegyi I, et al. Codon 129 prion protein genotype and sporadic Creutzfeldt-Jakob disease. *Lancet* 1999; 353: 1673–4.
- Bloom GS. Amyloid-beta and tau: the trigger and bullet in Alzheimer disease pathogenesis. *JAMA Neurol* 2014; 71: 505–8.
- Brandel JP, Heath CA, Head MW, Levavasseur E, Knight R, Laplanche JL, et al. Variant Creutzfeldt-Jakob disease in France and the United Kingdom: evidence for the same agent strain. *Ann Neurol* 2009; 65: 249–56.
- Choi IG, Woo SI, Kim HJ, Kim DJ, Park BL, Cheong HS, et al. Lack of association between *PRNP* M129V polymorphism and multiple sclerosis, mild cognitive impairment, alcoholism and schizophrenia in a Korean population. *Dis Markers* 2010; 28: 315–21.
- Croes EA, Dermaut B, Houwing-Duistermaat JJ, Van den Broeck M, Cruts M, Breteler MM, et al. Early cognitive decline is associated with prion protein codon 129 polymorphism. *Ann Neurol* 2003; 54: 275–6.
- de Bruijn RF, Akoudad S, Cremers LG, Hofman A, Niessen WJ, van der Lugt A, et al. Determinants, MRI correlates, and prognosis of mild cognitive impairment: the Rotterdam Study. *J Alzheimers Dis* 2014; 42: S239–49. Suppl
- de Bruijn RF, Bos MJ, Portegies ML, Hofman A, Franco OH, Koudstaal PJ, et al. The potential for prevention of dementia across two decades: the prospective, population-based Rotterdam Study. *BMC Med* 2015; 13: 132.
- Del Bo R, Scarlato M, Ghezzi S, Martinelli-Boneschi F, Fenoglio C, Galimberti G, et al. Is M129V of *PRNP* gene associated with Alzheimer's disease? A case-control study and a meta-analysis. *Neurobiol Aging* 2006; 27: 770.e1–5.
- Dermaut B, Croes EA, Rademakers R, Van den Broeck M, Cruts M, Hofman A, et al. *PRNP* Val129 homozygosity increases risk for early-onset Alzheimer's disease. *Ann Neurol* 2003; 53: 409–12.
- Golanska E, Sieruta M, Corder E, Gresner SM, Pfeffer A, Chodakowska-Zebrowska M, et al. The prion protein M129V polymorphism: longevity and cognitive impairment among Polish centenarians. *Prion* 2013; 7: 244–7.
- He J, Li X, Yang J, Huang J, Fu X, Zhang Y, et al. The association between the methionine/valine (M/V) polymorphism (rs1799990) in the *PRNP* gene and the risk of Alzheimer disease: an update by meta-analysis. *J Neurol Sci* 2013; 326: 89–95.
- Ikram MA, Brusselle GGO, Murad SD, van Duijn CM, Franco OH, Goedegeure A, et al. The Rotterdam Study: 2018 update on objectives, design and main results. *Eur J Epidemiol* 2017; 32: 807–50.
- Ikram MA, van der Lugt A, Niessen WJ, Koudstaal PJ, Krestin GP, Hofman A, et al. The Rotterdam Scan Study: design update 2016 and main findings. *Eur J Epidemiol* 2015; 30: 1299–315.
- Jeong BH, Lee KH, Jeong YE, Hwang KA, Lee YJ, Carp RI, et al. Polymorphisms at codons 129 and 219 of the prion protein gene (*PRNP*) are not associated with sporadic Alzheimer's disease in the Korean population. *Eur J Neurol* 2007; 14: 621–6.
- Kellett KA, Hooper NM. Prion protein and Alzheimer disease. *Prion* 2009; 3: 190–4.

- Kunkle BW, Grenier-Boley B, Sims R, Bis JC, Damotte V, Naj AC, et al.; Alzheimer Disease Genetics Consortium (ADGC). Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates Abeta, tau, immunity and lipid processing. *Nat Genet* 2019; 51: 414–30.
- Loh PR, Danecek P, Palamara PF, Fuchsberger C, Ar Y, H KF, et al. Reference-based phasing using the Haplotype Reference Consortium panel. *Nat Genet* 2016; 48: 1443–8.
- Mok T, Jaunmuktane Z, Joiner S, Campbell T, Morgan C, Wakerley B, et al. Variant Creutzfeldt-Jakob disease in a patient with heterozygosity at PRNP codon 129. *N Engl J Med* 2017; 376: 292–4.
- Niemeijer MN, van den Berg ME, Deckers JW, Aarnoudse AL, Hofman A, Franco OH, et al. ABCB1 gene variants, digoxin and risk of sudden cardiac death in a general population. *Heart* 2015; 101: 1973–9.
- Parchi P, de Boni L, Saverioni D, Cohen ML, Ferrer I, Gambetti P, et al. Consensus classification of human prion disease histotypes allows reliable identification of molecular subtypes: an inter-rater study among surveillance centres in Europe and USA. *Acta Neuropathol* 2012; 124: 517–29.
- Parchi P, Strammiello R, Notari S, Giese A, Langeveld JP, Ladogana A, et al. Incidence and spectrum of sporadic Creutzfeldt-Jakob disease variants with mixed phenotype and co-occurrence of PrP^{Sc} types: an updated classification. *Acta Neuropathol* 2009; 118: 659–71.
- Poleggi A, Bizzarro A, Acciarri A, Antuono P, Bagnoli S, Cellini E, et al. Codon 129 polymorphism of prion protein gene in sporadic Alzheimer's disease. *Eur J Neurol* 2008; 15: 173–8.
- Prusiner SB. Novel proteinaceous infectious particles cause scrapie. *Science* 1982; 216: 136–44.
- Rossi M, Kai H, Baiardi S, Bartoletti-Stella A, Carla B, Zenesini C, et al. The characterization of AD/PART co-pathology in CJD suggests independent pathogenic mechanisms and no cross-seeding between misfolded Abeta and prion proteins. *Acta Neuropathol Commun* 2019; 7: 53.
- Rujescu D, Hartmann AM, Gonnermann C, Moller HJ, Giegling I. M129V variation in the prion protein may influence cognitive performance. *Mol Psychiatry* 2003; 8: 937–41.
- Walker LC. Prion-like mechanisms in Alzheimer disease. *Handb Clin Neurol* 2018; 153: 303–19.
- Zhang W, Jiao B, Xiao T, Pan C, Liu X, Zhou L, et al. Mutational analysis of PRNP in Alzheimer's disease and frontotemporal dementia in China. *Sci Rep* 2016; 6: 38435.