

Scientific Foundation SPIROSKI, Skopje, Republic of Macedonia  
 Open Access Macedonian Journal of Medical Sciences. 2020 Mar 20; 8(B):114-118.  
<https://doi.org/10.3889/oamjms.2020.3895>  
 eISSN: 1857-9655  
 Category: B - Clinical Sciences  
 Section: Radiology and Radiotherapy



# Differentiating Benign from Suspicious Vertebral Marrow Lesions Detected with Conventional Magnetic Resonance Imaging Using Apparent Diffusion Coefficient and Diffusion-Weighted Image

Khaleel Ibraheem Mohson<sup>1\*</sup>, Qusay Tayser Naief<sup>2</sup>, Farah Abdul Jalil<sup>1</sup>

<sup>1</sup>National Cancer Research Center, University of Baghdad, Iraq; <sup>2</sup>Magnetic Resonance Unit, Oncology Teaching Hospital, Medical City Complex, Iraq

## Abstract

**Edited by:** Slavica Hristomanova-Mitkovska  
**Citation:** Mohson KI, Naief QT, Jalil FA. Differentiating Benign from Suspicious Vertebral Marrow Lesions Detected with Conventional Magnetic Resonance Imaging Using Apparent Diffusion Coefficient and Diffusion-Weighted Image. Open Access Maced J Med Sci. 2020 Mar 20; 8(B):114-118. <https://doi.org/10.3889/oamjms.2020.3895>  
**Keywords:** Vertebral metastasis; Magnetic resonance imaging; Apparent diffusion coefficient; Diffusion-weighted imaging  
**\*Correspondence:** Khaleel Ibraheem Mohson, National Cancer Research Center, University of Baghdad, Iraq. Mobile: +9647725297240. E-mail: [khaleelcabms@gmail.com](mailto:khaleelcabms@gmail.com)  
**Received:** Oct-18-2019  
**Revised:** Jan-11-2020  
**Accepted:** Mar-07-2020  
**Copyright:** © 2020 Khaleel Ibraheem Mohson, Qusay Tayser Naief, Farah Abdul Jalil  
**Funding:** This research did not receive any financial support  
**Competing Interests:** The authors have declared that no competing interests exist  
**Open Access:** This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**BACKGROUND:** Advanced magnetic resonance imaging (MRI) sequences, include the apparent diffusion coefficient (ADC) and diffusion-weighted imaging (DWI), are of great benefit in assessing troublesome vertebral marrow lesion in patients presented with primary malignancy outside the spines.

**OBJECTIVES:** The aim of the study was to evaluate the role of vertebral column MRI in characterization and evaluation of marrow lesion in patients with primary malignancy utilizing ADC and DWI in addition to conventional MRI sequences.

**METHODS:** This is a cross-sectional study which includes 80 patients referred to MR unit 58 of them referred from the oncology department in oncology teaching hospital having primary cancer outside the spine and also referred complaining of back pain and worry from metastasis while the remaining 22 are cancer-free patients referred to MRI because of lower back pain – the study was performed in Medical City, Baghdad, Iraq, during the period from the beginning of September 2018–October 2019.

**RESULTS:** The study sample consists of 55 females and 25 males, the breast cancer is main primary in females and represents 30/58 patients, while the prostate is the main cancer in males and represents 7/58 patients. When correlating the MR sequences (T1W, Fat sat and DWI) all together and after addition of ADC value ( $<9 \times 10^{-3} \text{ mm}^2/\text{sec}$ ) in diagnosis of malignancy there is decreasing frequency of positive cases from 39% before ADC to 22.4% after adding it which is close to histopathology results (17%). The sensitivity, specificity, positive predictive value, negative predictive value, and overall accuracy of ADC in detecting metastasis within the vertebral marrow are 94%, 95%, 88.9%, 97.5%, and 94.8%, respectively.

**CONCLUSION:** Conventional MRI using standard T1W, T2 weighted, and fat suppression sequences cannot discriminate between benign and pathological vertebral marrow lesions. Using DWI improves the recognition of pathological bony lesion(s) and this is strengthened when enforced by ADC value and by that, it can replace the need for intravenous contrast administration. DWI and ADC are beneficial in follow-up of previously detected restricted lesion and in assessment its response to treatment that was depicted by measuring its ADC that consequently elevated in healing phase.

## Introduction

Unluckily metastatic disease to bone which is mainly affect the axial skeleton (vertebral body, iliac bone, the proximal femora, and humeri) is a common squeal in many cancers as early or late presentation of the disease [1]. The presence of marrow metastasis has a great effect on the treatment plan, because its development was considered Stage IV, and here, only palliative treatment will be used in most cases [2]. Once bone metastasis occurs that there is dispersed morbidity from the neurological or stability problems and limitation of patient life from severe pain [3].

For that the characterization of the bony lesion in patients with primary malignancy is very important, this is broadly done using magnetic resonance imaging

(MRI), the conventional MRI can detect the lesion and delineate it clearly from nearby normal marrow, but has limitation in predicting whether this lesion is benign or malignant [4] so the introduction of functional MRI that includes apparent diffusion coefficient (ADC) maps and diffusion-weighted imaging (DWI) can do this job and predict the nature of the lesion, especially when calculating the ADC value of the lesion [5].

Primarily, the DWI was established in the imaging of the central nervous system's abnormalities [6]. With time, this MRI sequence has been used for variable body systems and examinations, mainly in the oncology patients [7].

Regarding our research, we focus the light on the appearance of variable bony lesions affecting the spine, some of them are common and are not neoplastic as typical hemangioma; here, the diagnosis is straight

forward while others are problematic as atypical hemangioma which leads to diagnostic dilemma in differentiating it from bony Mets [8].

The main principle of DWI technique is the regulations of free water transport in the cellular compartment, according to Brownian motion(9), by that DWI able to detect abnormal tissue by assessing the quantity of water within, the more compacted cells, the more limiting to water particles movement when compared with normal nearby tissues, this is opposing to the tissues with low cellular content or necrotic where disturbed cell membranes are predominant, diffusion is not restricted [10].

DWI is a spin-echo T2-weighted single-shot sequence that originates from two symmetric

motion gradient pulses about a 180° refocusing pulse, so the moving water particles do not realign and exhibit a signal drop on DWI, while the water molecules interposed by cellular membranes or intracellular component gain the signal and show high signal on DWI [11].

DWI and ADC values have been offered for the discernment between the malignant deposit, especially in patients with primary condition [12], in general, the bone marrow metastases appear as a high signal area on DWI within low signal normal vertebral marrow, while in ADC, it appears as low signal within relatively high signal normal vertebral marrow [13].

## Materials and Methods

This study included 80 patients referred to magnetic resonance (MR) unit from the oncology department in oncology teaching hospital – Medical City, Baghdad, Iraq, during the period from the beginning of September 3, 2019- July 7,2019.

All patients had a history of primary malignant tumor outside the bone marrow and complaining of back pain or symptoms related to spinal cord compression or having neurological deficit, those patients then subjected MRI examination of the spine, including the lumbar or thoracolumbar areas, using Avanto 1.5T closed MRI system Siemens (Germany).

The MRI protocols include sagittal T1-weighted (T1W) spin echo (SE), sagittal T2-weighted (T2W) SE, sagittal T2W fat suppression SE, and axial T2W SE.

After that, the DWIs were taken in the sagittal plane using a single-shot echo-planar sequence. The b values were 0, 500, and 800 s/mm<sup>2</sup>. DWIs were also obtained in sagittal plane.

After obtaining these images, full analysis of pathological findings was done by reviewing the T1 and T2 images and doing three-dimensional references

to allocated the position of the lesion in ADC and DW sequences, the signal of the lesion is recorded in all sequences, and the brightness of the suspicious lesion is seen on DW and its ADC value was calculated.

Diagnosis of metastasis is established either by biopsy or radioisotope using bone scan or positron emitting tomography, all these with Mets had follow-up by repeating the MRI examination 3–6 months after initiation of chemo- or radiotherapy.

## Statistical analysis

All the relevant data were entered using computerized statistical software; Statistical Package for the Social

Sciences (SPSS) version 20. Descriptive statistics presented as mean ± standard deviation and frequencies as percentages.

## Results

This is a cross-sectional study which includes 80 patients and four of them had follow-up study, 22 of them are cancer-free patients referred to MRI because of lower back pain, and the remaining 58 patients had primary cancer outside the spine and also referred complaining of back pain and worry from metastasis.

The study samples consist of 55 females and 25 males, the breast cancer is the main primary in females and represents 29/58 patients, while the prostate cancer is the main cancer in male and represents 6/58 patients. Details of age, gender, and cancer types are illustrated in Table 1.

The MR examination in the control group is normal in 18/22 patients, the remaining four reveal

**Table 1: Age, gender, and cancer types of the study population**

Age groups	Number (%)
Age	
<40 years	14 (17.5)
41–49	15 (18.75)
50–59	21 (26.25)
60–69	22 (27.5)
>70	8 (10)
Total	80 (100)
Gender	
Males	25 (31.25)
Females	55 (68.75)
Total	80 (100)
Cancer type	
Females	
Breast	29 (72.5)
Ovary	3 (7.5)
Uterus	3 (7.5)
Other	5 (12.5)
Total	40 (100)
Males	
Prostate	6 (46)
Lung	2 (15.5)
Other	5 (38.5)
Total	13(100)

typical hemangioma in two, atypical hemangioma in one, and Modic I end plate changes in the last one.

While the examination of the patients with primary cancer reveals metastasis in 14/58, the remaining findings are discussed briefly in Table 2.

**Table 2: MRI findings in control group and cancer patients**

MRI findings	Number (%)
Control group	
Normal	18 (81.8)
Typical hemangioma	2 (9.1)
Modic I	1 (4.54)
Atypical hemangioma	1 (4.54)
Total	22 (100)
Patients	
Metastasis	14 (24.1)
Free of metastasis	17 (29.3)
Inactive	9 (15.5)
Typical and atypical hemangioma	7 (12.06)
Modic changes	3 (5.17)
Fracture	5 (8.6)
Collection	1 (1.7)
Others	2 (3.4)
Total	58 (100)

MRI: Magnetic resonance imaging.

The correlation between T1W images and Fatsat sequence, on the one hand, with DWI results, on the other hand, in cases of malignant findings revealed that low signal lesions on T1, and high Fatsat and DWI seen in 23/58 (39.6%) of those only 10 (17%) patients are really metastasis, while when adding the low ADC value ( $< 9 \times 10^{-3} \text{ mm}^2/\text{s}$ ) to previous parameters, the metastasis percentage decreases from 39.6% to 22.4% (13/58).

The sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV) and overall accuracy of ADC in detecting metastasis within the vertebral marrow are illustrated in Table 3.

**Table 3: Sensitivity, specificity, positive predictive and negative value, and overall accuracy of ADC in detecting metastasis**

ADC value	Metastasis present	Metastasis absent	Total
Low	16/84*	2/84	18
Normal or high	1/84*	39/84	40
Total	17	41	

\*The total number of patients increased to 84 because four of the patients had two pathology, 1<sup>st</sup> is patient with Ewing sarcoma and had collection (abscess as well), two patients are the elderly with prostate cancer and had active and inert bony lesion, and finally, a patient with breast cancer and also had active and inert lesion.


	Value (%)	95% confidence interval (CI)
Sensitivity	94.12	71.31–99.85
Specificity	95.12	83.47–99.40
PPV	88.89	67.32–96.88
NPV	97.50	85.33–99.62
Accuracy	94.83	85.62–98.92

ADC: Apparent diffusion coefficient, PPV: Positive predictive value, NPV: Negative predictive value.

## Discussion


The introduction of diffusion-weighted MRI revolutionized the diagnosis of benign and malignant vertebral tumors in addition being non-invasive technique, this was achieved by considering the low signal lesion as benign and the high signal one as malignant, on the other hand, the high signal on ADC is more in keeping with a benign cause [14].

Before the DWI was introduced, it was difficult to diagnose the lesion, whether being metastatic or benign tumors using the T1W and T2W images and even with using the fat saturation sequence, the latter only highlighting the hidden lesion that could not clearly visualized in the T1W and T2W sequences. The intravenous contrast injection did not add anything to the diagnosis because many numbers of benign lesions took the contrast and by that, it was not solving this problem.



**Figure 1:** 55 years age female patient with history of breast cancer her MRI reveals (a) ADC image shows low signal in 4th and 5th lumbar vertebral bodies (arrows) ADC ( $6 \times 10^{-3} \text{ mm}^2/\text{sec}$ ) (b) DWI image shows restricted signal within the corresponding vertebral body ( $800\text{s}/\text{mm}^2$ )-histopathology diagnosis

However, after the entry of the ADC and the DW sequences, the problem was nearly solved, so when the lesion appeared of high signal on DWI, we compare that to it to corresponding ADC signal, if it is high on DWI and low on ADC, this raises the suspicious of malignant tumor.



**Figure 2:** 35 years age female patient with history of Ewing sarcoma, her MRI reveals paravertebral abscess (a) ADC sagittal image shows low signal rim collection around 5th lumbar and 1st sacral vertebral bodies (arrows) with ADC ( $3 \times 10^{-3} \text{ mm}^2/\text{sec}$ ) (b) DWI: restricted signal within the rim of collection.

In addition to the role of ADC mainly in the follow-up of cancer cases after treatment with chemotherapy, radiotherapy, or hormonal therapy, the response will be indicated by changing the low ADC value to high while the DWI signal either still high or reverse to low.

In our study, the ADC value in detection bony metastasis shows high sensitivity of 94%, specificity of 95%, and overall accuracy of 94% and this nearly in line with a study done by Tadros and Louka [15].

In 2016, a study showed that a sensitivity of 99%, a specificity of 98%, a PPV of 98%, a NPV of 96%, and an accuracy of 98% for the detection of bone metastatic deposits, and this is highly comparable to the results in our study [16].

Regarding the PPV and NPV of ADC value in detection of vertebral marrow secondary was 89% and 97% and these in concordance with a study done by Gong *et al.* [17].

## Conclusion

1. Conventional MRI using standard T1W, T2W, and fat suppression sequences cannot discriminate between benign and pathological vertebral marrow lesions.
2. Using DWI improves the recognition of pathological bony lesion and this is strengthened when combined with ADC value.
3. Using the above technique is nearly replacing the need intravenous contrast administration, especially in renal disease and when the contrast is not available.
4. DWI and ADC are beneficial in follow-up of previously detected restricted lesion and in assessment lesions response to the treatment.

## References

1. Coleman RE, Lipton A, Roodman GD, Guise TA, Boyce BF, Brufsky AM, *et al.* Metastasis and bone loss: Advancing treatment and prevention. *Cancer Treat Rev.* 2010;36(8):615-20. <https://doi.org/10.1016/j.ctrv.2010.04.003>  
PMid:20478658
2. Sathiakumar N, Delzell E, Morrissey MA, Falkson C, Yong M, Chia V, *et al.* Mortality following bone metastasis and skeletal-related events among women with breast cancer: A population-based analysis of U.S. Medicare beneficiaries, 1999-2006. *Breast Cancer Res Treat.* 2012;131(1):231-8. <https://doi.org/10.1007/s10549-011-1721-x>  
PMid:21842243
3. Tanaka R, Yonemori K, Hirakawa A, Kinoshita F, Takahashi N, Hashimoto J, *et al.* Risk factors for developing skeletal-related events in breast cancer patients with bone metastases undergoing treatment with bone-modifying agents. *Oncologist.* 2016;21(4):508-13. <https://doi.org/10.1634/theoncologist.2015-0377>  
PMid:26975863
4. Vande Berg BC, Lecouvet FE, Michaux L, Ferrant A, Maldague B,

- Malghem J. Magnetic resonance imaging of the bone marrow in hematological malignancies. *Eur Radiol.* 1998;8(8):1335-44. <https://doi.org/10.1007/s003300050548>  
PMid:9853210
5. Lecouvet FE, Larbi A, Pasoglou V, Omoumi P, Tombal B, Michoux N, *et al.* MRI for response assessment in metastatic bone disease. *Eur Radiol.* 2013;23(7):1986-97. <https://doi.org/10.1007/s00330-013-2792-3>  
PMid:23455764
6. Huisman TA. Diffusion-weighted imaging: Basic concepts and application in cerebral stroke and head trauma. *Eur Radiol.* 2003;13(10):2283-97. <https://doi.org/10.1007/s00330-003-1843-6>  
PMid:14534804
7. Subhawong TK, Jacobs MA, Fayad LM. Diffusion-weighted MR imaging for characterizing musculoskeletal lesions. *Radiographics.* 2014;34(5):1163-77. <https://doi.org/10.1148/rg.345140190>  
PMid:25208274
8. Matrawy KA, El-Nekeidy AA, El-Sheridy HG. Atypical hemangioma and malignant lesions of spine: Can diffusion weighted magnetic resonance imaging help to differentiate? *Egypt J Radiol Nucl Med.* 2013;44:259-263. <https://doi.org/10.1016/j.ejnm.2013.03.001>
9. Malayeri AA, El Khouli RH, Zaheer A, Jacobs MA, Corona-Villalobos CP, Kamel IR, *et al.* Principles and applications of diffusion-weighted imaging in cancer detection, staging, and treatment follow-up. *Radiographics.* 2011;31(6):1773-91. <https://doi.org/10.1148/rg.316115515>  
PMid:21997994
10. Koh DM, Collins DJ. Diffusion-weighted MRI in the body: Applications and challenges in oncology. *AJR Am J Roentgenol.* 2007;188:1622-235.
11. Neil JJ. Diffusion imaging concepts for clinicians. *J Magn Reson Imaging.* 2008;27(1):1-7.  
PMid:18050325
12. Wu LM, Gu HY, Zheng J, Xu X, Lin LH, Deng X, *et al.* Diagnostic value of whole-body magnetic resonance imaging for bone metastases: A systematic review and meta-analysis. *J Magn Reson Imaging.* 2011;34(1):128-35. <https://doi.org/10.1002/jmri.23697>  
PMid:21618333
13. Lecouvet FE, El Mouedden J, Collette L, Coche E, Danse E, Jamar F, *et al.* Can whole-body magnetic resonance imaging with diffusion-weighted imaging replace Tc 99m bone scanning and computed tomography for single-step detection of metastases in patients with high-risk prostate cancer? *Eur Urol.* 2012;62(1):68-75. <https://doi.org/10.1016/j.eururo.2012.02.020>  
PMid:22366187
14. Mubarak F, Akhtar W. Acute vertebral compression fracture: Differentiation of malignant and benign causes by diffusion weighted magnetic resonance imaging. *J Pak Med Assoc.* 2011;61(6):555-8.  
PMid:22204209
15. Tadros MY, Louka AL, between malignant marrow diffusion and shift DB. Discrimination between benign and malignant in vertebral marrow lesions with diffusion weighted MRI and chemical shift. *Egypt J Radiol Nucl Med.* 2016;47:557-69. <https://doi.org/10.1016/j.ejnm.2016.02.007>
16. Barchetti F, Stagnitti A, Megna V, Al Ansari N, Marini A, Musio D, *et al.* Unenhanced whole-body MRI versus PET-CT for the detection of prostate cancer metastases after primary treatment. *Eur Rev Med Pharmacol Sci.* 2016;20(18):3770-6.  
PMid:27735042

- 
17. Gong J, Cao W, Zhang Z, Deng Y, Kang L, Zhu P, *et al.* Diagnostic efficacy of whole-body diffusion-weighted imaging in the detection of tumour recurrence and metastasis by comparison with <sup>18</sup>F-2-fluoro-2-deoxy-D-glucose positron emission tomography o computed tomography in patients with gastrointestinal cancer. *Gastroenterol Rep (Oxf)*. 2015;3(2):128-35. <https://doi.org/10.1093/gastro/gou078> PMID:25406465