



# Delayed isolated peri-myocarditis in a Covid-19 patient with respiratory symptoms but without lung involvement

Giancarlo Spano<sup>1</sup> · Kady Fischer<sup>1</sup> · Cédric Maillat<sup>2</sup> · Grégory Vicario<sup>1,2</sup> · Adrian T. Huber<sup>1</sup> · Christoph Gräni<sup>1,3</sup> 

Received: 10 July 2020 / Accepted: 18 July 2020  
© The Author(s) 2020

A 49-year-old prior healthy male without any cardiovascular risk factors living in the western part of Switzerland, noted in mid of March 2020 anosmia and dysgeusia, similar to his wife and four people from his close family, with whom he had frequent contact and who were positive for SARS-CoV-2. Six weeks later, he presented to the hospital with new-onset of dyspnea NYHA 3, general weakness, intermittent epigastrical pain and nocturia without orthopnea nor fever. The retro-nasal SARS-CoV-2 PCR, 6 weeks after initial anosmia and dysgeusia was negative but the antibody IgG blood test for SARS-CoV-2 was positive. Computed tomography of the lungs showed no pulmonary embolism, no infiltrates but left heart congestion, suspected by previous thoracic X-ray (A) and pleural effusion (B). Echocardiography revealed diffuse hypokinesia with severely depressed left- and right-ventricular function. The patient showed elevated C-reactive protein, troponin and NT-proBNP. ECG showed dynamic T-wave changes (C) and after ruling out coronary artery disease,

he was diagnosed with isolated peri-myocarditis using multiparametric cardiac magnetic resonance imaging (CMR). CMR showed diffuse thickening of the myocardium and pericardium due to edema confirmed with T2 weighted imaging and T2 mapping (D). Further, pericardial effusion could be seen and tissue characterization revealed diffuse LGE, elevated T1 mapping values and an elevated extracellular volume fraction of 38% (normal value: < 30%), indicating diffuse fibrosis (E). Global myocardial strain of all heart chambers was diffusely impaired (i.e. peak global longitudinal RV-strain of - 11.6% and LV-strain of - 8.7; normal LV-strain value: < - 15%) (F). No other cause was found as an underlying reason for the peri-myocarditis. In the clinical setting of suspected Covid-19 with respiratory symptoms and negative pulmonary imaging, elevated C-reactive protein and troponin should lead to the suspicion of isolated peri-myocarditis. CMR is the primary noninvasive imaging tool to assess peri-myocarditis, also in Covid-19 patients [1, 2].

✉ Christoph Gräni  
[christoph.graeni@insel.ch](mailto:christoph.graeni@insel.ch)


<sup>1</sup> Bern University Hospital, Bern, Switzerland

<sup>2</sup> Hôpital du Jura Bernois St. Imier, Saint-Imier, Switzerland

<sup>3</sup> University Hospital Bern, University of Bern, Bern, Switzerland

Published online: 28 July 2020

 Springer



**Acknowledgements** Open access funding provided by University of Bern.

### Compliance with ethical standards

**Conflict of interest** There is no conflict of interest for any of the authors concerning this manuscript.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

### References

1. Gräni C, Eichhorn C, Bière L et al (2017) Prognostic value of cardiac magnetic resonance tissue characterization in risk stratifying patients with suspected myocarditis. *J Am Coll Cardiol* 17(16):1964–1976
2. Ferreira VM, Schulz-Menger J, Holmvang G et al (2018) Cardiovascular magnetic resonance in nonischemic myocardial inflammation: expert recommendations. *J Am Coll Cardiol* 72(24):3158–3176

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.