



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

## Journal Pre-proof

Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening

Olga Abian, David Ortega-Alarcon, Ana Jimenez-Alesanco, Laura Ceballos-Laita, Sonia Vega, Hugh T. Reyburn, Bruno Rizzuti, Adrian Velazquez-Campoy



PII: S0141-8130(20)33997-0

DOI: <https://doi.org/10.1016/j.ijbiomac.2020.07.235>

Reference: BIOMAC 16283

To appear in: *International Journal of Biological Macromolecules*

Received date: 17 June 2020

Revised date: 17 July 2020

Accepted date: 22 July 2020

Please cite this article as: O. Abian, D. Ortega-Alarcon, A. Jimenez-Alesanco, et al., Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening, *International Journal of Biological Macromolecules* (2020), <https://doi.org/10.1016/j.ijbiomac.2020.07.235>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening**

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5†</sup>, Ana Jimenez-Alesanco<sup>4,5†</sup>, Laura Ceballos-Laita<sup>2,4†</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

<sup>1</sup> Instituto Aragonés de Ciencias de la Salud (IACS), 50009, Zaragoza, Spain

<sup>2</sup> Instituto de Investigación Sanitaria de Aragón (IIS Aragón), Zaragoza, Spain

<sup>3</sup> Centro de Investigación Biomédica en Red en el Área Temática de Enfermedades Hepáticas Digestivas (CIBERehd), Madrid, Spain

<sup>4</sup> Institute for Biocomputation and Physics of Complex Systems (BIFI), Joint Units IQFR-CSIC-BIFI, and GBsC-CSIC-BIFI, Universidad de Zaragoza, 50009, Zaragoza, Spain

<sup>5</sup> Departamento de Bioquímica y Biología Molecular y Celular, Universidad de Zaragoza, 50009, Zaragoza, Spain

<sup>6</sup> Department of Immunology and Oncology, National Centre for Biotechnology (CNB), CSIC, 28049, Madrid, Spain

<sup>7</sup> CNR-NANOTEC, Licryl-UOS Cosenza and CEMIF.Cal, Department of Physics, University of Calabria, 87036, Rende, Italy

<sup>8</sup> Fundación ARAID, Gobierno de Aragón, 50018, Zaragoza, Spain

† Equal contribution

\* Corresponding authors:

Adrián Velázquez-Campoy, Institute BIFI - University of Zaragoza, Mariano Esquillor s/n, 50018, Zaragoza, Spain

Olga Abian, Instituto Aragonés de Ciencias de la Salud (IACS), San Juan Bosco 13, 50009, Zaragoza, Spain

**ABSTRACT**

The global health emergency generated by coronavirus disease 2019 (COVID-19) has prompted the search for preventive and therapeutic treatments for its pathogen, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). There are many potential targets for drug discovery and development to tackle this disease. One of these targets is the main protease, Mpro or 3CLpro, which is highly conserved among coronaviruses. 3CLpro is an essential player in the viral replication cycle, processing the large viral polyproteins and rendering the individual proteins functional. We report a biophysical characterization of the structural stability and the catalytic activity of 3CLpro from SARS-CoV-2, from which a suitable experimental *in vitro* molecular screening procedure has been designed. By screening of a small chemical library consisting of about 150 compounds, the natural product quercetin was identified as reasonably potent inhibitor of SARS-CoV-2 3CLpro ( $K_i \sim 7 \mu\text{M}$ ). Quercetin could be shown to interact with 3CLpro using biophysical techniques and bind to the active site in molecular simulations. Quercetin, with well-known pharmacokinetic and ADMET properties, can be considered as a good candidate for further optimization and development, or repositioned for COVID-19 therapeutic treatment.

**KEYWORDS**

SARS-CoV-2 3CL protease, structural stability, drug discovery

## INTRODUCTION

December 2019 was the starting point for a global pandemic, caused by a newly identified coronavirus, SARS-CoV-2, rising to dramatic numbers of infections and deaths worldwide[1, 2]. Much effort and many resources have been devoted to combat this health emergency on many fronts: 1) better diagnosis and patient management[3]; 2) improving protective and palliative care[4]; 3) reducing damage in the different organs affected by the virus and minimizing collateral harmful effects of the immune response[5]; and 4) developing preventive vaccines and therapeutic treatments[6], to mention the more relevant. In addition, very strict and severe social and economic measures, with the potential danger of causing a profound global crisis, have been taken[7]. In this context, even if an effective vaccine is promisingly close to becoming available, drugs are needed for infected people[8]. From experience with other pathogens, we have learnt that the best therapeutic strategy consists in administering a combination of several drugs acting through different mechanisms, in order to minimize the probability of drug resistance appearance[9]. At the same time, cost-efficient drugs (if possible, developed by repositioning or repurposing known drugs) are urgently needed to reaching all the population[10].

The SARS-CoV-2 genome is 82% identical to that of SARS-CoV, the causative virus of the 2002 coronavirus outbreak[11]. The genomic differences are reflected in a different infectivity and mortality rates for SARS-CoV-2[12]. Among all the potential protein targets within coronaviruses, the main protease (Mpro, or 3C-like protease, 3CLpro) stands out as associated to a highly conserved gene (96% sequence identity between the SARS-CoV-2 and SARS-CoV), making 3CLpro a good target for developing effective drugs against SARS-CoV-2 (and other future coronavirus variants)[13, 14]. Together with PLpro, 3CLpro is responsible for the processing of the viral polyproteins synthesized from the viral RNA after infection, rendering the individual viral proteins active and functional[15]. Remarkably, 3CLpro must process at least 11 cleavage sites on polyprotein 1ab (replicase 1ab), most of them sharing a common cleavage sequence LQ↓(S/A/G), quite unusual for human proteases[16]. Therefore, molecules able to inhibit SARS-CoV-2 3CLpro in a specific manner would hinder viral replication and represent appropriate candidates to develop low-toxicity drugs against this devastating pathogen[17].

Searching for drugs against a protein target always benefits from a comprehensive structural and functional characterization[18]. Experimental and computational knowledge gathered along this process is instrumental for identifying weak points and key interaction sites in the target, and developing suitable screening procedures for finding molecules interfering with or modulating its function[19]. Here we have characterized some structural and functional

features of SARS-CoV-2 3CLpro by employing biophysical techniques (spectroscopy and calorimetry), and we have implemented a fast *in vitro* screening procedure based on 3CLpro hydrolytic activity using a Förster resonance energy transfer (FRET) substrate, which allows the ready identification of small molecules blocking the enzymatic activity of 3CLpro. Starting from a small chemical library consisting of about 150 compounds, collected from successful screening programs previously carried out in our laboratory, a natural compound was identified as an inhibitor of SARS-CoV-2 3CLpro with enough potency (inhibition constant  $K_i \sim 7 \mu\text{M}$ ) to be considered a good candidate for further optimization and development. Quercetin, with known pharmacokinetic and ADMET properties, exhibits anti-oxidant, anti-allergic, anti-inflammatory, and anti-proliferative indications, and can be directed for drug repositioning.

## MATERIALS & METHODS

### Protein expression and purification

A pET22b plasmid containing the SARS-CoV-2 His-tagged 3CLPRO sequence was transformed into BL21 (DE3) Gold *E. coli* strain. Bacterial cultures were grown in 250 mL of LB/ampicillin (100  $\mu\text{g}/\text{mL}$ ) media at 37 °C overnight with gentle shaking. Then, 4 L of LB/ampicillin (100  $\mu\text{g}/\text{mL}$ ) were inoculated and incubated under the same conditions until reaching OD = 0.6 at a wavelength of 600nm. Protein expression was induced with 1 mM isopropyl 1-thio- $\beta$ -D-galactopyranoside (IPTG) at 18 °C for 5 h. Cells were harvested by centrifugation at 4 °C for 10 min at 10000 rpm (using a Beckman Coulter Avanti J-26 XP Centrifuge) and then resuspended in lysis buffer (sodium phosphate 50 mM, pH 8, sodium chloride 500 mM). Cell rupture was achieved by sonication (using a Sonics Vibra-Cell Ultrasonic Liquid Processor) in ice, and 20 U/mL of benzonase (Merck-Millipore) were added to remove nucleic acids. To remove cellular debris, the extract was centrifuged at 4 °C for 30 min at 20000 rpm, and filtered through a 0.45  $\mu\text{m}$ -pore membrane. After increasing imidazole concentration up to 10 mM, the protein was purified using immobilized metal ion affinity chromatography (IMAC) in a ÄKTA FPLC System (GE Healthcare Life Sciences) using a cobalt HiTrap TALON column (GE-Healthcare Life Sciences), eluting in an imidazole 10-250 mM gradient. Purity was checked by SDS-PAGE, and pure protein fractions were dialyzed to remove imidazole and reach the protein storage condition (sodium phosphate 50 mM, pH 8, sodium chloride 150 mM). The identity of the protein was assessed by mass spectrometry (LC-ESI-MS/MS). Potential DNA contamination was assessed by determining the UV absorption 260/280 ratio. An extinction coefficient of 32890  $\text{M}^{-1} \text{cm}^{-1}$  at 280 nm was employed for quantification. When needed, buffer exchange was performed using a PD-10 pre-packed Sephadex G-25 M (GE Healthcare Life Sciences).

**Circular dichroism and fluorescence spectroscopy**

Circular dichroism (CD) spectra were recorded in a Chirascan spectropolarimeter (Applied Photophysics) at 25 °C. Far-UV spectrum was recorded at wavelengths between 190 and 260 nm in a 0.1 cm path-length cuvette. Near-UV spectrum was recorded at wavelengths between 250 and 310 nm in a 1-cm path-length cuvette. Protein concentration was 10  $\mu$ M in all cases. Fluorescence measurements were performed in a Cary Eclipse fluorescence spectrophotometer (Agilent Technologies), monitoring the intrinsic tryptophan fluorescence of a protein solution at 2  $\mu$ M concentration. An excitation wavelength of 290 nm was used, with excitation and emission bandwidths of 5 nm, and recording fluorescence emission between 300 and 400 nm. All spectroscopic measurements were made in sodium phosphate 50 mM, pH 8.

Thermal denaturations were monitored by CD and fluorescence, employing a protein concentration of 10  $\mu$ M and 2  $\mu$ M, respectively, and performing thermal scans with a scanning rate of 60 °C/h.

**Dynamic light scattering**

Hydrodynamic radius of SARS-CoV-2 3CLpro was measured at different pH values for estimating the size of the oligomeric native state in a DynaPro NanoStar equipment (Wyatt Technology), employing a protein concentration of 3  $\mu$ M at pH 5, 7 and 8.

**Size-exclusion chromatography**

Self-association state of SARS-CoV-2 3CLpro was assessed employing an ÄKTA FPLC (GE Healthcare) using a Superdex Increase 75 10/30 chromatographic column, at pH 8 and 150 mM NaCl.

**Differential scanning calorimetry**

Thermal stability of SARS-CoV-2 3CLpro was assessed by temperature unfolding transitions monitored by high-precision differential scanning calorimetry (DSC). The partial molar heat capacity of the protein in solution was measured as a function of temperature in an Auto-PEAQ-DSC (MicroCal, Malvern-Panalytical). Experiments were performed with a protein solution at a concentration of 14  $\mu$ M in buffer, and scanning from 15 to 95 °C at a rate of 60 °C/h. Different buffers were employed to get protein stability information under different conditions: acetate 50 mM at pH 5, and phosphate at pH 7 and pH 8.

## Protein unfolding

### *Model-free calorimetric analysis*

The first approach in DSC experimental analysis consists of applying a model-free data analysis for discriminating between different possibilities: two-state unfolding, non-two-state unfolding, and oligomer unfolding of the protein. From the thermogram (excess molar heat capacity of the protein as a function of the temperature,  $\langle \Delta C_p(T) \rangle$ ), the calorimetric unfolding enthalpy,  $\Delta H_{cal}$ , the unfolding temperature,  $T_m$ , and the maximal unfolding heat capacity,  $C_{p,max}$ , can be readily evaluated, from which it is possible to calculate the van't Hoff enthalpy,  $\Delta H_{vH}$ :

$$\Delta H_{vH} = \frac{4RT_m^2 C_{p,max}}{\Delta H_{cal}} \quad (1)$$

From the ratio  $\Delta H_{vH}/\Delta H_{cal}$  different possibilities may arise: 1) if  $\Delta H_{vH}/\Delta H_{cal} = 1$ , the thermogram would reflect a single transition and the protein unfolds according to a two-state model (i.e., the protein contains a single energetic domain); 2) if  $\Delta H_{vH}/\Delta H_{cal} < 1$ , the thermogram would reflect at least two (partially) overlapping transitions and the protein unfolds according to a non-two-state model (i.e., the protein contains at least two domains which unfold in an independent manner); and 3) if  $\Delta H_{vH}/\Delta H_{cal} > 1$ , the thermogram would reflect an unfolding transition coupled to subunit dissociation (i.e., the protein is oligomeric and unfolds into monomers).

In the case of SARS-CoV-2 3CLpro, the thermograms under the different conditions assayed showed that  $\Delta H_{vH}/\Delta H_{cal} > 1$  and, therefore, the protein is not monomeric and dissociates upon thermal unfolding. In fact, according to the crystallographic structure of the unliganded species, the protein is dimeric in its native state, although it has been reported that SARS-CoV-1 3CLpro can populate tetramers and octamers.

### *Cooperative homo-oligomeric protein unfolding into isolated unfolded monomers*

The conformational equilibrium is governed by the equilibrium constant,  $K_U$ :



where  $N_n$  is the oligomeric native state constituted of  $n$  monomeric subunits, and  $U$  is the unfolded state of each identical monomeric subunit. The total protein concentration can be expressed (per monomer) as:

$$[P]_T = n[N_n] + [U] = \frac{n}{K_U} [U]^n + [U] \quad (3)$$

from which the molar fractions of the different species can be calculated:

$$F_N = \frac{n[U]^n}{K_U [P]_T} = \frac{n[P]_T^{n-1}}{K_U} F_U^n \quad (4)$$

$$F_U = \frac{[U]}{[P]_T}$$



Given the equilibrium constant and the total protein concentration, Eq. 3 can be solved numerically for the unknown [U], and the molar fraction of both protein species can be calculated at any temperature, from which the excess average molar unfolding enthalpy,  $\langle\Delta H\rangle(T)$  can be determined:

$$\langle\Delta H\rangle(T) = F_U(T)\Delta H_U(T) \quad (5)$$

where  $\Delta H_U$  is the enthalpy of the unfolded state, considering that of the native state as a reference. The temperature derivative of the excess molar unfolding enthalpy is the excess unfolding heat capacity at constant pressure,  $\langle\Delta C_p\rangle(T)$ :

$$\langle\Delta C_p\rangle(T) = \left(\frac{\partial\langle\Delta H\rangle(T)}{\partial T}\right)_p \simeq \frac{\langle\Delta H\rangle(T+\Delta T) - \langle\Delta H\rangle(T)}{\Delta T} \quad (6)$$

taking into account the temperature dependence of the different magnitudes:

$$\begin{aligned} \Delta H_U(T) &= \Delta H_U(T_0) + \Delta C_{p,U}(T - T_0) \\ \Delta S_U(T) &= \Delta S_U(T_0) + \Delta C_{p,U} \ln \frac{T}{T_0} = \frac{\Delta H_U(T_0)}{T_0} + \Delta C_{p,U} \ln \frac{T}{T_0} \\ \Delta G_U(T) &= \Delta H_U(T) - T\Delta S_U(T) \\ K_U &= e^{-n\Delta G_U(T)/RT} \end{aligned} \quad (7)$$

where  $\Delta S_U$ ,  $\Delta G_U$  and  $\Delta C_{p,U}$  are the entropy, the Gibbs energy and the heat capacity of the unfolded state (taking the native state as a reference), respectively, and  $T_0$  is an appropriate reference temperature. In this case,  $T_0$  is the temperature at which the unfolding Gibbs energy is zero, and does not coincide with the temperature for maximal heat capacity,  $T_{\max}$ , nor with the temperature for half denaturation,  $T_{1/2}$  (i.e., the temperature at which  $F_U = 0.5$ , or median temperature in the thermogram). This model particularized to  $n = 1$  corresponds to the two-state model, for which  $T_0$  is the apparent unfolding temperature,  $T_m$ , which is very close to both  $T_{\max}$  and  $T_{1/2}$ .

For a spectroscopically monitored unfolding of a homo-oligomeric protein, the procedure is similar, but the final observable quantity is the spectroscopic signal,  $S(T)$ , which can be calculated as:

$$S(T) = F_N(T)S_N(T) + F_U(T)S_U(T) \quad (8)$$

where  $S_N(T)$  and  $S_U(T)$  are the intrinsic spectroscopic signal values for the native and unfolded protein states, and can be considered linear functions of the temperature:

$$\begin{aligned} S_N(T) &= A_N + B_N T \\ S_U(T) &= A_U + B_U T \end{aligned} \quad (9)$$

### Proteolytic activity assay

*In vitro* catalytic activity of 3CLpro was determined using a fluorescence resonance energy transfer (FRET) cleavage continuous assay with the peptide substrate (Dabcyl)KTSAVLQSGFRKME(Edans)-NH<sub>2</sub> (Biosyntan GmbH)[20-22]. Briefly, enzyme at 0.2-2  $\mu\text{M}$

final concentration in sodium phosphate 50 mM, NaCl 150 mM, at pH 8, and the enzymatic reaction was initiated by adding the substrate at 20  $\mu$ M final concentration, in a final volume of 100  $\mu$ L. Fluorescence was continuously measured in a FluoDia T70 microplate reader (Photon Technology International) for 20 min (excitation wavelength, 380 nm; emission wavelength, 500 nm). Enzyme activity was quantified as the initial slope of the time evolution curve of the fluorescence signal. The slope ratio between the activity in the presence and absence of a given compound provides the percentage of inhibition. For comparison, the hydrolytic activity was also measured at different pH values. The Michaelis-Menten constant,  $K_m$ , and the catalytic rate constant or turnover number,  $k_{cat}$ , were estimated by measuring the initial enzymatic rate at different substrate concentrations at fixed 0.2 enzyme concentration and applying non-linear regression data analysis.

#### Activity-based screening

Screening of a small chemical library was performed based on the catalytic activity continuous FRET assay. In a 96-well microplate, enzyme at 1  $\mu$ M final concentration was incubated in the presence of chemical compounds (at 125  $\mu$ M final concentration) solubilized previously at 5 mM concentration in DMSO. The final concentration of DMSO was 2.5% v/v. The same amount of DMSO was added to control wells (no compounds added). Several hits were identified as those compounds reducing the enzymatic activity below a threshold (average enzymatic activity of controls minus twice their standard deviation). In order to determine the inhibition constants of the selected compounds, 2-fold serial dilutions of each compound were tested (ranging from 0 to 125  $\mu$ M) under the same conditions described above.

#### Inhibition assay

To assess the *in vitro* inhibition potency of the selected compounds, the inhibition constants were estimated from the experimental curves. Inhibition curves were obtained by measuring the enzyme activity as a function of compound concentration: monitoring the substrate fluorescence emission as a function of time, fixing the enzyme concentration at 2  $\mu$ M, the substrate concentration at 20  $\mu$ M, and varying the compound concentration from 0 to 125  $\mu$ M, while maintaining constant the percentage of DMSO. The enzymatic activity was quantitated as the initial slope of the substrate fluorescence emission time curve, and was plotted as a function of compound concentration. Non-linear regression analysis employing a simple inhibition model allowed us to estimate the apparent inhibition constant for both compounds, according to this equation:

$$\frac{v([I])}{v([I]=0)} = 1 - \frac{\frac{[I]}{K_i^{app}}}{1 + \frac{[I]}{K_i^{app}}} = \frac{1}{1 + \frac{[I]}{K_i^{app}}} \quad (10)$$

where  $v$  is the initial slope of the enzymatic activity trace at a (free) compound concentration  $[I]$ ,  $K_m$  is the Michaelis-Menten constant for the enzyme-substrate interaction,  $[S]$  is the substrate concentration, and  $K_i^{app}$  is the apparent inhibition constant for the compound. If the inhibitor acts through a purely competitive mechanism, the previous equation can be substituted by the following one:

$$\frac{v([I])}{v([I]=0)} = \frac{K_m + [S]}{K_m \left(1 + \frac{[I]}{K_i}\right) + [S]} = \frac{1}{1 + \frac{[I]}{K_i \left(1 + \frac{[S]}{K_m}\right)}} \quad (11)$$

where  $K_i$  is the intrinsic (i.e., substrate concentration-independent) inhibition constant.

### Thermal shift assay

For the thermal shift assay (TSA), SYPRO Orange (Thermo Fisher Scientific) was employed as an extrinsic fluorescent probe for reporting protein unfolding. A reaction mixture containing 3CLpro at 1  $\mu$ M of final concentration in sodium phosphate 50 mM, pH 8, and SYPRO Orange at a final concentration of 5X, was freshly prepared and dispensed into 96-well microplates (96-well PCR plate, non-skirted, from 4titude) at a final volume of 100  $\mu$ L. Compound from a 5 mM stock solution in pure DMSO were added to the mixture of protein and fluorescence probe to a final concentration of 250  $\mu$ M each. Controls were considered by adding the same volume of 100% DMSO instead of any compound. Unfolding curves were registered from 25  $^{\circ}$ C to 99  $^{\circ}$ C at a 1  $^{\circ}$ C/min scan rate in a Stratagene Mx3005P qPCR real-time thermal cycler (Agilent Technologies). Thermal denaturation was followed by fluorescence emission of the fluorescent probe, using 492 and 610 nm as excitation and emission wavelength filters, respectively. The apparent midpoint unfolding temperature,  $T_m$ , was calculated in each well as the temperature of maximum slope (i.e., the inflection point), and compared to the controls. To assess the concentration dependence of the stability change induced by the selected compound, 2-fold serial dilutions (ranging from 0 to 125  $\mu$ M) for each compound were assayed by following the same protocol described above.

### Isothermal titration calorimetry

The interaction between the compound and 3CLpro was further assessed by isothermal titration calorimetry. Calorimetric titrations were performed using a high precision Auto-iTC200 calorimeter (MicroCal, Malvern-Panalytical). Protein at 15  $\mu$ M and compound at 150  $\mu$ M were dissolved in sodium phosphate 50 mM, pH 8, with and without NaCl 150 mM.

Compound was titrated into protein solution by performing 19 injections of 2  $\mu\text{L}$  each, and mixed using a stirring speed of 750 rpm, maintaining a spacing between injections of 150 s, and applying a reference power of 10  $\mu\text{cal/s}$ . The association constant,  $K_a$ , the binding enthalpy,  $\Delta H$ , and the binding stoichiometry,  $N$ , were estimated through non-linear least squares regression analysis of the data, by using a model considering a single ligand binding site, implemented in Origin 7.0 (OriginLab). The dissociation constant,  $K_d$ , the binding Gibbs energy,  $\Delta G$ , and the binding entropy,  $-\Delta S$ , were obtained from basic thermodynamic relationships.

### **Molecular docking**

Molecular docking was performed by using the simulation software AutoDock Vina[23] and the supporting suite AutoDock Tools[24]. The structure of 3CLpro was extracted from the entries 6Y2E and 6Y2F[22] of the Protein Data Bank (PDB), which contain the crystallographic conformation of the protein in unliganded form and with an  $\alpha$ -ketoamide inhibitor bound in the active site, respectively. The protein structures were considered in all cases free from any ligand and water molecules. A few missing residues in the entry 6Y2F were reconstructed *in silico*, and the structure of quercetin was also built by using the molecular editor Avogadro[25]. A blind search was carried out on the whole protein surface, by considering a volume of size 50  $\text{\AA} \times 60 \text{\AA} \times 60 \text{\AA}$ . Extensive sampling was performed in all cases, using an exhaustiveness 16 times larger than the value normally recommended[26].

## **RESULTS**

### **SARS-CoV-2 3CLpro is properly folded**

Recombinant 3CLpro was expressed in *E. coli* at sufficiently high yield for initiating a biophysical characterization and an activity-based molecular screening (Figure S1). The far-UV circular dichroism (CD) spectrum showed negative bands around 208 and 222 nm, typical of proteins with  $\alpha$ -helical and  $\beta$ -sheet content (Figure 1A), in agreement with the crystallographic structure[22]. The near-UV CD spectrum indicated aromatic residues, especially tryptophans, are enclosed within an asymmetric environment (Figure 1B). The fluorescence spectrum monitoring the intrinsic emission of the tryptophans (there are three tryptophan residues, two located in the  $\alpha$ -helical domain and one in the  $\beta$ -sheet domain) showed a maximum around 330, confirming that some of them are partially exposed to the solvent (Figure 1C). These results confirmed 3CLpro adopted a folded conformation with well-defined secondary and tertiary structures.

**SARS-CoV-2 3CLpro self-associates**

3CLpro from other coronaviruses (SARS-CoV and MERS) dimerizes, and such dimerization is known to be essential for the enzymatic activity; even higher-level oligomerization structures (tetramers and octamers) have been reported[27]. Among other experimental techniques providing information about protein oligomerization (such as analytical ultracentrifugation and gel filtration chromatography), thermal denaturation monitored by spectroscopy or calorimetry can be used to detect the self-association of proteins, since the thermal unfolding of monomeric and oligomeric proteins show markedly different features[28].

Thermal unfolding of 3CLpro followed by far-UV CD showed a single transition with an apparent unfolding temperature close to 51 °C (Figure 2A). The same process followed by fluorescence also showed a single transition with an apparent unfolding temperature close to 48.5 °C (Figure 2B). The small difference in the unfolding temperature could be justified by the use of a different protein concentration in CD and fluorescence assays and considering the oligomer-monomer equilibrium coupled to the unfolding process. Non-linear regression analysis of the unfolding traces according to different two-state models for a monomeric ( $n = 1$ , see equation (1)), a dimeric ( $n = 2$ ), a tetrameric protein ( $n = 4$ ), and an octameric protein ( $n = 8$ ) were performed (see Materials & Methods section). These experiments confirmed 3CLpro adopts a well-folded conformation under native conditions and becomes unstructured upon temperature stress. However, it was not possible to discriminate which model reproduces better the experimental data, because the fitting curves from the different models overlapped. The fitting was marginally better (according to the residual sum of squares; see Table S1) for the two-state monomer model in the case of CD unfolding, and for the two-state tetramer model in the fluorescence unfolding. Therefore, spectroscopic unfolding traces were not sensitive enough for discriminating the self-association nature of 3CLpro native state.

Thermal unfolding monitored by differential scanning calorimetry (DSC) also showed a single apparent unfolding transition with a marked dependence on the pH (Figure 3A). The lower thermal stability observed at pH 5 and 8, compared to that at pH 7, may reflect the protonation/deprotonation equilibrium of ionizable sidechains titratable around pH 6-7 (e.g., histidines). Very importantly, the transitions are quite asymmetric, a clear indication of the oligomeric nature of the native state of 3CLpro. A model-free analysis (see Materials & Methods section) allowed to estimate an apparent unfolding enthalpy of 106 kcal/mol and an unfolding temperature of 54.2 °C, at pH 8. Again, the difference in the unfolding temperature compared to those obtained by using spectroscopic techniques could be justified by the use of a different protein concentration in the DSC assays and considering the oligomer-monomer equilibrium coupled to the unfolding process. From this model-free analysis, the ratio between

the van't Hoff enthalpy and the calorimetric enthalpy was estimated to be  $> 1$ , which is another clear indication of the oligomeric nature of the native state of the protein. A small influence of the ionic strength of the solution on the thermal stability of the protein was observed (Figure S2A). The increase in NaCl concentration to 150 mM caused a reduction in the apparent unfolding temperature of 0.6 °C, which suggested the uptake of salt ions upon protein unfolding.

The scan rate along the DSC assay may distort the stability parameters (i.e., the DSC profiles may show a scan rate dependence) if the unfolding kinetic equilibration for the transitions between the conformational states is sufficiently slow. In this case, irreversible kinetic models should be applied instead of equilibrium models. In our experiments on SARS-CoV-2 3CLpro, a small scan rate dependence could be observed (Figure S2B). The data analysis according to the Lumry-Eyring model for protein thermal denaturation (Figure S2B, inset) allowed estimating a high kinetic activation energy barrier for the irreversible denaturation process, close to 200 kcal/mol[29]. This energy barrier corresponds to a half-life for irreversible denaturation extremely long (close to 30 years at 37 °C, and much larger at 25 °C). These results suggested the analysis employing equilibrium models may be appropriate.

The two-state tetramer ( $n = 4$ ) unfolding model reproduced significantly better the experimental data at pH 7 and 8 (Figure 3B), while the two-state dimer ( $n = 2$ ) unfolding model reproduced better the experimental data at pH 5 (according to the residual sum of squares; see Table S1). Application of a more complex model considering the coexistence of tetramers and dimers ( $T \rightarrow 2D \rightarrow 4U$ ) did not improve the results (Table S1). In addition, other models considering a two-step unfolding process where the dimeric or tetrameric states undergo a first transition with no dissociation ( $N_n \rightarrow N_n^*$ ) followed by a second transition with coupled unfolding-dissociation ( $N_n^* \rightarrow nU$ ) did not improve the results either. This indicates that any unfolding model considering more than one step lacks the sufficient cooperativity to reproduce the experimental results. Therefore, calorimetric unfolding traces were sensitive enough for discriminating the self-association nature of 3CLpro native state, and very likely there is a pH dependence of the quaternary rearrangement in 3CLpro.

#### **SARS-CoV-2 3CLpro activity profile correlates with the thermal stability profile**

The enzymatic activity of 3CLpro was measured using a FRET substrate containing Edans/Dabcyl as the donor/acceptor FRET pair. Continuous assays allowed us monitoring the time evolution of the substrate fluorescence emission as a function of time at different pH values. The enzymatic activity was higher at pH 7, compared to pH 5 and 8 (Figure 4), which

correlates with the pH dependence of the structural stability observed by DSC assays. Among other factors, the pH dependence of the enzyme activity may reflect structural changes (either secondary, tertiary or quaternary, or a combination of them) and, very importantly, the ionization state of titratable sidechains from the catalytic dyad residues (Cys-145 and His-41). Very likely, the catalytic histidine is responsible for this pH effect.

The enzymatic parameters were estimated from the initial enzymatic rate dependence on the substrate concentration at pH 8 (Figure S3):  $K_m = 11 \mu\text{M}$  and  $k_{\text{cat}} = 0.040 \text{ s}^{-1}$ , resulting in a catalytic efficiency,  $k_{\text{cat}}/K_m$  of  $3640 \text{ M}^{-1} \text{ s}^{-1}$ , comparable to reported values considering the differences in the experimental conditions[21, 22].

### Screening of a small chemical library

The continuous FRET assay for measuring the activity of 3CLpro was adapted to an *in vitro* experimental screening procedure. In a first step, the screening was performed using a small chemical library consisting of 150 compounds, which also included approved drugs with known therapeutic indications. Hits consisted of compounds reducing the enzymatic activity of 3CLpro below a specified threshold, defined as the average activity for the controls (enzyme with no compound) minus twice their corresponding standard deviation (Figure 5). Several hits were selected and confirmed by the inhibition curve analysis.

### Quercetin inhibits SARS-CoV-2 3CLpro *in vitro*

To assess the *in vitro* potency of the compounds selected through the small-scale screening, their inhibition constants was estimated from the inhibition curve. Inhibition curves were obtained by measuring the enzymatic activity, applying the previously described protocol for the continuous FRET assay, by fixing the enzyme and the substrate concentrations while varying the amount of compound (Figure 6A). Enzymatic activity at each compound concentration was calculated as the initial slope in each curve, and a dose-dependent effect of the compound on the enzyme activity was observed (Figure 6B). Non-linear regression analysis employing a simple inhibition model, according to equation (10) (see Materials & Methods), allowed estimating an apparent inhibition constant,  $K_i^{\text{app}}$ , of  $21 \mu\text{M}$  for quercetin, the most active compound. Because it can be assumed this compound might bind to the active site, thus, functioning as a competitive inhibitor, this apparent inhibition constant can be transformed into an intrinsic inhibition constant according to:  $K_i = K_i^{\text{app}}/(1+[S]/K_m)$ , or, equivalently, the fitting can be carried out using equation (11), which accounts for competitive inhibition and the substrate concentration and  $K_m$  appear explicitly (see Materials & Methods). With a  $K_m$  value of  $11 \mu\text{M}$  recently reported in the literature[21, 22], an intrinsic inhibition

constant  $K_i$  of 7.4  $\mu\text{M}$  could be estimated for quercetin. This potency is similar to some of the first inhibitors found for SARS-CoV 3CLpro[30].

#### **Quercetin affects SARS-CoV-2 3CLpro thermal stability**

In general, the interaction of a ligand with a protein results in a change in the stability of the protein against thermal or chemical denaturation. More precisely, the preferential interaction of the ligand with certain conformational state stabilizes that state, resulting in a net global stabilization effect if the preferential interaction occurs with the native state or a net destabilization if the preferential interaction occurs with the non-native state. Thermal shift assay (TSA), also called differential scanning fluorimetry, consists in detecting ligand-induced stability changes in proteins by fluorescence[31, 32]. Using an extrinsic fluorescent probe reporting the progression of the protein unfolding process, it is possible to assess the thermal stability of the protein by measuring its apparent  $T_m$  in the presence of increasing concentrations of a given interacting compound.

The identified compound, quercetin, altered the thermal stability of 3CLpro causing a destabilization (Figure 7A). Although ligand-induced stabilizing effects on a protein are more common, it is not unusual to observe destabilizing effects[33-35]. In addition, the effect of the compounds was concentration-dependent (Figure 7B).

#### **Quercetin interacts with SARS-CoV-2 3CLpro**

Isothermal titration calorimetry (ITC) is considered the gold-standard for binding affinity determination, as well as target engagement in drug discovery[36, 37]. Careful experimental design in ITC provides invaluable and detailed information about the drug-target interaction[38]. ITC was employed to assess target engagement for quercetin and get a direct estimation of the dissociation constant for its interaction with 3CLpro. According to the experiments, quercetin interacts with 3CLpro with a dissociation constant of 2.7  $\mu\text{M}$  in the absence of NaCl, and a dissociation constant of 10  $\mu\text{M}$  in the presence of NaCl 150 mM (Figure 8), in good agreement with the inhibition constant estimated through the inhibition curve (which was determined in the presence of NaCl 150 mM).

#### **Quercetin binds to SARS-CoV-2 3CLpro active site**

Molecular docking was used to clarify the interaction of quercetin with 3CLpro. Two crystallographic structures were considered, corresponding to the protein in unliganded form or with an inhibitor bound in the active site, reported as entry 6Y2E and 6Y2F in the PDB[22], respectively. Figure 9 summarizes the overall findings obtained in the simulation with



quercetin. For both molecular receptors, the most favorable binding modes predicted through a blind search carried out on the whole protein surface were found in the known protein active site. The binding affinity ranged from -7.5 to -7.2 kcal/mol, corresponding to a dissociation constant of  $\sim 3\text{-}5\ \mu\text{M}$ , in reasonably good agreement with the value obtained experimentally ( $K_d$  of  $2.7\ \mu\text{M}$ ). A variety of slightly different binding modes were obtained, all distributed in the same binding location. This observation is consistent with the pseudo-symmetric chemical structure of quercetin, with five hydroxyl groups determining a number of possibilities in the formation of hydrogen bonds as donors.

As detailed in the Figure 9 (see Insets b and c), the two binding modes of quercetin most distant from each other still occupied the same protein crevice. In particular, they mostly overlapped with the position of a single larger known ligand, the  $\alpha$ -ketoamide inhibitor co-crystallized with 3CLpro in the PDB entry 6Y2F[22]. Interactions with a number of protein residues contribute to bind quercetin into the 3CLpro binding site. The key residue is Met165, which can form both hydrogen bonds and hydrophobic interactions with quercetin. Additional hydrogen bonds can be formed with the two residues Ser144 and Met165. Energetic contributions are also due to supplementary hydrophobic interactions with Met49, Phe140 and Leu141, and to electrostatic interaction formed with the polar residue His164 and the charged residue Glu166.

## DISCUSSION

The need for pharmacological agents acting against SARS-CoV-2 is urgent. The development of drugs acting through different mechanisms provides a good starting point for a combined use in a therapy with low susceptibility to drug resistance. 3CLpro (or Mpro) is an appropriate target, given its high conservation among coronaviruses, as well as a suitable experimental tractability and druggability. Drug discovery can be initiated from experimental or computational screenings, by employing small or large, highly diverse or focused (target-derived) chemical libraries. On the basis of an experimental pipeline for drug screening successfully employed in the past[35, 39-45], we have applied the same methodology to identify small molecules blocking the activity of SARS-CoV-2 3CLpro.

The first step in this screening program was to assess the stability and the *in vitro* activity of SARS-CoV-2 3CLpro. According to our results, 3CLpro is well folded in solution and exhibits a large kinetic stability (activation energy barrier  $\sim 200\ \text{kcal/mol}$ ), which allowed us to apply equilibrium models for unfolding. It has been established that monomeric 3CLpro is inactive[46] and the functional state of this protein is the homodimer. We observed that 3CLpro not only populates dimeric states in solution, but also tetrameric states, depending on

the pH. Interestingly, even tetrameric and highly-active octameric states have been previously reported for 3CLpro[22, 27]. In addition, the effect of pH on the structural stability parallels that of the enzymatic activity, suggesting some titratable residues in the range of pH 5-8 are instrumental to guarantee both these properties. Hydrodynamic radius measurements by dynamic light scattering showed a predominant population of particles compatible with dimeric 3CLpro (hydrodynamic radius of 3.89, 3.33, and 3.94 nm at pH 5, 7, and 8, respectively), with an expected smaller size around its isoelectric point ( $pI \sim 6$ ) due to smaller net charge and lower electrostatic repulsions (Figure S4). Additional experiments using size-exclusion chromatography were also in agreement with a predominantly dimeric 3CLpro (Figure S4). It may be hypothesized that, along the thermal denaturations, temperature could trigger a transient dimerization of dimers into tetramers (tetrameric intermediate) at moderate temperatures, a temperature-driven oligomerization process that has been observed in other proteins[47, 48], before reaching the unfolding temperature and ending in monomer unfolding/dissociation; work on this matter is in progress.

The screening test for 3CLpro could be based on detecting small molecules interfering with the enzymatic activity (which it is fairly easy for a protease, by using a continuous enzymatic assay and a peptidic FRET substrate) or based on detecting the effect of small molecules on altering the thermal stability of the protein against thermal denaturation (by using fluorescence TSA). The TSA-based screening has the advantage of being quite general for any protein, but it has some disadvantages: 1) binding affinity does not correlate with stability changes; 2) false negatives are somewhat common, because ligands may bind to unfolded or non-native protein states besides binding to the native state; 3) stability changes are highly dependent on the binding enthalpy and binding heat capacity; and 4) the ligand binding does not always translates into an inhibitory effect[49]. In contrast, the techniques based on detection of the protein activity provide direct evidence for inhibition during the screening test, but the activity of the protein target must be measurable through a quick and appropriate procedure, which is not straightforward in general.

Through the application of a small-scale activity-based screening to SARS-CoV-2 3CLpro, the quercetin was selected and further studied. Direct evidence for target engagement was gathered by different biophysical techniques, TSA and ITC, as well as through molecular docking simulations. Quercetin showed an inhibition constant of about 7  $\mu\text{M}$ , which corresponds to a binding affinity sufficiently favourable to be considered as a good candidate for optimization or preclinical studies.

The fact that quercetin exerts a destabilizing effect may be related to its physico-chemical properties and the type of interaction with the protein target. The extent of the stabilization

effect induced by a ligand depends on the specific set of conformational states, within the whole conformational landscape of the protein, that interact preferentially with the ligand. If the ligand interacts with most of the conformational states with similar binding affinities, the effect on the protein stability will be small. If the ligand interacts not only with the native state, but also with partially unfolded states (which could be the case for quercetin, due to its hydrophobic moiety), a destabilization effect could be expected if the affinity (or the binding stoichiometry) for the non-native states is higher. These are some of the caveats (described above) to be born in mind when TSA is applied for *in vitro* molecular screening. In addition, unlike in the case of isothermal inhibition assays, where only native-like states are accessible, through TSA it is possible to gradually explore non-native states close to the native state through a temperature-driven process, and binding to non-native states can be detected. In any case, regardless of whether the effect of the ligand on the protein was destabilizing or stabilizing, quercetin interacted and inhibited the activity of 3CLpro.

The influence of quercetin on the structural properties of 3CLpro was assessed (Figure S5), and provided an additional piece of evidence for the interaction of quercetin with 3CLpro. The far-UV CD spectrum of the protein was not significantly distorted by quercetin, but considerable changes occurred in the near-UV CD and fluorescence spectra. In particular, a substantial decrease in tryptophan fluorescence could be observed in 3CLpro upon quercetin binding. Very likely these alterations in the tertiary structure of the protein are associated with the destabilizing effect of quercetin on the thermal stability of 3CLpro observed by TSA.

According to ITC experiments, quercetin binds with favorable enthalpic ( $\Delta H = -3.6$  kcal/mol) and entropic ( $-T\Delta S = -4.0$  kcal/mol) contributions to the Gibbs energy of binding ( $\Delta G = -7.6$  kcal/mol). The hydrophobicity, the compactness and the rigidity of the molecule, and the ability to adopt multiple configurations within the binding site seem to be responsible for a large desolvation entropy gain and small conformational, configurational and rotational entropy losses upon binding, resulting in a global favorable entropy gain. Quercetin would be an appropriate scaffold to engineer new functional groups for developing new inhibitor compounds for SARS-CoV-2 3CLpro. The lower affinity at high ionic strength (Figure 8) indicates that the binding of quercetin to 3CLpro is coupled to the release of salt ions.

Quercetin compares quite well to SARS-CoV-2 3CLpro inhibitors already reported. An intrinsic inhibition constant  $K_i$  of  $0.19 \mu\text{M}$  could be estimated for an alpha-ketoamide inhibitor 13b reported by Zhang et al[22]. In order to compare the inhibition potency of ligands, their molecular size must be considered. The ligand efficiency or binding efficiency index ( $\text{BEI} = \text{p}K_i/\text{MW}$ ) normalizes the inhibitor potency ( $\text{p}K_i = -\log K_i$ ) with its molecular weight (MW)[50].

Because quercetin and inhibitor 13b have a MW of 0.302 kDa and 0.598 kDa, respectively, their ligand efficiencies are 17.0 and 11.2, respectively. Thus, although the alpha-ketoamide inhibitor has a much lower inhibition constant, which is not surprising given its larger MW of the alpha-ketoamide inhibitor; in fact, quercetin is close to be considered as a fragment (MW < 300 Da) from a screening perspective. However, BEI is significantly higher for quercetin, suggesting it is a good candidate establishing good interactions with the target and possessing high potential for optimization and development.

It is worth to note that quercetin is a natural product with well-known pharmacokinetics and ADMET properties and some proposed therapeutic indications, which points to a potential repurposing for COVID-19 treatment. Interestingly, and in agreement with our blind assays, experimental evidences had been reported in the past for an inhibitory effect of quercetin on the 3CLpro from the coronavirus variant MERS-CoV[51]. The work presented here provides a direct evidence for quercetin ability to target SARS-CoV-2 3CLpro on solid experimental bases. Furthermore, it represents a successful starting point that guarantees the possibility to extend our experimental approach for identifying SARS-CoV-2 3CLpro drug candidates to larger chemical libraries, in order to find small molecules to be used (or further optimized) as potential COVID-19 drugs.

#### ACKNOWLEDGMENTS

This work was supported by Fundación hna; Miguel Servet Program from Instituto de Salud Carlos III (CPII13/00017 to O.A.); Fondo de Investigaciones Sanitarias from Instituto de Salud Carlos III, and European Union (ERDF/ESF, 'Investing in your future') (PI18/00343 to O.A. and a FIS Research Contract to L.C.L.); Spanish Ministry of Economy and Competitiveness (BFU2016-78232-P to A.V.C. and SAF2017-83265-R to H.T.R.); Spanish Ministry of Science, Innovation and Universities (FPI Predoctoral Research Contract BES-2017-080739 to D.O.A.); the Spanish National Research Council (CSIC, project 202020E079); Diputación General de Aragón (Predoctoral Research Contract 2019 to A.J.A., 'Protein Targets and Bioactive Compounds Group' E45\_20R to A.V.C., 'Digestive Pathology Group' B25\_20R to O.A.); Centro de Investigación Biomédica en Red en Enfermedades Hepáticas y Digestivas (CIBERehd). The proteomic analysis was performed in the Proteomics Facility of The Spanish National Center for Biotechnology (CNB-CSIC) that belongs to ProteoRed, PRB3-ISCI, supported by grant PT17/0019.

**CONFLICTS OF INTEREST**

The authors declare no conflict of interest.

**SUPPLEMENTARY INFORMATION**

Supplementary Figures S1, S2, S3, S4 and S5, and Tables S1 and S2.

**REFERENCES**

- [1] N. Zhu, D. Zhang, W. Wang, X. Li, B. Yang, J. Song, X. Zhao, B. Huang, W. Shi, R. Lu, P. Niu, F. Zhan, X. Ma, D. Wang, W. Xu, G. Wu, G.F. Gao, W. Tan, A Novel Coronavirus from Patients with Pneumonia in China, 2019, *New England Journal of Medicine* 382(8) (2020) 727-733.
- [2] W.H.O. (WHO), World Health Organization. *Novel coronavirus (COVID-19) situation* . <https://covid19.who.int>.
- [3] C. Huang, Y. Wang, X. Li, L. Ren, J. Zhao, Y. Hu, L. Zhang, G. Fan, J. Xu, X. Gu, Z. Cheng, T. Yu, J. Xia, Y. Wei, W. Wu, X. Xie, W. Yin, H. Li, M. Liu, Y. Xiao, H. Gao, L. Guo, J. Xie, G. Wang, R. Jiang, Z. Gao, Q. Jin, J. Wang, B. Cao, Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China, *Lancet* 395(10223) (2020) 497-506.
- [4] J.G. Adams, R.M. Walls, Supporting the Health Care Workforce During the COVID-19 Global Epidemic, *JAMA*, United States, 2020.
- [5] K. Renu, P.L. Prasanna, A. Valsala Gopalakrishnan, Coronaviruses pathogenesis, comorbidities and multi-organ damage - A review, *Life Sci* 255 (2020) 117839.
- [6] M. Catanzaro, F. Fagiani, M. Racchi, E. Corsini, S. Govoni, C. Lanni, Immune response in COVID-19: addressing a pharmacological challenge by targeting pathways triggered by SARS-CoV-2, *Signal Transduction and Targeted Therapy* 5(1) (2020) 84.
- [7] L. Clarke, An introduction to economic studies, health emergencies, and COVID-19, *J Evid Based Med* 13(2) (2020) 161-167.
- [8] C.L. Atzrodt, I. Maknojia, R.D.P. McCarthy, T.M. Oldfield, J. Po, K.T.L. Ta, H.E. Stepp, T.P. Clements, A Guide to COVID-19: a global pandemic caused by the novel coronavirus SARS-CoV-2, *FEBS J* (2020).
- [9] V. Pirrone, N. Thakkar, J.M. Jacobson, B. Wigdahl, F.C. Krebs, Combinatorial approaches to the prevention and treatment of HIV-1 infection, *Antimicrobial agents and chemotherapy* 55(5) (2011) 1831-1842.
- [10] S. Pushpakom, F. Iorio, P.A. Eyers, K.J. Escott, S. Hopper, A. Wells, A. Doig, T. Williams, J. Latimer, C. McNamee, A. Norris, P. Sanseau, D. Cavalla, M. Pirmohamed, Drug repurposing: progress, challenges and recommendations, *Nature Reviews Drug Discovery* 18(1) (2019) 41-58.
- [11] K.G. Andersen, A. Rambaut, W.I. Lipkin, E.C. Holmes, R.F. Garry, The proximal origin of SARS-CoV-2, *Nature Medicine* 26(4) (2020) 450-452.
- [12] M.Z. Tay, C.M. Poh, L. Rénia, P.A. MacAry, L.F.P. Ng, The trinity of COVID-19: immunity, inflammation and intervention, *Nature reviews. Immunology* 20(6) (2020) 363-374.
- [13] Y.A. Helmy, M. Fawzy, A. Elasad, A. Sobieh, S.P. Kenney, A.A. Shehata, The COVID-19 Pandemic: A Comprehensive Review of Taxonomy, Genetics, Epidemiology, Diagnosis, Treatment, and Control, *Journal of clinical medicine* 9(4) (2020) 1225.
- [14] H. Yang, M. Yang, Y. Ding, Y. Liu, Z. Lou, Z. Zhou, L. Sun, L. Mo, S. Ye, H. Pang, G.F. Gao, K. Anand, M. Bartlam, R. Hilgenfeld, Z. Rao, The crystal structures of severe acute respiratory syndrome virus main protease and its complex with an inhibitor, *Proceedings of the National Academy of Sciences* 100(23) (2003) 13190.

- [15] T. Muramatsu, Y.T. Kim, W. Nishii, T. Terada, M. Shirouzu, S. Yokoyama, Autoprocessing mechanism of severe acute respiratory syndrome coronavirus 3C-like protease (SARS-CoV 3CLpro) from its polyproteins, *FEBS J* 280(9) (2013) 2002-13.
- [16] A. Wu, Y. Wang, C. Zeng, X. Huang, S. Xu, C. Su, M. Wang, Y. Chen, D. Guo, Prediction and biochemical analysis of putative cleavage sites of the 3C-like protease of Middle East respiratory syndrome coronavirus, *Virus Res* 208 (2015) 56-65.
- [17] J.S. Morse, T. Lalonde, S. Xu, W.R. Liu, Learning from the Past: Possible Urgent Prevention and Treatment Options for Severe Acute Respiratory Infections Caused by 2019-nCoV, *Chembiochem : a European journal of chemical biology* 21(5) (2020) 730-738.
- [18] L. Zhang, D. Lin, Y. Kusov, Y. Nian, Q. Ma, J. Wang, A. von Brunn, P. Leyssen, K. Lanko, J. Neyts, A. de Wilde, E.J. Snijder, H. Liu, R. Hilgenfeld,  $\alpha$ -Ketoamides as Broad-Spectrum Inhibitors of Coronavirus and Enterovirus Replication: Structure-Based Design, Synthesis, and Activity Assessment, *J Med Chem* 63(9) (2020) 4562-4578.
- [19] S.A. Khan, K. Zia, S. Ashraf, R. Uddin, Z. Ul-Haq, Identification of chymotrypsin-like protease inhibitors of SARS-CoV-2 via integrated computational approach, *J Biomol Struct Dyn* (2020) 1-10.
- [20] H. Lee, A. Mittal, K. Patel, J.L. Gatuz, L. Truong, J. Torres, D.C. Mulhearn, M.E. Johnson, Identification of novel drug scaffolds for inhibition of SARS-CoV 3-Chymotrypsin-like protease using virtual and high-throughput screenings, *Bioorganic & medicinal chemistry* 22(1) (2014) 167-177.
- [21] C.J. Kuo, Y.H. Chi, J.T. Hsu, P.H. Liang, Characterization of SARS main protease and inhibitor assay using a fluorogenic substrate, *Biochem Biophys Res Commun* 318(4) (2004) 862-7.
- [22] L. Zhang, D. Lin, X. Sun, U. Curth, C. Drosten, L. Sauerhering, S. Becker, K. Rox, R. Hilgenfeld, Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved  $\alpha$ -ketoamide inhibitors, *Science* 368(6489) (2020) 409-412.
- [23] O. Trott, A.J. Olson, AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading, *Journal of computational chemistry* 31(2) (2010) 455-461.
- [24] G.M. Morris, D.S. Goodsell, R.S. Halliday, R. Huey, W.E. Hart, R.K. Belew, A.J. Olson, Automated docking using a Lamarckian genetic algorithm and an empirical binding free energy function, *Journal of Computational Chemistry* 19(14) (1998) 1639-1662.
- [25] M.D. Hanwell, D.E. Curtis, D.C. Lonie, T. Vandermeersch, E. Zurek, G.R. Hutchison, Avogadro: an advanced semantic chemical editor, visualization, and analysis platform, *J Cheminform* 4(1) (2012) 17.
- [26] F. Grande, B. Rizzuti, M.A. Occhiuzzi, G. Ioele, T. Casacchia, F. Gelmini, R. Guzzi, A. Garofalo, G. Statti, Identification by Molecular Docking of Homoisoflavones from *Leopoldia comosa* as Ligands of Estrogen Receptors, *Molecules* 23(4) (2018).
- [27] S. Zhang, N. Zhong, F. Xue, X. Kang, X. Ren, J. Chen, C. Jin, Z. Lou, B. Xia, Three-dimensional domain swapping as a mechanism to lock the active conformation in a super-active octamer of SARS-CoV main protease, *Protein Cell* 1(4) (2010) 371-383.
- [28] J.W. Shriver, S.P. Edmondson, Defining the stability of multimeric proteins, *Methods Mol Biol* 490 (2009) 57-82.
- [29] J.M. Sanchez-Ruiz, Theoretical analysis of Lumry-Eyring models in differential scanning calorimetry, *Biophys J* 61(4) (1992) 921-35.
- [30] U. Bacha, J. Barrila, A. Velazquez-Campoy, S.A. Leavitt, E. Freire, Identification of Novel Inhibitors of the SARS Coronavirus Main Protease 3CLpro, *Biochemistry* 43(17) (2004) 4906-4912.
- [31] U.B. Ericsson, B.M. Hallberg, G.T. Detitta, N. Dekker, P. Nordlund, Thermofluor-based high-throughput stability optimization of proteins for structural studies, *Anal Biochem, United States*, 2006, pp. 289-98.



- [32] M.W. Pantoliano, E.C. Petrella, J.D. Kwasnoski, V.S. Lobanov, J. Myslik, E. Graf, T. Carver, E. Asel, B.A. Springer, P. Lane, F.R. Salemme, High-density miniaturized thermal shift assays as a general strategy for drug discovery, *J Biomol Screen* 6(6) (2001) 429-40.
- [33] P. Cimpmperman, L. Baranauskiene, S. Jachimoviciūte, J. Jachno, J. Torresan, V. Michailoviene, J. Matuliene, J. Sereikaite, V. Bumelis, D. Matulis, A quantitative model of thermal stabilization and destabilization of proteins by ligands, *Biophys J* 95(7) (2008) 3222-31.
- [34] A. González, J. Casado, E. Chueca, S. Salillas, A. Velázquez-Campoy, J. Sancho, Á. Lanás, Small Molecule Inhibitors of the Response Regulator ArsR Exhibit Bactericidal Activity against *Helicobacter pylori*, *Microorganisms* 8(4) (2020).
- [35] R. Villanueva, S. Romero-Tamayo, R. Laplaza, J. Martínez-Olivan, A. Velázquez-Campoy, J. Sancho, P. Ferreira, M. Medina, Redox- and Ligand Binding-Dependent Conformational Ensembles in the Human Apoptosis-Inducing Factor Regulate Its Pro-Life and Cell Death Functions, *Antioxid Redox Signal* 30(18) (2019) 2013-2029.
- [36] J.R. Lakowicz, R. Jayaweera, H. Szmackinski, W. Wiczak, Resolution of multicomponent fluorescence emission using frequency-dependent phase angle and modulation spectra, *Anal Chem* 62(18) (1990) 2005-12.
- [37] W.H. Ward, G.A. Holdgate, Isothermal titration calorimetry in drug discovery, *Prog Med Chem*, Netherlands, 2001, pp. 309-76.
- [38] R. Claveria-Gimeno, S. Vega, O. Abian, A. Velazquez-Campoy, A look at ligand binding thermodynamics in drug discovery, *Expert Opin Drug Discov* 12(4) (2017) 363-377.
- [39] A. González, S. Salillas, A. Velázquez-Campoy, V. Espinosa Angarica, M.F. Fillat, J. Sancho, Á. Lanás, Identifying potential novel drugs against *Helicobacter pylori* by targeting the essential response regulator HsrA, *Sci Rep* 9(1) (2019) 11294.
- [40] P. Santofimia-Castano, Y. Xia, W. Lan, Z. Zhou, C. Huang, L. Peng, P. Soubeyran, A. Velazquez-Campoy, O. Abian, B. Rizzuti, J.L. Neira, J. Iovanna, Ligand-based design identifies a potent NUPR1 inhibitor exerting anticancer activity via necroptosis, *J Clin Invest* 130 (2019).
- [41] J.L. Neira, J. Bintz, M. Arruebo, B. Rizzuti, T. Bonacci, S. Vega, A. Lanás, A. Velázquez-Campoy, J.L. Iovanna, O. Abián, Identification of a Drug Targeting an Intrinsically Disordered Protein Involved in Pancreatic Adenocarcinoma, *Sci Rep* 7 (2017) 39732.
- [42] J. Hidalgo, P. Latorre, J.A. Carrodeguas, A. Velázquez-Campoy, J. Sancho, P. López-Buesa, Inhibition of Pig Phosphoenolpyruvate Carboxykinase Isoenzymes by 3-Mercaptopicolinic Acid and Novel Inhibitors, *PLoS One* 11(7) (2016) e0159002.
- [43] O. Abian, S. Vega, J. Sancho, A. Velazquez-Campoy, Allosteric Inhibitors of the NS3 Protease from the Hepatitis C Virus, *Plos One* 8(7) (2013).
- [44] A.L. Pey, M. Ying, N. Cremades, A. Velazquez-Campoy, T. Scherer, B. Thöny, J. Sancho, A. Martinez, Identification of pharmacological chaperones as potential therapeutic agents to treat phenylketonuria, *J Clin Invest* 118(8) (2008) 2858-67.
- [45] N. Cremades, A. Velázquez-Campoy, M. Martínez-Júlvez, J.L. Neira, I. Pérez-Dorado, J. Hermoso, P. Jiménez, A. Lanás, P.S. Hoffman, J. Sancho, Discovery of specific flavodoxin inhibitors as potential therapeutic agents against *Helicobacter pylori* infection, *ACS Chem Biol* 4(11) (2009) 928-38.
- [46] J. Shi, Z. Wei, J. Song, Dissection study on the severe acute respiratory syndrome 3C-like protease reveals the critical role of the extra domain in dimerization of the enzyme: defining the extra domain as a new target for design of highly specific protease inhibitors, *J Biol Chem* 279(23) (2004) 24765-73.
- [47] L.A. Campos, R. Sharma, S. Alvira, F.M. Ruiz, B. Ibarra-Molero, M. Sadqi, C. Alfonso, G. Rivas, J.M. Sanchez-Ruiz, A. Romero Garrido, J.M. Valpuesta, V. Muñoz, Engineering protein assemblies with allosteric control via monomer fold-switching, *Nature Communications* 10(1) (2019) 5703.
- [48] M.R. Kasimova, S.J. Milstein, E. Freire, The conformational equilibrium of human growth hormone, *J Mol Biol*, 1998 Academic Press Limited., England, 1998, pp. 409-18.

- [49] A. Velazquez-Campoy, J. Sancho, O. Abian, S. Vega, Biophysical Screening for Identifying Pharmacological Chaperones and Inhibitors Against Conformational and Infectious Diseases, *Curr Drug Targets* 17(13) (2016) 1492-505.
- [50] C. Abad-Zapatero, J.T. Metz, Ligand efficiency indices as guideposts for drug discovery, *Drug Discov Today*, England, 2005, pp. 464-9.
- [51] S. Jo, H. Kim, S. Kim, D.H. Shin, M.-S. Kim, Characteristics of flavonoids as potent MERS-CoV 3C-like protease inhibitors, *Chemical Biology & Drug Design* 94(6) (2019) 2023-2030.

## CAPTIONS

**Figure 1. Spectroscopic properties of SARS-CoV-2 3CLpro.** (a) Far-UV CD at 10  $\mu\text{M}$  protein concentration. (b) Near-UV CD at 10  $\mu\text{M}$  protein concentration. (c) Fluorescence at 2  $\mu\text{M}$  protein concentration. All the spectra were recorded at pH 8.

**Figure 2. Spectroscopic thermal unfolding of SARS-CoV-2 3CLpro.** (a) Far-UV CD thermal unfolding monitoring the (raw) ellipticity at 222 nm, at 10  $\mu\text{M}$  protein concentration. (b) Fluorescence thermal unfolding monitoring the intrinsic tryptophan fluorescence emission at 330 nm, at 2  $\mu\text{M}$  protein concentration. Both experiments were recorded at pH 8. The experimental data (circles) and the non-linear regression analysis fitting curves (continuous lines) according to two-state monomeric, dimeric and tetrameric protein are shown (indistinguishable in the plot).

**Figure 3. Calorimetric thermal unfolding of SARS-CoV-2 3CLpro.** (a) Molar excess heat capacity as a function of temperature at pH 5 (black), pH 7 (red), and pH 8 (green), at a 14  $\mu\text{M}$  protein concentration. (b) Molar excess heat capacity as a function of temperature at pH 8. The experimental data (circles) and the non-linear regression analysis fitting curves (continuous lines) according to two-state monomeric (gray), dimeric (green), tetrameric (red), and octameric (blue) protein unfolding models are shown.

**Figure 4. Catalytic activity of SARS-CoV-2 3CLpro.** Time evolution of the substrate fluorescence emission as a function of time at pH 5 (black), pH 7 (red), and pH 8 (green), at protein concentration of 1  $\mu\text{M}$  and substrate concentration of 20  $\mu\text{M}$ . The increase in fluorescence emission reflects the decrease in Dabcyl quenching on Edans emission (i.e., reduction of FRET phenomenon) due to the increase in the spatial separation of the donor-acceptor pair as a result of the catalytic cleavage of the substrate molecule.



**Figure 5. Screening of a small chemical library.** Enzyme activity, in the presence and the absence of compound, was quantitated as the initial slope of the fluorescence emission as a function of time. Control wells (red squares) contained enzyme and substrate in the presence of an identical concentration of DMSO as that in the wells with compounds (black squares). Hits were selected as those compounds lowering the activity below a certain threshold (average activity of controls minus twice the standard deviation of those controls, lower dotted line). The blue square corresponds to quercetin.

**Figure 6. Hit confirmation through inhibition curve.** (a) Substrate fluorescence emission as a function of time, fixing the enzyme concentration at 2  $\mu\text{M}$ , the substrate concentration at 20  $\mu\text{M}$ , and varying quercetin concentration from 0 to 125  $\mu\text{M}$  (while maintaining constant the percentage of DMSO). The increase in quercetin concentration resulted in the decrease in the enzymatic activity. (b) Experimental inhibition data for quercetin (squares), together with the non-linear regression analysis fitting curve (black line) to a simple inhibition, from which the apparent inhibition constant ( $K_i^{\text{app}} = 21 \mu\text{M}$ ) or the intrinsic inhibition constant ( $K_i = 9.6 \mu\text{M}$ ) could be estimated.

**Figure 7. Hit confirmation through thermal shift assay (TSA).** (a) Unfolding traces of the compound-free protein (black) and the protein in the presence of 62.5  $\mu\text{M}$  quercetin (red). (b) Apparent unfolding temperature for the protein at different quercetin concentrations.

**Figure 8. Hit confirmation through isothermal titration calorimetry (ITC).** Interaction of quercetin with 3CLpro assessed by isothermal titration calorimetry in sodium phosphate 50 mM, pH 8, without (left) and with (right) NaCl 150 mM. The upper plot shows the thermogram (thermal power required to maintain a null temperature difference between sample and reference cells as a function of time) and the lower plot shows the binding isotherm (ligand-normalized heat effect per injection as a function of the molar ratio, the quotient between the ligand and protein concentrations in the cell). The fitting curve corresponds to the single ligand binding site model (continuous line). According to the data analysis, in the absence of NaCl quercetin interacts with 3CLpro with favorable enthalpic ( $\Delta H = -3.6 \text{ kcal/mol}$ ) and entropic ( $-T\Delta S = -4.0 \text{ kcal/mol}$ ) contributions to the Gibbs energy of binding ( $\Delta G = -7.6 \text{ kcal/mol}$ ), corresponding to a dissociation constant  $K_d$  of 2.7  $\mu\text{M}$ . In the presence of NaCl quercetin interacts with 3CLpro with favorable enthalpic ( $\Delta H = -4.3 \text{ kcal/mol}$ ) and entropic ( $-T\Delta S = -2.5$

kcal/mol) contributions to the Gibbs energy of binding ( $\Delta G = -6.8$  kcal/mol), corresponding to a dissociation constant  $K_d$  of  $10 \mu\text{M}$ . In both cases, the percentage of active (or binding-competent) protein is 0.75.

**Figure 9. Molecular docking of quercetin to 3CLpro.** Structure of (a) unliganded 3CLpro (from entry 6Y2E of the PDB; in ribbon representation, yellow) and  $\alpha$ -ketoamide inhibitor bound in the active site (from entry 6Y2F; in stick representation, yellow)[22]. The liganded protein was superimposed by a least square fit on the  $C^\alpha$  atoms of the liganded one, and only the latter is shown. (b) and (c) Detail of the 3CLpro active site, with the two best docking poses obtained using as receptor the structure 6Y2E (cyan) and 6Y2F (purple). Polar hydrogen atoms are explicitly shown (white) and oxygen atoms are also coloured (red).

## Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>





Figure 1

**Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening**

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

**Figure 2**

## Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>





Figure 3


**Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening**

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

**Figure 4**


**Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening**

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

**Figure 5**

**Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening**


Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

**Figure 6**




**Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening**

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

**Figure 7**


**Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening**

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

**Figure 8**

**Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening**

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

**Figure 9**

## Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

<sup>1</sup> Instituto Aragonés de Ciencias de la Salud (IACS), 50009, Zaragoza, Spain

<sup>2</sup> Instituto de Investigación Sanitaria de Aragón (IIS Aragón), Zaragoza, Spain

<sup>3</sup> Centro de Investigación Biomédica en Red en el Área Temática de Enfermedades Hepáticas Digestivas (CIBERehd), Madrid, Spain

<sup>4</sup> Institute for Biocomputation and Physics of Complex Systems (BIFI), Joint Units IQFR-CSIC-BIFI, and GBsC-CSIC-BIFI, Universidad de Zaragoza, 50009, Zaragoza, Spain

<sup>5</sup> Departamento de Bioquímica y Biología Molecular y Celular, Universidad de Zaragoza, 50009, Zaragoza, Spain

<sup>6</sup> Department of Immunology and Oncology, National Centre for Biotechnology (CNB), CSIC, 28049, Madrid, Spain

<sup>7</sup> CNR-NANOTEC, Licryl-UOS Cosenza and CEMIF.Cal, Department of Physics, University of Calabria, 87036, Cosenza, Italy

<sup>8</sup> Fundación ARAID, Gobierno de Aragón, 50018, Zaragoza, Spain

### AUTHOR CONTRIBUTIONS

Conceptualization, O.A. and A.V.C.; Methodology, O.A., A.V.C. and B.R.; Software, O.A., A.V.C. and B.R.; Validation, O.A., A.V.C., D.O.A., A.J.A., L.C.L., S.V., B.R. and H.T.R.; Formal Analysis, O.A., A.V.C., D.O.A., A.J.A., L.C.L., S.V., B.R. and H.T.R.; Investigation, D.O.A., A.J.A., L.C.L., S.V., H.T.R. and B.R.; Resources, O.A., A.V.C., B.R. and H.T.R.; Data Curation, O.A., A.V.C., B.R. and H.T.R.; Writing – Original Draft Preparation, O.A., A.V.C. and B.R.; Writing – Review & Editing, O.A., A.V.C., D.O.A., A.J.A., L.C.L., S.V., B.R. and H.T.R.; Visualization, O.A., A.V.C., D.O.A., A.J.A., L.C.L., S.V., and B.R.; Supervision, O.A., A.V.C.; Project Administration, O.A. and A.V.C.; Funding Acquisition, O.A. and A.V.C.

## Structural stability of SARS-CoV-2 3CLpro and identification of quercetin as an inhibitor by experimental screening

Olga Abian<sup>1,2,3,4,5\*</sup>, David Ortega-Alarcon<sup>4,5</sup>, Ana Jimenez-Alesanco<sup>4,5</sup>, Laura Ceballos-Laita<sup>2,4</sup>, Sonia Vega<sup>4</sup>, Hugh T. Reyburn<sup>6</sup>, Bruno Rizzuti<sup>7</sup>, Adrian Velazquez-Campoy<sup>2,3,4,5,8\*</sup>

<sup>1</sup> Instituto Aragonés de Ciencias de la Salud (IACS), 50009, Zaragoza, Spain

<sup>2</sup> Instituto de Investigación Sanitaria de Aragón (IIS Aragón), Zaragoza, Spain

<sup>3</sup> Centro de Investigación Biomédica en Red en el Área Temática de Enfermedades Hepáticas Digestivas (CIBERehd), Madrid, Spain

<sup>4</sup> Institute for Biocomputation and Physics of Complex Systems (BIFI), Joint Units IQFR-CSIC-BIFI, and GBsC-CSIC-BIFI, Universidad de Zaragoza, 50009, Zaragoza, Spain

<sup>5</sup> Departamento de Bioquímica y Biología Molecular y Celular, Universidad de Zaragoza, 50009, Zaragoza, Spain

<sup>6</sup> Department of Immunology and Oncology, National Centre for Biotechnology (CNB), CSIC, 28049, Madrid, Spain

<sup>7</sup> CNR-NANOTEC, Licryl-UOS Cosenza and CEMIF.Cal, Department of Physics, University of Calabria, 87036, Cosenza, Italy

<sup>8</sup> Fundación ARAID, Gobierno de Aragón, 50018, Zaragoza, Spain

### HIGHLIGHTS

- SARS-CoV-2 3CLpro shows cooperative unfolding in which dimers or tetramers predominate depending on the pH.
- SARS-CoV-2 3CLpro shows a pH-dependent hydrolytic activity that correlates with its structural stability.
- SARS-CoV-2 3CLpro is an appropriate target for drug discovery given its high sequence identity among different coronaviruses.
- Quercetin has been identified as a SARS-CoV-2 3CLpro inhibitor by an activity-based experimental screening.
- Experimental and computational evidences for quercetin target engagement have been obtained by ITC, spectroscopy, TSA, and molecular docking.