

1

1 **Dust and bullets: stable isotopes and GPS tracking disentangle lead sources for a**
2 **large avian scavenger**

3 ENEKO ARRONDO*¹, JOAN NAVARRO², JUAN MANUEL PEREZ-GARCÍA^{3,4},
4 RAFAEL MATEO⁵, PABLO R. CAMARERO⁵, ROSA C. RODRÍGUEZ MARTIN-
5 DOIMEADIOS⁵, MARÍA JIMÉNEZ-MORENO⁶, AINARA CORTÉS-AVIZANDA⁷,
6 ISABEL NAVAS⁸, ANTONIO JUAN GARCÍA-FERNÁNDEZ⁸, JOSÉ ANTONIO
7 SÁNCHEZ-ZAPATA³, JOSÉ ANTONIO DONÁZAR¹

8

9 ¹*Department of Conservation Biology, Doñana Biological Station-CSIC, Avda. Américo Vespucio, 26,*
10 *41092, Seville, Spain*

11 ²*Institut de Ciències del Mar Passeig Marítim de la Barceloneta, 37-49. E-08003 Barcelona*

12 ³*Department of Applied Biology, Miguel Hernández University, Avda. de la Universidad, s/n, 03202*
13 *Elche, Alicante, Spain*

14 ⁴*Department of Animal Science, Faculty of Life Sciences and Engineering, University of Lleida, Plaza de*
15 *Victor Siurana, 1, 25198, Lleida, Spain.*

16 ⁵*Instituto de Investigación en Recursos Cinegéticos Ronda de Toledo, 12 13071 Ciudad Real España*

17 ⁶*Departamento De Química Analítica y Tecnología de Alimentos, Instituto de Ciencias Ambientales,*
18 *Universidad de Castilla-LaMancha, Avda. Carlos III s/n, 45071 Toledo, Spain*

19 ⁷*Animal Demography and Ecology Unit, IMEDEA CSIC-UIB, C. Miquel Marqués 21, 07190, Esporles,*
20 *Mallorca, Spain.*

21 ⁸*Area of Toxicology, Department of Health Sciences, IMIB-Arrixaca, University of Murcia, Campus de*
22 *Espinardo, 30100 Murcia, Spain.*

23

24 ***Correspondence author**

25 *Eneko Arrondo, Doñana Biological Station-CSIC, [Avda. Américo Vespucio, 26, 41092 Sevilla](https://www.doñana.csic.es/), Spain*

26 *Tel: +34 954 232 340 / +34 954466700*

27 *E-mail: bioeaf@gmail.com*

28

30 Abstract

31 Lead intoxication is an important threat to human health and a large number of
32 wildlife species. Animals are exposed to several sources of lead highlighting hunting
33 ammunition and lead that is bioavailable in topsoil. Disentangling the role of each in
34 lead exposure is an important conservation issue, particularly for species potentially
35 affected by lead poisoning, such as vultures. The identification of lead sources in
36 vultures and other species has been classically addressed by means of stable-isotope
37 comparisons, but the extremely varied isotope signatures found in ammunition hinders
38 this identification when it overlaps with topsoil signatures. In addition, assumptions
39 related to the exposure of individual vultures to lead sources have been made without
40 knowledge of the actual feeding grounds exploited by the birds. Here, we combine lead
41 concentration analysis in blood, novel stable isotope approaches to assign the origin of
42 the lead and GPS tracking data to investigate the main foraging grounds of two Iberian
43 griffon vulture populations (N=58) whose foraging ranges differ in terms of topsoil lead
44 concentration and intensity of big game hunting activity. We found that the lead
45 signature in vultures was closer to topsoil than to ammunition, but this similarity
46 decreased significantly in the area with higher big game hunting activity. In addition,
47 attending to the individual home ranges of the tracked birds, models accounting for the
48 intensity of hunting activity better explained the higher blood lead concentration in
49 vultures than topsoil exposure. In spite of that, our finding also show that lead exposure
50 from topsoil is more important than previously thought.

51 Key words:

52 Lead, ammunition, ecotoxicology, GPS, vultures, stable isotopes

53 1. **Introduction**

54 Lead is a heavy metal whose toxic effects in humans have been known for millennia
55 (Papanikolaou et al., 2005). Its consequences in wildlife, however, were not described
56 until the 19th century (Calvert 1876). Since then, direct mortality due to lead toxicity
57 has been frequently reported for many avian species (Pain et al., 2019). There are,
58 nonetheless, more subtle and barely detectable sub-lethal effects that often go
59 unnoticed, such as alterations in behavior, morphology, and breeding success or
60 physiological functions ([Espín et al., 2015](#); Golden et al., 2016; Vallverdú-Coll et al.,
61 2016). Consequently, the study of the impact of lead pollution on wildlife has become
62 an extremely active field in conservation of threatened populations (Pain et al., 2019).

63

64 Vultures are one of the bird groups most sensitive to lead intoxication to the extent
65 that it has been noted as a significant conservation problem for many vulture species
66 worldwide (Golden et al., 2016; Plaza & Lambertucci 2019), threatening entire
67 populations and compromising the success of costly conservation programs (Finkelstein
68 et al., 2012). The obligate scavenging habits of vultures make them very prone to
69 ingesting ammunition from big game hunting remains (Mateo et al., 1997; García-
70 Fernández et al., 2005; Krone 2018). Carcasses and remains of shot animals are
71 frequently abandoned in nature ([Hunt et al., 2006](#); Legagneux et al., 2014) and can
72 contain up to hundreds of fragments of metallic lead that can be bioavailable for
73 vultures because of the characteristic extremely acidic gastric fluid of these species
74 (Hunt et al., 2006; Hunt et al., 2009; Knot et al., 2010).

75 Ammunition is not the only source of lead that could affect vultures. Alternative
76 sources of lead such as paint, contaminated water or soils have also been described as
77 possible causes of intoxication in wildlife (Katzner et al 2018). Some of them, such as

78 lead-based paint, are of little relevance to scavengers because of their low exposure
79 occurrence (Finkelstein et al., 2102). On the contrary, lead in soil is naturally
80 widespread, and mining activities have led to its bioavailability to wildlife. This is
81 relevant because wild and domestic ungulates, whose carcasses are the main food
82 source for vultures, accumulate lead from the soil in their tissues triggering potential
83 trophic transfer processes affecting higher trophic levels (García-Fernández, 2014;
84 Mateo-Tomás et al., 2016; Naidoo et al., 2017).

85 Starting from this scenario, it is crucial to identify the role that ammunition and
86 topsoil lead play in vulture intoxication, not only to counteract resistance to global
87 regulations on lead hunting ammunition (Cromie et al., 2014), but also to rule out
88 possible underestimates of the risk posed by topsoil lead. Thus far, the most direct
89 approaches have made use of stable isotope signatures (Church et al 2006; Mateo-
90 Tomás et al., 2016; Naidoo et al., 2017). In addition, the application of stable isotope
91 mixing models goes one step further, allowing a detailed assessment of the contribution
92 of potential lead sources (Longman et al., 2018). This approach alone, however, is
93 incomplete. It is well known that large avian scavengers perform huge long-distance
94 movements (Alarcón & Lambertucci 2018), which makes it difficult to determine where
95 the individuals may have been exposed to lead in topsoil and/or game carcasses
96 (Binkowski et al., 2016). In addition, from a population point of view, individual
97 foraging decisions are highly variable (Alarcón & Lambertucci 2018), which implies the
98 possibility that different birds in the same breeding area could be unequally exposed to
99 different lead sources. Recent studies have tried to deal with this but have been based on
100 direct observations (Church et al., 2006; Mateo-Tomás et al., 2016; Naidoo et al., 2017),
101 which can introduce important biases when the home ranges are very large or include
102 poorly accessible areas.

103 Here, taking advantage of GPS tracking of 58 griffon vultures of two Spanish
104 populations differently exposed to topsoil and ammunition, we aim to identify the
105 contribution of topsoil and ammunition sources to lead concentrations in the blood of
106 the tracked birds. Spain is an excellent place to address this issue because it holds 90%
107 of the European population and shows a high prevalence of abnormal blood lead levels
108 (García-Fernández et al., 2005; Mateo-Tomás et al. 2016; Descalzo and Mateo 2018).
109 Moreover, Spanish vultures are exposed to both target lead sources. Whereas elevated
110 lead exposure has been reported in wild ungulates, as well as in livestock, because of
111 topsoil contamination in some Spanish regions (Reglero et al., 2009, Taggart et al.,
112 2011, Pareja-Carrera et al., 2014), the populations of these game species are recovering
113 across most of the country, with the number of animals hunted being one of the largest
114 in Europe (Apollonio et al 2010). Our aim is to estimate for the first time, linkages
115 between sources of lead in the environment and that found in griffon vultures and the
116 spatial scale at which this species may be exposed to lead. We specifically predict that
117 1) blood lead in individual vultures derives from two different sources, ammunition and
118 topsoil; 2) lead in the blood of vultures differs between populations based on the
119 individual level of exposure to topsoil and ammunition; and 3) exposure to big game
120 hunting is the major driver of high levels of blood lead concentration.

121

122 2. **Methods**

123 2.1. *Focus species and study area*

124 The European griffon vulture is a large body-sized (up to 12 kg) obligate scavenger.
125 It is the most abundant European vulture (Margalida et al., 2010). The bulk (90%) of the
126 European populations are concentrated in Spain (Margalida et al., 2010) where a 2018
127 census estimated 30.946 breeding pairs (del Moral and Molina 2018). They nest on

128 cliffs and their main source of food is domestic and wild ungulates (Margalida et al.,
129 2011). They feed over areas covering thousands of square kilometers (Arrondo et al.,
130 2018) and thus rely on social information (Cortés-Avizanda et al 2014).

131 We captured and tagged 58 adult (more than seven years old) griffon vultures in two
132 distant populations (hereafter “southern” and “northern”) of the Iberian Peninsula (see
133 Arrondo et al 2019). Captures were done at baited sites by means of cannon-nets. Thirty
134 birds were trapped in Sierra de Segura Cazorla y las Villas Natural Park, Southern Spain
135 (Figure 1) in December 2014. The movements of these vultures extend mainly
136 westwards to the Portuguese border (see Arrondo et al., 2018). This area is dominated
137 by Mediterranean woodlands and “dehesas”, which are traditional silvopastoral
138 landscapes where two of the main economic activities are traditional livestock
139 (including free-ranging herds of sheep and pigs) and big game hunting (Acevedo et al.,
140 2011). In addition, this area has hosted significant lead-mining operations for centuries
141 (Reglero et al., 2009). The other 28 vultures were captured in Bardenas Reales Natural
142 Park, Northern Spain (Figure 1) in December 2015. In this area, griffon vultures are
143 mainly concentrated around Ebro Valley, a relatively flat area mainly characterized by
144 irrigated crops and intensive livestock farms and surrounded by mountain ranges with
145 Mediterranean woodlands and pastures (Lecina et al., 2005; Martín-Queller et al.,
146 2010). Big game hunting is common but less intense than in the southern area (Acevedo
147 et al., 2014). In addition, there is no history of lead mining activity, but natural lead is
148 present at high concentrations in mountain topsoil (Locutura et al., 2012). Additionally,
149 griffon vultures from this population also travel long distances to Southwestern Iberia,
150 where they share some foraging zones with vultures from the southern population
151 (Arrondo et al., 2018, 2020 and Figure 1).

152 Trapping and handling were carried out with the proper permits and bioethical
153 authorizations. During handling, safety protocols were followed to avoid stressing the
154 animals. Until the moment of the tagging, the individuals were isolated and safe. The
155 tagging was always done by at least two people and never lasted more than twenty
156 minutes.

157 All the individuals were tagged with 90 g GPS/GPRS-GSM backpack devices from
158 E-OBS Digital Telemetry (<https://www.e-obs.de/http://www.e-obs.de>). Devices were
159 equipped as backpacks using a Y type harness made of Teflon following the procedures
160 described in Kenward (2000). The devices were programmed to record variable
161 numbers of locations depending on weather conditions and the power level of the
162 batteries. During spring and summer, devices recorded one fix every 5 minutes if the
163 battery was full, every 20 minutes if the battery was half-full and every 30 minutes if
164 the battery was close to empty. In autumn and winter, the devices recorded every 10
165 minutes if the battery was full, every 30 minutes if the battery was half-full and every
166 60 minutes if the battery was close to empty. Throughout the year, if the batteries were
167 discharged below the safety level, the device would only record one fix per day. We
168 compiled movement data for all birds since the capture until December 2018 unless the
169 animal died or the device failed (Table S.1.).

170

171 2.2. *Lead analysis and isotopic determination*

172 We took blood samples by brachial puncture from all of the individuals. Whole blood
173 without anticoagulant was stored at -20 °C until the analyses of blood lead concentration
174 and isotope composition. Blood samples were also used to determine the sex of the
175 birds by molecular procedures (Wink et al., 1998).

176 Blood samples (0.4-1.0 g) were digested with 3 ml of HNO₃ (69% Analytical
177 Grade), 1ml of H₂O₂ (30% v/v Suprapur) and 4 ml of H₂O (Milli-Q grade) with a
178 microwave oven (Ethos E, Milestone) (Reglero et al., 2009). Lead concentrations and
179 the proportion of the stable isotopes ²⁰⁶Pb, ²⁰⁷Pb, and ²⁰⁸Pb were measured in the
180 digested solutions by inductively coupled plasma quadrupole mass spectrometry (ICP-
181 MS) following Martínez-Haro et al., (2011).

182 Stable lead isotope composition was also analyzed in the topsoil of mining sites of
183 Sierra Madrona-Valle de Alcudia (Table S.2). Here, elevated lead concentrations have
184 been detected in soil (average values of different sites: 7.78-8897 µg/g; Reglero et al.,
185 2008; Rodríguez-Estival et al., 2014) and in wild ungulates (red deer and wild boar
186 muscle showed geometric means with 95%CI of 0.483 (0.32-0.73) and 2.63 (1.13-6.15)
187 µg/g) and the livestock (sheep liver showed 6.16 (4.12-9.23) µg/g; Reglero et al., 2009,
188 Taggart et al., 2011, Pareja-Carrera et al., 2014). Additionally, vultures from both
189 populations usually forage in this area (Figure 1). The isotope ratios in topsoil lead of
190 this region are very similar to that found in the northern study area (Monna et al 2004).
191 Soil samples (≈ 100 g) were taken at a depth of 0-5 cm using a shovel and stored in
192 independent ziplock polyethylene bags. Soil samples were oven-dried, disaggregated in
193 a mortar and sieved through a 250 µm-aperture nylon mesh before being acid-digested
194 (0.2 g) as described above.

195 We also determined the isotopic composition of the most frequently used lead-
196 based bullets in Spain. For this purpose, we obtained 17 bullets and 3 cartridges of 6
197 commercial brands (Table S.2).

198 Blanks and a certified reference material of lobster hepatopancreas (TORT-2) with
199 0.39 µg/g of lead were processed in each batch of digestions. The limit of detection
200 (LODs) of lead in blood was 0.32 µg/dl. We calculated blood lead concentration in

201 $\mu\text{g}/\text{dl}$ considering blood density at 1.06 g/ml to make our results more comparable with
202 the available literature. The mean (\pm %RSD) lead recovery in the reference material
203 TORT-2 was 94.7% (\pm 5.8%, $n = 12$). The precision expressed as %RSD was lower
204 than 5.5% for lead concentration data ($n=12$).

205 Key operating conditions for isotope determination were quadrupole dwell time (10
206 ms for ^{206}Pb and ^{207}Pb and 5 ms for ^{208}Pb), number of scans per sample (800 sweeps),
207 and dead time correction factor (35 ns). Both internal ($^{203}\text{Tl}/^{205}\text{Tl}$ ratio) and external
208 (NIST SRM 981, certified isotopic composition (mean \pm 95%) of $24.144 \pm 0.006\%$ for
209 ^{206}Pb , of $22.083 \pm 0.003\%$ for ^{207}Pb , and of $52.347 \pm 0.009\%$ for ^{208}Pb) standards were
210 used for mass discrimination correction. All isotope ratios determined for SRM 981
211 during analysis were within an uncertainty $<1\%$ of the certified value (before a nominal
212 rolling correction was applied to all data). For isotopic analysis, six replicates of each
213 sample were run. Variability in isotopic data expressed as %RSD ($n=6$) was in all cases
214 lower than 0.28%. Detailed values for each lead isotope ratio and type of sample are
215 shown in Table S.3.

216

217 2.3. *Spatial variables*

218 We estimated the home ranges of GPS-tracked griffon vultures exclusively during
219 the big game hunting period (October to March). Since the birds were captured in the
220 middle of this period (in December, see above), we assume that the lead concentration
221 levels recorded are representative of the lead exposure during whole hunting period. To
222 ensure that core and foraging areas do not show a significant spatial variation during the
223 study period, we assessed the stability of home ranges. According to Fieberg and
224 Kochanny (2005), we used the Bhattacharyya's affinity (BA) index and the home range
225 estimators overlap (HRE).

226 Before performing home range estimations, we standardized our data by resampling
227 the dataset until we obtained for each individual a fix every 30 minutes. Home range
228 and overlapping analyses were done by means of bivariate kernel functions using the
229 adehabitatHR package (Calenge & Fortmann-Roe 2013) run in R version 3.5.1 (R
230 development core team 2018). Fixed 95% and 50% kernel density contours were
231 calculated to estimate the majority of the foraging areas, KDE 95%, and the core
232 (intensive use) areas, KDE 50%. We used as a smoothing parameter the ad hoc method
233 with a resolution of one ha (Margalida et al., 2016).

234 Potential topsoil and ammunition exposures were estimated by means of proxy
235 variables. In the first case, and on the basis of the national geochemical atlas (resolution
236 1x1m) elaborated by the Spanish Geological and Mining Institute (Locutura et al.,
237 2012), we calculated the median lead concentration (mg/kg) at the superficial ground
238 inside the KDE 50 and KDE 95 areas of each individual. Exposure to ammunition was
239 estimated in relation to hunting statistics. We defined the hunting intensity in KDE50
240 and KDE95 as the sum of wild boars (*Sus scrofa*) and red deer (*Cervus elaphus*) culled
241 (<https://www.mapa.gob.es/es/desarrollo-rural/estadisticas/>) in a 10 x10 km cell covering
242 all of peninsular Spain ([https://www.miteco.gob.es/es/biodiversidad/temas/inventarios-
243 nacionales/inventario-especies-terrestres/inventario-nacional-de-biodiversidad/bdn-ieet-
244 atlas-vert-mamif.aspx](https://www.miteco.gob.es/es/biodiversidad/temas/inventarios-nacionales/inventario-especies-terrestres/inventario-nacional-de-biodiversidad/bdn-ieet-atlas-vert-mamif.aspx)[https://www.miteco.gob.es/es/biodiversidad/temas/inventarios-
245 nacionales/inventario-especies-terrestres/inventario-nacional-de-biodiversidad/bdn-ieet-
246 atlas-vert-mamif.aspx](https://www.miteco.gob.es/es/biodiversidad/temas/inventarios-nacionales/inventario-especies-terrestres/inventario-nacional-de-biodiversidad/bdn-ieet-atlas-vert-mamif.aspx)).

247 Statistical details of both variables are described in Table S.4.

248

249

250 2.4. *Statistical analyses*

251 2.4.1. Lead sources in individual vultures

252 To infer the potential origin of the lead present in the blood of vultures, we applied
253 stable isotopic Bayesian mixing models (MixSIAR, Stock et al., 2018) using three lead
254 stable isotopic ratios, $^{206}\text{Pb}/^{207}\text{Pb}$, $^{207}\text{Pb}/^{208}\text{Pb}$, and $^{206}\text{Pb}/^{208}\text{Pb}$, in the blood of each GPS-
255 tracked vultures. We avoided the use of ^{204}Pb isotope because of its low presence in the
256 isotopic signature, which could introduce analytical biases in the calculations of isotope
257 ratios in biological samples with low lead levels. MixSIAR Bayesian isotopic mixing
258 models estimate the potential contribution of each isotopically distinct potential origin
259 of lead (in our case topsoil and ammunition sources) in the diet of the consumer (in our
260 case griffon vultures) based on the lead isotopic values of the consumer and its potential
261 source. MixSIAR estimates probability density functions using Markov chain Monte
262 Carlo methods, and each model was run with identical parameters. Model convergence
263 was determined using Gelman-Rubin and Geweke diagnostic tests (Stock & Semmens,
264 2016; Stock et al., 2018). Bayesian mixing models have been developed to allow
265 flexible model specification in a rigorous Bayesian statistical framework (Phillips et al.,
266 2014). We did not use trophic enrichment factors between vulture's blood and sources
267 of lead because no trophic enrichment factor occurs with lead as occurs with nitrogen
268 (Longman et al 2018).

269

270 2.4.2. Factors associated with blood lead concentration in vultures

271 We related the blood lead concentration, transformed by logarithm in base 10, to
272 the explanatory variables using General Linear Models (Gaussian error distribution and
273 identity linkage). The explanatory variables selected were: a) median topsoil lead
274 concentration at KDE 50; b) median topsoil lead concentration at KDE 95; c) big game

275 hunting intensity at KDE 50; d) big game hunting intensity at KDE 95, e) area of
276 KDE50, f) area of KDE95 and g) sex.
277 The two spatial scales analyzed (KDE50 and KDE95) were highly correlated in all
278 variables (topsoil lead concentration: $t = 6.94$, $df = 58$, $p < 0.001$, $r = 0.67$; big game
279 hunting intensity: $t = 50.90$, $df = 58$, $p < 0.001$, $r = 0.99$; area: $t = 9.82$, $df = 58$, $p < 0.001$,
280 $r = 0.79$). In addition, topsoil lead concentration and big game hunting intensity were
281 correlated at both scales KDE50 and KDE95 (KDE50: $t = 6.52$, $df = 58$, $p < 0.001$, $r =$
282 0.65 ; KDE95: $t = 21.02$, $df = 58$, $p < 0.001$, $r = 0.96$). All correlated variables were
283 modeled independently.

284 Model selection was done by means of the Akaike's information criterion corrected
285 for small sample size (AICc). Models with $\Delta AICc < 2$ were considered equivalents. We
286 discarded models including uninformative parameters, i.e. parameters whose 85%
287 confidence interval overlapped with 0 (Burnham and Anderson, 2002).

288

289 3. Results

290 Lead values above the background and toxic levels ($> 20 \mu\text{g}/\text{dl}$ and $> 50 \mu\text{g}/\text{dl}$, Pain
291 et al. 2019) appeared in 93.3% and 78.6 % of individuals from the southern population
292 and 66.7% and 28.6 % of individuals from the northern population, respectively (Table
293 1). Vultures from the southern population showed significantly higher mean lead
294 concentrations than those from the northern population (mean \pm SD respectively: $64.0 \pm$
295 29.9 vs. $40.1 \pm 25.3 \mu\text{g}/\text{dl}$; $t = -3.324$, $df = 54.718$, $p = 0.002$). Females tended to show
296 higher frequencies of toxic ($> 50 \mu\text{g}/\text{dl}$) lead concentrations than males: 72.7% vs.
297 63.2% and 37.5% vs. 28.6% of the birds in southern and northern populations,
298 respectively (Table 1).

299

300 3.1. *Stable isotopic results*

301 We found higher stable isotope ratios of $^{207}\text{Pb}/^{208}\text{Pb}$ and $^{206}\text{Pb}/^{208}\text{Pb}$ in vultures
302 sampled in the southern compared to the northern area (Figure 2; T-Student Tests;
303 $^{207}\text{Pb}/^{208}\text{Pb}$, $t=2.21$, $p=0.03$; $^{206}\text{Pb}/^{208}\text{Pb}$, $t=2.26$, $p=0.02$). In contrast, both populations
304 showed similar stable isotope ratios of $^{206}\text{Pb}/^{207}\text{Pb}$ (Figure 2; $t=2.21$, $p=0.03$). In the case
305 of lead sources, ammunition always showed higher stable isotope ratios of $^{207}\text{Pb}/^{208}\text{Pb}$,
306 $^{206}\text{Pb}/^{208}\text{Pb}$ and $^{206}\text{Pb}/^{207}\text{Pb}$ than topsoil (Figure 2).

307 Lead source estimates derived from isotopic mixing models revealed that, for both
308 populations the isotopic signature seems to be closer to topsoil than ammunition (Figure
309 3; T-Student tests; topsoil vs. ammunition; northern population, $t=-9.33$, $p<0.001$;
310 southern population, $t=-4.96$, $p<0.001$). However, the importance of ammunition was
311 higher in the southern than in the northern population (Figure 3; southern vs. northern
312 population; topsoil, $t=7.38$, $p<0.001$; ammunition, $t=-7.36$, $p<0.001$).

313

314 3.2. *Modeling blood lead concentration*

315 Overlap between years of utilization distribution areas was high ($60 \pm 25\%$, BA index),
316 while both KDE50% and KDE95% showed high stability (HRE: index showed 42.41%
317 $\pm 31.69\%$ and $47.93\% \pm 32.60\%$ (indiv= 46; indiv-year=286)).

318 We obtained three AIC-equivalent models explaining blood lead concentrations in
319 vultures (Table S5 and Table 2). Two models showed an effect of exposure to
320 ammunition from big game hunting based on the KDE50 and KDE95 with an additive
321 effect of sex. The third model selected included an effect of topsoil lead concentration at
322 KDE50. In spite of the equivalence of the three models, the one that included big game
323 hunting intensity at KDE50 and sex presented a weight of 44%, more than double the
324 models that included big game hunting intensity at KDE595 and topsoil lead

325 concentration whose weights were 20% and 17%, respectively. That is, those
326 individuals, especially females whose core areas are in areas with high intensities of big
327 game hunting, have higher levels of lead in blood (Table. S.5)

328

329 4. Discussion

330 Our results reveal that both topsoil and ammunition are important sources of lead
331 found in the blood of griffon vultures, but their relative contribution is clearly
332 asymmetric. Most of the vultures were exposed to background lead levels probably
333 derived from both direct topsoil exposure (e.g. contaminated dust inhalation or
334 ingestion) and a transfer between trophic levels. Toxic levels of lead is mainly
335 explained, however, by the ingestion of hunting ammunition. Thus, our study, with the
336 combination of GPS and isotopic signatures of blood lead analyses, is the first to
337 provide a fine-tuned approach to disentangling how fine-scale foraging patterns
338 determine individual variations in the contribution of different sources of lead.

339

340 4.1. *Sources of lead exposure in griffon vultures*

341 Our results showed that topsoil could has an important contribution to the lead
342 found in vultures which could be explained by chronic exposure to this source
343 compared to the exceptional exposure to ammunition. Topsoil lead is widely present in
344 foraging areas of both northern and southern populations. The bulk of the vultures' diet
345 is domestic and wild ungulates that are consistently exposed to lead from the topsoil,
346 especially in mining areas (Reglero et al., 2009; Taggart et al., 2011; Pareja-Carrera et
347 al., 2014). Apart from this, the remains of hunted wild ungulates in regions with topsoil
348 lead would contain lead from both sources (topsoil and ammunition). It should also be
349 noted, that the average concentration of lead in the muscle of ungulates from mining

350 areas is relatively low (0.08-2.6 $\mu\text{g/g}$; Taggart et al., 2011; Pareja-Carrera et al., 2014),
351 whereas a single piece of ammunition from a wounded animal can weigh more than 1
352 mg (Nadjafzadeh et al., 2015). Consequently, vultures would be continuously
353 incorporating small amounts of lead from the topsoil and exceptionally, large quantities
354 from ammunition.

355 This idea is reinforced by modeling procedures that showed that high levels of
356 blood lead concentrations were related to exposure to ammunition lead. It is well known
357 that ammunition is an agent of clinical lead intoxications in birds of prey (García-
358 Fernández, 2014; Naidoo et al., 2017; Garbett et al., 2018; Krone et al., 2018). More
359 recently, the presence of lead from topsoil and ammunition in griffon vultures has been
360 described (Mateo-Tomás et al., 2016). Nevertheless, to our knowledge, this is the first
361 time the relative contribution of both sources has been studied by integrating stable
362 isotope analysis with fine-scale GPS monitoring.

363 It could be argued that hunting intensity and topsoil lead exposure show high
364 spatial covariance. These results could be obscuring an additive effect between topsoil
365 and ammunition and can explain the striking differences found in lead concentrations
366 between the two populations, which confirms the findings of Mateo-Tomás et al.,
367 (2016) in another region in Spain. Thus, the southern population would be more
368 exposed not only to ammunition (Figure 1) but also to lead in the topsoil. In fact, tissues
369 from wild and domestic ungulates from our southern study area showed high
370 concentrations of lead in contrast to the levels found in these species in other Spanish
371 areas not affected by mining pollution (Santiago et al., 1998; Taggart et al., 2011;
372 Pareja-Carrera et al., 2014). For example, red deer and wild boar from southern study
373 area have lead in muscle of 0.48 and 2.63 respectively. This contrasts with the levels
374 found in these species in other areas not affected by lead mining pollution, where red

375 deer and wild boar showed 0.12 (0.08-0.19) and 0.32 (0.12-0.80) $\mu\text{g/g}$ d.w. of lead in
376 muscle, respectively (Taggart et al., 2011). Similarly, lead concentrations in liver of red
377 deer and wild boar from the mining sites were higher in the southern area (0.43 and 1.92
378 $\mu\text{g/g}$) than in control sites (0.11 and 0.39 $\mu\text{g/g}$) These differences are also noticeable in
379 domestic ungulates. Sheep southern area showed lead levels in liver and muscle of 6.16
380 (4.12-9.23) and 0.08 (0.07-0.09) $\mu\text{g/g}$ d.w., respectively, which are well above the levels
381 found in sheep from control sites of 0.21 (0.13-0.35) and 0.04 (0.03-0.05) $\mu\text{g/g}$ d.w., in
382 liver and muscle, respectively (Pareja-Carrera et al., 2014). All of this means that
383 griffons feeding on carrion from the southern area can be exposed to lead levels 2 to
384 8.3-fold greater through a diet of muscle and 4.9 to 29.3-fold higher from liver
385 consumption, which may well partially explain the higher background blood lead
386 concentrations found in griffon vultures from the southern area.

387 Our models showed that female vultures had higher lead levels that match previous
388 studies in this species (Mateo-Tomás et al., 2016). Our blood samples were taken in
389 winter, coinciding with the beginning of the breeding season and thanks to GPS, we
390 were able to verify that at least 78% of the females and 65% of the males tagged bred
391 during the season in which they were equipped with GPS. Thus, it is reasonable to
392 hypothesize that the sex-based differences could be due to the mobilization of lead from
393 bones occurring during eggshell formation (Gangoso et al., 2009) but certainly further
394 studies would be required to test this hypothesis.

395

396 4.2. *Ecological/Physiological Consequences of high lead exposure in vultures*

397 Almost 80% of the individuals from the southern population and 30% from the
398 northern population were above the threshold value limit established for clinical toxicity
399 (50 $\mu\text{g/dl}$; Pain et al., 2019). These high lead concentrations are probably related to the

400 fact that the studied vultures were captured in winter, during the big game hunting
401 season (Espín et al., 2014; Hernández & Margalida 2009; Mateo-Tomás et al., 2016;
402 Krone 2018; Garbett et al 2018). In any case, these lead values were above the
403 concentrations described in other species of large avian scavengers (Plaza &
404 Lambertucci 2019; Krüger & Amar 2018) and were comparable to those found in the
405 California condor (*Gymnogyps californianus*) undergoing chelation therapy to counter
406 lead poisoning (Finkelstein et al., 2012). However, we did not detect any deaths
407 attributable to lead intoxication (Arrondo et al., 2020), nor did we perceive intoxication
408 symptoms such as anorexia, dropping head or vomiting in the sampled individuals
409 during the handling process (Krone et al., 2018). This confirms the already described
410 high resistance of griffon vultures to lead exposure (García-Fernandez et al., 2005;
411 Espín et al., 2014). In fact, deaths due to lead exposure are known but seem
412 comparatively rare in relation to other vultures and large body-sized facultative
413 scavenger species (Mateo et al., 1997; Mateo 2009; Horowitz et al., 2014). Beyond the
414 absence of direct mortality and visible symptoms of intoxication, we cannot discard
415 hidden negative effects derived from chronic exposure such as alterations in bone
416 mineralization (Gangoso et al., 2009), physiological effects such as the suppression of
417 δ -ALAD (Espín et al., 2015) or behavioral alterations derived from sub lethal
418 exposures.

419

420 5. Further remarks

421 Topsoil lead can be found naturally (Locutura et al., 2012) but pollution derived
422 from mining activity as occurs in our southern study area is a major problem for wildlife
423 and ecosystems, largely because lead mining activity in Europe has been occurring for
424 millennia (Reglero et al., 2009, Taggart et al., 2011). Although for our target species, no

425 consequences were detected, it is possible to hypothesize that other sensitive threatened
426 species such as Egyptian vultures (*Neophron percnopterus*), red kites (*Milvus milvus*) or
427 Spanish imperial eagles (*Aquila adalberti*) can be affected if their territories and home
428 ranges include highly contaminated mining areas. Consequently, detailed information
429 on topsoil contamination at the level of the entire Iberian Peninsula is necessary to be
430 able to predict damage to wildlife, livestock and human health.

431 Our results also reinforce the idea that ammunition is the main cause of toxic lead
432 concentration in scavenger birds, such as vultures (García-Fernández 2014; Krone et al.,
433 2018; Pain et al., 2019). This finding is especially relevant in the current context of rural
434 abandonment in which wild ungulates are spreading across Europe as part of a passive
435 rewilding process (Apollonio et al., 2010). In parallel to the growth of wild ungulates
436 populations, hunting pressure is also increasing (Herruzo & Martínez-Jauregui 2013).
437 This inevitably entails a greater exposure to lead and more risk of intoxication for
438 vultures and other scavenger species that consume both the discarded remains of killed
439 animals and the carcasses of mortally injured animals not collected by hunters. In
440 addition, based on our results, exposure to ammunition could be occurring hundreds of
441 kilometers away from the breeding colonies. This is especially relevant for large body-
442 sized scavenging species, which can fly long distances daily crossing administrative
443 boundaries that expose them to different, and sometimes contradictory, legislation
444 (Arrondo et al., 2018). Therefore, the decision to ban lead ammunition partially or at
445 the local scale (Avery & Watson 2009; Mateo & Kanstrup 2019) may be insufficient. It
446 is obvious that a change in legislation regarding the replacement of lead with other
447 materials requires European regulations to develop integral conservation strategies
448 (Lambertucci et al., 2014; Arrondo et al., 2018). This might also contribute to

449 promoting hunting as a more sustainable activity within a rewilding Europe (Kanstrup
450 et al., 2018).

451

452 **Acknowledgements**

453 The research was funded by Comunidad de Bardenas Reales de Navarra the Project
454 RNM-1925 (Junta de Andalucía), Project CGL2015-66966-C2-1-2-R (Spanish Ministry
455 of Economy and Competitiveness and EU/ERDF) and Project PPII-2014-028-P (Junta
456 de Comunidades de Castilla-La Mancha). EA was supported by La Caixa-Severo Ochoa
457 International PhD Program 2015. JN was funded by the Spanish National Program
458 Ramón y Cajal (RYC-2015-17809).

459

460 **References**

- 461 1. Acevedo, P., Farfán, M. A., Márquez, A. L., Delibes-Mateos, M., Real, R., &
462 Vargas, J. M. (2011). Past, present and future of wild ungulates in relation to
463 changes in land use. *Landscape Ecology*, 26(1), 19-31.
- 464 2. Acevedo, P., Quirós-Fernández, F., Casal, J., & Vicente, J. (2014). Spatial
465 distribution of wild boar population abundance: Basic information for spatial
466 epidemiology and wildlife management. *Ecological Indicators*, 36, 594-600.
- 467 3. Alarcón, P. A., & Lambertucci, S. A. (2018). A three-decade review of telemetry
468 studies on vultures and condors. *Movement Ecology*, 6(1), 13.
- 469 4. Apollonio, M., Andersen, R., & Putman, R. (Eds.). (2010). *European ungulates*
470 *and their management in the 21st century*. Cambridge University Press.
- 471 5. Arrondo, E., Moleón, M., Cortés-Avizanda, A., Jiménez, J., Beja, P., Sánchez-
472 Zapata, J. A., & Donazar, J. A. (2018). Invisible barriers: Differential sanitary

- 473 regulations constrain vulture movements across country borders. *Biological*
474 *Conservation*, 219, 46-52.
- 475 6. Arrondo, E., Morales-Reyes, Z., Moleón, M., Cortés-Avizanda, A., Donázar, J. A.,
476 & Sánchez-Zapata, J. A. (2019). Rewilding traditional grazing areas affects
477 scavenger assemblages and carcass consumption patterns. *Basic and Applied*
478 *Ecology*, 41, 56-66.
- 479 7. Arrondo, E., Sanz-Aguilar, A., Pérez-García, J. M., Cortés-Avizanda, A., Sánchez-
480 Zapata, J. A., & Donázar, J. A. Landscape anthropization shapes the survival of a
481 top avian scavenger. *Biodiversity and Conservation*, 1-15.
- 482 8. Avery, D., and R. T. Watson. (2009). Regulation of lead-based ammunition around
483 the world. Pages 161-168 in *Ingestion of Lead from Spent Ammunition:*
484 *Implications for Wildlife and Humans* (R. T. Watson, M. Fuller, M. Pokras, and
485 W. G. Hunt, Eds.). The Peregrine Fund, Boise, Idaho
- 486 9. Berny, P., Vilagines, L., Cugnasse, J. M., Mastain, O., Chollet, J. Y., Joncour, G.,
487 & Razin, M. (2015). VIGILANCE POISON: Illegal poisoning and lead
488 intoxication are the main factors affecting avian scavenger survival in the Pyrenees
489 (France). *Ecotoxicology and Environmental Safety*, 118, 71-82.
- 490 10. Binkowski, Ł. J., Meissner, W., Trzeciak, M., Izevbekhai, K., & Barker, J. (2016).
491 Lead isotope ratio measurements as indicators for the source of lead poisoning in
492 Mute swans (*Cygnus olor*) wintering in Puck Bay (northern Poland).
493 *Chemosphere*, 164, 436-442.
- 494 11. Burnham, K.P. & Anderson, D.R., (2002). *Model Selection and Multimodel*
495 *Inference: A Practical Information-Theoretic Approach (2^a)*, *Ecological Modelling*.
496 Springer Science & Business Media. doi:10.1016/j.ecolmodel.2003.11.004

- 497 12. Calenge, C., & Fortmann-Roe, S. (2013). adehabitatHR: home range estimation. R
498 package version 0.4, 7.
- 499 13. Calvert JH (1876) Pheasant poisoning by swallowing shot. The Field 47 no. 1208,
500 Feb 19, p. 189
- 501 14. Church, M. E., Gwiazda, R., Risebrough, R. W., Sorenson, K., Chamberlain, C. P.,
502 Farry, S., Heinrich, W., Rideout, B.A. & Smith, D. R. (2006). Ammunition is the
503 principal source of lead accumulated by California condors re-introduced to the
504 wild. Environmental Science & Technology, 40(19), 6143-6150.
- 505 15. Cortés-Avizanda, A., R. Jovani, J. A. Donázar, and V. Grimm. 2014. Bird sky
506 networks: How do avian scavengers use social information to find carrion?
507 Ecology 95:1799–1808.
- 508 16. Cromie, R.L., A. Loram, L. Hurst, M. O'Brien, J. Newth, M.J. Brown, and J.P.
509 Harradine. 2010. Compliance with the Environmental Protection (Restrictions on
510 Use of Lead Shot) (England) Regulations 1999. DEFRA. Bristol, UK. Available
511 at: [http://randd.defra.gov.uk/Document.aspx?](http://randd.defra.gov.uk/Document.aspx?Document=WC0730_9719_FRP.pdf)
512 [Document=WC0730_9719_FRP.pdf](http://randd.defra.gov.uk/Document.aspx?Document=WC0730_9719_FRP.pdf).
- 513 17. Del Moral, J. C. y Molina, B. (Eds.) 2018. El buitre leonado en España, población
514 reproductora en 2018 y método de censo. SEO/BirdLife. Madrid.
- 515 18. Descalzo E, Mateo R (2018) La contaminación por munición de plomo en Europa:
516 el plumbismo aviar y las implicaciones en la seguridad de la carne de caza.
517 Instituto de Investigación en Recursos Cinegéticos (IREC), Ciudad Real, Spain. 82
518 pp. <http://doi.org/10.18239/aten.13.2018>
- 519 19. Espín, S., Martínez-López, E., Jiménez, P., María-Mojica, P., & García-Fernández,
520 A. J. (2014). Effects of heavy metals on biomarkers for oxidative stress in Griffon
521 vulture (*Gyps fulvus*). Environmental Research, 129, 59-68.

- 522 20. Espín, S., Martínez-López, E., Jiménez, P., María-Mojica, P., & García-Fernández,
523 A. J. (2015). Delta-aminolevulinic acid dehydratase (δ ALAD) activity in four free-
524 living bird species exposed to different levels of lead under natural conditions.
525 *Environmental Research*, 137, 185-198.
- 526 21. Fieberg, J., & Kochanny, C. O. (2005). Quantifying home-range overlap: the
527 importance of the utilization distribution. *The Journal of Wildlife Management*,
528 69(4), 1346-1359.
- 529 22. Finkelstein, M. E., Doak, D. F., George, D., Burnett, J., Brandt, J., Church, M.,
530 Grantham, J. & Smith, D. R. (2012). Lead poisoning and the deceptive recovery of
531 the critically endangered California condor. *Proceedings of the National Academy*
532 *of Sciences*, 109(28), 11449-11454.
- 533 23. Gangoso, L., Alvarez-Lloret, P., Rodríguez-Navarro, A. A., Mateo, R., Hiraldo, F.,
534 & Donazar, J. A. (2009). Long-term effects of lead poisoning on bone
535 mineralization in vultures exposed to ammunition sources. *Environmental*
536 *Pollution*, 157(2), 569-574.
- 537 24. García-Fernández, A. J., Martínez-López E., Romero D., María-Mojica P., Godino
538 A., & Jiménez P. (2005). High levels of blood lead in griffon vultures (*Gyps*
539 *fulvus*) from Cazorla Natural Park (Southern Spain). *Environmental Toxicology*
540 20(4): 459-463.
- 541 25. García-Fernández, A. J. (2014). *Ecotoxicology, Avian*. In: Wexler, P. (Ed.),
542 *Encyclopedia of Toxicology*, 3rd edition, vol 2. Elsevier Inc., Academic Press, pp.
543 289–294. ISBN: 9780123864543.
- 544 26. Garbett, R., Maude, G., Hancock, P., Kenny, D., Reading, R., & Amar, A. (2018).
545 Association between hunting and elevated blood lead levels in the critically

- 546 endangered African white-backed vulture *Gyps africanus*. *Science of the Total*
547 *Environment*, 630, 1654-1665.
- 548 27. Gochfeld, J. B. M. (2000). Effects of lead on birds (Laridae): a review of
549 laboratory and field studies. *Journal of Toxicology and Environmental Health Part*
550 *B: Critical Reviews*, 3(2), 59-78.
- 551 28. Golden, N. H., Warner, S. E., & Coffey, M. J. (2016). A review and assessment of
552 spent lead ammunition and its exposure and effects to scavenging birds in the
553 United States. In *Reviews of Environmental Contamination and Toxicology*
554 *Volume 237* (pp. 123-191). Springer, Cham.
- 555 29. González, F., López, I., Suárez, L., Moraleda, V., & Rodríguez, C. (2017). Levels
556 of blood lead in Griffon vultures from a Wildlife Rehabilitation Center in Spain.
557 *Ecotoxicology and Environmental Safety*, 143, 143-150.
- 558 30. Hernández, M., & Margalida, A. (2009). Poison-related mortality effects in the
559 endangered Egyptian vulture (*Neophron percnopterus*) population in Spain.
560 *European Journal of Wildlife Research*, 55(4), 415-423.
- 561 31. Herruzo, A. C., & Martínez-Jauregui, M. (2013). Trends in hunters, hunting
562 grounds and big game harvest in Spain. *Forest Systems*, 22(1), 114-122.
- 563 32. Horowitz, I. H., Yanco, E., Nadler, R. V., Anglister, N., Landau, S., Elias, R.,
564 Lublin, A., Perl, S., Edery, N. & Rosenzweig, A. B. (2014). Acute lead poisoning
565 in a griffon vulture (*Gyps fulvus*) in Israel. *Israel Journal of Veterinary Medicine*,
566 69(3), 163-168.
- 567 33. Hunt, W. G., Burnham, W., Parish, C. N., Burnham, K. K., Mutch, B. & Oaks, J.
568 L. (2006). Bullet fragments in deer remains: implications for lead exposure in
569 avian scavengers. *Wildlife Society Bulletin*, 34(1), 167-170.

- 570 34. Hunt, W. G., Watson, R. T., Oaks, J. L., Parish, C. N., Burnham, K. K., Tucker, R.
571 L., Belthoff, J.R. & Hart, G. (2009). Lead bullet fragments in venison from rifle-
572 killed deer: potential for human dietary exposure. PLoS One, 4(4), e5330.
- 573 35. Kanstrup, N., Swift, J., Stroud, D. A., & Lewis, M. (2018). Hunting with lead
574 ammunition is not sustainable: European perspectives. *Ambio*, 47(8), 846-857.
- 575 36. Katzner, T. E., Stuber, M. J., Slabe, V. A., Anderson, J. T., Cooper, J. L., Rhea, L.
576 L., & Millsap, B. A. (2018). Origins of lead in populations of raptors. *Animal*
577 *Conservation*, 21(3), 232-240.
- 578 37. Knott J., Gilbert J, Hoccom D.G. & Green RE (2010) Implications for wildlife and
579 humans of dietary exposure to lead from fragments of lead rifle bullets in deer shot
580 in the UK. *Science of the Total Environment* 409:95–99.
- 581 38. Krone, O. (2018). Lead Poisoning in Birds of Prey. In *Birds of Prey* (pp. 251-272).
582 Springer, Cham.
- 583 39. Krüger, S. C., & Amar, A. (2018). Lead Exposure in the Critically Endangered
584 Bearded Vulture (*Gypaetus barbatus*) Population in Southern Africa. *Journal of*
585 *Raptor Research*, 52(4), 491-500.
- 586 40. Lambertucci, S. A., Alarcón, P. A., Hiraldo, F., Sánchez-Zapata, J. A., Blanco, G.,
587 & Donazar, J. A. (2014). Apex scavenger movements call for transboundary
588 conservation policies. *Biological Conservation*, 170, 145-150.
- 589 41. Lecina, S., Playán, E., Isidoro, D., Dechmi, F., Causapé, J., & Faci, J. M. (2005).
590 Irrigation evaluation and simulation at the Irrigation District V of Bardenas
591 (Spain). *Agricultural Water Management*, 73(3), 223-245.
- 592 42. Legagneux, P., Suffice, P., Messier, J. S., Lelievre, F., Tremblay, J. A.,
593 Maisonneuve, C., Saint-Louis, R. & Bêty, J. (2014). High risk of lead

- 594 contamination for scavengers in an area with high moose hunting success. PLoS
595 One, 9(11), e111546.
- 596 43. Locutura, J., Bel-lan A., García-Cortés, A. & Martínez, S. (2012) Atlas
597 Geoquímico de España. Publicado por el IGME (Instituto Geológico y Minero de
598 España) ISBN: 978-84-7840-875-7. 592 pag. Madrid (España)
599 [http://www.igme.es/actividadesIGME/lineas/CartoGeo/geoquimica/geoquimicaEs](http://www.igme.es/actividadesIGME/lineas/CartoGeo/geoquimica/geoquimicaEsp.htm)
600 [p.htm](http://www.igme.es/actividadesIGME/lineas/CartoGeo/geoquimica/geoquimicaEsp.htm)
- 601 44. Longman, J., Veres, D., Ersek, V., Phillips, D. L., Chauvel, C., & Tamas, C. G.
602 (2018). Quantitative assessment of Pb sources in isotopic mixtures using a
603 Bayesian mixing model. *Scientific Reports*, 8(1), 6154.
- 604 45. Margalida, A., Donázar, J. A., Carrete, M., & Sánchez-Zapata, J. A. (2010).
605 Sanitary versus environmental policies: fitting together two pieces of the puzzle of
606 European vulture conservation. *Journal of Applied Ecology*, 47(4), 931-935.
- 607 46. Margalida, A., Colomer, M. À., & Sanuy, D. (2011). Can wild ungulate carcasses
608 provide enough biomass to maintain avian scavenger populations? An empirical
609 assessment using a bio-inspired computational model. *PLoS One*, 6(5), e20248.
- 610 47. Margalida, A., Pérez-García, J. M., Afonso, I., & Moreno-Opo, R. (2016). Spatial
611 and temporal movements in Pyrenean bearded vultures (*Gypaetus barbatus*):
612 Integrating movement ecology into conservation practice. *Scientific reports*, 6,
613 35746.
- 614 48. Martín-Queller, E., Moreno-Mateos, D., Pedrocchi, C., Cervantes, J., & Martínez,
615 G. (2010). Impacts of intensive agricultural irrigation and livestock farming on a
616 semi-arid Mediterranean catchment. *Environmental Monitoring and Assessment*,
617 167(1-4), 423-435.

- 618 49. Martinez-Haro, M., Taggart, M. A., Martín-Doimeadios, R.C. R., Green, A. J., &
619 Mateo, R. (2011). Identifying sources of Pb exposure in waterbirds and effects on
620 porphyrin metabolism using noninvasive fecal sampling. *Environmental Science &*
621 *Technology*, 45(14), 6153-6159.
- 622 50. Mateo, R., Molina, R., Grífols, J., & Guitart, R. (1997). Lead poisoning in a free
623 ranging griffon vulture (*Gyps fulvus*). *The Veterinary Record*, 140, 47-48.
- 624 51. Mateo, R. (2009). Lead poisoning in wild birds in Europe and the regulations
625 adopted by different countries. Pages 78–91 in *Ingestion of Lead from Spent*
626 *Ammunition: Implications for Wildlife and Humans* (Watson, R. T. , Fuller M. ,
627 Pokras M. , and Hunt W. G. , Eds.). The Peregrine Fund, Boise, Idaho.
- 628 52. Mateo, R., & Kanstrup, N. (2019). Regulations on lead ammunition adopted in
629 Europe and evidence of compliance. *Ambio*, 48, 989-998.
- 630 53. Mateo-Tomás, P., Olea, P. P., Jiménez-Moreno, M., Camarero, P. R., Sánchez-
631 Barbudo, I. S., Martín-Doimeadios, R. C. R., & Mateo, R. (2016). Mapping the
632 spatio-temporal risk of lead exposure in apex species for more effective mitigation.
633 *Proceedings of the Royal Society B*, 283(1835), 20160662.
- 634 54. Morales-Reyes, Z., Pérez-García, J. M., Moleón, M., Botella, F., Carrete, M.,
635 Donázar, J. A., Cortés-Avizanda, A., Arrondo, E., Moreno-Opo, R., Jiménez, J.,
636 Margalida, A., & Sánchez-Zapata, J. A. (2017). Evaluation of the network of
637 protection areas for the feeding of scavengers in Spain: from biodiversity
638 conservation to greenhouse gas emission savings. *Journal of Applied Ecology*, 54
639 (4), 1120–1129.
- 640 55. Nadjafzadeh, M., Hofer, H., & Krone, O. (2015). Lead exposure and food
641 processing in white-tailed eagles and other scavengers: an experimental approach

- 642 to simulate lead uptake at shot mammalian carcasses. *European Journal of Wildlife*
643 *Research*, 61(5), 763-774.
- 644 56. Naidoo, V., Wolter, K., Espie, I., & Kotze, A. (2012). Lead toxicity: consequences
645 and interventions in an intensively managed (*Gyps coprotheres*) vulture colony.
646 *Journal of Zoo and Wildlife Medicine*, 43(3), 573-578.
- 647 57. Naidoo, V., Wolter, K., & Botha, C. J. (2017). Lead ingestion as a potential
648 contributing factor to the decline in vulture populations in Southern Africa.
649 *Environmental Research*, 152, 150-156.
- 650 58. Núñez, O., Fernández-Navarro, P., Martín-Méndez, I., Bel-Lan, A., Rupérez, J. F.
651 L., & López-Abente, G. (2017). Association between heavy metal and metalloid
652 levels in topsoil and cancer mortality in Spain. *Environmental Science and*
653 *Pollution Research*, 24(8), 7413-7421.
- 654 59. Monna, F., Galop, D., Carozza, L., Tual, M., Beyrie, A., Marembert, F., Chateu, C.
655 & Grousset, F. E. (2004). Environmental impact of early Basque mining and
656 smelting recorded in a high ash minerogenic peat deposit. *Science of the Total*
657 *Environment*, 327(1-3), 197-214.
- 658 60. Pain DJ, Fisher IJ, Thomas VG (2009) A global update of lead poisoning in
659 terrestrial birds from ammunition sources. In: Watson RT, Fuller M, Pokras M,
660 Hunt G, editors. *Ingestion of Lead from Spent Ammunition: Implications for*
661 *Wildlife and Humans*. pp. 99–118.
- 662 61. Pain, D.J, Mateo, R., & Green, R. E. (2019). Effects of lead from ammunition on
663 birds and other wildlife: A review and update. *Ambio*, 1-19.
664 <https://doi.org/10.1007/s13280-019-01159-0>

- 665 62. Papanikolaou, N. C., Hatzidaki, E. G., Belivanis, S., Tzanakakis, G. N., &
666 Tsatsakis, A. M. (2005). Lead toxicity update. A brief review. *Medical Science*
667 *Monitor*, 11(10), RA329-RA336.
- 668 63. Pareja-Carrera, J., Mateo, R., & Rodríguez-Estival, J. (2014). Lead (Pb) in sheep
669 exposed to mining pollution: Implications for animal and human health.
670 *Ecotoxicology and Environmental Safety*, 108, 210-216.
- 671 64. Parnell, A. C., Inger, R., Bearhop, S., & Jackson, A. L. (2010). Source partitioning
672 using stable isotopes: coping with too much variation. *PloS One*, 5(3), e9672.
- 673 65. Phillips, D. L., Inger, R., Bearhop, S., Jackson, A. L., Moore, J. W., Parnell, A. C.
674 & Ward, E. J. (2014). Best practices for use of stable isotope mixing models in
675 food-web studies. *Canadian Journal of Zoology*, 92(10), 823-835. doi:10.1139/cjz-
676 2014-0127
- 677 66. Plaza, P.I., & Lambertucci, S. A. (2019). What do we know about lead
678 contamination in wild vultures and condors? A review of decades of research.
679 *Science of the Total Environment*, 654, 409-417.
- 680 67. R Core Team (2018). R: A language and environment for statistical
681 computing. R Foundation for Statistical Computing, Vienna, Austria.
682 URL<https://www.R-project.org/>.
- 683 68. Reglero, M. M., Monsalve-González, L., Taggart, M. A., & Mateo, R. (2008).
684 Transfer of metals to plants and red deer in an old lead mining area in Spain.
685 *Science of the Total Environment*, 406(1-2), 287-297.
- 686 69. Reglero, M. M., Taggart, M. A., Monsalve-Gonzalez, L., & Mateo, R. (2009).
687 Heavy metal exposure in large game from a lead mining area: effects on oxidative
688 stress and fatty acid composition in liver. *Environmental Pollution*, 157(4), 1388-
689 1395.

- 690 70. Rodríguez-Estival, J., Álvarez-Lloret, P., Rodríguez-Navarro, A. B., & Mateo, R.
691 (2013). Chronic effects of lead (Pb) on bone properties in red deer and wild boar:
692 relationship with vitamins a and D3. *Environmental Pollution*, 174, 142-149.
- 693 71. Rodríguez-Estival, J., Pareja-Carrera, J., Mateo, R., 2014. Lead poisoning in a calf
694 from the mining area of Sierra Madrona and Alcudia Valley. *Revista de*
695 *Toxicología*, 31, 47-49.
- 696 72. Santiago., Motas-Guzmán, M., Reja, A., María-Mojica, P, Rodero, B., & García-
697 Fernández, A. J. (1998). Lead and Cadmium in Red Deer and Wild Boar from
698 Sierra Morena Mountains (Andalusia, Spain). *Bulletin of Environmental*
699 *Contamination & Toxicology* 61: 730-737
- 700 73. Stock, C., & Semmens, B. X. (2016). Unifying error structures in commonly used
701 biotracer mixing models. *Ecology*, 97(10), 2562-2569.
- 702 74. Stock, B. C., Jackson, A. L., Ward, E. J., Parnell, A. C., Phillips, D. L., &
703 Semmens, B. X. (2018). Analyzing mixing systems using a new generation of
704 Bayesian tracer mixing models. *PeerJ*, 6, e5096.
- 705 75. Taggart, M. A., Reglero, M. M., Camarero, P. R., & Mateo, R. (2011). Should
706 legislation regarding maximum Pb and Cd levels in human food also cover large
707 game meat?. *Environment international*, 37(1), 18-25.
- 708 76. Vallverdú-Coll, N., Mougeot, F., Ortiz-Santaliestra, M. E., Rodríguez-Estival, J.,
709 López-Antia, A., & Mateo, R. (2016). Lead exposure reduces carotenoid-based
710 coloration and constitutive immunity in wild mallards. *Environmental Toxicology*
711 *and Chemistry*, 35(6), 1516-1525.
- 712 77. Wink, M., Sauer-Gürth, H., Martinez, F., Doval, G., Blanco, G., & Hatzofe, O.
713 (1998). The use of (GACA) 4 PCR to sex Old World vultures (Aves:
714 *Accipitridae*). *Molecular Ecology*, 7(6), 779-782.

716 **Table 1:** Number and percentage of individuals from the two study areas in each of the
717 categories of lead exposure defined by Pain et al. (2019).

718

719 **Table 2:** Results of the Generalized Linear Models (Gaussian family) performed to
720 determine sources of blood lead concentration in GPS-tagged vultures.

721

722 **Table S.1:** Individual characteristics of the birds included in this study. In column Sex
723 are represented males (M) and females (F). Areas of KDE50 and KDE95 are expressed
724 in km². Column Alive indicates if the animals were alive at the end of the study period
725 (December 2018). NA represents those birds whose GPS device failed.

726 **Table S.2:** Ammunition and topsoil (control and contaminated) samples used to
727 determine proportion of both lead sources in vultures blood.

728 **Table S.3:** Mean and range values of %RSD for replicate analyses (n=6) of vulture
729 blood, topsoil, and ammunition samples.

730 **Table S.4:** Statistical summary of the layers used to estimate ammunition and topsoil
731 lead exposure.

732 **Table S.5:** AIC-based model selection to assess the lead concentration in vultures. Only
733 models with informative variables are included.

734

735 **Figure 1:** Upper and lower panels show the KDE95 used by all the individuals from
736 northern (blue contour) and southern populations (red contour). Left maps represent the
737 lead concentration in the superficial topsoil (according to Locutura et al., 2012). Right
738 panels show the number of animals hunted per year in 10x10 km² cells including the
739 two species most commonly hunted, wild boar and red deer (see methods). Black stars
740 show trapping sites.

741 **Figure 2:** Lead isotope ratios (A: $^{207}\text{Pb}/^{208}\text{Pb}$ - $^{206}\text{Pb}/^{207}\text{Pb}$; B: $^{206}\text{Pb}/^{208}\text{Pb}$ - $^{206}\text{Pb}/^{207}\text{Pb}$; C:
742 $^{206}\text{Pb}/^{208}\text{Pb}$ - $^{207}\text{Pb}/^{208}\text{Pb}$) in blood of griffon vultures from northern and southern
743 populations. Red and blue dots represent southern and northern population individuals,
744 respectively. Mean and standard deviation of lead isotope ratios of the two lead sources
745 (ammunition and topsoil) are also shown.

746 **Figure 3:** Mean and 95% Confidence Interval of the estimated contribution of lead from
747 ammunition and topsoil to blood lead concentration in vultures from both northern and
748 southern populations, based on the results of the MixSIAR models.

749 Table 1

750

Population	Sex	Blood Pb concentration ($\mu\text{g}/\text{dl}$)			
		<20 Background	20-50 Sublethal effects	50-100 Clinical effects	>100 Potentially lethal
Northern	Female	2(12.5)	8(50.0)	5(31.3)	1(6.3)
	Male	4(28.6)	6(42.9)	2(14.3)	0(0.0)
	Total	6(21.4)	14(50.0)	7(25.0)	1(3.6)
Southern	Female	0(0.0)	3(27.3)	5(45.5)	3(27.3)
	Male	2(10.5)	5(26.3)	11(57.9)	1(5.3)
	Total	2(6.7)	8(26.7)	16(53.3)	4(13.3)
Both	Female	2(7.4)	11(40.7)	10(37.0)	4(14.8)
	Male	6(18.2)	11(33.3)	13(39.4)	1(3.0)
	Total	8(13.8)	22(37.9)	23(39.7)	5(8.6)

751

752

753

754

755

756

757

758

759

760

761

762

763

764 Table 2

Variables	Estimate±Std. Error	p-value
(Intercept)	3.438±0.159	<0.001
exposure to ammunition from big game hunting at KDE 50	0.004±0.001	<0.001
males	-0.337±0.153	0.003
(Intercept)	3.438±0.167	<0.001
exposure to ammunition from big game hunting at KDE 95	0.004±0.001	<0.001
males	-0.327±0.155	0.004
(Intercept)	3.141±0.196	<0.001
exposure to topsoil lead at KDE95	0.022±0.006	<0.001

765

766

767

36

768

769

770 Table S.1.

771

Individual	Alive	Population	Sex	KDE50	KDE95
L73	YES	Southern	M	70.88	1124.7916
L8J	YES	Southern	M	243.41	4668.9676
T00	YES	Southern	F	207.61	3426.2532
T01	YES	Southern	F	318.63	5675.5555
T02	YES	Southern	M	172.82	2492.7401
T03	YES	Southern	M	519.83	6090.678
T05	YES	Southern	M	78.10	5457.2433
T06	NA	Southern	M	84.64	2768.1997
T07	NO	Southern	M	59.07	2666.8988
T08	YES	Southern	F	196.74	3326.8942
T09	YES	Southern	M	81.93	4137.2157
T0A	YES	Southern	M	334.30	2798.9276
T0C	YES	Southern	M	105.33	4678.9345
T0H	YES	Southern	M	307.49	3596.8582
T0J	YES	Southern	F	964.21	8335.799
T0L	NA	Southern	M	55.26	3110.0696
T0U	NO	Southern	F	187.80	3960.7735
T0V	YES	Southern	M	231.85	2922.7373
T0W	NA	Southern	F	57.60	2701.8798
T0X	NA	Southern	M	164.98	5371.7382
T10	NA	Southern	F	464.65	5438.5837
T11	YES	Southern	M	51.72	5323.4727

T12	NA	Southern	M	83.62	2455.8705
T14	YES	Southern	M	226.10	4117.9899
T15	YES	Southern	F	780.49	8514.226
T16	YES	Southern	M	92.43	1898.6661
T17	YES	Southern	M	247.02	7765.4193
T19	NO	Southern	F	309.69	3856.2093
T1C	NA	Southern	F	462.86	3382.1542
T1J	YES	Southern	F	780.26	7626.0972
T1L	YES	Northern	M	57.71	1336.3249
T1N	YES	Northern	F	86.82	871.3995
T1R	YES	Northern	M	205.56	2111.6737
T1U	NO	Northern	F	134.95	891.8856
T1W	NO	Northern	M	46.04	1760.9609
T1X	YES	Northern	F	209.66	2653.92
T21	YES	Northern	M	38.42	543.4831
T22	NO	Northern	M	25.55	233.7521
T24	YES	Northern	F	88.19	1029.8596
T25	YES	Northern	M	698.48	7775.0831
T2C	NO	Northern	M	88.75	995.9196
T2F	YES	Northern	F	73.39	791.1746
T2H	YES	Northern	M	79.03	678.0347
T2L	YES	Northern	F	191.61	3106.063
T2M	YES	Northern	M	142.04	1890.6208
T2N	NO	Northern	M	34.25	536.0871
T2R	YES	Northern	F	738.00	7057.0544
T2T	YES	Northern	F	128.09	908.3614
T2U	YES	Northern	M	210.53	2475.1838
T2V	YES	Northern	F	152.63	2338.3821
T2W	YES	Northern	F	302.94	4039.008

T2X	NO	Northern	F	54.22	1311.9559
T30	YES	Northern	F	66.13	1967.6696
T31	YES	Northern	F	362.49	5181.7917
T33	YES	Northern	F	271.52	2665.2463
T35	NO	Northern	F	132.29	2478.3662
T36	YES	Northern	M	20.05	142.9229
T3T	NO	Northern	F	71.20	518.6238

772

773

774 Table S.2.

775

ID	N	Kind of lead	206/207	207/208	206/208
Topsoil T05_10 A	1	Control topsoil	1.1934	0.4004	0.4778
Topsoil T05_10 B	1	Control topsoil	1.1945	0.3996	0.4773
Topsoil T05_10 C	1	Control topsoil	1.1711	0.4059	0.4754
Topsoil T05_9	1	Control topsoil	1.1931	0.4015	0.4790
Topsoil NM2	1	Contaminated topsoil	1.1711	0.4073	0.4770
Topsoil Pto 122 B	1	Contaminated topsoil	1.1626	0.4106	0.4774
Topsoil Pto 123	1	Contaminated topsoil	1.1580	0.4097	0.4744
Topsoil Pto 124	1	Contaminated topsoil	1.1536	0.4112	0.4743
Topsoil Pto 125	1	Contaminated topsoil	1.1542	0.4098	0.4730
Topsoil Pto 133	1	Contaminated topsoil	1.1549	0.4103	0.4738
Topsoil Pto 74	1	Contaminated topsoil	1.1526	0.4119	0.4748
Norma 3006 A	3	Bullet	1.1455	0.4141	0.4743
Norma 3006 B	3	Bullet	1.1544	0.4130	0.4768
S&B 3006	2	Bullet	1.1473	0.4118	0.4724

REMINGTON 270	1	Bullet	1.2323	0.4051	0.4992
REMINGTON 300 A	2	Bullet	1.2241	0.4058	0.4968
REMINGTON 300 B	3	Bullet	1.2003	0.4087	0.4905
WINCHESTER 270 WSM	2	Bullet	1.2000	0.4091	0.4910
HORNADY 300	2	Bullet	1.2018	0.4094	0.4920
BROWING 12	3	Cartridge	1.1529	0.4132	0.4764

776

777

778

779

780 Table S.3

Sample	206/207	207/208	206/208
Vulture blood	0.124 (0.029-0.229)	0.147 (0.035-0.279)	0.140 (0.004-0.271)
Topsoil	0.140 (0.080-0.245)	0.134 (0.049-0.231)	0.127 (0.057-0.205)
Ammunition	0.138 (0.097-0.183)	0.138 (0.064-0.277)	0.119 (0.067-0.193)

781

782

783

784

785

786 Table S.4

Variables	Minimum	Maximum	Mean	Standard Deviation
Hunting exposure	0	320	70.07	59.90
Topsoil lead exposure	1	8545	27.22	54.32

787


788

789 Table S.5

Model	AICc	Δ AICc	Cumulative weight	Weight	R ²
exposure to ammunition from big game hunting at KDE 50+sex	104.77	0	0.44	0.44	23.8
exposure to ammunition from big game hunting at KDE 95+sex	106.44	1.67	0.19	0.63	21.9
exposure to topsoil lead at KDE95	106.63	1.86	0.17	0.8	18.1
exposure to ammunition from big game hunting at KDE 50	107.36	2.59	0.12	0.92	
exposure to ammunition from big game hunting at KDE 95+sex	108.66	3.89	0.06	0.98	
exposure to topsoil lead at KDE50	111.53	6.76	0.01	1	
NULL	116.02	11.25	0	1	

790


791 Figure 1



792

793

794 Figure 2



795

796

797

798

799

800

801

802

803

804

805


806

807

808

809 Figure 3

810



811