

Education as important predictor for successful employment in adults with congenital heart disease worldwide

Maayke A. Sluman MD, PhD^{1,2}  | Silke Apers PhD^{3,4} | Judith K. Sluiter PhD^{1*} | Karen Nieuwenhuijsen PhD¹ | Philip Moons PhD^{4,5} | Koen Luyckx PhD^{6,7} | Adrienne H. Kovacs PhD^{8,9} | Corina Thomet MSc¹⁰ | Werner Budts MD, PhD¹¹ | Junko Enomoto PhD¹² | Hsiao-Ling Yang PhD¹³ | Jamie L. Jackson PhD¹⁴  | Paul Khairy MD, PhD¹⁵ | Stephen C. Cook MD¹⁶ | Raghavan Subramanian MD¹⁷ | Luis Alday MD¹⁸ | Katrine Eriksen MSc¹⁹ | Mikael Dellborg MD, PhD^{20,21} | Malin Berghammer PhD^{5,22} | Eva Mattsson MD, PhD²³ | Andrew S. Mackie MD²⁴ | Samuel Menahem MD²⁵ | Maryanne Caruana MD, PhD²⁶  | Kathy Gosney MSSW²⁷ | Alexandra Soufi MD²⁸ | Susan M. Fernandes PA²⁹ | Kamila S. White PhD³⁰ | Edward Callus PhD³¹ | Shelby Kutty MD, PhD³² | Berto J. Bouma MD, PhD³³ | Barbara J.M. Mulder MD, PhD³³ on behalf of the APPROACH-IS consortium, the International Society for Adult Congenital Heart Disease (ISACHD)

¹Coronel Institute of Occupational Health, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands

²Department of Cardiology, Jeroen Bosch Hospital, 's Hertogenbosch, The Netherlands

³Department of Development and Regeneration, KU Leuven, University of Leuven, Leuven, Belgium

⁴Department of Public Health and Primary Care, KU Leuven, University of Leuven, Leuven, Belgium

⁵Center for Person-Centered Care (GPCC), University of Gothenburg, Gothenburg, Sweden

⁶School Psychology and Development in Context, KU Leuven, University of Leuven, Leuven, Belgium

⁷Department of Psychology, UNIBS, University of the Free State, Bloemfontein, South Africa

⁸Department of Psychology, University Health Network, Toronto, Canada

⁹The Knight Cardiovascular Institute, Oregon Health & Science University, Portland, Oregon

¹⁰Center for Congenital Heart Disease, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

¹¹Congenital and Structural Cardiology, Department of Cardiovascular Sciences, University Hospitals of Leuven, KU Leuven, Leuven, Belgium

¹²Department of Adult Congenital Heart Disease, Chiba Cardiovascular Center, Chiba, Japan

¹³School of Nursing, College of Medicine, National Taiwan University, Taipei, Taiwan

¹⁴Center for Biobehavioral Health, Nationwide Children's Hospital, Columbus, Ohio

¹⁵Montreal Heart Institute, University of Montreal, Montreal, Canada

¹⁶Adult Congenital Heart Disease Center, Helen DeVos Children's Hospital, Grand Rapids, Michigan

¹⁷Frontier Lifeline Hospital, Dr. K. M. Cherian Heart Foundation, Chennai, India

¹⁸Division of Cardiology, Hospital de Niños, Córdoba, Argentina

¹⁹Department of Cardiology, Oslo University Hospital, Oslo, Norway

²⁰Institute of Medicine, Sahlgrenska Academy, Gothenburg University, Gothenburg, Sweden

*Judith K. Sluiter passed away in May 2018 (after this manuscript was finished).

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors. *Congenital Heart Disease* Published by Wiley Periodicals, Inc.

²¹Adult Congenital Heart Unit, Sahlgrenska University Hospital/Östra, Gothenburg, Sweden

²²Department of Health Sciences, University West, Trollhättan, Sweden

²³Department of Cardiology, Karolinska University Hospital, Stockholm, Sweden

²⁴Department of Pediatric Cardiology, University of Alberta, Edmonton, Canada

²⁵Monash Medical Center, Melbourne, Australia

²⁶Department of Cardiology, Mater Dei Hospital, Msida, Malta

²⁷Adult Congenital Heart Disease Center, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio

²⁸Hospital Louis Pradel, Lyon, France

²⁹Adult Congenital Heart Disease Program at Stanford, Lucile Packard Children's Hospital and Stanford Health Care, Palo Alto, California

³⁰Adult Congenital Heart Disease Center, Washington University and Barnes Jewish Heart & Vascular Center, University of Missouri, Saint Louis, Missouri

³¹Clinical Psychology Service, IRCCS Policlinico San Donato Hospital, Milan, Italy

³²Adult Congenital Heart Disease Center, University of Nebraska Medical Center/Children's Hospital & Medical Center, Omaha, Nebraska

³³Amsterdam UMC, Department of Cardiology, University of Amsterdam, Amsterdam, The Netherlands

Correspondence

Maayke Sluman, Coronal Institute of Occupational Health, Amsterdam UMC, University of Amsterdam, The Netherlands.
Email: m.a.sluman@amc.uva.nl

Abstract

Background: Conflicting results have been reported regarding employment status and work ability in adults with congenital heart disease (CHD). Since this is an important determinant for quality of life, we assessed this in a large international adult CHD cohort.

Methods: Data from 4028 adults with CHD (53% women) from 15 different countries were collected by a uniform survey in the cross-sectional APPROACH International Study. Predictors for employment and work limitations were studied using general linear mixed models.

Results: Median age was 32 years (IQR 25-42) and 94% of patients had at least a high school degree. Overall employment rate was 69%, but varied substantially among countries. Higher education (OR 1.99-3.69) and having a partner (OR 1.72) were associated with more employment; female sex (OR 0.66, worse NYHA functional class (OR 0.67-0.13), and a history of congestive heart failure (OR 0.74) were associated with less employment. Limitations at work were reported in 34% and were associated with female sex (OR 1.36), increasing age (OR 1.03 per year), more severe CHD (OR 1.31-2.10), and a history of congestive heart failure (OR 1.57) or mental disorders (OR 2.26). Only a university degree was associated with fewer limitations at work (OR 0.62).

Conclusions: There are genuine differences in the impact of CHD on employment status in different countries. Although the majority of adult CHD patients are employed, limitations at work are common. Education appears to be the main predictor for successful employment and should therefore be encouraged in patients with CHD.

KEYWORDS

adult, congenital heart defects, disability, education, employment, work ability

1 | INTRODUCTION

With survival rates to adulthood over 90%, most children with congenital heart disease (CHD) go to school and subsequently obtain employment.^{1,2} Employment and work ability contribute to well-being and are nowadays crucial in daily life for most adults with CHD.³ Employment has been reported as one of the most

important aspects of quality of life (QOL) and, inversely, lack of employment with lower QOL.^{4,5} Aspects influencing work such as concentration and fatigue, are important patient-reported outcomes that receive increasing attention. However, in contrast to previous guidelines,⁶ education and employment are no longer mentioned in the latest European guidelines for the management of grown-up CHD.⁷

Higher rates of unemployment, disease-related work absences, and limitations at work have been reported in adults with CHD compared to the general population in The Netherlands, Germany, Denmark, the United Kingdom, and the United States.⁸⁻¹³ In contrast, studies from Finland, Sweden, and Malta, have reported employment rates similar or even above those of the general population.¹⁴⁻¹⁶ Research in other chronic diseases (eg, rheumatoid arthritis, asthma, chronic obstructive pulmonary disease, diabetes mellitus, and ischemic heart disease) has consistently shown that employment and work ability are negatively influenced by these conditions. Patients with other chronic diseases are less often employed, work fewer hours when they work and report limitations at work.¹⁷ These findings were associated with older age, female sex, perceived health complaints, and limitations in daily physical activities caused by the disease.¹⁸ It remains unknown whether these contributing factors for work disability are applicable to adults with CHD.

The conflicting findings on employment and limitations at work and contributing factors in adults with CHD, suggest that they might be influenced by other factors than solely the CHD itself. Therefore, the aim of this study was to explore employment, work ability, and the presence of limitations at work in a large international adult CHD cohort in a uniform way, to investigate differences between countries and to identify predictors for employment and work limitations.

2 | METHODS

2.1 | Study design and setting

Data were collected from April 2013 to March 2015 in the "Assessment of Patterns of Patient-Reported Outcomes in Adults with Congenital Heart disease—International Study" (APPROACH-IS). APPROACH-IS is an international cross-sectional multicenter study conducted in partnership with the International Society for Adult Congenital Heart Disease (ISACHD). The rationale and methodology of this study have been previously described.¹⁹ In summary, inclusion criteria were age 18 years or above, known CHD with continuing follow-up and the ability to complete self-report questionnaires. The study was approved by the institutional review board of the coordinating center (University Hospitals Leuven/KU Leuven, Belgium) and local institutional review boards when required. All participating patients provided written informed consent. Data from 5 different continents were collected in 15 participating countries: Belgium, France, Italy, Malta, Norway, Sweden, Switzerland, and The Netherlands (Europe); Canada and the United States (North America); India, Japan, and Taiwan (Asia); Argentina (South America); and Australia.

2.2 | Variables and measurements

A uniform survey developed by the research team consisting of questions on demographic, medical and QOL items was sent to either a random or consecutively approached selection of patients at each participating center. The first question of the Work Ability

Index, inquiring about current work ability compared to lifetime best, was used to assess work ability. This question is considered a valid and reliable predictor of work disability and is frequently used to measure work ability.^{20,21} Patients were asked to rate their current work ability on a scale from 0 (not being able to work at all) to 10 (equivalent of lifetime best work ability). A Work Ability Score (WAS) of 8 or higher was considered good to excellent, a score of 6 or 7 was considered moderate, and any score of 5 or below was considered poor. Limitations at work were collected through several questions on experiencing limitations at work attributed to the CHD, eg, having symptoms at work, having to slow down due to the symptoms, or having to work part-time because of the disease. Country-specific data on employment and unemployment were drawn from the International Monetary Fund (IMF) and the Organisation for Economic Co-operation and Development (OECD) and used as reference data.^{22,23}

2.3 | Definitions

CHD was diagnosed according to the European Paediatric Cardiac Code Short List coding scheme.²⁴ To categorize complexity of CHD, a prespecified hierarchical scheme was used, classifying type of CHD as either mild, moderate, or severe.²⁵ Employment status was classified by the patient as either full- or part-time, unemployed (including job seeking), disabled/government financial assistance, homemaker, student, retired, or other. From all patients that had chosen "other," additional free text was individually verified. For all analyses regarding work ability, only patients with working experiences were analyzed. Therefore, since most of the participating countries have a retirement age of 65 years and older, patients were censored from that age and homemakers, early retired participants, and students were all excluded from work ability analyses. Unemployed or disabled patients who considered themselves entirely unable to work were also excluded from all analyses on working experiences. Limitations at work were considered when "yes" was answered on one or more questions regarding work limitations. Only employed patients were included for the analyses on work limitations.

2.4 | Statistical methods

Categorical data were presented as numbers and percentages and continuous data as medians with interquartile ranges (IQR) since they were not normally distributed. Clinical characteristics were compared using chi-square statistics for dichotomous and categorical variables and Mann-Whitney *U* tests for continuous variables. The differences in parameters for employment and work limitations were first tested by univariate analyses. The association of patient-specific characteristics of employment and work limitations was estimated with the use of general linear mixed models. By this multilevel structure, patients were nested within countries, to account for as much influence from cultural differences as possible. Odds

TABLE 1 Characteristics of the study population

| Variable | Respondents | APPROACH-IS (N = 4028) | | | P value |
|-----------------------------------|-------------|------------------------|-----------------------|--------------------------------|---------|
| | | Men (N = 1897 (47)) | Women (N = 2115 (53)) | Total (N = 4012 ^a) | |
| Age, y (IQR) | 4013 | 31 (24-43) | 32 (25-41) | 32 (25-42) | .649 |
| Origin, N (%) ^b | 4028 | | | | |
| Europe | | 869 (46) | 866 (41) | 1 735 (43) | .003 |
| North America | | 560 (29) | 700 (33) | 1 260 (32) | .016 |
| Asia | | 330 (17) | 377 (18) | 707 (18) | .761 |
| South America | | 68 (4) | 110 (5) | 178 (4) | .018 |
| Australia | | 70 (4) | 62 (3) | 132 (3) | .177 |
| Marital status | 4008 | | | | |
| Never married | | 911 (48) | 839 (40) | 1750 (44) | <.001 |
| Married/living with partner | | 901 (48) | 1134 (54) | 2035 (51) | <.001 |
| Divorced/widowed | | 73 (4) | 129 (6) | 202 (5) | .001 |
| Having children | 4004 | 696 (37) | 881 (42) | 1577 (40) | .002 |
| Educational level | 3989 | | | | |
| Less than high school | | 109 (6) | 114 (5) | 223 (6) | .623 |
| High school | | 834 (44) | 873 (42) | 1707 (43) | .086 |
| College degree | | 390 (21) | 455 (22) | 845 (21) | .459 |
| University degree | | 542 (29) | 657 (31) | 1199 (30) | .085 |
| CHD severity | 4028 | | | | |
| Mild | | 420 (22) | 618 (29) | 1038 (26) | <.001 |
| Moderate | | 924 (49) | 1021 (48) | 1945 (49) | .783 |
| Severe | | 553 (29) | 476 (23) | 1029 (25) | <.001 |
| NYHA I and II | 3927 | 1665 (90) | 1778 (88) | 3443 (89) | <.001 |
| NYHA III and IV | | 186 (10) | 251 (12) | 437 (11) | .037 |
| Comorbidity | | | | | |
| Congestive heart failure | 3959 | 193 (10) | 240 (12) | 435 (11) | .471 |
| Cognitive impairment | 3998 | 29 (2) | 19 (1) | 48 (1) | .066 |
| History of mental disorder | 4012 | 149 (8) | 265 (13) | 418 (10) | <.001 |
| Employment status | 3993 | | | | |
| Employed | | 1381 (74) | 1364 (65) | 2745 (69) | <.001 |
| Full-time | | 1151 (83) | 893 (65) | 2044 (74) | <.001 |
| Part-time | | 230 (17) | 471 (35) | 701 (26) | <.001 |
| Homemaker or retired | | 79 (4) | 255 (12) | 334 (8) | <.001 |
| Unemployed ^c | | 192 (10) | 201 (10) | 393 (10) | .511 |
| Disabled | | 127 (7) | 173 (8) | 300 (8) | .074 |
| Student or other | | 101 (5) | 105 (5) | 206 (5) | .606 |
| Good WAS (≥8) ^d | 3227 | 1248 (78) | 1247 (77) | 2495 (77) | .596 |
| Poor WAS (≤5) | | 135 (8) | 148 (9) | 283 (9) | .467 |
| Any work limitations ^e | 2745 | 441 (32) | 499 (37) | 940 (34) | .010 |

Abbreviations: CHD, congenital heart disease; NYHA, New York Heart Association (functional class).

^aSex unknown in 16 patients.

^bPercentages within columns.

^cUnemployed also includes job seeking.

^dWAS = work ability score = only from patients aged below 65 years who were currently employed or had experiences with employment (N = 3283, from 56 patients WAS unknown).

^eWork limitations among employed patients (N = 2756, from 11 patients sex unknown).

ratios (ORs) and 95% confidence intervals (CIs) were calculated using multivariable logistic regression. The level of statistical significance was set at $P \leq .05$ and all reported P values were two-tailed. All statistical analyses were performed using SPSS statistical software for Windows (version 23; IBM, Armonk, New York).

3 | RESULTS

A total of 4028 adults with CHD were enrolled in APPROACH-IS. Characteristics of the study population are described in Table 1. Median age was 32 years (IQR 25–42). Slightly more women than men (53% vs 47%, $P = .002$) were included and CHD distribution was more severe in men. The majority of patients originated from Europe or North America, had at least a high school degree, were married or living with a partner, had no children and had NYHA functional class I symptoms. Fifty-one percent of all patients had a college or university degree.

Data on employment status were missing from 35 (<1%) participants. In addition, 135 (3%) patients 65 years or older, 334 (8%) homemaker or retired, 157 (4%) students, and 180 (5%) unemployed or disabled patients who considered themselves entirely unable to work, were excluded. Exclusion criteria were partly overlapping.

Factors influencing employment status and work ability were assessed on the remaining 3283 (82%) patients.

3.1 | Employment status

As shown in Table 1, 69% of all patients were employed, varying from 43% in India to 80% in Belgium. Employment status per country is depicted in Figures 1 and 2 (see Tables S1, S2, and S3, and Table 1, for exact numbers). Women were generally less often employed than men (65% vs 74%, $P < .001$), varying from 23% in India to 77% in Belgium. Disability rates increased by age and CHD severity (from 4% in mild to 11% in severe CHD, $P < .001$). Additional data on employment status, WAS and work limitations according to CHD complexity can be found online in Table 2. As shown in Figure 1, employment rates were lower than expected in 8 of 12 countries, particularly in India and Switzerland, and higher than expected in the remaining 4 countries. There was notable variation between countries, ranging from an employment rate of 18% above the general population in Belgium to 18% below the general population in India. Overall, unemployment (including job seeking) was observed in 10% of patients, equally in men and women. In all countries except for Belgium and The Netherlands, unemployment rates were higher in CHD patients than in the general population (Table 1). The proportion of patients




FIGURE 1 Distribution of employment status in ACHD patients per country. Since reference data are not available, employment rates for the general population per country (available for 12 countries) were derived from the Organisation for Economic Co-operation and Development (OECD). They are highlighted with a dark blue bar




FIGURE 2 Distribution of employment and work limitations (work limitations among employed patients) among participating countries. This map shows the amount of employment and work limitations per country, with darker colors reflecting less unemployment and more limitations. Participating countries with total number of participants and percentage employed: Argentina ($N = 174$, 52% employed), Australia ($N = 132$, 67%), Belgium ($N = 275$, 80%), Canada ($N = 517$, 75%), France ($N = 95$, 63%), India ($N = 199$, 43%), Italy ($N = 64$, 74%), Japan ($N = 257$, 67%), Malta ($N = 116$, 74%), Norway ($N = 172$, 70%), Sweden ($N = 468$, 73%), Switzerland ($N = 276$, 72%), Taiwan ($N = 250$, 64%), The Netherlands ($N = 254$, 70%), and United States ($N = 744$, 70%)

TABLE 2 Factors associated with employment in adult CHD patients ($N = 3283$ with $N = 2756$ employed patients)

| Variable | OR | 95% CI | P value |
|-------------------------------------|------|-----------|---------|
| Female gender | .66 | .57–.76 | <.001 |
| Age (per year) | .99 | .97–1.00 | .080 |
| High school education | 1.99 | 1.37–2.90 | <.001 |
| College degree | 3.52 | 2.25–5.50 | <.001 |
| University degree | 3.69 | 2.65–5.14 | <.001 |
| NYHA II | .67 | .57–.78 | <.001 |
| NYHA III | .30 | .21–.41 | <.001 |
| NYHA IV | .13 | .06–.26 | <.001 |
| Moderate CHD | 1.06 | .89–1.26 | .514 |
| Severe CHD | .88 | .74–1.05 | .166 |
| Having a partner | 1.72 | 1.48–1.99 | <.001 |
| Having children | .97 | .77–1.23 | .807 |
| History of congestive heart failure | .74 | .60–.92 | .008 |
| History of mental disorder | .85 | .65–1.11 | .228 |

Abbreviations: NYHA = New York Heart Association (functional class); CHD = congenital heart disease.

working part-time varied between countries from 8% in India to 51% in Australia. Women were more likely to work part-time than men (35% vs 17%, $P < .001$, Table 1). Overall, part-time employment did not differ according to CHD severity (Table 2), but patterns in part-time workers varied strongly between countries. For example, in

Australia, part-time workers were mostly patients with more severe CHD, whereas in The Netherlands, mainly women worked part-time.

Factors associated with less employment in multivariate logistic regression analyses by generalized linear models were female sex, worse NYHA functional class, and a history of congestive heart

failure (but not CHD severity) (Table 2). Higher education (ORs ranging from 1.99, 95% CI 1.37-2.90 for only high school to 3.69, 95% CI 2.65-5.14, for university degree) and having a partner (OR 1.72, 95% CI 1.48-1.99) were positively associated with being employed.

3.2 | Work ability

Most employed patients (77%) found themselves well capable of working, expressed in a WAS of 8 or higher. This varied from 57% in Taiwan to 94% in Malta (data per country shown online in Table 1). In total, only 283 patients (9%) with working experiences scored a WAS of 5 or below (ranging from 0% in Malta to 19% in India). Among all employed patients, 5% considered themselves not at all capable of working, varying from 0% in Argentina, Malta, and The Netherlands to 11% of employed patients in India and Taiwan.

3.3 | Work limitations

Among currently employed patients, 34% experienced some degree of limitations (varying from 16% in Malta to 50% in India and Taiwan, Table 1). Figure 2 reflects the employment situation including limitations per country. Patients that experienced working limitations were slightly older (35 ± 11 vs 34 ± 11 years, $P < .001$) and more likely to be female: limitations were reported by 37% of working women vs 32% of working men ($P = .010$). Part-time workers reported more limitations than patients who worked full-time (43% vs 31% limitations among full-time workers, $P < .001$). Table 3 shows that female sex (OR 1.36, 95% CI 1.17-1.58), increasing age (OR 1.03, 95% CI 1.02-1.04; for each increasing year of age, 3% more limitations were seen), more severe CHD (OR 1.31, 95% CI 1.09-1.58 for moderate to 2.10, 95% CI 1.68-2.62 severe CHD), history of

congestive heart failure (OR 1.57, 95% CI 1.19-2.08), and mental illness (OR 2.26, 95% CI 1.67-3.06) were all associated with limitations at work. Only 4% of patients with work limitations had none of these associated factors. A university degree was the only factor associated with fewer limitations (OR 0.62, 95% CI 0.41-0.93). Patients with more severe CHD reported more work limitations (presented in Table 2, together with the reported symptoms). The impact of severe CHD (compared to nonsevere CHD) on work limitations varied greatly by country, but was strongly predictive in Japan, India, Sweden, Switzerland, Canada, and the United States (ORs ranging from 1.9 to 5.2, see Table 3).

4 | DISCUSSION

In the present study, employment was studied in a large international cohort using a uniform methodology. The majority of CHD patients were employed, despite experiencing severe limitations. The most significant factors positively associated with employment were education and male gender.

To date, there are only a few prior studies investigating employment in adults with CHD and—despite increasing focus on patient-reported outcomes—they are mostly limited to the assessment of employment status. Previous studies indicated that adults with CHD from Western Europe and the United States were less likely to be employed than healthy controls, particularly in cyanotic but even in mild CHD.^{8-10,12,26} However, two Scandinavian studies reported employment levels higher than observed in the general population.^{14,15} In our study, employment rates were lower than in the general population in most countries, but there was substantial variation between countries. Since the same questionnaires and

TABLE 3 Risk factors associated with work limitations among employed patients ($N = 941$ in $N = 2756$ employed patients)

| Variable | OR | 95% CI | P value |
|-------------------------------------|------|-----------|---------|
| Female sex | 1.36 | 1.17–1.58 | <.001 |
| Age (per year) | 1.03 | 1.02–1.04 | <.001 |
| Moderate CHD | 1.31 | 1.09–1.58 | .005 |
| Severe CHD | 2.10 | 1.68–2.62 | <.001 |
| High school education | 1.02 | .69–1.52 | .915 |
| College degree | .79 | .54–1.17 | .243 |
| University degree | .62 | .41–.93 | .021 |
| History of congestive heart failure | 1.57 | 1.19–2.08 | .002 |
| History of mental disorder | 2.26 | 1.67–3.06 | <.001 |
| Having a partner | .86 | .71–1.03 | .105 |
| Having children | .88 | .71–1.09 | .244 |

Abbreviation: CHD = congenital heart disease.

reference data were used for all participating countries, we assume that there are indeed genuine differences in the impact of CHD on employment status in different continents and countries.

Although the majority of patients were employed and self-reported as well capable of working, one third experienced limitations at work. More work limitations and a poor self-declared work ability were observed in economically disadvantaged countries. This may be explained by insufficient social security with less financial support for disabled patients. For example in India, a considerable number of employed patients still considered themselves not capable of working and 50% reported limitations. However, even in a wealthy country like Australia, patients with cardiovascular disease have been reported to be at risk of living in poverty when unemployed due to their condition.²⁷

4.1 | Predictors for employment and limitations at work

Factors associated with employment in adults with CHD were in great part similar to factors associated with employment in the general population and in other chronic diseases.¹⁷ Higher educational levels and male gender were associated with more employment. Our previous qualitative study on barriers and limitations at work showed that less physical work was beneficial for our patients.¹¹ Higher education generally leads to jobs with less physical work, more internal and external recovery possibilities, and, often, better job conditions. However, patients with CHD are at increased risk for neurodevelopmental disorders.²⁸ On the whole, patients with CHD attain lower educational levels than the general population.^{10,29-32} Since this study demonstrates a positive effect on employment and limitations at work from higher education, efforts should be undertaken to maximize academic attainment in this population. In a scientific statement, Marino et al. called for vocational planning early in adolescence to maximize appropriate educational options.²⁸ Previous studies have indicated that patients with CHD lack advice on education and career.^{8,11} Although they were less likely to consider offered advice helpful than healthy controls, the chance to return to work in patients on sick leave can be influenced by the treating physician.³³ Since most patients with CHD are in lifelong cardiac care, the potential impact of advice from health care providers should not be taken lightly.

As in the general population, female gender was found to be a risk factor for less employment.²³ This factor might therefore be more of a cultural reflection of specific gender roles or social stereotyping than a specific risk factor for patients with CHD. However, in prior studies comparing adult CHD patients to the general population, the effect of CHD on employment was larger for men, meaning that employment patterns in male CHD patients differed more from the patterns in the male general population than female patterns did.^{9,10,34} It would be interesting, but beyond the scope of this study, to compare male and female working patterns per country.

What could promote successful employment in CHD patients besides good education? Besides early developmental interventions and governmental disability support, hypothetically, working

part-time could be seen as a way to stay employed despite limitations. In our study, patients working part-time reported more limitations than those working full-time. However, in countries where a large proportion of patients worked part-time, disability, or unemployment rates were not specifically lower than countries with fewer part-time workers. Working part-time requires flexibility from the employer to accommodate part-time employment and requires alternative sources for the patient for the remaining income. The lack of the latter may explain that for some patients in certain countries, working part-time was not an option.

4.2 | Strengths and limitations

The relatively large sample size is a major strength of this study. However, the number of participants (as well as countries per continent) varied between countries, which did not allow for detailed analyses on a national or continental level. Furthermore, the most important limitation, is the lack of reference data on employment on a national level. Therefore, the influence of socioeconomic and cultural factors is unknown. The study was based on uniform questionnaires assessing patient-reported outcomes. In hindsight, definitions—favorably country-specific—could have been made more explicit, since some descriptions such as “disability” may have different meanings or regulations between countries. Similarly, the Work Ability Score consists of a “ladder” scale system and the way in which patients interpret this scale and are likely to choose extreme values, could be influenced by country or culture. Furthermore, data on work limitations were only studied in employed patients. This selected sample is likely healthier than the larger CHD population (known as the “healthy worker effect”), thereby limiting generalizability of our findings.³⁵ However, we prioritized internal validity by including only employed patients. Any bias from a “healthy worker effect” is likely in the direction of underestimation of work-related limitations.

5 | CONCLUSIONS

A history of CHD has consequences beyond the medical field. Although most adults with CHD are employed, CHD continues to have a negative impact on employment. Further, in this global study with uniform questionnaires, employment status and difficulties varied widely between countries. Differences could only be partly explained by economic status. Education is the main predictor for successful employment with higher education appearing to even protect against limitations at work. Since the Work Ability Score (WAS) has proven to be a useful tool to predict future disability in other chronic diseases, a longitudinal follow-up study is required to determine whether it can also predict disability in CHD patients. In the meantime, a poor WAS in an employed patient should be considered a poor prognostic sign that needs to be addressed and may indicate consultation of an occupational physician or therapist. In general, our results support advocating for patients with CHD to reach their full educational potential.

REGISTRATION

ClinicalTrials.gov: NCT02150603.

AUTHOR CONTRIBUTIONS

On behalf of the APPROACH-IS consortium and the International Society for Adult Congenital Heart Disease (ISACHD)

Maayke A. Sluman: Concept/Design; Data collection; Data analysis/interpretation; Statistics; Drafting article; Critical revision of article; Approval of article.

Silke Apers: Data collection; Critical revision of article; Approval of article.

Judith K. Sluiter: Concept/Design; Data analysis/interpretation; Statistics; Critical revision of article; Approval of article.

Karen Nieuwenhuijsen: Critical revision of article; Approval of article.

Philip Moons: Concept/Design; Critical revision of article; Funding secured by; Approval of article.

Koen Luyckx: Critical revision of article; Approval of article.

Adrienne H. Kovacs: Critical revision of article; Approval of article.

Corina Thomet: Data collection; Critical revision of article; Approval of article.

Werner Budts: Critical revision of article; Approval of article.

Junko Enomoto: Data collection; Critical revision of article; Approval of article.

Hsiao-Ling Yang: Data collection; Approval of article.

Jamie L. Jackson: Critical revision of article; Approval of article.

Paul Khairy: Critical revision of article; Approval of article.

Stephen C. Cook: Critical revision of article; Approval of article.

Raghavan Subramanyan: Critical revision of article; Approval of article.

Luis Alday: Data collection; Critical revision of article; Approval of article.

Katrine Eriksen: Data collection; Approval of article.

Mikael Dellborg: Critical revision of article; Approval of article.

Malin Berghammer: Critical revision of article; Approval of article.

Eva Mattsson: Data collection; Critical revision of article; Approval of article.

Andrew S. Mackie: Critical revision of article; Approval of article.

Samuel Menahem: Critical revision of article; Approval of article.

Maryanne Caruana: Data collection; Critical revision of article; Approval of article

Kathy Gosney: Data collection; Approval of article.

Alexandra Soufi: Data collection; Critical revision of article; Approval of article.

Susan M. Fernandes: Critical revision of article; Approval of article.

Kamila S. White: Critical revision of article; Approval of article.

Edward Callus: Critical revision of article; Approval of article.

Shelby Kutty: Data collection; Critical revision of article; Approval of article.

Berto J. Bouma: Concept/Design; Data analysis/interpretation; Statistics; Critical revision of article; Approval of article.

Barbara JM Mulder: Concept/Design; Data analysis/interpretation; Statistics; Critical revision of article; Approval of article.

All authors take responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

ORCID

Maayke A. Sluman  <https://orcid.org/0000-0002-1199-6179>

Jamie L. Jackson  <https://orcid.org/0000-0002-1752-1377>

Maryanne Caruana  <https://orcid.org/0000-0003-3947-3350>

REFERENCES

1. Moons P, Bovijn L, Budts W, Belmans A, Gewillig M. Temporal trends in survival to adulthood among patients born with congenital heart disease from 1970 to 1992 in Belgium. *Circulation*. 2010;122:2264-2272.
2. Mandalenakis Z, Rosengren A, Skoglund K, Lappas G, Eriksson P, Dellborg M. Survivorship in children and young adults with congenital heart disease in Sweden. *JAMA Intern Med*. 2016;177:224-230.
3. Vigl M, Niggemeyer E, Hager A, Schwedler G, Kropf S, Bauer U. The importance of socio-demographic factors for the quality of life of adults with congenital heart disease. *Qual Life Res*. 2011;20:169-177.
4. Moons P, Van Deyk K, Marquet K, et al. Individual quality of life in adults with congenital heart disease: a paradigm shift. *Eur Heart J*. 2005;26:298-307.
5. Apers S, Kovacs AH, Luyckx K, et al. Quality of life of adults with congenital heart disease in 15 countries: evaluating country-specific characteristics. *J Am Coll Cardiol*. 2016;67:2237-2245.
6. Deanfield J, Thaulow E, Warnes C, et al. Management of grown up congenital heart disease. *Eur Heart J*. 2003;24:1035-1084.
7. Baumgartner H, Bonhoeffer P, De Groot NM, et al. ESC Guidelines for the management of grown-up congenital heart disease (new version 2010). *Eur Heart J*. 2010;31:2915-2957.
8. Crossland DS, Jackson SP, Lyall R, Burn J, O'Sullivan JJ. Employment and advice regarding careers for adults with congenital heart disease. *Cardiol Young*. 2005;15:391-395.
9. Geyer S, Norozi K, Buchhorn R, Wessel A. Chances of employment in women and men after surgery of congenital heart disease: comparisons between patients and the general population. *Congenit Heart Dis*. 2009;4:25-33.
10. Zomer AC, Vaartjes I, Uiterwaal CS, et al. Social burden and lifestyle in adults with congenital heart disease. *Am J Cardiol*. 2012;109:1657-1663.
11. Sluman MA, de Man S, Mulder BJ, Sluiter JK. Occupational challenges of young adult patients with congenital heart disease. *Neth Heart J*. 2014;22:216-224.
12. Simko LC, McGinnis KA, Schembri J. Educational needs of adults with congenital heart disease. *J Cardiovasc Nurs*. 2006;21:85-94.
13. Bygstad E, Pedersen LC, Pedersen TA, Hjortdal VE. Tetralogy of Fallot in men: quality of life, family, education, and employment. *Cardiol Young*. 2012;22:417-423.
14. Nieminen H, Sairanen H, Tikanoja T, et al. Long-term results of pediatric cardiac surgery in Finland: education, employment, marital status, and parenthood. *Pediatrics*. 2003;112:1345-1350.
15. Ternstedt BM, Wall K, Oddsson H, Riesenfeld T, Groth I, Schollin J. Quality of life 20 and 30 years after surgery in patients operated

- on for tetralogy of Fallot and for atrial septal defect. *Pediatr Cardiol*. 2001;22:128-132.
16. Caruana M, Grech V. Congenital heart disease has no negative impact on educational achievements and employment among Maltese adult patients under clinical follow-up. *Inter Cardiovasc Forum J*. 2016;8:41-46.
 17. Vooijs M, Leensen MC, Hoving JL, Daams JG, Wind H, Frings-Dresen MH. Disease-generic factors of work participation of workers with a chronic disease: a systematic review. *Int Arch Occup Environ Health*. 2015;88:1015-1029.
 18. Detaille SI, Heerkens YF, Engels JA, van der Gulden JW, van Dijk FJ. Common prognostic factors of work disability among employees with a chronic somatic disease: a systematic review of cohort studies. *Scand J Work Environ Health*. 2009;35:261-281.
 19. Apers S, Kovacs AH, Luyckx K, et al. Assessment of patterns of patient-reported outcomes in adults with congenital heart disease—International Study (APPROACH-IS): rationale, design, and methods. *Int J Cardiol*. 2015;179:334-342.
 20. Tuomi K, Jahkola A, et al. *Work Ability Index*. Helsinki: Finnish Institute of Occupational Health; 1998.
 21. Alavinia SM, de Boer AG, van Duivenbooden JC, Frings-Dresen MH, Burdorf A. Determinants of work ability and its predictive value for disability. *Occup Med (Lond)*. 2009;59:32-37.
 22. World economic outlook database. Available at: <https://www.imf.org/external/pubs/ft/weo/2016/01/weodata/download.aspx>. Accessed January 15, 2019.
 23. Employment rates per country. Available at: <https://data.oecd.org/emp/employment-rate.htm>. Accessed January 15, 2019.
 24. Franklin RC, Anderson RH, Daniëls O, et al. Report of the Coding Committee of the Association for European Paediatric Cardiology. *Cardiol Young*. 2002;12:611-618.
 25. Warnes CA, Liberthson R, Danielson GK, et al. Task force 1: the changing profile of congenital heart disease in adult life. *J Am Coll Cardiol*. 2001;37:1170-1175.
 26. Kamphuis M, Vogels T, Ottenkamp J, Van Der Wall EE, Verloove-Vanhorick SP, Vliegen HW. Employment in adults with congenital heart disease. *Arch Pediatr Adolesc Med*. 2002;156:1143-1148.
 27. Schofield DJ, Callander EJ, Shrestha RN, Percival R, Kelly SJ, Passey ME. Labour force participation and the influence of having CVD on income poverty of older workers. *Int J Cardiol*. 2012;156:80-83.
 28. Marino BS, Lipkin PH, Newburger JW, et al. Neurodevelopmental outcomes in children with congenital heart disease: evaluation and management: a scientific statement from the American Heart Association. *Circulation*. 2012;126:1143-1172.
 29. Olsen M, Hjortdal VE, Mortensen LH, Christensen TD, Sørensen HT, Pedersen L. Educational achievement among long-term survivors of congenital heart defects: a Danish population-based follow-up study. *Cardiol Young*. 2011;21:197-203.
 30. Mulkey SB, Bai S, Luo C, et al. School-age test proficiency and special education after congenital heart disease surgery in infancy. *J Pediatr*. 2016;178:47-54.
 31. Mulkey SB, Swearingen CJ, Melguizo MS, et al. Academic proficiency in children after early congenital heart disease surgery. *Pediatr Cardiol*. 2014;35:344-352.
 32. Özcan EE, Küçük A. Impact of severity of congenital heart diseases on university graduation rate among male patients. *Türk Kardiyol Dern Ars*. 2012;40:229-234.
 33. Dunstan DA, Covic T, Tyson GA. What leads to the expectation to return to work? Insights from a Theory of Planned Behavior (TPB) model of future work outcomes. *Work*. 2013;46:25-37.
 34. Sluman MA, Zomer AC, Vaartjes I, Bouma BJ, Mulder BJ. Congenital heart disease may hurt men more than women in job participation. *Int J Cardiol*. 2014;172:230-232.
 35. Shah D. Healthy worker effect phenomenon. *Indian J Occup Environ Med*. 2009;13:77-79.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

TABLE S1 Employment status, work ability and work limitations per country, N=3993

TABLE S2 Employment status, WAS and work limitations per CHD severity

TABLE S3 Impact of severe CHD as risk factor for work limitations (N=941 among N=2756 employed participants), adjusted for age and sex

How to cite this article: Sluman MA, Apers S, Sluiter JK, et al. Education as important predictor for successful employment in adults with congenital heart disease worldwide. *Congenital Heart Disease*. 2019;14:362-371. <https://doi.org/10.1111/chd.12747>



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Sluman, MA; Apers, S; Sluiter, JK; Nieuwenhuijsen, K; Moons, P; Luyckx, K; Kovacs, AH; Thomet, C; Budts, W; Enomoto, J; Yang, H-L; Jackson, JL; Khairy, P; Cook, SC; Subramanyan, R; Alday, L; Eriksen, K; Dellborg, M; Berghammer, M; Mattsson, E; Mackie, AS; Menahem, S; Caruana, M; Gosney, K; Soufi, A; Fernandes, SM; White, KS; Callus, E; Kutty, S; Bouma, BJ; Mulder, BJM

Title:

Education as important predictor for successful employment in adults with congenital heart disease worldwide

Date:

2019-05-01

Citation:

Sluman, M. A., Apers, S., Sluiter, J. K., Nieuwenhuijsen, K., Moons, P., Luyckx, K., Kovacs, A. H., Thomet, C., Budts, W., Enomoto, J., Yang, H. -L., Jackson, J. L., Khairy, P., Cook, S. C., Subramanyan, R., Alday, L., Eriksen, K., Dellborg, M., Berghammer, M. ,... Mulder, B. J. M. (2019). Education as important predictor for successful employment in adults with congenital heart disease worldwide. CONGENITAL HEART DISEASE, 14 (3), pp.362-371. <https://doi.org/10.1111/chd.12747>.

Persistent Link:

<http://hdl.handle.net/11343/249791>

File Description:

published version

License:

CC BY-NC