

Note di Matematica
Note Mat. 40 (2020) no. 1, 13–25.

ISSN 1123-2536, e-ISSN 1590-0932
doi:10.1285/i15900932v40n1p13

A new subclass of meromorphic functions with positive coefficients defined by Bessel function

B. Venkateswarlu

*Department of Mathematics, GSS, GITAM University,
Doddaballapur- 562 163, Bengaluru Rural, India.
bvmaths@gmail.com*

P.Thirupathi Reddy

*Department of Mathematics, Kakatiya University,
Warangal- 506 009, Telangana, India.
reddypt2@gmail.com*

Chao Meng

*Department of Mathematics, School of Science,
Shenyang Aerospace University, P.O.Box 110136, Shenyang, China.
mengchaosau@163.com*

R. Madhuri Shilpa

*Department of Mathematics, GSS, GITAM University,
Doddaballapur- 562 163, Bengaluru Rural, India.
madhuri.4073@gmail.com*

Received: 11.11.2019; accepted: 21.12.2019.

Abstract. In this paper, we introduce and study a new subclass of meromorphic univalent functions defined by Bessel function. We obtain coefficient inequalities, extreme points, radius of starlikeness and convexity. Finally we obtain partial sums and neighborhood properties for the class $\sigma_p^*(\eta, k, \lambda, \tau, c)$.

Keywords: meromorphic; extreme point; partial sums; neighborhood.

MSC 2000 classification: 30C45 .

1 Introduction

Let Σ denote the class of meromorphic functions of the form

$$f(z) = \frac{1}{z} + \sum_{n=1}^{\infty} a_n z^n \quad (1.1)$$

which are analytic in the punctured disc $U^* = \{z : z \in C \text{ and } 0 < |z| < 1\} = U \setminus \{0\}$.

A function $f \in \Sigma$ is said to be meromorphically starlike of order α if it satisfies

$$\operatorname{Re} \left\{ -\frac{zf'(z)}{f(z)} \right\} > \alpha$$

for some $\alpha(0 \leq \alpha < 1)$ and for all $z \in U$. Further, a function $f \in \Sigma$ is said to be meromorphically convex of order α if it satisfies

$$\operatorname{Re} \left\{ -\left(1 + \frac{zf''(z)}{f'(z)}\right) \right\} > \alpha,$$

for some $\alpha(0 \leq \alpha < 1)$ and for all $z \in U$.

Some subclasses of Σ were introduced and studied by Pommerenke [12], Miller [10], Mogra et al. [11], Cho [6], Cho et al. [7], Aouf [1, 2] and see also [17, 18].

For functions $f \in \Sigma$ given by (1.1) and $g \in \Sigma$ given by

$$g(z) = \frac{1}{z} + \sum_{n=1}^{\infty} b_n z^n,$$

we define the Hadamard product (or convolution) of f and g by

$$(f * g)(z) = \frac{1}{z} + \sum_{n=1}^{\infty} a_n b_n z^n.$$

We recall here the generalized Bessel function of first kind of order γ (see [8]), denoted by

$$W(z) = \sum_{n=0}^{\infty} \frac{(-c)^n}{n! \Gamma(\gamma + n + \frac{b+1}{2})} \left(\frac{z}{2}\right)^{2n+\gamma} \quad (z \in U)$$

(where Γ stands for the Gamma Euler function) which is the particular solution of the second order linear homogeneous differential equation (see, for details, [19])

$$z^2 w''(z) + bz w'(z) + [cz^2 - \gamma^2 + (1-b)\gamma]w(z) = 0$$

where $c, \gamma, b \in C$.

We introduce the function φ defined, in terms of the generalized Bessel function w by

$$\varphi(z) = 2^\gamma \Gamma\left(\gamma + \frac{b+1}{2}\right) z^{-(1+\frac{\gamma}{2})} w(\sqrt{z}).$$

By using the well-known Pochhammer symbol $(x)_\mu$ defined, for $x \in C$ and in terms of the Euler gamma function, by

$$(x)_\mu = \frac{\Gamma(x + \mu)}{\Gamma(x)} = \begin{cases} 1, & (\mu = 0); \\ x(x + 1) \cdots (x + \mu - 1), & (\mu = n \in N = \{1, 2, 3, \dots\}). \end{cases}$$

We obtain the following series representation for the function $\varphi(z)$

$$\varphi(z) = \frac{1}{z} + \sum_{n=0}^{\infty} \frac{(-c)^{n+1}}{4^{n+1}(n+1)!(\tau)_{n+1}} z^n$$

(where $\tau = \gamma + \frac{b+1}{2} \notin Z_0^- = \{0, -1, -2, \dots\}$).

Corresponding to the function φ define the Bessel operator S_τ^c by the following Hadamard product

$$\begin{aligned} S_\tau^c f(z) = (\varphi * f)(z) &= \frac{1}{z} + \sum_{n=0}^{\infty} \frac{\left(\frac{-c}{4}\right)^{n+1} a_n}{(n+1)!(\tau)_{n+1}} z^n \\ &= \frac{1}{z} + \sum_{n=1}^{\infty} \phi(n, \tau, c) a_n z^n, \end{aligned} \tag{1.2}$$

where $\phi(n, \tau, c) = \frac{\left(\frac{-c}{4}\right)^n}{(n)!(\tau)_n}$.

It easy to verify from the definition (1.2) that

$$z[S_{\tau+1}^c f(z)] = \tau S_\tau^c f(z) - (\tau + 1)S_{\tau+1}^c f(z). \tag{1.3}$$

Motivated by Sivaprasad Kumar et al. [16], Atshan et al. [5] and Venkateswarlu et al. [17, 18], now we define a new subclass $\sigma_p^*(\eta, k, \lambda, \tau, c)$ of Σ .

Definition 1. For $0 \leq \eta < 1, k \geq 0, 0 \leq \lambda < \frac{1}{2}$, we let $\sigma_p^*(\eta, k, \lambda, \tau, c)$ be the subclass of Σ_p consisting of functions of the form (1.1) and satisfying the analytic criterion

$$\begin{aligned} -Re \left(\frac{z(S_\tau^c f(z))'}{S_\tau^c f(z)} + \lambda z^2 \frac{(S_\tau^c f(z))''}{S_\tau^c f(z)} + \eta \right) \\ > k \left| \frac{z(S_\tau^c f(z))'}{S_\tau^c f(z)} + \lambda z^2 \frac{(S_\tau^c f(z))''}{S_\tau^c f(z)} + 1 \right|. \end{aligned} \tag{1.4}$$

In order to prove our results we need the following lemmas [4].

Lemma 1. *If η is a real number and $\omega = -(u + iv)$ is a complex number then*

$$Re(\omega) \geq \eta \Leftrightarrow |\omega + (1 - \eta)| - |\omega - (1 - \eta)| \geq 0.$$

Lemma 2. *If $\omega = u + iv$ is a complex number and η is a real number then*

$$-Re(\omega) \geq k|\omega + 1| + \eta \Leftrightarrow -Re\left(\omega(1 + ke^{i\theta}) + ke^{i\theta}\right) \geq \eta, -\pi \leq \theta \leq \pi.$$

The main object of this paper is to study some usual properties of the geometric function theory such as the coefficient bounds, extreme points, radii of meromorphic starlikeness and convexity for the class $\sigma_p^*(\eta, k, \lambda, \tau, c)$. Further, we obtain partial sums and neighborhood properties for the class also.

2 Coefficient estimates

In this section we obtain necessary and sufficient condition for a function f to be in the class $\sigma_p^*(\eta, k, \lambda, \tau, c)$.

Theorem 1. *Let $f \in \Sigma$ be given by (1.1). Then $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ if and only if*

$$\sum_{n=1}^{\infty} [n(k+1)(1+(n-1)\lambda) + (k+\eta)] \phi(n, \tau, c) a_n \leq (1-\eta) - 2\lambda(1+k). \quad (2.1)$$

Proof. Let $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$. Then by Definition 1 and using Lemma 2, it is enough to show that

$$-Re \left\{ \left(\frac{z(S_\tau^c f(z))'}{S_\tau^c f(z)} + \lambda z^2 \frac{(S_\tau^c f(z))''}{S_\tau^c f(z)} \right) (1 + ke^{i\theta}) + ke^{i\theta} \right\} > \eta, -\pi \leq \theta \leq \pi. \quad (2.2)$$

For convenience

$$\begin{aligned} C(z) &= - [z(S_\tau^c f(z))' + \lambda z^2 (S_\tau^c f(z))''] (1 + ke^{i\theta}) - ke^{i\theta} S_\tau^c f(z) \\ D(z) &= S_\tau^c f(z) \end{aligned}$$

that is, the equation (2.2) is equivalent to

$$-Re \left(\frac{C(z)}{D(z)} \right) \geq \eta.$$

In view of Lemma 1, we only need to prove that

$$|C(z) + (1-\eta)D(z)| - |C(z) - (1-\eta)D(z)| \geq 0.$$

Therefore

$$\begin{aligned}
& |C(z) + (1 - \eta)D(z)| \\
& \geq (2 - \eta - 2\lambda(k + 1)) \frac{1}{|z|} \\
& \quad - \sum_{n=1}^{\infty} [n(k + 1)(1 + (n - 1)\lambda) + (k + \eta - 1)] \phi(n, \tau, c) a_n |z|^n \\
\text{and } & |C(z) - (1 - \eta)D(z)| \\
& \leq (\eta + 2\lambda(k + 1)) \frac{1}{|z|} \\
& \quad + \sum_{n=1}^{\infty} [n(k + 1)(1 + (n - 1)\lambda) + (k + \eta + 1)] \phi(n, \tau, c) a_n |z|^n.
\end{aligned}$$

It is to show that

$$\begin{aligned}
& |C(z) + (1 - \eta)D(z)| - |C(z) - (1 + \eta)D(z)| \\
& \geq (2(1 - \eta) - 4\lambda(k + 1)) \frac{1}{|z|} \\
& \quad - 2 \sum_{n=1}^{\infty} [n(k + 1)(1 + (n - 1)\lambda) + (k + \eta)] \phi(n, \tau, c) a_n |z|^n \\
& \geq 0, \text{ by the given condition (2.1).}
\end{aligned}$$

Conversely, suppose $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$. Then by Lemma 1, we have (2.2).

Choosing the values of z on the positive real axis the inequality (2.2) reduces to

$$\operatorname{Re} \left\{ \frac{[1 - \eta - 2\lambda(1 + ke^{i\theta})] \frac{1}{z^2} + \sum_{n=1}^{\infty} [n(1 + (n - 1)\lambda)(1 + ke^{i\theta}) + (\eta + ke^{i\theta})] \phi(n, \tau, c) z^{n-1}}{\frac{1}{z^2} + \sum_{n=1}^{\infty} \phi(n, \tau, c) a_n z^{n-1}} \right\} \geq 0.$$

Since $\operatorname{Re}(-e^{i\theta}) \geq -|e^{i\theta}| = -1$, the above inequality reduces to

$$\operatorname{Re} \left\{ \frac{[1 - \eta - 2\lambda(1 + k)] \frac{1}{r^2} + \sum_{n=1}^{\infty} [n(1 + k)(1 + (n - 1)\lambda) + (\eta + k)] \phi(n, \tau, c) a_n r^{n-1}}{\frac{1}{r^2} + \sum_{n=1}^{\infty} \phi(n, \tau, c) r^{n-1}} \right\} \geq 0.$$

Letting $r \rightarrow 1^-$ and by the mean value theorem, we have obtained the inequality (2.1). \square

Corollary 1. *If $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ then*

$$a_n \leq \frac{(1 - \eta) - 2\lambda(k + 1)}{[n(1 + k)(1 + (n - 1)\lambda) + (\eta + k)]\phi(n, \tau, c)}. \quad (2.3)$$

By taking $\lambda = 0$ in Theorem 1, we get the following corollary.

Corollary 2. *If $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ then*

$$a_n \leq \frac{2 - \eta}{[n(1 + k) + (\eta + k)]\phi(n, \tau, c)}. \quad (2.4)$$

Theorem 2. *If $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ then for $0 < |z| = r < 1$,*

$$\frac{1}{r} - \frac{(1 - \eta) - 2\lambda(k + 1)}{(2k + \eta + 1)\phi(1, \tau, c)} r \leq |f(z)| \leq \frac{1}{r} + \frac{(1 - \eta) - 2\lambda(k + 1)}{(2k + \eta + 1)\phi(1, \tau, c)} r. \quad (2.5)$$

This result is sharp for the function

$$f(z) = \frac{1}{z} + \frac{(1 - \eta) - 2\lambda(k + 1)}{(2k + \eta + 1)\phi(1, \tau, c)} z. \quad (2.6)$$

Proof. Since $f(z) = \frac{1}{z} + \sum_{n=1}^{\infty} a_n z^n$, we have

$$|f(z)| = \frac{1}{r} + \sum_{n=1}^{\infty} a_n r^n \leq \frac{1}{r} + r \sum_{n=1}^{\infty} a_n. \quad (2.7)$$

Since $n \geq 1$, $(2k + \eta + 1) \leq n(k + 1)(k + \eta)\phi(n, \tau, c)$, using Theorem 1, we have

$$\begin{aligned} (2k + \eta + 1) \sum_{n=1}^{\infty} a_n &\leq \sum_{n=1}^{\infty} n(k + 1)(k + \eta)\phi(n, \tau, c) \\ &\leq (1 - \eta) - 2\lambda(k + 1) \\ &\Rightarrow \sum_{n=1}^{\infty} a_n \leq \frac{(1 - \eta) - 2\lambda(k + 1)}{(2k + \eta + 1)\phi(1, \tau, c)}. \end{aligned}$$

Using the above inequality in (2.7), we have

$$\begin{aligned} |f(z)| &\leq \frac{1}{r} + \frac{(1 - \eta) - 2\lambda(k + 1)}{(2k + \eta + 1)\phi(1, \tau, c)} r \\ \text{and } |f(z)| &\geq \frac{1}{r} - \frac{(1 - \eta) - 2\lambda(k + 1)}{(2k + \eta + 1)\phi(1, \tau, c)} r. \end{aligned}$$

The result is sharp for the function $f(z) = \frac{1}{z} + \frac{(1 - \eta) - 2\lambda(k + 1)}{(2k + \eta + 1)\phi(1, \tau, c)} z$. \square

Corollary 3. *If $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ then*

$$\frac{1}{r^2} - \frac{(1-\eta) - 2\lambda(k+1)}{(2k+\eta+1)\phi(1, \tau, c)} \leq |f'(z)| \leq \frac{1}{r^2} + \frac{(1-\eta) - 2\lambda(k+1)}{(2k+\eta+1)\phi(1, \tau, c)}.$$

The result is sharp for the function given by (2.6).

3 Extreme points

Theorem 3. *Let $f_0(z) = \frac{1}{z}$ and*

$$f_n(z) = \frac{1}{z} + \sum_{n=1}^{\infty} \frac{(1-\eta) - 2\lambda(k+1)}{[n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c)} z^n, \quad n \geq 1. \quad (3.1)$$

Then $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ if and only if it can be expressed in the form

$$f(z) = \sum_{n=0}^{\infty} u_n f_n(z), \quad u_n \geq 0 \quad \text{and} \quad \sum_{n=1}^{\infty} u_n = 1. \quad (3.2)$$

Proof. Suppose $f(z)$ can be expressed as in (3.2). Then

$$\begin{aligned} f(z) &= \sum_{n=0}^{\infty} u_n f_n(z) = u_0 f_0(z) + \sum_{n=1}^{\infty} u_n f_n(z) \\ &= \frac{1}{z} + \sum_{n=1}^{\infty} u_n \frac{(1-\eta) - 2\lambda(k+1)}{[n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c)} z^n. \end{aligned}$$

Therefore

$$\begin{aligned} &\sum_{n=1}^{\infty} u_n \frac{\{(1-\eta) - 2\lambda(k+1)\} \{ [n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c) \}}{\{ [n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c) \} \{ (1-\eta) - 2\lambda(k+1) \}} \\ &= \sum_{n=1}^{\infty} u_n = 1 - u_0 \leq 1. \end{aligned}$$

So by Theorem 1, $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$.

Conversely, suppose that $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$. Since

$$a_n \leq \frac{(1-\eta) - 2\lambda(k+1)}{[n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c)} \quad n \geq 1.$$

We set $u_n = \frac{[n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c)}{(1-\eta) - 2\lambda(k+1)} a_n$, $n \geq 1$ and $u_0 = 1 - \sum_{n=1}^{\infty} u_n$.

Then we have $f(z) = \sum_{n=0}^{\infty} u_n f_n(z) = u_0 f_0(z) + \sum_{n=1}^{\infty} u_n f_n(z)$.

Hence the result follows. \square

4 Radii of meromorphically starlike and meromorphically convexity

Theorem 4. *Let $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$. Then f is meromorphically starlike of order δ , ($0 \leq \delta \leq 1$) in the unit disc $|z| < r_1$, where*

$$r_1 = \inf_n \left[\frac{(1-\delta)}{(n+2-\delta)} \frac{[n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c)}{(1-\eta) - 2\lambda(k+1)} \right]^{\frac{1}{n+1}}, \quad n \geq 1.$$

The result is sharp for the extremal function $f(z)$ given by (2.6).

Proof. The function $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ of the form (1.1) is meromorphically starlike of order δ in the disc $|z| < r_1$ if and only if it satisfies the condition

$$\left| \frac{zf'(z)}{f(z)} + 1 \right| < (1-\delta). \quad (4.1)$$

Since

$$\left| \frac{zf'(z)}{f(z)} + 1 \right| \leq \left| \frac{\sum_{n=1}^{\infty} (n+1)a_n z^{n+1}}{1 + \sum_{n=1}^{\infty} a_n z^{n+1}} \right| \leq \frac{\sum_{n=1}^{\infty} (n+1)|a_n||z|^{n+1}}{1 - \sum_{n=1}^{\infty} |a_n||z|^{n+1}}.$$

The above expression is less than $(1-\delta)$ if $\sum_{n=1}^{\infty} \frac{(n+2-\delta)}{(1-\delta)} a_n |z|^{n+1} < 1$.

Using the fact that $f(z) \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ if and only if

$$\sum_{n=1}^{\infty} \frac{[n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c)}{(1-\eta) - 2\lambda(k+1)} a_n \leq 1.$$

Thus, (4.1) will be true if $\frac{(n+2-\delta)}{(1-\delta)} |z|^{n+1} < \frac{[n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c)}{(1-\eta) - 2\lambda(k+1)}$

or equivalently $|z|^{n+1} < \frac{(1-\delta)}{(n+2-\delta)} \frac{[n(1+k)(1+(n-1)\lambda) + (\lambda+k)]\phi(n, \tau, c)}{(1-\eta) - 2\lambda(k+1)}$

which yields the starlikeness of the family. □ QED

The proof of the following theorem is analogous to that of Theorem 4, hence we omit the proof.

Theorem 5. *Let $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$. Then f is meromorphically convex of order δ , ($0 \leq \delta \leq 1$) in the unit disc $|z| < r_2$, where*

$$r_2 = \inf_n \left[\frac{(1-\delta)}{n(n+2-\delta)} \frac{[n(1+k)(1+(n-1)\lambda) + (\eta+k)]\phi(n, \tau, c)}{(1-\eta) - 2\lambda(k+1)} \right]^{\frac{1}{n+1}}, \quad n \geq 1.$$

The result is sharp for the extremal function $f(z)$ given by (2.6).

5 Partial Sums

Let $f \in \Sigma$ be a function of the form (1.1). Motivated by Silverman [14] and Silvia [15] and also see [3], we define the partial sums f_m defined by

$$f_m(z) = \frac{1}{z} + \sum_{n=1}^m a_n z^n, (m \in N). \quad (5.1)$$

In this section we consider partial sums of function from the class $\sigma_p^*(\eta, k, \lambda, \tau, c)$ and obtain sharp lower bounds for the real part of the ratios of f to f_m and f' to f'_m .

Theorem 6. *Let $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ be given by (1.1) and define the partial sums $f_1(z)$ and $f_m(z)$ by*

$$f_1(z) = \frac{1}{z} \text{ and } f_m(z) = \frac{1}{z} + \sum_{n=1}^m |a_n| z^n, (m \in N \setminus \{1\}). \quad (5.2)$$

Suppose also that $\sum_{n=1}^{\infty} d_n |a_n| \leq 1$, where

$$d_n \geq \begin{cases} 1, & \text{if } n = 1, 2, \dots, m \\ \frac{[n(1+k)(1+(n-1)\lambda) + (\eta+k)]\phi(n, \tau, c)}{(1-\eta) - 2\lambda(k+1)}, & \text{if } n = m+1, m+2, \dots \end{cases}. \quad (5.3)$$

Then $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$. Furthermore

$$\operatorname{Re} \left(\frac{f(z)}{f_m(z)} \right) > 1 - \frac{1}{d_{m+1}} \quad (5.4)$$

$$\text{and } \operatorname{Re} \left(\frac{f_m(z)}{f(z)} \right) > \frac{d_{m+1}}{1 + d_{m+1}}. \quad (5.5)$$

Proof. For the coefficient d_n given by (5.3) it is not difficult to verify that

$$d_{m+1} > d_m > 1. \quad (5.6)$$

Therefore we have

$$\sum_{n=1}^m |a_n| + d_{m+1} \sum_{n=m+1}^{\infty} |a_n| \leq \sum_{n=1}^{\infty} |a_n| d_m \leq 1 \quad (5.7)$$

by using the hypothesis (5.3). By setting

$$g_1(z) = d_{m+1} \left(\frac{f(z)}{f_m(z)} - \left(1 - \frac{1}{d_{m+1}} \right) \right) = 1 + \frac{d_{m+1} \sum_{n=m+1}^{\infty} a_n z^{n-1}}{1 + \sum_{n=1}^{\infty} |a_n| z^{n-1}}$$

then it sufficient to show that

$$\operatorname{Re}(g_1(z)) \geq 0, (z \in U) \quad \text{or} \quad \left| \frac{g_1(z) - 1}{g_1(z) + 1} \right| \leq 1, (z \in U)$$

and applying (5.7), we find that

$$\begin{aligned} \left| \frac{g_1(z) - 1}{g_1(z) + 1} \right| &\leq \frac{d_{m+1} \sum_{n=m+1}^{\infty} |a_n|}{2 - 2 \sum_{n=1}^m |a_n| - d_{m+1} \sum_{n=m+1}^{\infty} |a_n|} \\ &\leq 1, (z \in U) \end{aligned}$$

which ready yields the assertion (5.4) of Theorem 6. In order to see that

$$f(z) = \frac{1}{z} + \frac{z^{m+1}}{d_{m+1}} \quad (5.8)$$

gives sharp result, we observe that for

$$z = re^{\frac{i\pi}{m}} \quad \text{that} \quad \frac{f(z)}{f_m(z)} = 1 - \frac{r^{m+2}}{d_{m+1}} \rightarrow 1 - \frac{1}{d_{m+1}} \quad \text{as} \quad r \rightarrow 1^-.$$

Similarly, if we takes $g_2(z) = (1 + d_{m+1}) \left(\frac{f_m(z)}{f(z)} - \frac{d_{m+1}}{1+d_{m+1}} \right)$ and making use of (5.7), we denote that

$$\left| \frac{g_2(z) - 1}{g_2(z) + 1} \right| < \frac{(1 + d_{m+1}) \sum_{n=m+1}^{\infty} |a_n|}{2 - 2 \sum_{n=1}^m |a_n| - (1 - d_{m+1}) \sum_{n=m+1}^{\infty} |a_n|}$$

which leads us immediately to the assertion (5.5) of Theorem 6.

The bound in (5.5) is sharp for each $m \in N$ with extremal function $f(z)$ given by (5.8). \square

The proof of the following theorem is analogous to that of Theorem 6, so we omit the proof.

Theorem 7. *If $f \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ be given by (1.1) and satisfies the condition (2.1) then*

$$\begin{aligned} \operatorname{Re} \left(\frac{f'(z)}{f'_m(z)} \right) &> 1 - \frac{m+1}{d_{m+1}} \\ \text{and} \quad \operatorname{Re} \left(\frac{f'_m(z)}{f'(z)} \right) &> \frac{d_{m+1}}{m+1+d_{m+1}}, \end{aligned}$$

$$\text{where } d_n \geq \begin{cases} n, & \text{if } n = 2, 3, \dots, m \\ \frac{[n(1+k)(1+(n-1)\lambda)+(\eta+k)]\phi(n,\tau,c)}{(1-\eta)-2\lambda(k+1)}, & \text{if } n = m+1, m+2, \dots \end{cases}$$

The bounds are sharp with the extremal function $f(z)$ of the form (2.6).

6 Neighbourhoods for the class $\sigma_p^{*\xi}(\eta, k, \lambda, \tau, c)$

In this section, we determine the neighborhood for the class $\sigma_p^{*\xi}(\eta, k, \lambda, \tau, c)$ which we define as follows:

Definition 2. A function $f \in \Sigma$ is said to be in the class $\sigma_p^{*\xi}(\eta, k, \lambda, \tau, c)$ if there exists a function $g \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ such that

$$\left| \frac{f(z)}{g(z)} - 1 \right| < 1 - \xi, \quad (z \in E, 0 \leq \xi < 1). \quad (6.1)$$

Following the earlier works on neighbourhoods of analytic functions by Goodman [9] and Ruscheweyh [13], we define the δ -neighbourhoods of function $f \in \Sigma$ by

$$N_\delta(f) = \left\{ g \in \Sigma : g(z) = \frac{1}{z} + \sum_{n=1}^{\infty} b_n z^n \text{ and } \sum_{n=1}^{\infty} n|a_n - b_n| \leq \delta \right\}. \quad (6.2)$$

Theorem 8. If $g \in \sigma_p^*(\eta, k, \lambda, \tau, c)$ and

$$\xi = 1 - \frac{\delta(2k + \eta + 1)\phi(1, \tau, c)}{(2k + \eta + 1)\phi(1, \tau, c) - (1 - \eta) + 2\lambda(k + 1)} \quad (6.3)$$

then $N_\delta(g) \subset \sigma_p^{*\xi}(\eta, k, \lambda, \tau, c)$.

Proof. Let $f \in N_\delta(g)$. Then we find from (6.2) that

$$\sum_{n=1}^{\infty} n|a_n - b_n| \leq \delta \quad (6.4)$$

which implies the coefficient inequality

$$\sum_{n=1}^{\infty} |a_n - b_n| \leq \delta \quad (n \in N). \quad (6.5)$$

Since $g \in \sigma_p^*(\eta, k, \lambda, \tau, c)$, we have

$$\sum_{n=1}^{\infty} b_n \leq \frac{(1 - \eta) - 2\lambda(k + 1)}{(2k + \eta + 1)\phi(1, c, \tau)}. \quad (6.6)$$

So that

$$\begin{aligned} \left| \frac{f(z)}{g(z)} - 1 \right| &< \frac{\sum_{n=1}^{\infty} |a_n - b_n|}{1 - \sum_{n=1}^{\infty} b_n} \\ &= \frac{\delta(2k + \eta + 1)L(1, c, \tau)}{(2k + \eta + 1)\phi(1, \tau, c) - (1 - \eta) + 2\lambda(k + 1)} \\ &= 1 - \xi \end{aligned}$$

provided ξ is given by (6.3). Hence by definition, $f \in \sigma_p^{*\xi}(\eta, k, \lambda, \tau, c)$ for ξ given by which completes the proof. \square

Acknowledgements. The authors express their sincere thanks to the esteemed referee(s) for their careful readings, valuable suggestions and comments, which helped them to improve the presentation of the paper.

References

- [1] M. K. AOUF : *A certain subclass of meromorphically starlike functions with positive coefficients*, Rend. Mat., **9** (1989), 225-235.
- [2] M. K. AOUF : *On certain class of meromorphically univalent functions with positive coefficients*, Rend. Mat., **11** (1991), 209-219.
- [3] M. K. AOUF AND H. SILVERMAN: *Partial sums of certain meromorphic p -valent functions*, J. Inequal. Pure Appl. Math., **7** (2006), no. 4, art. 119, 1-7 .
- [4] E. AQLAN , J. M. JHANGIRI AND S. R. KULKARNI : *Class of k -uniformly convex and starlike functions*, Tamkang J. Math., **35** (2004), 1-7.
- [5] W. G ATSHAN AND S. R. KULKARNI : *Subclass of meromorphic functions with positive coefficients defined by Ruscheweyh derivative*, I. J. Rajasthan Acad. Phys. Sci., **6** (2007), no. 2, 129-140.
- [6] N. E. CHO : *On certain class of meromorphic functions with positive coefficients*, J. Inst. Math. Comput.Sci., **3** (1990), no. 2, 119-125.
- [7] N. E. CHO AND H. M. SRIVASTAVA : *Argument estimates of certain analytic functions defined by a class of multiplier transformations*, Math. Comput. Model, **37** (2003), 39-49 .
- [8] E. DENIZ, H. ORHAN AND H. M. SRIVASTAVA: *Some sufficient conditions for univalence of certain families of integral operators involving generalized Bessel functions*, Taiwanese J. Math., **15** (2011), no. 2, 883-917.
- [9] A. W. GOODMAN: *Univalent functions and non-analytic curves*, Proc. Amer. Math. Soc., **8** (1957), 598-601.
- [10] J. E. MILLER : *Convex meromorphic mapping and related functions*, Proc. Am. Math. Soc., **25** (1970), 220-228.

- [11] M. L. MOGRA, T. R. REDDY AND O.P. JUNEJA : *Meromorphic univalent functions with positive coefficients*, Bull. Aust. Math. Soc., **32** (1985), 161-176.
- [12] CH. POMMERENKE : *On meromorphic starlike functions*, Pac. J. Math., **13** (1963), 221-235.
- [13] ST. RUSCHEWEYH : *Neighbourhoods of univalent functions*, Proc. Amer. Math. Soc., **81** (1981), 521-527.
- [14] H. SILVERMAN : *Partial sums of starlike and convex functions*, J. Math. Anal. Appl., **209** (1997), no. 2, 221-227.
- [15] E. M. SILVIA : *On partial sums of convex functions of order α* , Houston J. Math., **11** (1985), no. 2, 397-404.
- [16] S. SIVAPRASAD KUMAR, V. RAVICHANDRAN AND G. MURUGUSUNDARAMOORTHY : *Classes of meromorphic p -valent parabolic starlike functions with positive coefficients*, Aust. J. Math. Anal. Appl., **2** (2005), no. 2, art. 3, 1- 9.
- [17] B. VENKATESWARLU, P. THIRUPATHI REDDY AND N. RANI : *Certain subclass of meromorphically uniformly convex functions with positive coefficients* , Mathematica, **61 (84)** (2019), no. 2, 85-97 .
- [18] B. VENKATESWARLU, P. THIRUPATHI REDDY AND N. RANI: *On new subclass of meromorphically convex functions with positive coefficients*, Surv. Math. Appl., **14** (2019), 49 - 60.
- [19] G. N. WATSON : A Treatise on the Theory of Bessel Functions, (2nd edn.), Cambridge University Press, Cambridge, 1944.

