



TITLE:

Update on the efficacy and safety of sodium–glucose cotransporter 2 inhibitors in Asians and non-Asians

AUTHOR(S):

Fujita, Yoshihito; Inagaki, Nobuya

CITATION:

Fujita, Yoshihito ...[et al]. Update on the efficacy and safety of sodium–glucose cotransporter 2 inhibitors in Asians and non-Asians. *Journal of Diabetes Investigation* 2019, 10(6): 1408-1410

ISSUE DATE:

2019-11

URL:

<http://hdl.handle.net/2433/255850>

RIGHT:

© 2019 The Authors. *Journal of Diabetes Investigation* published by Asian Association for the Study of Diabetes (AASD) and John Wiley & Sons Australia, Ltd This is an open access article under the terms of the Creative Commons Attribution - NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Update on the efficacy and safety of sodium–glucose cotransporter 2 inhibitors in Asians and non-Asians

Sodium–glucose cotransporter 2 (SGLT2) inhibitors are a novel class of antidiabetic drugs for the treatment of diabetes; they reduce plasma glucose concentrations and bodyweight by inhibiting glucose reabsorption in the kidney¹. As the action of SGLT2 inhibitors is unique, they have been receiving attention regarding their efficacy and safety from well before their launch. In clinical use, further insights on the efficacy and cautions on the safety of SGLT2 inhibitors have been accumulating. Meta-analysis suggests that SGLT2 inhibitors, as well as glucagon-like peptide 1 analogs, lead to comparable bodyweight decreases in type 2 diabetes patients². Bodyweight loss is a major target for treatment of patients with type 2 diabetes, especially because overweight is associated with insulin resistance and cardiovascular risk factors.

Based on the evidence of several major mega-trials, which showed that SGLT2 inhibitors also exert preventive effects on major adverse cardiovascular events, heart failure hospitalization and progression of renal impairment, they were prioritized among other second-line oral drugs after metformin for patients with cardiovascular disease and chronic kidney disease in the consensus reports of the American Diabetes Association and the European Association for the Study of Diabetes³.

These consensus reports were edited based on evidence targeted primarily to Caucasians. It is widely recognized that

type 2 diabetes in Asians, especially East Asians, is characterized by β -cell dysfunction and less obesity compared with Caucasians⁴. These pathophysiological differences might have an impact on the therapeutic consideration of SGLT2 inhibitors. Cai *et al.*⁵ carried out a meta-analysis to compare the efficacy and adverse effects of SGLT2 inhibitors between Asian and non-Asian type 2 diabetes patients. They found no significant difference in the reduction of glycated hemoglobin or bodyweight. Furthermore, no disparity was found in the risk of all-cause mortality or hypoglycemia⁵. Subgroup analysis in the Empagliflozin Cardiovascular Outcome Event Trial in Type 2 Diabetes Mellitus Patients—Removing Excess Glucose (EMPA-REG OUTCOME) showed that the reductions in the risk of cardiovascular events and mortality by empagliflozin were consistent between Asians and the overall population⁶. These reductions were also consistent between East Asians and the overall population; however, a statistical test was not carried out, because the number of East Asian patients with cardiovascular events was small.

Clinical findings using SGLT2 inhibitors in Asian patients have been accumulating. Luseogliflozin added to liraglutide improves glycemic control and leads to a reduction in bodyweight (especially fat mass) in Japanese patients⁷. This suggests that combination therapy could be an attractive treatment approach for overweight Japanese patients with type 2 diabetes. Tofogliflozin showed better improvement of glycemic control and an increase in insulin secretion, especially in subgroups with high insulin levels at baseline⁸. These data suggest that SGLT2

inhibitors might facilitate recovery from impaired β -cell dysfunction in patients in whom insulin secretion capacity is preserved to a certain extent. Furthermore, administration of SGLT2 inhibitors with or without dipeptidyl peptidase-4 inhibitor decreased nocturnal hypoglycemia when taking basal–bolus insulin therapy, suggesting that SGLT2 inhibitors might increase the insulin counter-regulatory hormone concentrations, including glucagon, growth hormone, cortisol and catecholamine⁹. SGLT2 inhibitors also have the effect of reducing blood pressure comparable to that of low-dose thiazide diuretics in patients with both type 2 diabetes and hypertension¹⁰.

Regarding safety concerns, empagliflozin was found to be well tolerated in pooled analysis of East Asian patients with type 2 diabetes, which is consistent with results from the safety data in overall population pooled analysis from placebo-controlled trials¹¹. In Japan, shortly after six SGLT2 inhibitors were launched, safety became a major concern due to reports of several serious adverse events. These led to a “recommendation on the appropriate usage of SGLT2 inhibitors” by a committee of Japanese experts in June 2014. This statement recommended that all patients aged ≥ 65 years starting the drugs within 3 months after the launch should be registered in a post-marketing study. The clinical incidence of adverse reactions to tofogliflozin in patients aged ≥ 65 years differed little from that noted in preapproval clinical trials¹². Recently, prevention of sarcopenia, which might accompany aging in patients with diabetes, has been given special attention¹³. Luseogliflozin was found to produce beneficial changes in

*Corresponding author. Nobuya Inagaki

Tel: +81-75-751-3560

Fax: +81-75-751-4244

E-mail address: inagaki@kuhp.kyoto-u.ac.jp

Received 11 September 2019; revised 17

September 2019; accepted 18 September 2019

Table 1 | Major benefits and risks of sodium–glucose cotransporter 2 inhibitors use

Benefits	Risks
<ul style="list-style-type: none"> • Plasma glucose ↓ • Bodyweight ↓ • Body fat ↓ • Blood pressure ↓ • MACE ↓ • HF hospitalization ↓ • Renal protection 	<ul style="list-style-type: none"> • GTI • UTI • DKA (especially used in type 1 diabetes) • Hypoglycemia (with insulin or SU) • Volume depletion • Skin lesion

DKA, diabetic ketoacidosis; GTI, genital tract infection; HF, heart failure; MACE, major adverse cardiovascular events; SGLT2, sodium–glucose cotransporter 2; SU, Sulfonylureas; UTI, urinary tract infection.


body composition in moderately obese Japanese type 2 diabetes patients, including body fat reduction with minimal muscle reduction, and the signs of sarcopenia were not observed after treatment¹³. The latest edition of “recommendations” updated in July 2019 notes safety concerns, such as diabetic ketoacidosis, especially when the drug is used in type 1 diabetes patients, hypoglycemia when used with sulfonylurea or insulin, volume depletion, skin lesion and urogenital infections.

One of the most common adverse effects of SGLT2 inhibitors is genital tract infection. The rate of developing positive colonization and symptomatic vaginitis by the use of SGLT2 inhibitors in women who attended outpatient clinics in Japan appears to be higher in real-world practice than in clinical trials¹⁴. Meta-analysis of the efficacy and safety profiles of SGLT2 inhibitors in East Asian patients with type 2 diabetes showed that the drugs significantly increased the risk of genital tract infection, but did not significantly increase the risk of hypoglycemia, urinary tract infection, hypotension or mortality¹⁵.

The major benefits and risks of using SGLT2 inhibitors are listed in Table 1; they should be carefully considered for appropriate use of the drugs^{3,5,15}. Reports regarding the clinical use of SGLT2 inhibitors are accumulating for both Asians and non-Asians, and are expected to further clarify the risks and benefits of these drugs.

DISCLOSURE

NI received research grants, honoraria for lectures, and scholarship grants from Mitsubishi Tanabe Pharma, Daiichi Sankyo, AstraZeneca, Kowa Pharmaceutical, Kissei Pharmaceutical, Taisho Toyama Pharmaceutical, Sanofi, Daiichi Sankyo, Mitsubishi Tanabe Pharma, Takeda, Japan Tobacco, Kyowa Kirin, Sumitomo Dainippon Pharma, Astellas Pharma, MSD, Eli Lilly Japan, Ono Pharmaceutical, Sanwa Kagaku Kenkyusho, Nippon Boehringer Ingelheim, Novo Nordisk Pharma, Novartis Pharma and Teijin Pharma. YF received research grants from Japan Diabetes Society, AstraZeneca, Ono Pharmaceutical and FFDR.

Yoshihito Fujita, Nobuya Inagaki* 
Department of Diabetes, Endocrinology
and Nutrition, Graduate School of
Medicine, Kyoto University, Kyoto, Japan

REFERENCES

1. Fujita Y, Inagaki N. Renal sodium glucose cotransporter 2 inhibitors as a novel therapeutic approach to treatment of type 2 diabetes: clinical data and mechanism of action. *J Diabetes Investig* 2014; 5: 265–275.
2. Cai X, Ji L, Chen Y, *et al.* Comparisons of weight changes between sodium-glucose cotransporter 2 inhibitors treatment and glucagon-like peptide-1 analogs treatment in type 2 diabetes patients: A meta-analysis. *J Diabetes Investig* 2017; 8: 510–517.

3. Davies MJ, D'Alessio DA, Fradkin J, *et al.* Management of hyperglycemia in type 2 diabetes, 2018. A Consensus Report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* 2018; 2018(41): 2669–2701.
4. Yabe D, Seino Y, Fukushima M, *et al.* β cell dysfunction versus insulin resistance in the pathogenesis of type 2 diabetes in East Asians. *Curr Diab Rep* 2015; 15: 602.
5. Cai X, Gao X, Yang W, *et al.* No disparity of the efficacy and all-cause mortality between Asian and non-Asian type 2 diabetes patients with sodium-glucose cotransporter 2 inhibitors treatment: a meta-analysis. *J Diabetes Investig* 2018; 9: 850–861.
6. Kaku K, Lee J, Mattheus M, *et al.* Empagliflozin and cardiovascular outcomes in Asian patients with type 2 diabetes and established cardiovascular disease - results From EMPA-REG OUTCOME®. *Circ J* 2017; 81: 227–234.
7. Seino Y, Yabe D, Sasaki T, *et al.* Sodium-glucose cotransporter-2 inhibitor luseogliflozin added to glucagon-like peptide 1 receptor agonist liraglutide improves glycemic control with bodyweight and fat mass reductions in Japanese patients with type 2 diabetes: a 52-week, open-label, single-arm study. *J Diabetes Investig* 2018; 9: 332–340.
8. Tobe K, Suganami H, Kaku K. Sodium-glucose cotransporter 2 inhibitor, tofogliflozin, shows better improvements of blood glucose and insulin secretion in patients with high insulin levels at baseline. *J Diabetes Investig* 2018; 9: 862–869.
9. Okajima F, Nagamine T, Nakamura Y, *et al.* Preventive effect of ipragliflozin on nocturnal hypoglycemia in patients with type 2 diabetes treated with basal-bolus insulin therapy: an open-label, single-center, parallel, randomized control study. *J Diabetes Investig* 2017; 8: 341–345.
10. Kimura T, Sanada J, Shimoda M, *et al.* Switching from low-dose thiazide

- diuretics to sodium-glucose cotransporter 2 inhibitor improves various metabolic parameters without affecting blood pressure in patients with type 2 diabetes and hypertension. *J Diabetes Investig* 2018; 9: 875–881.
11. Yabe D, Yasui A, Ji L, *et al.* Safety and tolerability of empagliflozin in East Asian patients with type 2 diabetes: pooled analysis of phase I-III clinical trials. *J Diabetes Investig* 2019; 10: 418–428.
 12. Utsunomiya K, Shimmoto N, Senda M, *et al.* Safety and effectiveness of tofogliflozin in elderly Japanese patients with type 2 diabetes mellitus: a post-marketing study (J-STEP/EL Study). *J Diabetes Investig* 2017; 8: 766–775.
 13. Sasaki T, Sugawara M, Fukuda M. Sodium-glucose cotransporter 2 inhibitor-induced changes in body composition and simultaneous changes in metabolic profile: 52-week prospective LIGHT (Luseogliflozin: the Components of Weight Loss in Japanese Patients with Type 2 Diabetes Mellitus) Study. *J Diabetes Investig* 2019; 10: 108–117.
 14. Yokoyama H, Nagao A, Watanabe S, *et al.* Incidence and risk of vaginal candidiasis associated with sodium-glucose cotransporter 2 inhibitors in real-world practice for women with type 2 diabetes. *J Diabetes Investig* 2019; 10: 439–445.
 15. Yang L, Zhang L, He H, *et al.* Efficacy and safety of sodium-glucose cotransporter 2 inhibitors in East Asians with type 2 diabetes: a systematic review and meta-analysis. *Diabetes Ther*; 2019; 10: 1921–1934.

Doi: 10.1111/jdi.13150