

LARGE TIME BEHAVIOR FOR A NONLOCAL DIFFUSION EQUATION WITH ABSORPTION AND BOUNDED INITIAL DATA: THE SUBCRITICAL CASE

ARIEL SALORT, JOANA TERRA, AND NOEMI WOLANSKI

ABSTRACT. In this paper we continue our study of the large time behavior of the bounded solution to the nonlocal diffusion equation with absorption

$$\begin{cases} u_t = \mathcal{L}u - u^p & \text{in } \mathbb{R}^N \times (0, \infty), \\ u(x, 0) = u_0(x) & \text{in } \mathbb{R}^N, \end{cases}$$

where $p > 1$, $u_0 \geq 0$ and bounded and

$$\mathcal{L}u(x, t) = \int J(x - y) (u(y, t) - u(x, t)) dy$$

with $J \in C_0^\infty(\mathbb{R}^N)$, radially symmetric, $J \geq 0$ with $\int J = 1$.

Our assumption on the initial datum is that $0 \leq u_0 \in L^\infty(\mathbb{R}^N)$ and

$$|x|^\alpha u_0(x) \rightarrow A > 0 \quad \text{as } |x| \rightarrow \infty$$

This problem was studied in [17, 18] in the supercritical and critical cases $p \geq 1 + 2/\alpha$.

In the present paper we study the subcritical case $1 < p < 1 + 2/\alpha$. More generally, we consider bounded non-negative initial data such that

$$|x|^{\frac{2}{p-1}} u_0(x) \rightarrow \infty \quad \text{as } |x| \rightarrow \infty$$

and prove that

$$t^{\frac{1}{p-1}} u(x, t) \rightarrow \left(\frac{1}{p-1}\right)^{\frac{1}{p-1}} \quad \text{as } t \rightarrow \infty$$

uniformly in $|x| \leq k\sqrt{t}$, for every $k > 0$.

Of independent interest is our study of the positive eigenfunction of the operator \mathcal{L} in the ball B_R in the L^∞ setting that we include in Section 3.

1. INTRODUCTION

In this paper we continue our study of the large time behavior of the solution to the nonlocal diffusion equation with absorption

$$(1.1) \quad \begin{cases} u_t = \mathcal{L}u - u^p & \text{in } \mathbb{R}^N \times (0, \infty), \\ u(x, 0) = u_0(x) & \text{in } \mathbb{R}^N, \end{cases}$$

where $p > 1$, $u_0 \geq 0$ and bounded and

$$(1.2) \quad \mathcal{L}u(x, t) = \int J(x - y) (u(y, t) - u(x, t)) dy$$

with $J \in C_0^\infty(\mathbb{R}^N)$, radially symmetric, $J \geq 0$ with $\int J = 1$.

Our assumption on the initial datum is that $0 \leq u_0 \in L^\infty(\mathbb{R}^N)$ and

$$(1.3) \quad |x|^\alpha u_0(x) \rightarrow A > 0 \quad \text{as } |x| \rightarrow \infty$$

These kind of nonlocal diffusions appear in several applications such as population dynamics, disease propagation, image enhancement, etc (see, for instance, [2, 3, 4, 5, 6, 9, 12, 19]).

2010 *Mathematics Subject Classification.* 35R09, 45K05, 35K57 35B40 .

Key words and phrases. Nonlocal diffusion, Large time behavior.

Supported by CONICET PIP625 Res. 960/12, ANPCyT PICT-2012-0153 and UBACYT X117.

When the kernel J in the nonlocal operator (1.2) satisfies the hypotheses in this paper, the long time behavior of the solutions is closely related to that of the corresponding problem for the heat operator with a diffusivity related to the kernel J (see, for instance, [7, 10, 14, 16, 17, 18]).

In [16] the authors started the study of (1.1) when $u_0 \in L^1(\mathbb{R}^N)$, in the supercritical case $p > 1 + 2/N$. Then, in [17, 18] we studied this problem under assumption (1.3).

The main question we address is what is the interplay between the parameters p , α and the dimension N in the large time behavior of the solution.

In [17, 18] the critical and supercritical cases were studied. This is, we assumed that, either $u_0 \in L^1(\mathbb{R}^N)$ and $p \geq 1 + 2/N$ (completing the results of [16] by considering the critical case), or $0 < \alpha < N$ and $p \geq 1 + 2/\alpha$. Also some intermediate asymptotics for u_0 involving logarithms were considered in [18], always in the supercritical case.

In the present paper we complete our study by considering the subcritical case $1 < p < 1 + 2/\alpha$ that was left open in the previous articles.

The critical value $p_c = 1 + 2/\alpha$ is the one that makes diffusion and absorption of the “same size”. It is interesting to observe that this critical value depends on the size of the initial condition at infinity.

In the supercritical case, diffusion wins and the reaction component disappears in the long run. In the critical case, both diffusion and reaction remain in the time asymptotics (see [16, 17, 18]).

In the present paper we show that, in the subcritical case, only reaction remains in the large time behavior and the solution behaves as that of the equation

$$u_t = -u^p, \quad u(1) = \left(\frac{1}{p-1}\right)^{\frac{1}{p-1}}.$$

This is

$$(1.4) \quad t^{\frac{1}{p-1}} u(x, t) \rightarrow \left(\frac{1}{p-1}\right)^{\frac{1}{p-1}} \quad \text{as } t \rightarrow \infty \quad \text{uniformly in } \{|x| \leq k\sqrt{t}\}$$

for every $k > 0$.

It is interesting to observe that the final profile is independent of the initial datum u_0 as long as it is bounded and satisfies (1.3). In the critical and supercritical cases, both the constant A and the exponent α in (1.3) intervene in the time asymptotics.

Our result is similar to the one obtained by Gmira and Veron in [13] for the heat equation with absorption. As in [13], we get this behavior for any nonnegative and bounded initial datum u_0 such that

$$(1.5) \quad |x|^{\frac{2}{p-1}} u_0(x) \rightarrow \infty \quad \text{as } |x| \rightarrow \infty$$

thus allowing a more general behavior of u_0 at infinity than the one in (1.3).

In this paper we follow the ideas of [13] where the authors constructed subsolutions of separated variables with the right asymptotic behavior. These subsolutions involve the positive eigenfunctions h_R of the laplacian in the balls B_R , normalized so that the $\|h_R\|_{L^\infty(B_R)} = 1$.

The authors make use of the scaling invariance of the laplacian so that $h_R(x) = h_1(x/R)$ and the principal eigenvalue $\lambda_R = R^{-2}\lambda_1$.

One of the main differences when dealing with problem (1.1) is the lack of any scaling invariance of the problem. Nevertheless, a parabolic scaling leads –in the limit of the scaling parameter going to infinity– to the heat equation with diffusivity $A(J) = \frac{1}{2N} \int J(z)|z|^2 dz$. And this fact explains, in a way, the interplay between the time asymptotics of the nonlocal diffusion equation and that of the heat equation with diffusivity $A(J)$, as was made clear in [18].

This scaling property was also the basis for the understanding of the behavior as $|x| \rightarrow \infty$ of the solution to

$$\begin{cases} \mathcal{L}\phi = 0 & \text{in } \mathbb{R}^N \setminus \Omega \\ \phi = 1 & \text{in } \Omega \\ \phi(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty. \end{cases}$$

with Ω an open bounded set, studied in [8].

One of the main contributions of the present paper is a thorough study of the positive eigenfunction H_R to the nonlocal operator \mathcal{L} in the ball B_R with Dirichlet boundary conditions $H_R = 0$ in $\mathbb{R}^N \setminus B_R$, normalized so that $\|H_R\|_{L^\infty} = 1$.

This study was initiated in [11] where the existence of a principal eigenvalue Λ_R associated to a positive eigenfunction was proved. Moreover, in [11] the authors proved that, asymptotically the principal eigenvalue behaves as that of the laplacian with diffusivity $A(J)$. This is,

$$R^2 \Lambda_R \rightarrow A(J) \lambda_1 \quad \text{as } R \rightarrow \infty.$$

In [11] the authors also studied the associated eigenfunction in the L^2 setting and they proved that, after rescaling to the unit ball with an L^2 normalization, one gets convergence in L^2 to the positive eigenfunction of the laplacian in the unit ball with Dirichlet boundary conditions and unit L^2 - norm.

In the present paper, due to the application to the study of the asymptotics of (1.1) we have in mind, we are interested in a different normalization and convergence. Namely, we normalize so that the L^∞ - norm is preserved and prove uniform convergence in the unit ball.

In order to get this kind of compactness, the arguments in [11] cannot be applied. Instead, we get uniform bounds for the derivatives of the rescaled eigenfunctions $\tilde{H}_R(x) = H_R(Rx)$, on smaller balls B_r with $0 < r < 1$, by using an integral representation formula for H_R and a precise decay in terms of R of H_R in a neighborhood of the boundary of B_R . To this end, we construct an upper barrier. This barrier also allows to get uniform smallness of the rescaled eigenfunctions and their limits in a neighborhood of ∂B_1 that gives, in particular, uniform convergence in the whole ball. The uniform limit is then identified as being h_1 , the positive eigenfunction of the laplacian in the unit ball with Dirichlet boundary conditions and unit L^∞ - norm.

We believe that the results concerning the eigenfunctions H_R are of independent interest.

The paper is organized as follows. In Section 2 we state the results of [11] on the principal eigenvalue of the operator \mathcal{L} with Dirichlet boundary conditions set in the ball B_R . Then, in Section 3 we perform our study of the eigenfunctions associated to the principal eigenvalues in the L^∞ setting. In Section 4 we construct a subsolution to (1.1) by following the ideas of [13] for the heat equation. Due to the lack of any regularizing effect of the nonlocal operator, we need to prove that $\inf_{B_R} u(\cdot, t) > 0$ for every $R > 0$, $t > 0$ (Lemma 4.1). Finally, in Section 5 we prove our main result, namely that (1.4) is satisfied.

2. DEFINITIONS AND PRELIMINARY RESULTS

In this section we discuss notation and basic definitions. Moreover we state some previous results on the first eigenvalue of the nonlocal problem with Dirichlet boundary conditions in a ball.

Let $R > 0$ and define the ball of radius R as

$$B_R = \{x \in \mathbb{R}^n : |x| < R\}$$

We denote by λ_R the first eigenvalue of the laplacian in B_R . That is, λ_R verifies that there is a solution to the following problem,

$$(2.1) \quad \begin{cases} -\Delta u = \lambda_R u & \text{in } B_R, \\ u = 0 & \text{on } \partial B_R, \\ u > 0 & \text{in } B_R. \end{cases}$$

We know that λ_R is simple. Let us call h_R the associated eigenfunction satisfying

$$(2.2) \quad 0 < h_R(x) \leq 1 = \max_{x \in B_R} h_R(x) \quad \text{in } B_R.$$

It is well known that, due to the scaling of the laplacian there holds that, $\lambda_R = R^{-2} \lambda_1$.

We now consider the nonlocal eigenvalue problem,

$$(2.3) \quad \begin{cases} -\mathcal{L}u(x) = \Lambda_R u(x) & \text{in } B_R \\ u = 0 & \text{in } \mathbb{R}^N \setminus B_R, \\ u > 0 & \text{in } B_R. \end{cases}$$

where $\mathcal{L}u(x) = \int J(x-y)(u(y,t) - u(x,t)) dy$ and $J \in C_0^\infty(\mathbb{R}^N)$ is radially symmetric and $J \geq 0$ with $\int J = 1$.

It was proved in [11] that such an eigenvalue exists, it is simple and moreover,

$$(2.4) \quad \Lambda_R \sim A(J) \frac{\lambda_1}{R^2} \quad \text{as } R \rightarrow +\infty$$

with

$$(2.5) \quad A(J) = \frac{1}{2N} \int_{\mathbb{R}^N} J(z)|z|^2 dz.$$

This is,

$$\Lambda_R = A(J)(1 + o(1)) \frac{\lambda_1}{R^2} \quad \text{as } R \rightarrow +\infty.$$

Consequently, the first eigenvalue Λ_R for the nonlocal problem (2.3) behaves asymptotically as the first eigenvalue λ_R of the laplacian (2.1), as R tends to infinity.

Moreover, in [11] the authors proved that Λ_R is given variationally as

$$\Lambda_R = \inf_{\substack{0 \neq u \in L^2(B_R) \\ u=0 \text{ in } B_R^c}} \frac{1}{2} \frac{\int \int J(x-y)(u(x) - u(y))^2 dx dy}{\int u^2(x) dx}.$$

3. SOME RESULTS ON THE EIGENFUNCTIONS

In this section we study the eigenfunctions of the nonlocal problem in the ball B_R .

The eigenfunction problem was studied in [11] in the L^2 setting. This is, in [11] the authors consider the family of eigenfunctions normalized as to have the $L^2(B_R)$ -norm equal to 1 and prove that, when properly rescaled, they converge to the unique positive eigenfunction of the laplacian in the unit ball with $L^2(B_1)$ -norm equal to 1.

In the present paper we are interested in the family H_R of positive eigenfunctions normalized so that the $L^\infty(B_R)$ -norm is 1. We prove that, when properly rescaled, they converge to the unique positive eigenfunction of the laplacian in the unit ball with the same normalization. The convergence is uniform in the unit ball.

In order to get our result, we cannot use the compactness argument of [11] that holds only in L^p for $p < \infty$. Instead, we use Arzelà-Ascoli. To this end, we get uniform estimates of the derivatives of the rescaled eigenfunctions on compact subsets of the unit ball. The argument is delicate and uses a precise decay, in terms of R , of H_R in a neighborhood of the boundary of B_R . This decay is obtained by comparison with a supersolution that we construct to this end. In this way we obtain uniform convergence on compact subsets of the unit ball.

The supersolution also allows us to prove that the rescaled eigenfunctions \tilde{H}_R are smaller than any positive constant in a neighborhood of ∂B_1 if R is large. This, in turn, gives that the convergence is uniform in the unit ball to a function that is continuous in the closure and vanishes on the boundary. This limit function is therefore h_1 .

In this way, we get our main result in this section. Namely,

Theorem 3.1. *Let \mathcal{L} the operator in (1.2). Let $\Lambda_R \in \mathbb{R}$, $H_R \in C(\overline{B_R})$, $H_R = 0$ in $\mathbb{R}^N \setminus B_R$, be the unique solution to*

$$\begin{cases} -\mathcal{L}H(x) = \Lambda_R H(x) & \text{in } B_R, \\ H(x) = 0 & \text{in } \mathbb{R}^N \setminus B_R, \\ H(x) > 0 & \text{in } B_R \end{cases}$$

obtained in [11], with the normalization $0 < H(x) \leq 1 = \|H_R\|_{L^\infty(B_R)}$ for $x \in B_R$.

Let $\tilde{H}_R(x) = H_R(Rx)$ for $x \in B_1$ and $h_1 \in C^\infty(B_1) \cap C(\bar{B}_1)$ the positive eigenfunction of the laplacian in the unit ball such that $\|h_1\|_{L^\infty(B_1)} = 1$. Then,

$$\tilde{H}_R \rightarrow h_1 \quad (R \rightarrow \infty) \quad \text{uniformly in } B_1.$$

For the proof of this theorem we need a couple of lemmas.

Lemma 3.2. *Let $R_n \rightarrow \infty$ be such that $\tilde{H}_{R_n} \rightarrow H$ in $L^1_{loc}(B_1)$. Then H is a solution to,*

$$\begin{cases} -\Delta H = \lambda_1 H, & \text{in } B_1 \\ H = 0 & \text{on } \partial B_1. \end{cases}$$

Proof. In order to prove that H is a weak solution to the equation, we let $\phi \in C_0^\infty(B_1)$ and,

$$\begin{aligned} A(J) \int_{B_1} H(x) \Delta \phi(x) dx &= \\ &= A(J) \int_{B_1} \tilde{H}_{R_n}(x) \Delta \phi(x) dx + A(J) \int_{B_1} (H(x) - \tilde{H}_{R_n}(x)) \Delta \phi(x) dx \\ &= \int_{B_1} \tilde{H}_{R_n}(x) R^2 (J_R * \phi(x) - \phi(x)) dx + A(J) \int_{B_1} (H(x) - \tilde{H}_{R_n}(x)) \Delta \phi(x) dx \\ &\quad - \int_{B_1} \tilde{H}_{R_n}(x) (R^2 (J_R * \phi(x) - \phi(x)) - A(J) \Delta \phi(x)) dx \\ &= R^2 \int_{B_1} (J_R * \tilde{H}_{R_n}(x) - \tilde{H}_{R_n}(x)) \phi(x) dx \\ &\quad + A(J) \int_{B_1} (H(x) - \tilde{H}_{R_n}(x)) \Delta \phi(x) dx - \int_{B_1} \tilde{H}_{R_n}(x) O(R^{-3}) dx \\ &= -R^2 \Lambda_R \int_{B_1} \tilde{H}_{R_n}(x) \phi(x) dx + A(J) \int_{B_1} (H(x) - \tilde{H}_{R_n}(x)) \Delta \phi(x) dx \\ &\quad - \int_{B_1} \tilde{H}_{R_n}(x) O(R^{-3}) dx. \end{aligned}$$

Since $\tilde{H}_{R_n} \rightarrow H$ strongly in $L^2(B_1)$ and $R^2 \Lambda_R \rightarrow A(J) \lambda_1$, by taking limit as n tends to infinity we obtain,

$$A(J) \int_{B_1} H(x) \Delta \phi(x) dx = -\lambda_1 A(J) \int_{B_1} H(x) \phi(x) dx,$$

that is, H satisfies the equation $-\Delta H = \lambda_1 H$, in B_1 . □

Our next result is the construction of a barrier for H_R .

Lemma 3.3. *Let h_1 be the positive eigenfunction corresponding to the first eigenvalue λ_1 of the laplacian in B_1 with Dirichlet boundary conditions and the normalization $1 = \max_{x \in B_1} h_1(x)$. Let us consider the function*

$$v(x) = h_1\left(\frac{x}{2R}\right) \quad \text{for } x \in B_{2R}.$$

There exists $C > 0$, $R_0 > 0$ such that

$$C \mathcal{L}v(x) \leq \mathcal{L}H_R \quad \text{in } B_R \quad \text{if } R \geq R_0.$$

Proof. Assume $R \geq 1$. By using Taylor's expansion and the symmetry of J we get for $x \in B_R$,

$$\mathcal{L}v(x) = A(J) \Delta v(x) + O\left(\max_{|\beta|=4} \|D^\beta v\|_{L^\infty(B_{R+1})}\right).$$

Then,

$$\begin{aligned}
(3.1) \quad \mathcal{L}v &= \frac{A(J)}{4} R^{-2} \Delta h_1 \left(\frac{x}{2R} \right) + O(R^{-4}) \\
&= -\lambda_1 \frac{A(J)}{4} R^{-2} h_1 \left(\frac{x}{2R} \right) + O(R^{-4}) \\
&\leq -\frac{1}{8} \lambda_1 A(J) R^{-2} h_1 \left(\frac{x}{2R} \right)
\end{aligned}$$

if R is large.

Here we have used that there exists a positive constant c such that,

$$(3.2) \quad c < h_1 \left(\frac{x}{2R} \right) \quad x \in B_R.$$

Finally, since $\lambda_1 A(J) R^{-2} = \Lambda_R + o(1)$, we get for R large enough,

$$\mathcal{L}v \leq -\frac{1}{16} \Lambda_R h_1 \left(\frac{x}{2R} \right) \leq -\frac{c}{16} \Lambda_R \leq -\frac{c}{16} \Lambda_R H_R = \frac{c}{16} \mathcal{L}H_R$$

since $0 \leq H_R \leq 1$. □

Recall that h_1 is radially symmetric, radially decreasing, smooth with $h_1(0) = 1$. Let η such that $h_1(x) = \eta(|x|)$.

Now we use the supersolution constructed in Lemma 3.3 in order to bound H_R . There holds,

Lemma 3.4. *Let $\eta(|x|) = h_1(x)$ with h_1 as in Lemma 3.3. There exist constants $C, C_0 > 0$ and $R_0 > 0$ such that,*

$$H_R(x) \leq C \left\{ \eta \left(\frac{|x|}{2R} \right) - \eta \left(\frac{1}{2} \right) + \frac{C_0}{R} \right\} \quad \text{if } R \geq R_0.$$

Proof. In Lemma 3.3 we found a constant $C > 0$ and $R_0 > 0$ such that, for any $C_0 \in \mathbb{R}$, $R \geq R_0$, the function

$$w(x) = C \left\{ \eta \left(\frac{|x|}{2R} \right) - \eta \left(\frac{1}{2} \right) + \frac{C_0}{R} \right\}$$

satisfies

$$\mathcal{L}w \leq \mathcal{L}H_R \quad \text{in } B_R.$$

In order to be able to apply the comparison principle we need to show that, for some constant C_0 , there holds that

$$(3.3) \quad w \geq 0 \quad \text{in } \{x \in \mathbb{R}^N \setminus B_R / \text{dist}(x, B_R) < 1\} = \{R \leq |x| < R+1\}.$$

And, in fact (3.3) holds if $C_0 \geq \|\eta'\|_{L^\infty(0,1)}$.

Finally, by applying the Comparison Principle, the lemma is proved. □

From this lemma we get the following corollary that will be used to bound the derivatives of \tilde{H}_R .

Corollary 3.5. *There exists a constant $K > 0$ such that*

$$J * H_R \leq \frac{K}{R} \quad \text{in } \{R \leq |x| < R+1\}.$$

Proof. Let $R \leq |x| < R+1$. Then, if $J(x-y)H_R(y) \neq 0$, there holds that $R-1 \leq |y| < R$. Therefore,

$$H_R(y) \leq w(y) = C \left\{ \eta \left(\frac{|y|}{2R} \right) - \eta \left(\frac{1}{2} \right) + \frac{C_0}{R} \right\} \leq \frac{K}{R}$$

for a certain constant $K > 0$ and,

$$(J * H_R)(x) = \int J(x-y)H_R(y) dy \leq \frac{K}{R}.$$

□

In order to prove our main result in this section, we will use an integral representation formula for H_R . To this end, let us recall some results on the fundamental solution to the operator $\partial_t - \mathcal{L}$.

In [7] the authors found that the fundamental solution of the nonlocal operator $\partial_t - L$ in the whole space, is

$$F(x, t) = e^{-t}\delta(x) + \omega(x, t)$$

where δ is the Dirac mass at the origin in \mathbb{R}^N and ω is a smooth function.

Then, in [18] pointwise and integral estimates for w and its derivatives were obtained. In particular,

$$(3.4) \quad |\nabla\omega(x, t)| \leq C \frac{t}{|x|^{N+3}},$$

and,

$$(3.5) \quad \int_{\mathbb{R}^N} |\nabla\omega(x, t)| \leq Ct^{-\frac{1}{2}}.$$

We can now prove our main result in this section.

Proof of Theorem 3.1. The proof follows from the Arzelá-Ascoli Theorem.

In order to get uniform estimates of the derivatives of \tilde{H}_R let us observe that the first eigenfunction of (2.3) is the unique bounded solution of the following non-homogeneous equation defined in the whole \mathbb{R}^N ,

$$(3.6) \quad \begin{cases} w_t - \mathcal{L}w = \Lambda_R w - \mathcal{X}_{B_R^c}(J * w) & \text{in } \mathbb{R}^N \times (0, \infty), \\ w(x, 0) = H_R(x) & \text{in } \mathbb{R}^N. \end{cases}$$

As the solution of (3.6) is defined in the whole space, it can be expressed in terms of the fundamental solution $F = F(x, t)$ by means of the variation of constants formula. Thus, for $t \geq 0$ we have

$$(3.7) \quad \begin{aligned} H_R(x) &= e^{-t}H_R(x) + \int_{\mathbb{R}^N} \omega(x-y, t)H_R(y) dy + \Lambda_R \left(\int_0^t e^{-(t-s)} ds \right) H_R(x) \\ &- \left(\int_0^t e^{-(t-s)} ds \right) \mathcal{X}_{B_R^c}(x)(J * H_R)(x) + \Lambda_R \int_0^t \int_{\mathbb{R}^N} \omega(x-y, t-s)H_R(y) dy ds \\ &- \int_0^t \int_{B_R^c} \omega(x-y, t-s)(J * H_R)(y) dy ds. \end{aligned}$$

For $x \in B_R$, there holds that $\mathcal{X}_{B_R^c}(x) = 0$. Thus, we can rewrite (3.7) for $x \in B_R$ as,

$$(3.8) \quad \begin{aligned} (1 - e^{-t})(1 - \Lambda_R)H_R(x) &= \\ &= \int_{\mathbb{R}^N} \omega(x-y, t)H_R(y) dy + \Lambda_R \int_0^t \int_{\mathbb{R}^N} \omega(x-y, t-s)H_R(y) dy ds \\ &- \int_0^t \int_{B_R^c} \omega(x-y, t-s)(J * H_R)(y) dy ds. \end{aligned}$$

Observe that we are free to select the parameter t in expression (3.8).

Let us now rescale the identity (3.8). We have,

$$\begin{aligned} (1 - e^{-t})(1 - \Lambda_R)\tilde{H}_R(x) &= \\ &= \int_{\mathbb{R}^N} \omega(Rx-y, t)H_R(y) dy + \Lambda_R \int_0^t \int_{\mathbb{R}^N} \omega(Rx-y, t-s)H_R(y) dy ds \\ &- \int_0^t \int_{B_R^c} \omega(Rx-y, t-s)(J * H_R)(y) dy ds \\ &:= (i) + (ii) - (iii). \end{aligned}$$

In order to bound the derivatives of (i), (ii) and (iii) we will choose the value $t = R^2$. First, let us estimate the derivative of (i). By (3.5), since $0 \leq H_R \leq 1$, it follows that

$$(3.9) \quad \begin{aligned} \left| \nabla \int_{\mathbb{R}^N} \omega(Rx - y, t) H_R(y) dy \right| &= R \left| \int_{\mathbb{R}^N} \nabla \omega(Rx - y, t) H_R(y) dy \right| \\ &\leq R \int_{\mathbb{R}^N} |\nabla \omega(y, t)| dy \\ &\leq CRt^{-\frac{1}{2}} = C. \end{aligned}$$

Similarly, since $\Lambda_R \leq CR^{-2}$,

$$(3.10) \quad \begin{aligned} \Lambda_R \left| \nabla \int_0^t \int_{\mathbb{R}^N} \omega(Rx - y, t - s) H_R(y) dy ds \right| \\ \leq \Lambda_R R \int_0^t \int_{\mathbb{R}^N} |\nabla \omega(y, t - s)| dy ds \\ \leq CR^{-1} \int_0^t (t - s)^{-\frac{1}{2}} ds \\ \leq CR^{-1} t^{\frac{1}{2}} = C. \end{aligned}$$

Now, by using the pointwise estimate (3.4), Corollary 3.5 and the fact that $\text{supp}(J * H_R) = B_{R+1}$ we can bound the derivative of (iii) as

$$\begin{aligned} \left| \nabla \int_0^t \int_{B_R^c} \omega(Rx - y, t - s) (J * H_R)(y) dy ds \right| &= \\ &= R \left| \int_0^t \int_{B_R^c} \nabla \omega(Rx - y, t - s) (J * H_R)(y) dy ds \right| \\ &\leq C \left| \int_0^t \int_{R < |y| < R+1} \frac{t - s}{|Rx - y|^{N+3}} dy ds \right| \\ &\leq Ct^2 \int_{R < |y| < R+1} \frac{1}{|Rx - y|^{N+3}} dy. \end{aligned}$$

Assume now, $|x| \leq r$ with $0 < r < 1$. Then, if $|y| > R$ we get that $|Rx - y| \geq R(1 - r)$ and then,

$$(3.11) \quad \begin{aligned} \left| \nabla \int_0^t \int_{B_R^c} \omega(x - y, t - s) (J * H_R)(y) dy ds \right| &\leq \\ &\leq CR^{-N-3} t^2 \frac{1}{(1 - r)^{N+3}} |\{R < |y| < R + 1\}| \\ &\leq C_r R^{-4} t^2 = C_r. \end{aligned}$$

Thus, since $(1 - e^{-R^2})(1 - \Lambda_R) \geq \alpha_0 > 0$ for $R \geq R_0$, we conclude that for every $0 < r < 1$ there exists $C > 0$ such that,

$$\sup_{|x| \leq r} |\nabla \tilde{H}_R(x)| \leq C$$

if $R \geq R_0$.

We can apply Arzelà-Ascoli on every ball B_r with $0 < r < 1$ to get, for every sequence $R_n \rightarrow \infty$ a subsequence \tilde{H}_{R_n} uniformly convergent in B_r . Then, a diagonal argument gives a subsequence uniformly convergent on every compact subset of B_1 to a function H . By the previous lemmas, we know that H is a solution to

$$-\Delta H = \lambda_1 H \quad \text{in } B_1.$$

Moreover, $0 \leq H \leq 1$. Let us see that $H \in C(\overline{B_1})$ with $H = 0$ on ∂B_1 . In fact, we show that the subsequence $\tilde{H}_{R_{n_k}}$ converging to H uniformly on compact subsets of B_1 is actually uniformly convergent in B_1 . In fact, we use Lemma 3.4 to get for $\varepsilon > 0$,

$$\tilde{H}_R(x) \leq C \left(\eta \left(\frac{|x|}{2} \right) - \eta \left(\frac{1}{2} \right) + \frac{C_0}{R} \right) < \frac{\varepsilon}{2}$$

if $1 - |x| < \delta_0$ and $R \geq R_0$.

On the other hand, for every $x \in B_1$, by taking limit as $k \rightarrow \infty$ we find that,

$$H(x) \leq C \left(\eta \left(\frac{|x|}{2} \right) - \eta \left(\frac{1}{2} \right) \right) < \frac{\varepsilon}{2}$$

if $1 - |x| < \delta_0$.

Observe that, in particular, $H \in C(\overline{B_1})$ with $H = 0$ on ∂B_1 .

Then,

$$|\tilde{H}_{R_{n_k}}(x) - H(x)| \leq \varepsilon \quad \text{if } |x| > 1 - \delta_0, \quad k \geq k_0.$$

On the other hand, due to the uniform convergence of $\tilde{H}_{R_{n_k}}$ to H in $\overline{B_{1-\delta_0}}$,

$$|\tilde{H}_{R_{n_k}}(x) - H(x)| \leq \varepsilon \quad \text{if } |x| \leq 1 - \delta_0, \quad k \geq k_1.$$

So that, the convergence is uniform in B_1 , $H \in C(\overline{B_1})$ with $H = 0$ on ∂B_1 . So that, $H = h_1$ is independent of the subsequence and the theorem is proved. \square

4. BACK TO THE EVOLUTIONARY PROBLEM. CONSTRUCTION OF A BARRIER.

In this section we construct a barrier for the nonlocal problem which is similar to the one constructed in [13] for the laplacian. This barrier is a function of separated variables involving the eigenfunctions H_R studied in Section 3.

In order to be able to further analyze our solution u , we state a result that is needed because of the lack of a regularizing effect of the nonlocal diffusion equation.

Lemma 4.1. *Let $0 \leq u_0 \in L^\infty$, $u_0 \not\equiv 0$. Then, for every $R > 0$, $t > 0$,*

$$(4.1) \quad \inf_{x \in B_R} u(x, t) > 0.$$

Proof. We recall some results that can be found, for instance, in [17]. First, $u \in L^\infty$ and bounded by $\|u_0\|_\infty$. Moreover, $u \geq 0$ since $v \equiv 0$ is a solution to the equation and a comparison principle for bounded solutions holds (see, for instance [15]).

Moreover, $u(x, t) > 0$ for every $x \in \mathbb{R}^N$, $t > 0$. In fact, let $A \geq 1 + \|u_0\|_\infty^{p-1}$. Then, since $0 \leq u \leq \|u_0\|_\infty$,

$$u_t + Au \geq u_t + u + u^p = J * u.$$

Thus,

$$(4.2) \quad u(x, t) \geq e^{-At} u_0(x) + \int_0^t e^{-A(t-s)} (J * u(\cdot, s))(x) dx$$

so that, if $u(x, t) = 0$ for some $x \in \mathbb{R}^N$, $t > 0$ there holds,

$$0 \geq \int_0^t e^{-A(t-s)} (J * u(\cdot, s))(x) dx \geq 0.$$

We deduce that $u(y, s) = 0$ in $B_1(x) \times (0, t)$ and, since \mathbb{R}^N is connected, a continuation argument gives that $u = 0$ in $\mathbb{R}^N \times (0, t)$. But, by (4.2),

$$u(x, t) \geq e^{-At} u_0(x)$$

and $u_0 \not\equiv 0$.

Therefore, $u(x, t) > 0$ in $\mathbb{R}^N \times (0, \infty)$.

Let us now prove (4.1). In not, there exists a sequence $\{x_n\} \subset B_R$ such that $u(x_n, t) \rightarrow 0$. Without loss of generality we may assume that $x_n \rightarrow \bar{x} \in \overline{B_R}$. Going back to (4.2) and using that $J * u(\cdot, s)$ is a continuous function in \mathbb{R}^N we get

$$0 \leftarrow u(x_n, t) \geq \int_0^t e^{-A(t-s)} (J * u(\cdot, s))(x_n) dx \rightarrow \int_0^t e^{-A(t-s)} (J * u(\cdot, s))(\bar{x}) dx.$$

We deduce that $u = 0$ in $B_1(\bar{x}) \times (0, t)$, a contradiction. \square

Now, we construct the barrier.

Lemma 4.2. *Let Λ_R be the principal eigenvalue of (2.3) in the ball B_R and H_R the positive eigenfunction with the normalization $\|H_R\|_{L^\infty(B_R)} = 1$. Assume $0 \leq u_0 \in L^\infty(\mathbb{R}^N)$ and let u be the unique bounded solution of (1.1). Then, the following inequality holds in $B_R \times (0, +\infty)$:*

$$(4.3) \quad u(x, t) \geq \psi_R(t)H_R(x),$$

where ψ_R is the solution to

$$(4.4) \quad \begin{cases} \frac{d}{dt}\psi_R + \Lambda_R\psi_R + \psi_R^p = 0 & \text{in } (0, \infty) \\ \psi_R(0) = c = \inf_{x \in B_R} \frac{u_0(x)}{H_R(x)}. \end{cases}$$

Proof. We set $w(x, t) = \psi_R(t)H_R(x)$. Then, for $x \in B_R$,

$$\begin{aligned} w_t - \mathcal{L}w + w^p &= H_R \frac{d}{dt}\psi_R - \psi_R \mathcal{L}H_R + \psi_R^p H_R^p \\ &= H_R \frac{d}{dt}\psi_R + \psi_R \Lambda_R H_R + \psi_R^p H_R^p \\ &= H_R \left(\frac{d}{dt}\psi_R + \Lambda_R\psi_R + \psi_R^p \right) + H_R \psi_R ((H_R \psi_R)^{p-1} - \psi_R^{p-1}). \end{aligned}$$

Since ψ_R satisfies (4.4), $0 \leq H_R \leq 1$ and $p \geq 1$ we deduce that,

$$w_t - \mathcal{L}w + w^p \leq 0 \quad \text{for } x \in B_R.$$

As $w(x, 0) = \psi_R(0)H_R(x) \leq u_0(x)$ and $w(x, t) = 0$ in $B_R^c \times (0, \infty)$, we deduce by the comparison principle for sub- and super-solutions on bounded sets that,

$$w(x, t) \leq u(x, t).$$

\square

Remark 4.3. The function ψ can be computed explicitly (see [13]). In fact, if $c > 0$,

$$(4.5) \quad \psi_R(t) = \left(\frac{\Lambda_R}{(1 + c^{1-p}\Lambda_R)e^{\Lambda_R(p-1)t} - 1} \right)^{\frac{1}{p-1}}.$$

The following technical lemma was proved in [13]. This result will be used later on in Section 5 in order to obtain the region where we can identify the asymptotic behavior of u .

Lemma 4.4 (Gmira and Veron, Lemma 2.2, [13]). *Set $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that*

$$\lim_{y \rightarrow \infty} y^2 \varphi(y) = \infty.$$

Then, there exists a nondecreasing function R from \mathbb{R}^+ into \mathbb{R}^+ such that

$$(4.6) \quad \lim_{y \rightarrow \infty} \frac{y}{R^2(y)} = 0, \quad \lim_{y \rightarrow \infty} y \varphi(R(y)) = \infty.$$

Remark 4.5. In [13] the function φ was assumed continuous. But it is easy to see that this assumption is not needed.

Now, we prove a key lemma. Once again the goal is to establish a lower bound for $u(\cdot, t)$ by constructing an appropriate auxiliary function $\varphi(R)$. This function will be used as an initial condition for the function ψ_R from Lemma 4.2 in the proof of Theorem 5.1.

Proposition 4.6. *Suppose $0 \leq u_0 \in L^\infty(\mathbb{R}^N)$ is such that*

$$(4.7) \quad |x|^{\frac{2}{p-1}} u_0(x) \rightarrow \infty \quad \text{as } |x| \rightarrow \infty$$

and let u be the bounded solution to (1.1). Then, for any $t > 0$ the following equivalent properties hold:

- (i) $\lim_{|x| \rightarrow \infty} |x|^{\frac{2}{p-1}} u(x, t) = \infty$.
- (ii) $\lim_{R \rightarrow \infty} R^{\frac{2}{p-1}} \inf_{|x| \leq R} u(x, t) = \infty$.
- (iii) *There exists a positive, non-increasing, real-valued function φ such that*

$$(4.8) \quad \lim_{r \rightarrow \infty} r^{\frac{2}{p-1}} \varphi(r) = \infty$$

and,

$$u(x, t) \geq \varphi(R) H_R(x), \quad \forall x \in B_R.$$

Proof. By (4.2), for every $t > 0$,

$$|x|^{\frac{2}{p-1}} u(x, t) \geq e^{-At} |x|^{\frac{2}{p-1}} u_0(x) \rightarrow \infty \quad \text{as } |x| \rightarrow \infty.$$

Thus, (i) holds.

Let us see that (i) \Rightarrow (ii).

If not, there exist $R_n \rightarrow \infty$ and a constant $C > 0$ such that

$$R_n^{\frac{2}{p-1}} \inf_{B_{R_n}} u(\cdot, t) \leq C.$$

This in turn implies that there exists $x_n \in B_{R_n}$ such that

$$(4.9) \quad R_n^{\frac{2}{p-1}} u(x_n, t) \leq 2C.$$

If there exist $R_0 > 0$ and a subsequence R_{n_k} such that $\{x_{n_k}\} \subset B_{R_0}$ we would have, by (4.9) and Lemma 4.1,

$$2C \geq R_{n_k}^{\frac{2}{p-1}} u(x_{n_k}, t) \geq R_{n_k}^{\frac{2}{p-1}} \inf_{B_{R_0}} u(\cdot, t) \rightarrow \infty \quad \text{as } k \rightarrow \infty$$

which is a contradiction. Therefore, $|x_n| \rightarrow \infty$ as $n \rightarrow \infty$. But then, since $x_n \in B_{R_n}$, by (i),

$$2C \geq R_n^{\frac{2}{p-1}} u(x_n, t) \geq |x_n|^{\frac{2}{p-1}} u(x_n, t) \rightarrow \infty \quad \text{as } n \rightarrow \infty$$

which again is a contradiction. So, (ii) holds.

(ii) \Rightarrow (iii). We define for $R > 0$

$$(4.10) \quad \varphi(R) = \inf_{|x| \leq R} \frac{u(x, t)}{H_R(x)},$$

where H_R is the positive eigenfunction of (2.3) with $\|H_R\|_\infty = 1$. As $u(x, t)/H_R \geq u(x, t)$ in B_R , there holds that $\varphi(R)$ is positive.

From (4.10) we have in B_R ,

$$u(x, t) \geq \varphi(R) H_R(x),$$

and, as $0 \leq H_R(x) \leq 1$,

$$R^{\frac{2}{p-1}} \varphi(R) \geq R^{\frac{2}{p-1}} \inf_{|x| \leq R} u(x, t) \rightarrow \infty \quad \text{as } R \rightarrow \infty,$$

by (ii). So that, (iii) holds.

(iii) \Rightarrow (ii). In fact, by Theorem 3.1 we know that

$$\tilde{H}_R(x) \rightarrow h_1 \quad \text{uniformly in } B_{1/2}.$$

Since $h_1(x) \geq \beta > 0$ in $B_{1/2}$, there holds that

$$\tilde{H}_R(x) \geq \frac{\beta}{2} \quad \text{in } B_{1/2}$$

if $R \geq R_0$.

This is,

$$H_R(x) \geq \frac{\beta}{2} \quad \text{in } B_{R/2}$$

if $R \geq R_0$. Hence,

$$u(x, t) \geq \varphi(R)H_R(x) \geq \frac{\beta}{2}\varphi(R) \quad \text{in } B_{R/2}$$

if $R \geq R_0$.

Multiplying by $R^{\frac{2}{p-1}}$, taking infimum over $B_{R/2}$ and letting $R \rightarrow \infty$ gives (ii).

(ii) trivially implies (i). \square

Remark 4.7. Observe that the function $\varphi(R)$ depends on $t > 0$.

5. MAIN RESULT

In this section we prove our main result. This is, we obtain the large time behavior of u in the subcritical case $1 < p < 1 + 2/\alpha$.

Theorem 5.1. *Suppose $0 \leq u_0 \in L^\infty(\mathbb{R}^N)$ satisfies (4.7). Let $u(x, t)$ be the bounded solution of (1.1). Then,*

$$\lim_{t \rightarrow \infty} t^{\frac{1}{p-1}} u(x, t) = \left(\frac{1}{p-1} \right)^{\frac{1}{p-1}},$$

uniformly on the sets

$$E_k = \{x \in \mathbb{R}^N : |x| \leq k\sqrt{t}\},$$

where k is an arbitrary constant.

Proof. From Proposition 4.6, by considering $u(x, t)$ for $t \geq t_0 > 0$ we deduce that there is no loss of generality in assuming that there exists a nondecreasing function $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ satisfying (4.8) such that,

$$u_0(x) \geq \varphi(R)H_R(x) \quad \forall x \in B_R.$$

From Lemma 4.2 we have that $u(x, t) \geq H_R(x)\psi_R(t)$ in $B_R \times \mathbb{R}^+$, where ψ_R is the solution of

$$(5.1) \quad \begin{cases} \frac{d}{dt}\psi_R + \Lambda_R\psi_R + \psi_R^p = 0 & \text{in } (0, \infty) \\ \psi_R(0) = \varphi(R). \end{cases}$$

By (2.4) the principal eigenvalue of (2.3) we can written in the form

$$(5.2) \quad \Lambda_R = C_R \frac{\lambda_1}{R^2} \quad \text{as } R \rightarrow +\infty,$$

where λ_1 is the principal eigenvalue of (2.1) and $C_R \rightarrow A(J)$ as $R \rightarrow \infty$ with $A(J)$ given by (2.5). By using (5.2) and (4.5) we have

$$\begin{aligned} t^{\frac{1}{p-1}}\psi_R(t) &= t^{\frac{1}{p-1}} \left(\frac{\Lambda_R}{(1 + \varphi^{1-p}(R)\Lambda_R)e^{\Lambda_R(p-1)t} - 1} \right)^{\frac{1}{p-1}} \\ &= \frac{(C_R t \lambda_1 R^{-2})^{\frac{1}{p-1}} e^{-C_R \lambda_1 t R^{-2}}}{(1 + C_R \lambda_1 \varphi^{1-p}(R) R^{-2} - e^{-C_R \lambda_1 (p-1)t R^{-2}})^{\frac{1}{p-1}}}. \end{aligned}$$

By using the Taylor expansion for $e^{-C_R \lambda_1 (p-1)tR^{-2}}$ at the origin we get,

$$(5.3) \quad t^{\frac{1}{p-1}} \psi_R(t) = \left(\frac{\frac{C_R t \lambda_1}{R^2}}{\frac{C_R \lambda_1}{R^2 \varphi^{p-1}(R)} + \frac{C_R \lambda_1 (p-1)t}{R^2} + O(R^{-4}t^2)} \right)^{\frac{1}{p-1}} e^{-C_R \lambda_1 t R^{-2}}.$$

Now, as $\lim_{R \rightarrow \infty} R^2 \varphi^{p-1}(R) = \infty$ we deduce from Lemma 4.4 that there exists a nondecreasing function $t \mapsto R(t)$ such that

$$(5.4) \quad \lim_{t \rightarrow \infty} \frac{t}{R^2(t)} = 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} t \varphi^{p-1}(R(t)) = \infty.$$

Replacing R by $R(t)$ in (5.3) and using (5.4) yields

$$\lim_{t \rightarrow \infty} t^{\frac{1}{p-1}} \psi_{R(t)}(t) = \left(\frac{1}{p-1} \right)^{\frac{1}{p-1}}.$$

If we consider x such that $\frac{|x|}{\sqrt{t}} \leq k$ for some constant k , we have

$$(5.5) \quad \lim_{t \rightarrow \infty} \frac{|x|}{R(t)} = \lim_{t \rightarrow \infty} \frac{|x|}{\sqrt{t}} \frac{\sqrt{t}}{R(t)} = 0.$$

Since $w(x, t) = \left(\frac{1}{(p-1)t} \right)^{\frac{1}{p-1}}$ is a supersolution for $t > 0$, and u is bounded there holds that,

$$(5.6) \quad \left(\frac{1}{p-1} \right)^{\frac{1}{p-1}} \geq t^{\frac{1}{p-1}} u(x, t) \geq t^{\frac{1}{p-1}} \psi_{R(t)}(t) H_{R(t)}(x).$$

Now, let us prove that

$$(5.7) \quad \lim_{t \rightarrow \infty} H_{R(t)}(x) = 1$$

uniformly on $|x| \leq k\sqrt{t}$ for all $k > 0$. In fact, from the uniform convergence on B_1 obtained in Theorem 3.1, for every $\varepsilon > 0$ there exists $t_1 > 0$ such that,

$$|\tilde{H}_{R(t)}(y) - h_1(y)| < \varepsilon \quad \text{if} \quad |y| \leq 1 \quad \text{and} \quad t \geq t_1.$$

On the other hand, if $|x| \leq k\sqrt{t}$ and we put $x = R(t)y$, we get that

$$|y| \leq \frac{k\sqrt{t}}{R(t)} \leq 1 \quad \text{if} \quad t \geq t_2$$

and consequently, if $|x| \leq k\sqrt{t}$

$$(5.8) \quad \left| H_{R(t)}(x) - h_1\left(\frac{x}{R(t)}\right) \right| = |\tilde{H}_{R(t)}(y) - h_1(y)| < \varepsilon \quad \text{if} \quad t \geq \max\{t_1, t_2\}.$$

From the continuity of h_1 it follows that,

$$(5.9) \quad \left| h_1\left(\frac{x}{R(t)}\right) - h_1(0) \right| \leq \varepsilon \quad \text{if} \quad |x| \leq k\sqrt{t} \quad \text{and} \quad t \geq t_3.$$

Hence, from (5.8) and (5.9) we obtain that,

$$|H_{R(t)}(x) - h_1(0)| < 2\varepsilon \quad \text{if} \quad |x| \leq k\sqrt{t} \quad \text{and} \quad t \geq \{t_1, t_2, t_3\}.$$

Since $h_1(0) = 1$, (5.7) follows.

Taking limit as $t \rightarrow \infty$ in (5.6) and using (5.7) we obtain that

$$\lim_{t \rightarrow \infty} t^{\frac{1}{p-1}} u(x, t) = \left(\frac{1}{p-1} \right)^{\frac{1}{p-1}}$$

uniformly on $E_k = \{x \in \mathbb{R}^N : |x| \leq k\sqrt{t}\}$ and the proof is finished. \square

REFERENCES

- [1] F. Andreu, J. Mazón, J. D. Rossi, J. Toledo, *A nonlocal p -Laplacian evolution equation with nonhomogeneous Dirichlet boundary conditions*, SIAM J. Math. Anal. **40**(5), 2008/09, 1815–1851.
- [2] P. Bates, A. Chmaj, *An integrodifferential model for phase transitions: Stationary solutions in higher dimensions*, J. Statistical Phys. **95**, 1999, 1119–1139.
- [3] P. Bates, A. Chmaj, *A discrete convolution model for phase transitions*, Arch. Rat. Mech. Anal. **150**, 1999, 281–305.
- [4] P. Bates, P. Fife, X. Ren, X. Wang, *Travelling waves in a convolution model for phase transitions*, Arch. Rat. Mech. Anal. **138**, 1997, 105–136.
- [5] P. Bates, G. Zhao *Existence, Uniqueness and Stability of the Stationary Solution to a Nonlocal Evolution Equation Arising in Population Dispersal*, J. Math. Anal. Appl., **332**, 2007, 428–440.
- [6] C. Carrillo, P. Fife, *Spatial effects in discrete generation population models*, J. Math. Biol. **50**(2), 2005, 161–188.
- [7] M. Chaves, E. Chasseigne, J. D. Rossi, *Asymptotic behavior for nonlocal diffusion equations*, Adv. Differential Equations, **2**, 2006, 271–291.
- [8] C. Cortazar, M. Elgueta, F. Quiros, N. Wolanski, *Asymptotic Behavior for a Nonlocal Diffusion Equation in Domains with Holes*, Arch. Rational Mech. Anal. **205**, 2012, 673–697.
- [9] P. Fife, *Some nonclassical trends in parabolic and parabolic-like evolutions*, Trends in nonlinear analysis, 153–191, Springer, Berlin, 2003.
- [10] J. García-Melián, F. Quirós, *Fujita exponents for evolution problems with nonlocal diffusion*, J. Evol. Equ. **10**(1), 2010, 147–161.
- [11] J. García-Melián, J. D. Rossi, *On the principal eigenvalue of some nonlocal diffusion problems*, J. Differential Equations **246**(1), 2009, 21–38.
- [12] G. Gilboa, S. Osher, *Nonlocal operators with application to image processing*, Multiscale Model. Simul., **7**(3), 2008, 1005–1028.
- [13] A. Gmira, L. Veron, *Large Time Behaviour of the Solutions of a Semilinear Parabolic Equation in R^N* , Jour. Diff. Eqs. **53**, 1984, 258–276.
- [14] Ignat, J. D. Rossi, *Refined asymptotic expansions for nonlocal diffusion equations*, J. Evolution Equations. **8**, 2008, 617–629.
- [15] C. Lederman, N. Wolanski, *Singular perturbation in a nonlocal diffusion model*, Communications in PDE **31**(2), 2006, 195–241.
- [16] A. Pazoto, J. D. Rossi, *Asymptotic behavior for a semilinear nonlocal equation*. Asymptotic Analysis. **52**(1-2), 2007, 143–155.
- [17] J. Terra, N. Wolanski, *Asymptotic behavior for a nonlocal diffusion equation with absorption and nonintegrable initial data. The supercritical case*, Proc. Amer. Math. Soc., **139** (4), 2011, 1421–1432.
- [18] J. Terra, N. Wolanski, *Large time behavior for a nonlocal diffusion equation with absorption and bounded initial data*, Discrete Cont. Dyn. Syst. A, 31 (2011) 2, 581–605.
- [19] L. Zhang, *Existence, uniqueness and exponential stability of traveling wave solutions of some integral differential equations arising from neuronal networks*, J. Differential Equations **197**(1), 2004, 162–196.

ARIEL SALORT
 IMAS-CONICET AND
 DEPARTAMENTO DE MATEMÁTICA, FCEYN-UBA
 (1428) BUENOS AIRES, ARGENTINA.
E-mail address: `asalort@dm.uba.ar`

JOANA TERRA
 IMAS-CONICET AND
 DEPARTAMENTO DE MATEMÁTICA, FCEYN-UBA
 (1428) BUENOS AIRES, ARGENTINA.
E-mail address: `jterra@dm.uba.ar`

NOEMI WOLANSKI
 IMAS-CONICET AND
 DEPARTAMENTO DE MATEMÁTICA, FCEYN-UBA
 (1428) BUENOS AIRES, ARGENTINA.
E-mail address: `wolanski@dm.uba.ar`