

Essential oil of *Tagetes filifolia* against the flour beetle *Tribolium castaneum* and its relation to acetylcholinesterase activity and lipid peroxidation

Olmedo, R.; J.M. Herrera, E.I. Lucini, M.P. Zunino, R.P. Pizzolitto, J.S. Dambolena and J.A. Zygadlo

SUMMARY

The development of natural insecticides would help to decrease the negative impact of synthetic insecticides. Fumigant toxicity of essential oil of *Tagetes filifolia* Lag (Asterales: Asteraceae) and its major compounds were evaluated against *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). The essential oils were analyzed by gas chromatography–mass spectrometry. The essential oil and (E)-anethole were the most toxic at 24 h against adult insects (CL_{50} = 2.4 y 2.6 μ L/mL water, respectively). Changes in the products of lipid peroxidation were evidenced by an increase in malondialdehyde content. Acetylcholinesterase activity *in vitro*, of *T. castaneum* adults, was also reported. (E)-anethole and estragole showed similar acetylcholinesterase inhibition (54 and 63%, respectively at 5 mM). The present study constitutes one of the first contributions to understanding the relationship between the insecticidal action of essential oil and oxidative stress.

Keywords: fumigant toxicity, *Tagetes filifolia*, *Tribolium castaneum*, lipid peroxidation, acetylcholinesterase activity.

Olmedo, R.; J.M. Herrera, E.I. Lucini, M.P. Zunino, R.P. Pizzolitto, J.S. Dambolena y J.A. Zygadlo, 2015. Aceite esencial de *Tagetes filifolia* contra *Tribolium castaneum* y su relación con la actividad acetilcolinesterasa y peroxidación de lípidos. Agriscientia 32 (2): 113-121

RESUMEN

El desarrollo de insecticidas naturales podría ayudar a disminuir el impacto negativo de insecticidas sintéticos. La acción insecticida fumigante del aceite esencial (AEs) de *Tagetes filifolia* Lag (Asterales: Asteraceae) y sus compuestos principales fueron evaluados contra adultos de *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). Los aceites esenciales fueron analizados por cromatografía gaseosa-espectrometría de masas. El aceite esencial y (E)-anetol mostraron mayor toxicidad contra insectos adultos a las 24 h (CL_{50} = 2,4 y 2,6 μ L/mL water, respectivamente). Se evidenciaron

cambios en los productos de peroxidación lipídica por un incremento en el contenido de malondialdehído. También se estudió la actividad *in vitro* de la enzima acetilcolinesterasa de insectos adultos de *T. castaneum*. (E)-anetol y estragol mostraron similar inhibición de la enzima (54 y 63%, respectivamente a 5 mM). El presente estudio constituye una de las primeras contribuciones para comprender la relación entre la acción insecticida del aceite esencial y el estrés oxidativo.

Palabras clave: toxicidad fumigante, *Tagetes filifolia*, *Tribolium castaneum*, peroxidación de lípidos, actividad acetilcolinesterasa.

R. Olmedo y E.I. Lucini: Cátedra de Química Biológica, Facultad de Ciencias Agropecuarias, Universidad Nacional de Córdoba, CC 509, 5000 Córdoba, Argentina. J.M. Herrera, M.P. Zunino, R.P. Pizzolitto, J.S. Dambolena and J.A. Zygadlo: Cátedra de Química Orgánica y Productos Naturales. Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, IMBIV-CONICET-ICTA, Av. Vélez Sarsfield 1611, Córdoba, Argentina. Correspondence to: pauzun@efn.uncor.edu

INTRODUCTION

The flour beetle, *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) is a serious insect pest of stored products around the world. Synthetic insecticides such as methyl bromide and phosphine are used for stored-product protection. Current control measures for this species have several limitations.

Methyl bromide affects the ozone layer thus, it has been banned in many countries and there is a commitment to reduce its use in 2015 (Arthur, 2012; Rajashekar *et al.*, 2012). The inadequate and sustained use of phosphine enabled the emergence of resistance in insect populations (Jagadeesan *et al.*, 2012; Opit *et al.* 2012). Recent studies indicated high levels of phosphine resistance in Brazilian populations of *T. castaneum* (Lorini *et al.*, 2007; Pimentel *et al.*, 2007). Although synthetic pesticides are effective, its frequent use has led to the presence of toxic residues in the environment and in food, and then generates negative effects on human health. These problems have highlighted the need to develop new and safe alternatives based on reducing the use of organophosphate, carbamates and chlorinated insecticides and promote the use of new techniques and products designed to reduce the resistance of insects and their presence in food. Thus, essential oils (EOs) obtained from aromatic plants are recognized as an alternative to synthetic insecticides against insect pest in stored foods (Rajashekar *et al.*, 2012; Regnault-Roger *et al.*, 2012). Plant EOs are

promising in that they are easily biodegradable and have not described resistance. With a few exceptions, their mammalian toxicity is low and environmental persistence is short (Regnault-Roger *et al.*, 2012). The genus *Tagetes* L. includes species with medicinal properties and their EOs have insecticidal activity (Krishna *et al.*, 2005; Caballero-Gallardo *et al.*, 2011; López *et al.*, 2011; Stefanazzi *et al.*, 2011). Among *Tagetes* species, particularly *T. filifolia* (Asteraceae, Helenieae) showed different degree in their insecticidal activity between different groups of insects (Camarillo *et al.*, 2009; López *et al.* 2011).

The insecticidal activity of EOs and their components can affect different and specific targets (Enan, 2001, 2005; Jbilou *et al.*, 2008; Abdelgaleil *et al.*, 2009; Rattan, 2010; Waliwitiya *et al.*, 2010). However, the inhibitory effect of phenylpropenes on *T. castaneum* acetylcholinesterase (AChE) of adults, were not reported. Moreover, like synthetic pesticides, EOs could generate oxidative stress. The products of lipid degradation and decomposition are used in assessing oxidative stress, including conjugated dienes (CD), as well as the widely used product malondialdehyde (MDA (Zunino & Zygadlo, 2004).

Therefore, in the present paper, we investigated the insecticidal activity of *T. filifolia* EO and their major components against *T. castaneum*. Changes in the products of lipid peroxidation, MDA and CD as indicators of oxidative stress are also discussed. The inhibitory effect on AChE activity of the adult insect was also evaluated.

MATERIALS AND METHODS

Insect material

All experiments were conducted in the laboratory with colonies of the red flour beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). Insects were reared on wheat flour mixed with yeast (10:1, w/w) at controlled temperature and humidity (28 °C and 70% RH) and a light/dark regime of 12:12 h (FAO, 1974). Adult insects used in toxicity studies were 3-7 days post-emergence.

Essential oils and pure compounds

Plants of *Tagetes filifolia* L. (Asteraceae, Helenieae) were collected from fields around Sierra de Los Gigantes from Córdoba province, Argentina. Specimen plants (Voucher specimen N° 2644) were deposited at the herbarium of Facultad de Agronomía (UNC) Córdoba, Argentina (ACOR).

Plants were dried in the shade to approximately 10% moisture content. The EO was obtained by hydrodistillation (López *et al.*, 2011) and stored at -20 °C in airtight microtubes prior to analysis by gas chromatography-mass spectrometry (GC-MS).

The compounds 1-methoxy-4-[(1E)-prop-1-en-1-yl] benzene ((E)-anethole) and 1-methoxy-4-(prop-2-en-1-yl) benzene (estragole) were purchased from Sigma Aldrich Co (St. Louis, MO USA) (Figure 1). DDVP (dichlorvos or 2,2-dichlorovinyl dimethyl phosphate) (contains >98% of active material, Chemotécnica S.A, Buenos Aires, Argentina) was used as reference insecticide.




Figure 1. Chemical structures of natural compounds studied in the present work.

Analysis of the essential oil (GC and GC-MS analysis)

The EO was analyzed using a Perkin-Elmer Clarus 500 gas chromatograph equipped with a flame ionization detector (GC-FID). The gas chromatography analysis was performed using a DB-5 capillary column (60 m 0.25 mm i.d., film thickness 0.25 mm). The carrier gas was helium with a flow rate of 0.9 mL/min. The oven temperature was maintained at 30 °C for 5 min and then programmed at 5 °C/min to 250 °C. The injector and detector were maintained at 260 and 280 °C, respectively. The sample, 0.2 µL, was injected with a 1:100 split ratio.

GC/MS. EOs samples were diluted with hexane. The injection volume was 1 µL. The identification of the components of the EO was realized by GC-MS. A Perkin-Elmer Q 700 GC-MS coupled with an ion trap mass detector was employed for the identification. A capillary column DB-5 (30 m x 0.25 mm i.d. and 0.25 µm coating thickness) was used for the separation of the components. Helium was used as carrier gas with a flow rate of 0.9 mL/min. The temperature program for the oven and injector was the same as that for the GC-FID. Ionization was realized by electron impact at 70 eV. Mass spectral data were acquired in the scan mode in the m/z range 35-450. Identification of the oil components was based on the comparison of their Retention Indices (RI) and mass spectra with those of authentic samples and/or the Adams, Nist and Wiley Library spectra. RI was computed using a formula by Adams (2007).

Fumigant Toxicity Assay

Different amounts of EO and pure compounds were placed inside a glass vial (1.5 mL) onto Petri dish (9 cm diameter). Inside the glass vial was placed 1 mL of water with the EO or pure compounds dissolved in ethanol to achieve the appropriate concentration (0.25 - 3 µL/mL water) (Figure 2). The ethanol that was added to the medium did not exceed 1% v/v. Twenty adults of




Figure 2: Fumigant toxicity method. A: Petri dish (9 cm diameter) with 20 insects; B: glass vial (1.5 mL) with EO or pure compounds dissolved in ethanol/mL water.

T. castaneum were placed into each Petri dish (4 replicates/doses). The Petri dish was sealed with parafilm to avoid the compounds loss. Insect controls were kept under same conditions without EO and pure compounds. Insect mortality was checked at 24 and 72 h of treatment. At 72 h, the EO or pure compounds were evaporated completely and only the vapour phase of the compounds was present in the Petri dish. The mortality percentages and LC₅₀ values were calculated according to Finney (1971).

Lipid Extraction from *T. castaneum*

One hundred *T. castaneum* adults from the different treatments (after 72 h exposure) were homogenized in 6 mL of hexane using a glass rod homogenizer. Homogenates were agitated and allowed to stand for 24 h, and then they were filtered through filter paper to remove coarse solids and to rinse the filter system with three portions of 0.5 mL of hexane. Supernatants were used for determination of MDA and CD content.

Measurement of malondialdehyde (MDA)

The MDA level was used as an index of lipid peroxidation and was expressed as nmol g⁻¹ of fresh weight. The MDA content was evaluated using the methodology described by Cervera *et al.* (2003), with some modification. Briefly, 1 mL of the supernatant of the lipid extraction was added to 4 mL of 0.5% thiobarbituric acid (TBA) in 20% trichloroacetic acid (TCA). The mixture was incubated at 95 °C for 30 min followed by quick cooling over ice and then centrifugated at 10,000 × *g* for 10 min. The absorbance of the supernatant was read at 532 nm and corrected for non-specific absorbance at 600 nm. MDA amount was determined using the extinction coefficient of 155 mM⁻¹ cm⁻¹. All the experiments were done in triplicate.

Measurement of conjugates dienes (CD)

Lipid extraction aliquots (0.5 mL) were used for determination of CD content. Indices were analyzed spectrophotometrically at a wavelength of 232 nm (Zunino and Zygadlo, 2004). All the experiments were done in triplicate.

AChE Activity Assay

T. castaneum adults (0.5 g) were separately

homogenized in 5 mL of 0.1 M ice-cold phosphate buffer (pH 7.4) using a Teflon glass tissue homogenizer. Homogenates were centrifuged (5,000 rpm for 20 min at 0 °C), and supernatants were used as the enzyme source for determination of AChE activity. Inhibition of AChE was determined by the colorimetric method of Ellman *et al.* (1961) using acetylthiocholine iodide (ATChI) at 0.25 mM (Sigma Aldrich Co., St. Louis, MO USA) as substrate, according to Abdelgaleil *et al.* (2009), with some modifications. Enzyme aliquots (100 µL) and 5,5-dithio-bis (2-nitrobenzoic) acid (DTNB) (100 µL of 0.01 M) were added to 0.1 M phosphate buffer (pH 7.4; 600 µL) and to this mixture, phenylpropene test solutions (100 µL) prepared in absolute ethanol were added. Control treatments were prepared by the addition of absolute ethanol (100 µL) instead of a phenylpropene. Mixtures were incubated at 35°C for 15 min and reactions were started by adding ATChI (100 µL). Absorbance was measured at 412 nm using UV/VIS Spectrometer, Lambda 25, Perkin Elmer. Tested compounds were examined at two concentrations: 1 mM and 5 mM. Each test and control was corrected by blanks for nonenzymic hydrolysis. All the experiments were done in triplicate. Inhibition percentage of AChE activity was calculated as follows:

$$\text{AChE inhibition \%} = (\text{ODC} - \text{ODT} / \text{ODC}) \times 100$$

where ODC is the optical density of control and ODT is the optical density of treatment. AChE activity values were compared to the control AChE activity, which was assumed to be 100%.

Statistical analysis

The lethal concentration (LC₅₀) values were calculated according to Finney (1971) using SPSS Statistics program version 17.0 (SPSS Inc). A one-way analysis of variance was used ($P \leq 0.05$). Comparisons between treatments were performed by the Duncan's multiple range test. All statistical analyses were calculated by using the InfoStat software Professional 2010 (Di Rienzo *et al.*, 2010).

The data of AChE inhibition was analyzed by one-way analysis of variance. Mean separations were performed by Duncan test and differences at $P < 0.05$ were considered as significant.

RESULTS

Chemical composition of the essential oil

The chemical composition of the EO obtained from *T. filifolia* is shown in Table 1. The main

constituents of EO were (E)-anethole (74.6%) and estragole (or methyl chavicol) followed by (E)-isoeugenol and anisic aldehyde, as have been described previously by López *et al.* (2011).

Toxicity of essential oil and its main compounds

The EO of *T. filifolia* was toxic against *T. castaneum*. The toxicity of the EO (LC₅₀: 2.0 µL/mL water, at 72h) was higher than that the estragole (LC₅₀: 5.6 µL/mL water, at 72h). (E)-anethol was more toxic than estragole both at 24 and at 72 hours of treatment (LC₅₀: 2.6 and 2.2 µL/mL water,

respectively). However, DDVP was the most toxic compound (Table 2).

Effect of *T. filifolia* oil and its main components on MDA and CD content of *T. castaneum*

The MDA content was positively correlated with increase concentrations of EO and (E)-anethole, while CD content showed a decrease at 3.0 µL/mL water. Estragole did not affect significantly MDA and CD content at low concentrations. Only at high concentration (5.2 µL/mL water) it showed significant differences with control in MDA and CD content (Table 3).

Table 1: Chemical constituents of the essential oil from *Tagetes filifolia* plants collected from Córdoba province, Argentina.

RI ^a	Compound	Content (%)	Methods of identification
1291	(E)-anethole	74.6	GCMS - Co
1194	estragole	23.7	GCMS - Co
1455	(E)- isoeugenol	0.9	GCMS - Co
1252	anisic aldehyde	0.8	GCMS

^a Retention index on a DB-5 column relative to homologous series of *n*-alkanes. GCMS: peak identifications are based on MS comparison with file spectra. Co: peak identification is based on standard comparison with relative retention time.

Table 2: Fumigant toxicity of *Tagetes filifolia* essential oil and their mains components against *Tribolium castaneum* at 24 h and 72 h after exposure.

Compounds	Hours of treatment	LC ₅₀ (µL/mL water)	95% confidence limits (µL/mL water)	
			Lower	Upper
T. filifolia oil	24	2.4	1.6	3.5
	72	2.0	1.6	2.6
(E)-anethole	24	2.6	2.4	2.8
	72	2.2	2.1	2.3
estragole	24	6.4	5.5	8.1
	72	5.6	4.9	7.0
DDVP	24	0.00053	0.00049	0.00057
	72	0.00040	0.00036	0.00043

Table 3: Effect of *Tagetes filifolia* oil and its mains components on MDA and conjugated dienes content of *Tribolium castaneum* at 72 h after exposure.

Compounds	Concentration (µL/mL water)	Mortality (%)	MDA (mM g insect weight ⁻¹)	Conjugated Dienes (Abs 232 nm)
T. filifolia oil	0	0 a	0.309 ± 0.006 a	0,5234 ± 0,0200 b
	2.4	100 b	0.344 ± 0.006 b	0,5224 ± 0,0808 b
	3.0	100 b	0.366 ± 0.005 c	0,2363 ± 0,0144 a
(E)-anethole	0	0 a	0,297 ± 0,005 a	0,4585 ± 0,0081 b
	2.4	77±5 b	0,317 ± 0,007 a	0,4809 ± 0,0063 b
	3.0	97±3 c	0,377 ± 0,007 b	0,2458 ± 0,0105 a
estragole	0	0 a	0,297 ± 0,005 a	0,4585 ± 0,0081 bc
	2.4	0 a	0,293 ± 0,005 a	0,4905 ± 0,0065 c
	3.0	22±5 b	0,301 ± 0,002 a	0,4480 ± 0,0170 ab
	5.2	45±5 c	0,323 ± 0,002 b	0,4192 ± 0,0078 a

Values (means ± ES) having different letters in the same compound are significantly different from each other according to Duncan's multiple range test at P ≤ 0.05 (n=3).

Inhibitory effect of monoterpenes on AChE activity

In vitro inhibitory effect of phenylpropenes on AChE activity from adults *T. castaneum* was examined at 1 mM and 5 mM. (E)-anethole and estragole showed similar inhibition (%) at the two concentrations tested. Maximal inhibition was observed at 5 mM (54-63%) (Table 4).

DISCUSSION

A similar composition of the essential oil of *T. filifolia* from other regions was observed by other authors (Serrato-Cruz *et al.*, 2008; Ruiz *et al.*, 2010; Armas *et al.*, 2012).

This study showed that *T. castaneum* adults are susceptible to fumigant action of *T. filifolia* EO and its pure component (E)-anethole. Besides, EO and (E)-anethole were more effective than estragole. López *et al.* (2008) reported that anethole and estragole may have a synergized insecticide action when presented together as part of the EO. Nevertheless, in this study the EO and (E)-anethole have shown a similar activity (Table 2). The high content of (E)-anethole in the EO could be the responsible of the bioactivity of this oil (Table 1). Previous studies founded that (E)-anethole and estragole showed similar insecticidal activity against *T. castaneum* adults (Ho *et al.*, 1997; Chang *et al.*, 2009; Mondal & Khalequzzaman, 2010; Ebadollahi, 2011; Chu *et al.*, 2012).

Natural or synthetic insecticides have different modes of action; increasing evidence for their secondary effects suggests that they disturb cellular homeostasis by generating free radical intermediates that trigger lipid peroxidation (Bakkali *et al.*, 2008; Aslanturk *et al.*, 2011; Yu *et al.*, 2011; Leelaja & Rajini, 2012; Prakash, 2015). The most commonly used criteria of oxidative stress are those based on determination of the concentrations of lipid oxidation products, usually polyunsaturated fatty acids, which are susceptible to attack by free radicals. In the present study, oxidative stress was evidenced principally in the different treatments by a significant increase in MDA content. However,

concentrations of the different treatments did not have significant differences in CD content except at 3.0 $\mu\text{L}/\text{mL}$ water for EO and (E)-anethole, and at 5.2 $\mu\text{L}/\text{mL}$ water for estragole, in which they decreased (Table 3). This decrease may result from the fact that conjugated dienic hydroperoxides, the initial products of lipid peroxidation, are labile, unlike the more stable MDA (Aslanturk *et al.*, 2011) formed upon their cleavage. An increase in MDA content has been shown with the use of different synthetic insecticides (Bakkali *et al.*, 2008; Aslanturk *et al.*, 2011; Yu *et al.*, 2011; Leelaja & Rajini, 2012). On the other hand, is known that the insecticidal activity of many terpenes and phenylpropanoids is directly related to their ability to inhibit the enzyme acetylcholinesterase. In this study, we showed an inhibition of insect AChE (> 60%) at 5 mM in the main EO components. López *et al.* (2015) reported the inhibitory effect of anethole and estragole on *Electrophorus* and bovine erythrocyte AChE activity, respectively.

The insecticidal activity of EO and anethole at 24h could indicate that the central nervous system (AChE, octopamine, GABA receptor, etc.) is one of their possible targets. Thus, the effect of anethole on AChE activity showed 60% of inhibition. On the other hand, estragole revealed low insecticidal activity at 24 h and high AChE inhibition. Similarly, Lee *et al.* (2001) did not find a direct correlation between insect toxicity against *Sitophilus oryzae* and AChE inhibition by menthone. Herrera *et al.* (2015) reported that camphor showed low insecticidal activity and high AChE inhibition. The same was observed by Yeom *et al.* (2012) with α -pinene. Besides, Yeom *et al.* (2012) showed that (E)-anethole has not ability to inhibit this enzyme in cockroaches and López & Pascual Villalobos (2010) showed the same but using electric eel AChE. This could be to the low penetration rate in the target site of the insect treated with these compounds. Therefore, oxidative stress could be involved in insect mortality at 72 h by estragole; due to this mechanism, do not cause an immediate effect in insect mortality. In addition, the values of MDA and CD content showed a correlation with insecticidal activity. So, oxidative stress is one of the possible mechanisms of target toxicity. Similar results were

Table 4: In vitro inhibition of *Tribolium castaneum* adult acetylcholinesterase by phenylpropenes.

Compounds	AChE-inhibition (%)	
	1 mM	5 mM
(E)- anethole	22.5 \pm 11.5 a	54.0 \pm 13.1 b
estragole	8.3 \pm 7.9 a	63.0 \pm 1.5 b

Data (%) are means \pm SE of three replicates. Different letters are significantly different from each other according to Duncan's multiple range test at $P \leq 0.05$.

reported by Prakash (2015) who revealed that the toxicity of EO of *Rosmarinus officinalis* against *Sitophilus oryzae* and *Oryzaephilus surinamensis*, might be associated with oxidative imbalance (antioxidant enzymes activity increased) and inhibition of AChE activity.

In conclusion, the data of the present study suggest that EO from *T. filifolia* and (E)-anethole present insecticidal activity and showed oxidative stress altering levels of MDA. Although other assays of oxidative stress must be analyzed, our recent results contribute to understanding the effect underlying the action of EO of *T. filifolia* and its principal components as oxidizing agents on insect. These findings suggest that the phenylpropanoids could act not only on AChE activity but also trigger oxidative stress.

The development of natural insecticides would help to decrease the negative impact of synthetic insecticides; because of their low toxicity, they do not generate the development of resistant populations of insects and they are safe to the environment despite their high relative cost and elevated doses compared with synthetic pesticide.

Based on these findings, EO from *Tagetes filifolia* and its main compound (E)-anethole may serve as viable alternatives to increase the lethality and effectiveness of currently available insecticides.

ACKNOWLEDGEMENTS

This research was supported by grants from Secretaría de Ciencia y Técnica de la Universidad Nacional de Córdoba and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). J.M. Herrera and R. Olmedo hold a fellowship from CONICET. J.A. Zygadlo, M.P. Zunino, R.P. Pizzolitto and J.S. Dambolena are career members of CONICET.

REFERENCES

- Abdelgaleil, S.A.M.; M.I.E. Mohamed, M.E.I. Badawi and S.A.A. El Arami, 2009. Fumigant and contact toxicities of monoterpenes to *Sitophilus oryzae* (L.) and *Tribolium castaneum* (Herbst) and their inhibitory effects on acetylcholinesterase activity. *Journal of Chemical Ecology*, 35:518–525.
- Adams, P.R., 2007. Identification of essential oil components by gas chromatography/mass spectrometry. Allured Publishing Corp., Carol Stream, Illinois, USA.
- Armas, K.; J. Rojas, L. Rojas and A. Morales, 2012. Comparative study of the chemical composition of essential oils of five *Tagetes* species collected in Venezuela. *Natural Products Communication*, 7:1225-1226.
- Arthur, F.H., 2012. Aerosols and contact insecticides as alternatives to methyl bromide in flour mills, food production facilities, and food warehouses. *Journal of Pest Science*, 85: 323-329.
- Aslanturk, A.; S. Kalender, M. Uzunhisarcikli and Y. Kalender, 2011. Effects of methidathion on antioxidant enzyme activities and malondialdehyde level in midgut tissues of *Lymantria dispar* (Lepidoptera) larvae. *Journal of the Entomological Research Society*, 13: 27-38.
- Bakkali, F.; S. Averbeck, D. Averbeck and M. Idaomar, 2008. Biological effects of essential oils- A review. *Food and Chemical Toxicology*, 46: 446-475.
- Caballero-Gallardo, K.; J. Olivero-Verbel and E.E. Stashenko, 2011. Repellent activity of essential oils and some of their individual constituents against *Tribolium castaneum*. *Herbst. Journal of Agricultural Food and Chemistry*, 59:1690-1696.
- Camarillo, G.R.; L.D.A. Ortega, M.A.C. Serrato and C.H. Rodríguez, 2009. Biological activity of *Tagetes filifolia* (Asteraceae) on *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae). *Revista Colombiana de Entomología*, 35: 177-184.
- Cervera, C.; A.C. Maymó, R. Martínez-Pardo and M.D. Garcerá, 2003. Antioxidant enzymes in *Oncopeltus fasciatus* (Heteroptera: Lygaeidae) exposed to cadmium. *Environmental Entomology*, 32: 705–710.
- Chang, C.L.; I.L.K. Cho and Q.X. Li, 2009. Insecticidal activity of basil oil, trans-anethole, estragole, and linalool to adult fruit flies of *Ceratitis capitata*, *Bactrocera dorsalis*, and *Bactrocera cucurbitae*. *Journal of Economic Entomology*, 102: 203-209.
- Chu, S.S.; J. Cao, Q.Z. Liu, S.S. Du, Z.W. Deng and Z.L. Liu, 2012. Chemical composition and insecticidal activity of *Heracleum moellendorffii* Hance essential oil. *Chemija*, 23: 108-112.
- Di Rienzo, J.A.; F. Casanoves, M.G. Balzarini, L. Gonzalez, M. Tablada and C.W. Robledo, 2010. Infostat version 2010. Grupo Infostat, FCA, Universidad Nacional de Córdoba, Argentina.
- Ebadollahi, A., 2011. Chemical constituents and toxicity of *Agastache foeniculum* (Pursh) Kuntze essential oil against two stored-product insect pests. *Chilean Journal Agricultural Research*, 71:212-217.
- Ellman, G.L.; D. Courtney, V. Andres and R.M. Featherstone, 1961. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology*, 7:88–95.
- Enan, E.E., 2001. Insecticidal activity of essential

- oils: octopaminergic sites of action. *Comparative Biochemistry and Physiology*, 130: 325-337.
- Enan, E.E., 2005. Molecular response of *Drosophila melanogaster* tyramine receptor cascade to plant essential oils. *Insect Biochemistry and Molecular Biology*, 35: 309-321
- FAO, 1974. Boletín Fitosanitario de la FAO. Método provisional para gorgojos adultos importantes en cereales almacenados con malation o lindano. Método N° 15 de la FAO, 22:127-137
- Finney, D.J., 1971. *Probit Analysis*. Cambridge University Press, New York.
- Herrera, J.M.; M.P. Zunino, J.S. Dambolena, R.P. Pizzolitto, N.A. Gañan, E.I. Lucini and J.A. Zygadlo, 2015. Terpene ketones as natural insecticides against *Sitophilus zeamais*. *Industrial Crops and Products*, 70: 435-442.
- Ho, S.H.; Y. Ma and A. Huang, 1997. Anethole a potential insecticide from *Illicium verum* Hook f., against two stored product insects. *International Pest Control*, 39:50-51.
- Jagadeesan, R.; P.J. Collins, G.J. Darglish, P.R. Ebert and D.I. Schlipalius, 2012. Phosphine resistance in the rust red flour beetle, *Tribolium castaneum* (Coleoptera: Tenebrionidae): Inheritance, Gene Interactions and Fitness Costs. *PLoS ONE* 7: e31582.
- Jbilou, R.; H. Amri, N. Bouayad, N. Ghailani, A. Ennabili and F. Sayah, 2008. Insecticidal effects of extracts of seven plant species on larval development, α -amylase activity and offspring production of *Tribolium castaneum* (Herbst) (Insecta: Coleoptera: Tenebrionidae). *Bioresource Technology*, 99:959-964.
- Krishna, A.; V. Prajapati, S. Bhasney, A.K. Tripathi and S. Kumar, 2005. Potential toxicity of new genotypes of *Tagetes* (Asteraceae) species against stored grain insect pests. *International Journal of Tropical Insect Science*, 25:122-128.
- Lee, S.E.; B.H. Lee, W.S. Choi, B.S. Park, J.G. Kim and B.C. Campbell, 2001. Fumigant toxicity of volatile natural products from Korean spices and medicinal plants towards the rice weevil, *Sitophilus oryzae* (L). *Pest Management Science*, 57:548-553.
- Leelaja, B.C. and P.S. Rajini, 2012. Impact of phosphine exposure on development in *Caenorhabditis elegans*: Involvement of oxidative stress and the role of glutathione. *Pesticide Biochemistry and Physiology*, 104:38-43.
- López, M.D.; J. Contreras and M. J. Pascual Villalobos, 2008. Toxic compounds in essential oils of coriander, caraway and basil active against stored rice pests. *Journal of Stored Products Research*, 44:273-278.
- López, M.D.; F.J. Campoy, M.J. Pascual-Villalobos, E. Muñoz-Delgado, and C.J. Vidal. 2015. Acetylcholinesterase activity of electric eel is increased or decreased by selected monoterpenoids and phenylpropanoids in a concentration dependent manner. *Chemico-Biological Interactions*, 229: 36-43.
- López M.D. and M. J. Pascual Villalobos. 2010. Mode of inhibition of acetylcholinesterase by monoterpenoids and implications for pest control. *Industrial Crops and Products*, 31: 284-288.
- López, S.B.; M.L. López, L.M. Aragón, M.L. Tereschuk, A.C. Slanis, G.E. Feresin, J. A Zygadlo and A.A. Tapia, 2011. Composition and anti-insect activity of essential oils from *Tagetes* L species (Asteraceae, Helenieae) on *Ceratitis capitata* Wiedemann and *Triatoma infestans* Klug. *Journal of Agricultural Food and Chemistry*, 59:5286-5292.
- Lorini, I.; P.J. Collins, G.J. Darglish, M.K. Nayak and H. Pavic, 2007. Detection and characterization of strong resistance to phosphine in Brazilian *Rhyzopertha dominica* (F.) (Coleoptera: Bostrichidae). *Pest Management Science*, 63:358-364.
- Mondal, M. and M. Khalequzzaman, 2010. Toxicity of naturally occurring compounds of plant essential oil against *Tribolium castaneum* (Herbst). *Journal of Biological Science*, 10:10-17.
- Opit, G.P.; T.W. Phillips, M. J. Aikins and M.M. Hasan, 2012. Phosphine resistance in *Tribolium castaneum* and *Rhyzopertha dominica* from stored wheat in Oklahoma. *Journal of Economy Entomology*, 105:1107-1114.
- Pimentel, M.A.G., L.R.D.A. Faroni, M.R. Tótola and R.N.C. Guedes, 2007. Phosphine resistance, respiration rate and fitness consequences in stored-product insects. *Pest Management Science*, 63: 876-881.
- Prakash, K.S.B., 2015. Toxicity and biochemical efficacy of chemically characterized *Rosmarinus officinalis* essential oil against *Sitophilus oryzae* and *Oryzaephilus surinamensis*. *Industrial Crops and Products*, 74:817-823.
- Rajashekar, Y.; N. Bakthavatsalam and T. Shivanandappa, 2012. Botanicals as Grain Protectants. *Psyche* (New York), article: 646740.
- Rattan, R.S., 2010. Mechanism of action of insecticidal secondary metabolites of plant origin. *Crop Protection*, 29: 913 - 920.
- Regnault-Roger, C.; C. Vincent and J.T. Arnason, 2012. Essential oils in insect control: Low-risk products in a high-stakes world. *Annual Review Entomology*, 57:405-424.
- Ruiz, S.; O. Malagón, T. Zaragoza and E. Valarezo, 2010. Composition of the essential oils of *Artemisia sodiroi* Hieron., *Siparuna eggersii* Hieron., *Tagetes filifolia* Lag. and *Clinopodium nubigenum* (Kunth) Kuntze from Loja Ecuador. *Journal Essential Oil Bearing Plants*, 13:676-691.

- Serrato Cruz, M.Á.; F. Díaz Cedillo and J.S. Barajas Pérez, 2008. Composition of essential oil in germplasm of *Tagetes filifolia* Lag. from Central-South region of México. *Agrociencia*, 42: 277-285.
- SPPSS Inc, 2008. SPSS-vs. 17. User's Guide.
- Stefanazzi, N.; T. Stadler and A. Ferrero, 2011. Composition and toxic, repellent and feeding deterrent activity of essential oils against the stored-grain pests *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus oryzae* (Coleoptera: Curculionidae). *Pest Management Science*, 67, 639-646.
- Waliwitiya, R.; P. Belton, R A Nicholson, Lowenberger CA. 2010. Effects of the essential oil constituent thymol and other neuroactive chemicals on flight motor activity and wing beat frequency in the blowfly *Phaenicia sericata*. *Pest Management Science*, 66: 277-289.
- Yeom, H.J.; J.S. Kang, G.H. Kim and I.K. Park, 2012. Insecticidal and acetylcholine esterase inhibition activity of Apiaceae plant essential oils and their constituents against adults of German cockroach (*Blattella germanica*). *Journal of Agricultural Food and Chemistry*, 60:7194–7203.
- Yu Q.Y.; S.M. Fang, Z.D. Zuo, F.Y. Dai, Z. Zhang and C. Lu, 2011. Effect of organophosphate phoxim exposure on certain oxidative stress biomarkers in the silkworm. *Journal of Economy Entomology*, 104: 101-106.
- Zunino, M. P. and J.A. Zygadlo, 2004. Effect of monoterpenes on lipid oxidation of maize. *Planta*, 219: 303-309.