

Invariants for Continuous Linear Dynamical Systems

Shaull Almagor 

Department of Computer Science, Technion, Haifa, Israel
shaull@cs.technion.ac.il

Edon Kelmendi

Department of Computer Science, Oxford University, UK
edon.kelmendi@cs.ox.ac.uk

Joël Ouaknine

Max Planck Institute for Software Systems, Saarland Informatics Campus, Saarbrücken, Germany
Department of Computer Science, Oxford University, UK
joel@mpi-sws.org

James Worrell

Department of Computer Science, Oxford University, UK
jbw@cs.ox.ac.uk

Abstract

Continuous linear dynamical systems are used extensively in mathematics, computer science, physics, and engineering to model the evolution of a system over time. A central technique for certifying safety properties of such systems is by synthesising inductive invariants. This is the task of finding a set of states that is closed under the dynamics of the system and is disjoint from a given set of error states. In this paper we study the problem of synthesising inductive invariants that are definable in \mathcal{O} -minimal expansions of the ordered field of real numbers. In particular, assuming Schanuel's conjecture in transcendental number theory, we establish effective synthesis of \mathcal{O} -minimal invariants in the case of semi-algebraic error sets. Without using Schanuel's conjecture, we give a procedure for synthesising \mathcal{O} -minimal invariants that contain all but a bounded initial segment of the orbit and are disjoint from a given semi-algebraic error set. We further prove that effective synthesis of semi-algebraic invariants that contain the whole orbit, is at least as hard as a certain open problem in transcendental number theory.

2012 ACM Subject Classification Theory of computation \rightarrow Logic and verification; Computing methodologies \rightarrow Algebraic algorithms; Mathematics of computing \rightarrow Continuous mathematics; Mathematics of computing \rightarrow Continuous functions; Theory of computation \rightarrow Finite Model Theory; Software and its engineering \rightarrow Formal software verification

Keywords and phrases Invariants, continuous linear dynamical systems, continuous Skolem problem, safety, \mathcal{O} -minimality

Digital Object Identifier 10.4230/LIPIcs.ICALP.2020.107

Category Track B: Automata, Logic, Semantics, and Theory of Programming

Related Version A full version of the paper is available at [3] <https://arxiv.org/abs/2004.11661>.

Funding *Shaull Almagor*: Supported by a European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 837327.

Joël Ouaknine: Supported by ERC grant AVS-ISS (648701) and DFG grant 389792660 as part of TRR 248 (see <https://perspicuous-computing.science>).

James Worrell: Supported by EPSRC Fellowship EP/N008197/1.



© Shaull Almagor, Edon Kelmendi, Joël Ouaknine, and James Worrell;
licensed under Creative Commons License CC-BY

47th International Colloquium on Automata, Languages, and Programming (ICALP 2020).

Editors: Artur Czumaj, Anuj Dawar, and Emanuela Merelli; Article No. 107; pp. 107:1–107:15

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



1 Introduction

A *continuous linear dynamical system* (CDS) is a system whose evolution is governed by a differential equation of the form $\dot{\mathbf{x}}(t) = A\mathbf{x}(t)$, where A is a matrix with real entries. CDSs are ubiquitous in mathematics, physics, and engineering; they have been extensively studied as they describe the evolution of many types of systems (or abstractions thereof) over time. More recently, CDSs have become central in the study of cyber-physical systems (see, e.g., the textbook [4]).

In the study of CDSs, particularly from the perspective of control theory, a fundamental problem is *reachability* – namely whether the orbit $\{\mathbf{x}(t) : t \geq 0\}$ intersects a given target set $Y \subseteq \mathbb{R}^d$. For example, when $\mathbf{x}(t)$ describes the state of an autonomous car (i.e., its location, velocity, etc.). Y may describe situations where the car is not able to stop in time to respond to a hazard.

When Y is a singleton set, reachability is decidable [14, Theorem 2]. However, already when Y is a half-space it is open whether or not reachability is decidable. The latter decision problem is known in the literature as the *continuous Skolem problem*. Some partial positive results were given in [6] and [10]. The continuous Skolem problem is related to notoriously difficult problems in the theory of Diophantine approximation: specifically a procedure for the continuous Skolem problem would yield one for computing to arbitrary precision the *Diophantine-approximation types* of all real algebraic numbers [10].

In lieu of an algorithm to decide reachability, a popular approach to certifying non-reachability is by finding an *inductive invariant*, that is, a set X that is closed under the dynamics of the system and disjoint from the target Y . For such a set, it is immediate that the orbit of any point in X does not reach Y . In order to make this approach effective it is natural to seek invariants among suitably tame classes of sets (e.g., polyhedra, semi-algebraic sets, etc). In this work we take a very general approach – we consider *o-minimal invariants*, that is, invariants definable in *some* o-minimal expansion of the ordered set of real numbers. The class of o-minimal invariants includes those defined by Boolean combinations of inequalities involving polynomials, the real exponential function, and bounded versions of the trigonometric functions. One can potentially use many other functions to define o-minimal invariants, but the key to our approach is to show that the ingredients just listed suffice in the case of a CDS and semi-algebraic target Y .

The papers [2, 1] study o-minimal invariants for *discrete* linear dynamical systems. There it is proved that when the target Y is a semi-algebraic set, the question of whether there exists an o-minimal invariant disjoint from Y is decidable. Furthermore, if there is an o-minimal invariant then there is in fact a semi-algebraic invariant which can moreover be constructed effectively. The present paper uses similar ideas, although the case of continuous linear dynamical systems differs in several important ways.

Main Contributions. We consider the following problem: given a CDS by means of a matrix A with rational entries, an initial point $\mathbf{x}_0 = \mathbf{x}(0)$, and a semi-algebraic set Y of error states, decide whether there exists a set that is definable in some o-minimal expansion of the ordered real field and is (1) disjoint from Y , (2) invariant under the dynamics of the system, and (3) contains the initial point \mathbf{x}_0 . We show that in searching for such invariants it suffices to look among sets definable in the expansion of the reals with the real exponential function and trigonometric functions restricted to bounded domains. Moreover, assuming Schanuel’s conjecture (a unifying conjecture in transcendental number theory), we prove that the existence of such an invariant is decidable, and that invariants can effectively be constructed when they exist.

Without assuming Schanuel’s conjecture we can decide a related problem, namely the question of whether there exists a set that is definable in an o-minimal expansion of the real field and is (1) disjoint from Y , (2) invariant under the dynamics of the system, and (3) meets the orbit of the initial point \mathbf{x}_0 . Notice that such a set – which could be called an *eventual invariant* – must contain all but a bounded initial segment of the orbit. We show that when such a set exists, it can be effectively constructed and moreover that it can be chosen to be a semi-algebraic set. Such an invariant can serve as a certificate that the orbit does not enter the error set Y infinitely often. The latter is a very difficult problem to decide, even when the target set is a half-space [9].

As mentioned earlier, for discrete linear dynamical systems the question of whether there exists a semi-algebraic invariant that contains the *whole* orbit is decidable [2, 1]. We provide an explanation of why the analogous result for continuous systems is not easy to prove; this is by way of a reduction from a difficult problem that highlights the complications of continuous systems. The problem asks whether a given exponential polynomial of the form

$$f(t) = a_1 e^{b_1 t} + \dots + a_n e^{b_n t}$$

has zeros in a bounded interval, where a_i, b_i are real algebraic numbers. Deciding whether f has zeros in a bounded region seems to be difficult because all the zeros have to be transcendental (a consequence of Hermite-Lindemann Theorem), and they can be tangential, i.e., f never changes its sign, yet it has a zero.

Related Work. Invariant synthesis is a central technique for establishing safety properties of hybrid systems. It has long been known how to compute a strongest *algebraic* invariant [20] (i.e., a smallest algebraic set that contains the collection of reachable states) for an arbitrary CDS. Here an algebraic invariant is one that is specified by a conjunction of polynomial equalities. If one moves to the more expressive setting of semi-algebraic invariants, which allow inequalities, then there is typically no longer a strongest (or smallest) invariant, but one can still ask to decide the existence of an invariant that avoids a given target set of configurations. This is the problem that is addressed in the present paper.

Partial positive results are known, for example when strong restrictions on the matrix A are imposed, such as when all the eigenvalues are real and rational, or purely imaginary with rational imaginary part [15].

A popular approach in previous work has been to seek invariants that match a given syntactic *template*, which allows to reduce invariant synthesis to constraint solving [13, 23, 16]. While this technique can be applied to much richer classes of systems than those considered here (e.g., with discrete control modes and non-linear differential equations), it does not appear to offer a way to decide the existence of arbitrary semi-algebraic invariants. An alternative to the template approach for invariant generation involves obtaining candidate invariants from semi-algebraic abstractions of a system [21]. Another active area of current research lies in developing powerful techniques to check whether a given semi-algebraic set is actually an invariant [12, 16].

Other avenues for analysing dynamical systems in the literature include bisimulations [7], forward/backward reach-set computation [5], and methods for directly proving liveness properties [22]. The latter depends on constructing *staging sets*, which are essentially semi-algebraic invariants.

Often, questions about dynamical systems can be reduced to deciding whether a sentence belongs to the elementary theory of an appropriate expansion of the ordered field of real numbers. While the latter is typically undecidable, there are partial positive results, namely

quasi-decidability in bounded domains, see [11] and the references therein. This can be used to reason about the dynamics of a system in a bounded time interval, under the assumption that it does not tangentially approach the set that we want to avoid. However, it seems unlikely that such results can be easily applied to the problems considered here.

The rest of the paper is organised as follows. In Section 2, we give the necessary definitions and terminology. In Section 3, we define *cones*, which are over-approximations of the orbit, and prove that they are in a certain sense canonical. The positive results assuming Schanuel's conjecture are subsequently given in this section. Section 4 is devoted to the effective construction of the semi-algebraic invariants which allows us to state and prove the unconditional positive results. In Section 5, we give the aforementioned reduction, from finding zeros of exponential polynomials.

See [3] for the complete proofs.

2 Preliminaries

A *continuous-time linear dynamical system* is a pair

$$\langle A, \mathbf{x}_0 \rangle$$

where $A \in \mathbb{Q}^{d \times d}$ and $\mathbf{x}_0 \in \mathbb{Q}^d$. The system evolves in time according the function $x(t)$ which is the unique solution to the differential equation $\dot{\mathbf{x}}(t) = A\mathbf{x}(t)$ with $\mathbf{x}(0) = \mathbf{x}_0$. Explicitly this solution can be written as:

$$\mathbf{x}(t) = e^{At} \mathbf{x}_0.$$

The *orbit* of $\langle A, \mathbf{x}_0 \rangle$ from time t_0 is the set $\mathcal{O}(t_0) = \{e^{At} \mathbf{x}_0 : t \geq t_0\}$. An *invariant* for $\langle A, \mathbf{x}_0 \rangle$ from time t_0 is a set $\mathcal{I} \subseteq \mathbb{R}^d$ that contains $e^{At_0} \mathbf{x}_0$ and is stable under applications of e^{At} , i.e., $e^{At} \mathcal{I} \subseteq \mathcal{I}$ for every $t \geq 0$. Note that an invariant from time t_0 contains $\mathcal{O}(t_0)$. Given a set $Y \subseteq \mathbb{R}^d$ (referred to henceforth as an *error set*), we say that the invariant \mathcal{I} *avoids* Y if the two sets are disjoint.

We denote by \mathfrak{R}_0 the structure $\langle \mathbb{R}, 0, 1, +, \cdot, < \rangle$. This is the ordered field of real numbers with constants 0 and 1. A sentence in the corresponding first-order language is a quantified Boolean combination of atomic propositions of the form $P(x_1, \dots, x_n) > 0$, where P is a polynomial with integer coefficients and x_1, \dots, x_n are variables. In addition to \mathfrak{R}_0 , we also consider its following expansions:

- $\mathfrak{R}_{\text{exp}}$, obtained by expanding \mathfrak{R}_0 with the real exponentiation function $x \mapsto e^x$.
- \mathfrak{R}^{RE} , obtained by expanding \mathfrak{R}_0 with the *restricted elementary functions*, namely $x \mapsto e^x|_{[0,1]}$, $x \mapsto \sin x|_{[0,1]}$, and $x \mapsto \cos x|_{[0,1]}$.
- $\mathfrak{R}_{\text{exp}}^{\text{RE}}$, obtained by expanding $\mathfrak{R}_{\text{exp}}$ with the restricted elementary functions.

Tarski famously showed that the first-order theory of \mathfrak{R}_0 admits quantifier elimination, moreover the elimination is effective and therefore the theory is decidable [24, Theorem 37].

It is an open question whether the theory of the reals with exponentiation ($\mathfrak{R}_{\text{exp}}$) is decidable; however decidability was established subject to Schanuel's conjecture by MacIntyre and Wilkie [18, Theorem 1.1]. MacIntyre and Wilkie further showed in [18, Section 5] that decidability of the theory of $\mathfrak{R}_{\text{exp}}$ implies a weak form of Schanuel's conjecture.

Similarly, it is an open question whether \mathfrak{R}^{RE} and $\mathfrak{R}_{\text{exp}}^{\text{RE}}$ are decidable, but they are also known to be decidable subject to Schanuel's conjecture [17, Theorem 3.1]¹.

¹ More precisely, the decidability of $\mathfrak{R}_{\text{exp}}$ requires Schanuel's conjecture over \mathbb{R} , whereas that of $\mathfrak{R}_{\text{exp}}^{\text{RE}}$ requires it over \mathbb{C} .

Let \mathfrak{R} be an expansion of the structure \mathfrak{R}_0 . A set $S \subseteq \mathbb{R}^d$ is *definable* in \mathfrak{R} if there exists a formula $\phi(x_1, \dots, x_d)$ in \mathfrak{R} with free variables x_1, \dots, x_d such that $S = \{(c_1, \dots, c_d) \in \mathbb{R}^d \mid \mathfrak{R} \models \phi(c_1, \dots, c_d)\}$. For $\mathfrak{R} = \mathfrak{R}_0$, the ordered field of real numbers, \mathfrak{R}_0 -definable sets are known as *semi-algebraic* sets.

► **Remark 2.1.** There is a natural first-order interpretation of the field of complex numbers \mathbb{C} in the field of real numbers \mathbb{R} . We shall say that a set $S \subseteq \mathbb{C}^d$ is *\mathfrak{R} -definable* if the image $\{(x, y) \in \mathbb{R}^d \times \mathbb{R}^d \mid x + iy \in S\}$ of S under this interpretation is \mathfrak{R} -definable.

A totally ordered structure $\langle M, <, \dots \rangle$ is said to be *o-minimal* if every definable subset of M is a finite union of intervals. Tarski's result on quantifier elimination implies that \mathfrak{R}_0 is o-minimal. The o-minimality of $\mathfrak{R}_{\text{exp}}$ and \mathfrak{R}^{RE} is shown in [27], and the o-minimality of \mathfrak{R}^{RE} and $\mathfrak{R}_{\text{exp}}^{\text{RE}}$ is due to [25, 26].

A *semi-algebraic invariant* is one that is definable in \mathfrak{R}_0 . An *o-minimal invariant* is one that is definable in an o-minimal expansion of $\mathfrak{R}_{\text{exp}}$.

3 Orbit Cones

In this section we define orbit cones, an object that plays a central role in the subsequent results. They can be thought of as over-approximations of the orbit that has certain desirable properties, and moreover it is canonical in the sense that any other invariant must contain a cone.

3.1 Jordan Normal Form

Let $\langle A, \mathbf{x}_0 \rangle$ be a continuous linear dynamical system. The exponential of a square matrix A is defined by its formal power series as

$$e^A \stackrel{\text{def}}{=} \sum_{n=0}^{\infty} \frac{A^n}{n!}.$$

Let $\lambda_1, \dots, \lambda_k$ be the eigenvalues of A , and recall that when $A \in \mathbb{Q}^{d \times d}$, all the eigenvalues are algebraic. We can write A in *Jordan Normal Form* as $A = PJP^{-1}$ where $P \in \mathbb{C}^{d \times d}$ is an invertible matrix with algebraic entries, and $J = \text{diag}(B_1, \dots, B_k)$ is a block-diagonal matrix where each block B_l is a Jordan block that corresponds to eigenvalue λ_l , and it has the form

$$B_l = \begin{pmatrix} \lambda_l & 1 & 0 & \cdots & 0 \\ 0 & \lambda_l & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_l \end{pmatrix} \in \mathbb{C}^{d_l \times d_l}$$

with $\sum_{l=1}^k d_l = d$.

From the power series, we can write $e^{At} = Pe^{Jt}P^{-1}$. Further, $e^{Jt} = \text{diag}(e^{B_1 t}, \dots, e^{B_k t})$. For each $1 \leq l \leq k$, write $B_l = \Lambda_l + N_l$, where Λ_l is the $d_l \times d_l$ diagonal matrix $\text{diag}(\lambda_l, \dots, \lambda_l)$ and N_l is the $d_l \times d_l$ matrix $\text{diag}_2(1, \dots, 1)$; where $\text{diag}_j(\cdot)$ is the j -th diagonal matrix, with other entries zero.

The matrices Λ_l and N_l commute, since the former is a diagonal matrix. A fundamental property of matrix exponentiation is that if matrices A, B commute, then $e^{A+B} = e^A e^B$. Thus, we have

$$e^{Jt} = e^{\text{diag}(\Lambda_1 t + N_1 t, \dots, \Lambda_k t + N_k t)} = \text{diag}(e^{\lambda_1 t}, \dots, e^{\lambda_k t}) e^{\text{diag}(N_1 t, \dots, N_k t)},$$

where by $\text{diag}(e^{\lambda_1 t}, \dots, e^{\lambda_k t})$ we mean the $d \times d$ diagonal matrix that has the entry $e^{\lambda_1 t}$ written d_1 times, the entry $e^{\lambda_2 t}$ written d_2 times and so on. It will always be clear from the context whether we repeat the entries because of their multiplicity or not.

Matrices N_l are nilpotent, so its power series expansion is a finite sum, i.e. a polynomial in $N_l t$. More precisely, one can verify that:

$$e^{N_l t} = I + \text{diag}_2(t, \dots, t) + \text{diag}_3\left(\frac{t^2}{2}, \dots, \frac{t^2}{2}\right) + \dots + \text{diag}_{d_l}\left(\frac{t^{(d_l-1)}}{(d_l-1)!}\right).$$

Set $Q(t) \stackrel{\text{def}}{=} \text{diag}(e^{N_1 t}, \dots, e^{N_k t})$. From the equation above, the entries of $Q(t)$ are polynomials in t with rational coefficients.

Write the eigenvalues as $\lambda_l = \rho_l + i\omega_l$, so that

$$\text{diag}(e^{\lambda_1 t}, \dots, e^{\lambda_k t}) = \underbrace{\text{diag}(e^{\rho_1 t}, \dots, e^{\rho_k t})}_{E(t)} \cdot \underbrace{\text{diag}(e^{i\omega_1 t}, \dots, e^{i\omega_k t})}_{R(t)}$$

We have in this manner decomposed the orbit

$$\mathcal{O}(t_0) = \{P \ E(t) \ R(t) \ Q(t) \ P^{-1} \mathbf{x}_0 : t \geq t_0\},$$

into an exponential $E(t)$, a rotation $R(t)$, and a simple polynomial $Q(t)$ matrices that commute with one another. Having the orbit in such a form will facilitate the analysis done in the sequel.

3.2 Cones as Canonical Invariants

In a certain sense, the rotation matrix $R(t)$ is the most complicated, because of it, the orbit is not even definable in $\mathfrak{R}_{\text{exp}}$. The purpose of cones is to abstract away this matrix by a much simpler subgroup of the complex torus

$$\mathbb{T} \stackrel{\text{def}}{=} \{\mathbf{z} \in \mathbb{C}^k : |z_i| = 1, 1 \leq i \leq k\}.$$

To this end, consider the group of additive relations among the frequencies $\omega_1, \dots, \omega_k$:

$$S \stackrel{\text{def}}{=} \{\mathbf{a} \in \mathbb{Z}^k : a_1 \omega_1 + \dots + a_k \omega_k = 0\}.$$

The subgroup of the torus of interest, respects the additive relations as follows:

$$\mathbb{T}_\omega \stackrel{\text{def}}{=} \{(\tau_1, \dots, \tau_k) \in \mathbb{T} : \text{for all } \mathbf{a} \in S, \tau_1^{a_1} \dots \tau_k^{a_k} = 1\}.$$

Its desirable properties are summarised in the following proposition:

► **Proposition 3.1.** *For algebraic numbers $\omega_1, \dots, \omega_k$,*

1. \mathbb{T}_ω is semi-algebraic,
2. diagonals of $\{R(t) : t \geq 0\}$ form a dense subset of \mathbb{T}_ω .

Proof. Being an Abelian subgroup of \mathbb{Z}^k , S has a finite basis, moreover this basis can be computed because of effective bounds, [19, Section 3]. To check that (τ_1, \dots, τ_k) belongs to \mathbb{T}_ω , it suffices to check that $\tau_1^{a_1} \dots \tau_k^{a_k} = 1$ for (a_1, \dots, a_k) in the finite basis. This forms a finite number of equations, therefore \mathbb{T}_ω is semi-algebraic. The fact that this is a subset of vectors of complex numbers is not problematic in this case because of the simple first-order interpretation in the theory of reals, see Remark 2.1.

The second statement of the proposition is a consequence of Kronecker's theorem on inhomogeneous simultaneous Diophantine approximations, see [8, Page 53, Theorem 4]. The proof of a slightly stronger statement can also be found in [9, Lemma 4]. Examples can be found where the set of diagonals of $\{R(t) : t \geq 0\}$ is a strict subset of \mathbb{T}_ω . ◀

The orbit cone can now be defined by replacing the rotations with the subgroup of the torus. As it turns out, for our purposes this approximation is not too rough.

► **Definition 3.2.** *The orbit cone from $t_0 \geq 0$ is*

$$\mathcal{C}_{t_0} \stackrel{\text{def}}{=} \{P E(t) \text{diag}(\boldsymbol{\tau}) Q(t) P^{-1} \mathbf{x}_0 : \boldsymbol{\tau} \in \mathbb{T}_\omega, t \geq t_0\}.$$

We prove that the cone is an inductive invariant and also a subset of \mathbb{R}^d .

► **Lemma 3.3.** *For all $\delta, t_0 \geq 0$, $e^{A\delta} \mathcal{C}_{t_0} \subseteq \mathcal{C}_{t_0}$.*

Proof. Fix $t \geq t_0$ and $\boldsymbol{\tau} \in \mathbb{T}_\omega$, and consider the point

$$\mathbf{v} = P E(t) \text{diag}(\boldsymbol{\tau}) Q(t) P^{-1} \mathbf{x}_0 \in \mathcal{C}_{t_0},$$

then we can write $e^{A\delta} \mathbf{v}$ as

$$\begin{aligned} e^{A\delta} \mathbf{v} &= P E(\delta) R(\delta) Q(\delta) \cdot E(t) \text{diag}(\boldsymbol{\tau}) Q(t) P^{-1} \mathbf{x}_0 \\ &= P E(\delta + t) R(\delta) \text{diag}(\boldsymbol{\tau}) Q(\delta) Q(t) P^{-1} \mathbf{x}_0. \end{aligned}$$

The matrix $R(\delta) \text{diag}(\boldsymbol{\tau})$ is equal to $\text{diag}(\boldsymbol{\tau}')$ for some $\boldsymbol{\tau}' \in \mathbb{T}_\omega$. Otherwise said, the vector $(e^{\delta\omega_1 i} \tau_1, \dots, e^{\delta\omega_k i} \tau_k)$ belongs to \mathbb{T}_ω . Indeed this is the case because for any $\mathbf{a} \in S$ we have

$$e^{a_1 \delta \omega_1 i} \tau_1^{a_1} \dots e^{a_k \delta \omega_k i} \tau_k^{a_k} = e^{\delta i (a_1 \omega_1 + \dots + a_k \omega_k)} \cdot \tau_1^{a_1} \dots \tau_k^{a_k} = 1.$$

Finally, by induction on the dimension d one can verify that $Q(\delta)Q(t) = Q(\delta + t)$. ◀

The fact that cones are subsets of \mathbb{R}^d comes as a corollary of the following proposition whose proof can be found in Appendix A of the full version of the paper [3].

► **Proposition 3.4.** *Let $A = PJP^{-1}$ as above, and let $C_i \in \mathbb{C}^{d_i \times d_i}$ for $i = 1, \dots, k$, with dimensions compatible to the Jordan blocks of A , and such that for every i_1, i_2 , if $B_{i_1} = \overline{B_{i_2}}$, then $C_{i_1} = \overline{C_{i_2}}$. Then $P \text{diag}(C_1, \dots, C_k) P^{-1}$ has real entries.*

The matrix $E(t) \text{diag}(\boldsymbol{\tau}) Q(t)$ can be written as $\text{diag}(C_1, \dots, C_k)$ where the C_i matrices satisfy the conditions of Proposition 3.4, hence the following corollary.

► **Corollary 3.5.** *For all $t_0 \geq 0$ we have $\mathcal{C}_{t_0} \subseteq \mathbb{R}^d$.*

It is surprising that, already, the cones are a complete characterisation of o-minimal inductive invariants in the following sense.

► **Theorem 3.6.** *Let \mathcal{I} be an o-minimal invariant that contains the orbit $\mathcal{O}(u)$ from some time $u \geq 0$, then there exists $t_0 \geq u$ such that:*

$$\mathcal{C}_{t_0} \subseteq \mathcal{I}.$$

Proof sketch. Conceptually, the proof follows along the lines of its analogue in [1]. There are a few differences, namely that the entries of the matrix A in [1] are assumed to be algebraic, while this is not true for the entries of e^A .

We define rays of the cone, which are subsets where $\boldsymbol{\tau} \in \mathbb{T}_\omega$ is fixed. Then we prove that for every ray, all but a finite part of it, is contained in the invariant. This is done by contradiction: if a ray is not contained in the invariant, a whole dense subset of the cone can be shown not to be contained in the invariant, leading to a contradiction, since the invariant is assumed to contain the orbit. We achieve this using some results on the topology of o-minimal sets.

The complete proof can be found in Appendix B of [3]. ◀

Another desirable property of cones is that they are $\mathfrak{R}_{\text{exp}}$ -definable. Also, one can observe that for every t_0 , the set $\{e^{At}\mathbf{x}_0 : 0 \leq t \leq t_0\}$ is definable in $\mathfrak{R}_{\text{exp}}^{\text{RE}}$ (as we only need bounded restrictions of sin and cos to capture e.g. $e^{i\omega t}$ up to time t_0). As an immediate corollary of Theorem 3.6, we have the following theorems.

► **Theorem 3.7.** *Let $\langle A, \mathbf{x}_0 \rangle$ be a CDS. For every $t_0 \geq 0$, the set $\mathcal{C}_{t_0} \cup \{e^{At}\mathbf{x}_0 : 0 \leq t \leq t_0\}$ is an invariant that contains the whole orbit of $\langle A, \mathbf{x}_0 \rangle$. Moreover, this invariant is definable in $\mathfrak{R}_{\text{exp}}^{\text{RE}}$ (and in particular is o-minimal).*

► **Theorem 3.8.** *Let $\langle A, \mathbf{x}_0 \rangle$ be a CDS and let $Y \subseteq \mathbb{R}^d$ be an error set. There exists an o-minimal invariant \mathcal{I} that contains the orbit and is disjoint from Y if and only if there exists t_0 such that $\mathcal{C}_{t_0} \cup \{e^{At}\mathbf{x}_0 : 0 \leq t \leq t_0\}$ is such an invariant.*

Theorem 3.8 now allows us to provide an algorithm for deciding the existence of an invariant, subject to Schanuel's conjecture:

► **Theorem 3.9.** *Assuming Schanuel's conjecture, given a CDS $\langle A, \mathbf{x}_0 \rangle$ and an $\mathfrak{R}_{\text{exp}}^{\text{RE}}$ definable error set Y , it is decidable whether there exists an o-minimal invariant for $\langle A, \mathbf{x}_0 \rangle$ that avoids Y . Moreover, if such an invariant exists, we can compute a representation of it.*

Proof. By Theorem 3.8, there exists an o-minimal invariant \mathcal{I} that avoids Y if and only if there exists some $t_0 \in \mathbb{R}$ such that $\mathcal{C}_{t_0} \cup \{e^{At}\mathbf{x}_0 : 0 \leq t \leq t_0\}$ is such an invariant. Thus, the problem reduces to deciding the truth value of the following $\mathfrak{R}_{\text{exp}}^{\text{RE}}$ sentence:

$$\exists t_0 : (\mathcal{C}_{t_0} \cup \{e^{At}\mathbf{x}_0 : 0 \leq t \leq t_0\}) \cap Y = \emptyset$$

The theory of $\mathfrak{R}_{\text{exp}}^{\text{RE}}$ is decidable subject to Schanuel's conjecture, and therefore we can decide the existence of an invariant. Moreover, if an invariant exists, we can compute a representation of it by iterating over increasing values of t_0 , until we find a value for which the sentence $(\mathcal{C}_{t_0} \cup \{e^{At}\mathbf{x}_0 : 0 \leq t \leq t_0\}) \cap Y = \emptyset$ is true. ◀

4 Semi-algebraic Error Sets and Fat Trajectory Cones

In this section, we restrict attention to semi-algebraic invariants and semi-algebraic error sets, in order to regain unconditional decidability.

Substitute $s = e^t$ in the definition of the cone to get:

$$\mathcal{C}_{t_0} = \{P E(\log s) \text{diag}(\boldsymbol{\tau}) Q(\log s) P^{-1}\mathbf{x}_0 : \boldsymbol{\tau} \in \mathbb{T}_\omega, s \geq e^{t_0}\}.$$

Written this way, observe that $E(\log s) = \text{diag}(s^{\rho_1}, \dots, s^{\rho_k})$, which is almost semi-algebraic, apart from the fact that the exponents need not be rational.

4.1 Unconditional Decidability

We give the final, yet crucial property of the cones. When the error set is semi-algebraic, it is possible to decide, unconditionally, whether there exists some cone that avoids the error set. Moreover the proof is constructive, it will produce the cone for which this property holds.

► **Theorem 4.1.** *For a semi-algebraic error set Y , it is (unconditionally) decidable whether there exists $t_0 \geq 0$ such that $\mathcal{C}_{t_0} \cap Y = \emptyset$. Moreover, such a t_0 can be computed.*

Proof. Define the set

$$U \stackrel{\text{def}}{=} \{ \mathcal{V} \in \mathbb{R}^{d \times d} : \forall \tau \in \mathbb{T}_\omega, P \mathcal{V} \text{diag}(\tau) P^{-1} \mathbf{x}_0 \in \mathbb{R}^d \setminus Y \}.$$

The set U can be seen to be semi-algebraic and thus is expressed by a quantifier-free formula that is a finite disjunction of formulas of the form $\bigwedge_{l=1}^m R_l(\mathcal{V}) \sim_l 0$, where each R_l is a polynomial with integer coefficients, over $d \times d$ variables of the entries of the matrix \mathcal{V} , and $\sim_l \in \{>, =\}$. Define the matrix

$$\Lambda(s) \stackrel{\text{def}}{=} \text{diag}(s^{\rho_1}, \dots, s^{\rho_k}) Q(\log s) \in \mathbb{R}^{d \times d},$$

and notice that $\mathcal{C}_{t_0} \cap Y = \emptyset$ if and only if $\Lambda(s) \in U$ for every $s \geq e^{t_0}$. Thus, it is enough to decide whether there exists $s_0 \geq 1$ such that for every $s \geq s_0$, at least one of the disjuncts $\bigwedge_{l=1}^m R_l(\Lambda(s)) \sim_l 0$ is satisfied.

Since $R_l(\Lambda(s))$ are polynomials in entries of the form s^{ρ_i} and $\log(s)$, there is an effective bound s_0 such that for all $s \geq s_0$, none of the values $R_l(\Lambda(s))$ change sign for any $1 \leq l \leq m$. Hence we only need to decide whether there exists some $s'_0 \geq s_0$ such that for all $s \geq s'_0$ we have $R_l(\Lambda(s)) \sim_l 0$ for every $1 \leq l \leq m$.

Fix some l . The polynomial $R_l(v_1, \dots, v_D)$ has the form $\sum_i a_i v_1^{n_{i,1}} \dots v_D^{n_{i,D}}$. After identifying the matrix $\Lambda(s)$ with a vector in \mathbb{R}^D for $D = d^2$, we see that $R_l(\Lambda(s))$ is a sum of terms of the form

$$a_i s^{n'_{i,1}\rho_1 + \dots + n'_{i,k}\rho_k} \cdot Q_{i,1}(\log s) \cdots Q_{i,D}(\log s)$$

where the $n'_{i,j}$ are aggregations of the $n_{i,j}$ for identical entries of $\text{diag}(s_1^{\rho_1}, \dots, s_k^{\rho_k})$, and $Q_{i,j}(\log s)$ are polynomials obtained from the entries of $Q(\log s)$ under R_l . We can join the polynomials Q_1, \dots, Q_D into a single polynomial f_i , which would also absorb a_i . Thus, we rewrite R_l in the form $\sum_i s^{n'_{i,1}\rho_1 + \dots + n'_{i,k}\rho_k} f_i(\log s)$ where each f_i is a polynomial with rational coefficients (as the coefficients in $Q(\log s)$ are rational).

In order to reason about the sign of this expression as $s \rightarrow \infty$, we need to find the leading term of $R_l(\Lambda(s))$. This, however, is easy: the exponents $n'_{i,1}\rho_1 + \dots + n'_{i,k}\rho_k$ are algebraic numbers, and are therefore susceptible to effective comparison. Thus, we can order the terms by magnitude. Then, we can determine the asymptotic sign of each coefficient $f_i(\log s)$ by looking at the leading term in f_i .

We can thus determine the asymptotic behaviour of each $R_l(\Lambda(s))$, to conclude whether $\bigwedge_{l=1}^m R_l(\Lambda(s)) \sim_l 0$ eventually holds. Moreover, for rational s , every quantity above can be computed to arbitrary precision, therefore it is possible to compute a threshold s'_0 , after which, for all $s \geq s'_0$, $\bigwedge_{l=1}^m R_l(\Lambda(s)) \sim_l 0$ holds. This completes the proof. ◀

► **Theorem 4.2.** *For a semi-algebraic set Y , it is decidable whether there exists a 0-minimal invariant, disjoint from Y , that contains the orbit $\mathcal{O}(u)$ after some time $u \geq 0$. Moreover in the positive instances an invariant that is $\mathfrak{R}_{\text{exp}}$ -definable can be constructed.*

Proof. If there is an invariant \mathcal{I} that contains $\mathcal{O}(u)$, for some $u \geq 0$, then Theorem 3.6 implies that there exists some $t_0 \geq u$ such that \mathcal{C}_{t_0} is contained in \mathcal{I} . Consequently, the question that we want to decide is equivalent to the question of whether there exists a t_0 , such that $\mathcal{C}_{t_0} \cap Y = \emptyset$. The latter is decidable thanks to Theorem 4.1. The effective construction follows from the fact that such a t_0 is computable and that the cone is $\mathfrak{R}_{\text{exp}}$ -definable. ◀

4.2 Effectively Constructing the Semi-algebraic Invariant

We now turn to show that in fact, for semi-algebraic error sets Y , we can approximate \mathcal{C}_{t_0} with a semi-algebraic set such that if \mathcal{C}_{t_0} avoids Y , so does the approximation. Intuitively, this is done by relaxing the “non semi-algebraic” parts of \mathcal{C}_{t_0} in order to obtain a *fat cone*. This relaxation has two parts: one is to “rationalize” the (possibly irrational) exponents ρ_1, \dots, ρ_k , and the other is to approximate the polylogs in $Q(\log s)$ by polynomials.

Relaxing the exponents. We start by approximating the exponents ρ_1, \dots, ρ_k with rational numbers. We remark that naively taking rational approximations is not sound, as the approximation must also adhere to the additive relationships of the exponents.

Let $\boldsymbol{\ell} = (\ell_1, \dots, \ell_k)$ and $\mathbf{u} = (u_1, \dots, u_k)$ be tuples of rational numbers such that $\ell_i \leq \rho_i \leq u_i$ for $i = 1, \dots, k$. Define $\mathbb{S} \subseteq \mathbb{R}^k$ as:

$$\mathbb{S} \stackrel{\text{def}}{=} \left\{ (q_1, \dots, q_k) \in \mathbb{R}^k : \forall n_1, \dots, n_k \in \mathbb{Z}, \left(\sum_{i=1}^k n_i \rho_i = 0 \Rightarrow \sum_{i=1}^k n_i q_i = 0 \right) \right\}$$

Thus, \mathbb{S} captures the integer additive relationships among the ρ_i . Define

$$\text{Box}(\boldsymbol{\ell}, \mathbf{u}) \stackrel{\text{def}}{=} \{ \text{diag}(\mathbf{q}) : \boldsymbol{\ell} \leq \mathbf{q} \leq \mathbf{u}, \mathbf{q} \in \mathbb{S} \}.$$

Approximating polylogs. Let $\epsilon, \delta > 0$. We simply replace $\log s$ by r such that $\delta \leq r \leq s^\epsilon$. Note that it is not necessarily the case that $\delta \leq \log s \leq s^\epsilon$, so this replacement is a-priori not sound. However, for large enough s the inequalities do hold, which will suffice for our purposes.

We can now define the fat cone. Let $\epsilon, \delta > 0$ and $\boldsymbol{\ell} = (\ell_1, \dots, \ell_k)$ and $\mathbf{u} = (u_1, \dots, u_k)$ as above, the fat orbit cone $\mathcal{F}_{s_0, \epsilon, \delta, \boldsymbol{\ell}, \mathbf{u}}$ is the set:

$$\left\{ P \text{diag}(s^{q_1}, \dots, s^{q_k}) \text{diag}(\boldsymbol{\tau}) Q(r) P^{-1} \mathbf{x}_0 : \boldsymbol{\tau} \in \mathbb{T}_\omega, s \geq s_0, \delta \leq r \leq s^\epsilon, \mathbf{q} \in \text{Box}(\boldsymbol{\ell}, \mathbf{u}) \right\}.$$

That is, the fat cone is obtained from \mathcal{C}_{t_0} with the following changes:

- $R(\log s) = \text{diag}(s^{\rho_1}, \dots, s^{\rho_k})$ is replaced with $\text{diag}(s^{q_1}, \dots, s^{q_k})$, where the q_i are rational approximations of the ρ_i , and maintain the additive relationships.
- $Q(\log s)$ is replaced with $Q(r)$ where $\delta \leq r \leq s^\epsilon$.
- The variable s starts from s_0 (as opposed to e^{t_0}).

We first show that the fat cone is semi-algebraic (the proof is in [3] Appendix C), then proceed to prove that if there is a cone that avoids the error set, then there is a fat one that avoids it as well.

► **Lemma 4.3.** $\mathcal{F}_{s_0, \epsilon, \delta, \boldsymbol{\ell}, \mathbf{u}}$ is definable in \mathfrak{R}_0 , and we can compute a representation of it.

► **Lemma 4.4.** Let $Y \subseteq \mathbb{R}^d$ be a semi-algebraic error set such that $\mathcal{C}_{t_0} \cap Y = \emptyset$ for some $t_0 \in \mathbb{R}$, then there exists $\delta, \epsilon, s_0, \boldsymbol{\ell}, \mathbf{u}$ as above such that

1. $\mathcal{F}_{s_0, \epsilon, \delta, \boldsymbol{\ell}, \mathbf{u}} \cap Y = \emptyset$, and
2. for every $t \geq 0$ it holds that $e^{At} \cdot \mathcal{F}_{s_0, \epsilon, \delta, \boldsymbol{\ell}, \mathbf{u}} \subseteq \mathcal{F}_{s_0, \epsilon, \delta, \boldsymbol{\ell}, \mathbf{u}}$.

The result is constructive, so when t_0 is given, the constants $s_0, \epsilon, \delta, \boldsymbol{\ell}, \mathbf{u}$ can be computed. It follows that a corollary of this lemma, and Lemma 4.3, is a stronger statement than that of Theorem 4.2, namely one where $\mathfrak{R}_{\text{exp}}$ is replaced by \mathfrak{R}_0 . We state it here before moving on with the proof of Lemma 4.4.

► **Theorem 4.5.** *For a semi-algebraic set Y , it is decidable whether there exists a 0-minimal invariant, disjoint from Y , that contains the orbit $\mathcal{O}(u)$ after some time $u \geq 0$. Moreover in the positive instances an invariant that is \mathfrak{R}_0 -definable can be constructed.*

The proof of Lemma 4.4 is given by the two corresponding steps. The second step, proving the invariance of the fat cone, can be found in [3]. We turn our attention to the first step.

► **Lemma 4.6.** *Let $Y \subseteq \mathbb{R}^d$ be a semi-algebraic error set, and let $t_0 \in \mathbb{R}$ be such that $\mathcal{C}_{t_0} \cap Y = \emptyset$, then there exists $\delta, \epsilon, s_0, \ell, \mathbf{u}$ as above such that $\mathcal{F}_{s_0, \epsilon, \delta, \ell, \mathbf{u}} \cap Y = \emptyset$.*

Proof. We use the same analysis and definitions of $U, R_l, \sim_l, \Lambda(s)$ as in the proof of Theorem 4.1 and focus on a single polynomial R_l . Recall that we had

$$R_l(\Lambda(s)) = \sum_i s^{n_{i,1}\rho_1 + \dots + n_{i,k}\rho_k} f_i(\log s) \tag{1}$$

where each f_i is a polynomial with rational coefficients.

Denote $\boldsymbol{\rho} = (\rho_1, \dots, \rho_k)$. We show, first, how to replace the exponents vector $\boldsymbol{\rho}$ by any exponents vector in $\text{Box}(\ell, \mathbf{u})$ for appropriate ℓ, \mathbf{u} , and second, how to replace $\log s$ by r where $\delta \leq r \leq s^\epsilon$ for some appropriate δ and ϵ , while maintaining the inequality or equality prescribed by \sim_l .

Denote by N the set of vectors $\mathbf{n}_i = (n_{i,1}, \dots, n_{i,k})$ of exponents in (1). Let $\mu > 0$, such that for every $\mathbf{n}, \mathbf{n}' \in N$, if $\boldsymbol{\rho} \cdot (\mathbf{n} - \mathbf{n}') \neq 0$ then $|\boldsymbol{\rho} \cdot (\mathbf{n} - \mathbf{n}')| > \mu$. That is, μ is a lower bound on the minimal difference between distinct exponents in (1). Observe that we can compute a description of μ , as the exponents are algebraic numbers.

Let $M = \max_{\mathbf{n}, \mathbf{n}' \in N} \|\mathbf{n} - \mathbf{n}'\|$ (where $\|\cdot\|$ is the Euclidean norm in \mathbb{R}^k).

▷ **Claim 4.7.** Let $\mathbf{c} \in \mathbb{R}^k$ be such that $\|\boldsymbol{\rho} - \mathbf{c}\| \leq \frac{\mu}{2M}$, then, for all $\mathbf{n}, \mathbf{n}' \in N$, if $\boldsymbol{\rho} \cdot (\mathbf{n} - \mathbf{n}') > 0$ then $\mathbf{c} \cdot (\mathbf{n} - \mathbf{n}') > \frac{\mu}{2}$.

Proof of Claim 4.7. Suppose that $\boldsymbol{\rho} \cdot (\mathbf{n} - \mathbf{n}') > 0$, then by the above we have $\boldsymbol{\rho} \cdot (\mathbf{n} - \mathbf{n}') > \mu$, and hence

$$\mathbf{c} \cdot (\mathbf{n} - \mathbf{n}') = \boldsymbol{\rho} \cdot (\mathbf{n} - \mathbf{n}') + (\mathbf{c} - \boldsymbol{\rho}) \cdot (\mathbf{n} - \mathbf{n}') \geq \mu - \|\mathbf{c} - \boldsymbol{\rho}\| \cdot \|\mathbf{n} - \mathbf{n}'\| \geq \mu - \frac{\mu}{2M} M = \frac{\mu}{2}. \blacktriangleleft$$

We can now choose ℓ and \mathbf{u} such that $u_i - \ell_i \leq \frac{\mu}{2M\sqrt{k}}$ and for all $\mathbf{c} \in \text{Box}(\ell, \mathbf{u})$ we have

$$\|\boldsymbol{\rho} - \mathbf{c}\| \leq \sqrt{\sum_{i=1}^k (u_i - \ell_i)^2} \leq \sqrt{\frac{\mu^2}{(2M)^2}} = \frac{\mu}{2M}.$$

It follows from Claim 4.7 and from the definition of $\text{Box}(\ell, \mathbf{u})$ that, intuitively, every $\mathbf{c} \in \text{Box}(\ell, \mathbf{u})$ maintains the order of magnitude of the monomials $s^{n_{i,1}\rho_1 + \dots + n_{i,k}\rho_k}$ in $R_l(\Lambda(s))$.

More precisely, let $\Lambda'(s) = \text{diag}(s^{c_1}, \dots, s^{c_k})Q(\log s)$ for some $\mathbf{c} \in \text{Box}(\ell, \mathbf{u})$, then the exponent of the ratio of every two monomials in $R_l(\Lambda'(s))$ has the same (constant) sign as the corresponding exponent in $R_l(\Lambda(s))$. Moreover, the exponents of distinct monomials in $R_l(\Lambda(s))$ differ by at least $\frac{\mu}{2}$ in $R_l(\Lambda'(s))$.

We now turn our attention to the $\log s$ factor. First, let s_0 be large enough that $f_i(\log s)$ has constant sign for every $s \geq s_0$. We can now let δ be large enough such that for every $r \geq \delta$, the sign of $f_i(\log s)$ coincides with the sign of $f_i(r)$ for every $s \geq s_0$. It remains to give an upper bound on r of the form s^ϵ such that plugging $f_i(r)$ instead of $f_i(\log s)$ does not change the ordering of the terms (by their magnitude) in $R_l(\Lambda'(s))$.

107:12 Invariants for Continuous Linear Dynamical Systems

Let B be the maximum degree of all polynomials f_i in (1), and define $\epsilon = \frac{\mu}{3B}$ (in fact, any $\epsilon < \frac{\mu}{2B}$ would suffice), then we have that, for $s \geq s_0$, $f_i(r)$ has the same sign as $f_i(\log s)$ for every $\delta \leq r \leq s^\epsilon$ (by our choice of δ), and guarantees that plugging s^ϵ instead of s does not change the ordering of the terms (by their magnitude) in R_l . Since the exponents of the monomials in $R_l(\Lambda'(s))$ differ by at least $\frac{\mu}{2}$, it follows that their order is maintained when replacing $\log s$ by $\delta \leq r \leq s^\epsilon$.

Let $\Lambda''(s) = \text{diag}(s^{c_1}, \dots, s^{c_k})Q(r)$ for some $\mathbf{c} \in \text{Box}(\ell, \mathbf{u})$ and $\delta \leq r \leq s^\epsilon$, then by our choice of ϵ , the dominant term in $R_l(\Lambda''(s))$ is the same as that in $R_l(\Lambda(s))$. Therefore, for large enough s , the signs of $R_l(\Lambda''(s))$ and $R_l(\Lambda(s))$ are the same.

Note that since $\mathcal{C}_{t_0} \cap Y = \emptyset$, then w.l.o.g. $R_l(\Lambda(s)) \sim_l 0$ for every l . Thus, by repeating the above argument for each R_l , we can compute $s_0 \in \mathbb{R}$, $\epsilon > 0$, $\delta \in \mathbb{R}$, and $\ell, \mathbf{u} \in \mathbb{Q}^k$ such that $\mathcal{F}_{s_0, \epsilon, \delta, \ell, \mathbf{u}} \cap Y = \emptyset$, and we are done. \blacktriangleleft

5 A Reduction from Zeros of an Exponential Polynomial

In Theorem 4.5, we showed unconditional decidability for the question of whether there exists an invariant containing the orbit $\mathcal{O}(u)$, for some $u \geq 0$. Even though we construct such an invariant, it cannot be used as a certificate proving that the orbit never enters the error set; however it is a certificate that the orbit of the system does not enter Y after time u .

In this section we give indications that deciding whether there exists an invariant that takes into account the orbit $\leq u$ is difficult. More precisely, we will reduce a problem about zeros of a certain exponential polynomial to the question of whether there exists a semi-algebraic invariant disjoint from Y containing $\mathcal{O}(0)$.

► **Remark 5.1.** In the setting of discrete linear dynamical systems, the existence of a semi-algebraic invariant from time t_0 immediately implies the existence of one from time 0. This is because the system goes through finitely many points from 0 to t_0 , which can be added one by one to the semi-algebraic set. In this respect CDSs are more complicated to analyse.

The problem that we reduce from, can be stated as follows. We are given as input real algebraic numbers $a_1, \dots, a_n, \rho_1, \dots, \rho_n$, and $t_0 \in \mathbb{Q}$, and asked to decide whether the exponential function:

$$f(t) \stackrel{\text{def}}{=} a_1 e^{\rho_1 t} + \dots + a_n e^{\rho_n t},$$

has any zeros in the interval $[0, t_0]$. This is a special case of the so-called Continuous Skolem Problem [6, 10].

While there has been progress on characterising the asymptotic distribution of complex zeros of such functions, less is known about the real zeros, and we lack any effective characterisation, see [6, 10] and the references therein. The difficulty of knowing whether f has a zero in the specified region is because (a) all the zeros have to be transcendental (a consequence of Hermite-Lindemann Theorem) and (b) there can be tangential zeros, that is f has a zero but it never changes its sign. See the discussion in [6, Section 6]. Finding the zeros of such a polynomial is a special case of the *bounded* continuous Skolem problem. We note that when ρ_i are all rational the problem is equivalent to a sentence of \mathfrak{R}_0 (and hence decidable) by replacing $t = \log s$.

The rest of this section is devoted to the proof of the following theorem.

► **Theorem 5.2.** *For every exponential polynomial f we can construct a CDS $\langle A, \mathbf{x}_0 \rangle$ and semi-algebraic set Y such that the following two statements are equivalent:*

- *there exists a semi-algebraic invariant disjoint from Y that contains $\mathcal{O}(0)$,*
- *f does not have a zero in $[0, t_0]$.*

Fix the function f , *i.e.* real algebraic numbers $a_1, \dots, a_n, \rho_1, \dots, \rho_n$ and $t_0 \in \mathbb{Q}$. Without loss of generality we can assume that ρ_1, \dots, ρ_n are all nonnegative, since $e^{\rho t} f(t) = 0$ if and only if $f(t) = 0$ where ρ is larger than all ρ_1, \dots, ρ_n .

Since every ρ_i is algebraic, there is a minimal polynomial p_i , that has ρ_i as a simple root. Let A be the $d \times d$ companion matrix of the polynomial $p_1(x) \cdots p_n(x)x^2$. The numbers ρ_i are eigenvalues A of multiplicity one, and the latter also has zero as an eigenvalue of multiplicity two. In addition to those, the matrix A generally has other (complex) eigenvalues as well. We put A in Jordan normal form, $P^{-1}AP = J$ where J is made of two block diagonals: \tilde{A} and B , where

$$\tilde{A} \stackrel{\text{def}}{=} \begin{pmatrix} \text{diag}(\rho_1, \dots, \rho_n) & & \\ & 0 & 1 \\ & 0 & 0 \end{pmatrix},$$

and B is some $(d - n - 2) \times (d - n - 2)$ matrix. Define:

$$\tilde{\mathbf{x}}_0 \stackrel{\text{def}}{=} (\underbrace{1, \dots, 1}_{n+2}, 0, \dots, 0),$$

the vector that has $n + 2$ ones and the rest, $d - (n + 2)$ zeros, whose purpose is to ignore the contribution of the eigenvalues in matrix B in the system. To simplify notation, since $\tilde{\mathbf{x}}_0$ is ignoring the contribution of the matrix B , the dynamics of the system $\langle J, \tilde{\mathbf{x}}_0 \rangle$ can be assume to be the same as:

$$e^{\tilde{A}t}(1, \dots, 1) = (e^{\rho_1 t}, \dots, e^{\rho_n t}, t).$$

Focus on a single eigenvalue, *i.e.* on the graph $\{(e^{\rho t}, t) : t \geq 0\}$, as the analysis will easily generalise to the CDS in question. This is itself a CDS, so terminology such as orbits *etc.* make sense. The challenge is to find a family of *tubes* around this exponential curve such that (a) all the tubes together with $\{(y, t) : t \geq t_0\}$ are invariants and (b) the tubes are arbitrarily close approximations of the curve.

We achieve this by the following families of polynomials:

- under-approximations are given by the family indexed by $n \in \mathbb{N}$:

$$P_n(t) \stackrel{\text{def}}{=} \sum_{k=0}^n \frac{(\rho t)^k}{k!}.$$

- over-approximations are given by a family indexed by $n \in \mathbb{N}$ and $\mu > 1$:

$$Q_{n,\mu}(t) \stackrel{\text{def}}{=} P_n(\mu t).$$

Define:

$$\mathcal{I}_{n,\mu} \stackrel{\text{def}}{=} \{(y, t) : P_n(t) \leq y \leq Q_{n,\mu}(t) \text{ and } 0 \leq t \leq t_0\}.$$

It is clear from Taylor's theorem and the assumption that $\rho > 0$, that by taking $n \rightarrow \infty$, and $\mu \rightarrow 1^+$ the sets $\mathcal{I}_{n,\mu}$ are arbitrary precise approximations of the graph $\{(e^{\rho t}, t) : t \geq 0\}$, what remains to show is that they are invariant.

- **Lemma 5.3.** *For every $\mu > 1$ there exists $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$ the set*

$$\mathcal{I}_{n,\mu} \cup \{(y, t) : t > t_0\}$$

*is an invariant containing the whole orbit, *i.e.* $\{(e^{\rho t}, t) : t \geq 0\}$.*

107:14 Invariants for Continuous Linear Dynamical Systems

The proof is in Appendix D of [3].

We can construct such invariants for every curve $e^{\rho_i t}$, and thus build $\tilde{\mathcal{I}}_{n,\mu}$ for

$$\{(e^{\rho_1 t}, \dots, e^{\rho_n t}, t) : t \geq 0\}.$$

To prove Theorem 5.2 we define \tilde{Y} by the formula

$$\Phi(x_1, \dots, x_n, x_{n+1}) \stackrel{\text{def}}{=} a_1 x_1 + \dots + a_n x_n = 0 \text{ and } 0 \leq x_{n+1} \leq t_0.$$

Since the analysis was done on the CDS $\langle J, \tilde{\mathbf{x}}_0 \rangle$, whose entries are not rational in general, before proceeding with the proof of Theorem 5.2, we need the following lemma to say that changing basis does not have an effect in the decision problem at hand:

► **Lemma 5.4.** *For every \tilde{Y} semi-algebraic, there exists another semi-algebraic set Y and \mathbf{x}_0 with rational entries such that the following two statements are equivalent:*

- $\langle J, \tilde{\mathbf{x}}_0 \rangle$ has a semi-algebraic invariant disjoint from \tilde{Y} , containing the whole orbit,
- $\langle PJP^{-1}, \mathbf{x}_0 \rangle$ has a semi-algebraic invariant disjoint from Y , containing the whole orbit.

The proof can be found in [3]. Thanks to this lemma, we can prove Theorem 5.2 for the CDS $\langle J, \tilde{\mathbf{x}}_0 \rangle$ and the set \tilde{Y} instead. This is done as follows. The direct implication is trivial. For the converse, observe that $f(t)$ does not have a zero in $[0, t_0]$ if and only if the $\mathcal{O}(0)$ and \tilde{Y} are disjoint. Since both $\mathcal{O}(0)$ and \tilde{Y} are closed sets, we can find a tube that contains $\mathcal{O}(0)$ and is disjoint from \tilde{Y} , *i.e.* there exists some $\mu > 1$ and $n \in \mathbb{N}$ such that

$$\tilde{\mathcal{I}}_{n,\mu} \cup \{(y, t) : t > t_0\},$$

is an invariant that is disjoint from \tilde{Y} but contains $\mathcal{O}(0)$.

References

- 1 Shaull Almagor, Dmitry Chistikov, Joël Ouaknine, and James Worrell. O-minimal invariants for linear loops. In *45th International Colloquium on Automata, Languages, and Programming ICALP*, pages 1–14. Schloss Dagstuhl, 2018.
- 2 Shaull Almagor, Dmitry Chistikov, Joël Ouaknine, and James Worrell. O-minimal invariants for discrete-time dynamical systems, 2020.
- 3 Shaull Almagor, Edon Kelmendi, Joël Ouaknine, and James Worrell. Invariants for continuous linear dynamical systems, 2020. [arXiv:2004.11661](https://arxiv.org/abs/2004.11661).
- 4 Rajeev Alur. *Principles of cyber-physical systems*. MIT Press, 2015.
- 5 Hirokazu Anai and Volker Weispfenning. Reach set computations using real quantifier elimination. In *International Workshop on Hybrid Systems: Computation and Control*, pages 63–76. Springer, 2001.
- 6 Paul C. Bell, Jean-Charles Delvenne, Raphaël M. Jungers, and Vincent D. Blondel. The continuous skolem-pisot problem. *Theor. Comput. Sci.*, 411(40–42):3625–3634, September 2010.
- 7 Mireille Broucke. Reachability analysis for hybrid systems with linear dynamics. *Mathematical Theory of Networks and Systems (MTNS'02)*, 2002.
- 8 John W.S. Cassels. *An Introduction to Diophantine Approximation*. Cambridge University Press, 1965.
- 9 Ventsislav Chonev, Joël Ouaknine, and James Worrell. On recurrent reachability for continuous linear dynamical systems. In *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science*, pages 515–524, 2016.
- 10 Ventsislav Chonev, Joël Ouaknine, and James Worrell. On the skolem problem for continuous linear dynamical systems. In *43rd International Colloquium on Automata, Languages, and Programming (ICALP 2016)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2016.

- 11 Peter Franek, Stefan Ratschan, and Piotr Zgliczynski. Quasi-decidability of a fragment of the first-order theory of real numbers. *Journal of Automated Reasoning*, 57(2):157–185, 2016.
- 12 Khalil Ghorbal, Andrew Sogokon, and André Platzer. A hierarchy of proof rules for checking positive invariance of algebraic and semi-algebraic sets. *Comput. Lang. Syst. Struct.*, 47:19–43, 2017.
- 13 Sumit Gulwani and Ashish Tiwari. Constraint-based approach for analysis of hybrid systems. In *Computer Aided Verification, 20th International Conference, CAV 2008, Proceedings*, volume 5123 of *Lecture Notes in Computer Science*, pages 190–203. Springer, 2008.
- 14 Emmanuel Hainry. Reachability in linear dynamical systems. In Arnold Beckmann, Costas Dimitracopoulos, and Benedikt Löwe, editors, *Logic and Theory of Algorithms*, pages 241–250. Springer Berlin Heidelberg, 2008.
- 15 Gerardo Lafferriere, George J. Pappas, and Sergio Yovine. Symbolic reachability computation for families of linear vector fields. *J. Symb. Comput.*, 32(3):231–253, 2001.
- 16 Jiang Liu, Naijun Zhan, and Hengjun Zhao. Computing semi-algebraic invariants for polynomial dynamical systems. In *Proceedings of the 11th International Conference on Embedded Software, EMSOFT 2011*, pages 97–106. ACM, 2011.
- 17 Angus Macintyre. Turing meets schanuel. *Annals of Pure and Applied Logic*, 167(10):901–938, 2016.
- 18 Angus Macintyre and Alex J. Wilkie. On the decidability of the real exponential field. In Piergiorgio Odifreddi, editor, *Kreiseliana. About and Around Georg Kreisel*, pages 441–467. A K Peters, 1996.
- 19 David W Masser. Linear relations on algebraic groups. *New Advances in Transcendence Theory*, pages 248–262, 1988.
- 20 Enric Rodríguez-Carbonell and Ashish Tiwari. Generating polynomial invariants for hybrid systems. In *Hybrid Systems: Computation and Control, 8th International Workshop, HSCC 2005, Proceedings*, volume 3414 of *Lecture Notes in Computer Science*, pages 590–605. Springer, 2005.
- 21 Andrew Sogokon, Khalil Ghorbal, Paul B. Jackson, and André Platzer. A method for invariant generation for polynomial continuous systems. In *Verification, Model Checking, and Abstract Interpretation - 17th International Conference, VMCAI, Proceedings*, volume 9583 of *Lecture Notes in Computer Science*, pages 268–288. Springer, 2016.
- 22 Andrew Sogokon and Paul B Jackson. Direct formal verification of liveness properties in continuous and hybrid dynamical systems. In *International Symposium on Formal Methods*, pages 514–531. Springer, 2015.
- 23 Thomas Sturm and Ashish Tiwari. Verification and synthesis using real quantifier elimination. In *Symbolic and Algebraic Computation, International Symposium, ISSAC 2011, Proceedings*, pages 329–336. ACM, 2011.
- 24 Alfred Tarski. A decision method for elementary algebra and geometry. *RAND Corporation, R-109*, 1951.
- 25 Lou van den Dries, Angus Macintyre, and David Marker. The elementary theory of restricted analytic fields with exponentiation. *Annals of Mathematics*, 140(1):183–205, 1994.
- 26 Lou Van den Dries, Chris Miller, et al. Geometric categories and o-minimal structures. *Duke Math. J.*, 84(2):497–540, 1996.
- 27 A. J. Wilkie. Model completeness results for expansions of the ordered field of real numbers by restricted pfaffian functions and the exponential function. *Journal of the American Mathematical Society*, 9(4):1051–1094, 1996.