

# A Tight $(3/2 + \varepsilon)$ Approximation for Skewed Strip Packing

Waldo Gálvez 

Technical University of Munich, Germany  
galvez@in.tum.de

Fabrizio Grandoni 

IDSIA, USI-SUPSI, Manno, Switzerland  
fabrizio@idsia.ch

Afrouz Jabal Ameli 

IDSIA, USI-SUPSI, Manno, Switzerland  
afrouz@idsia.ch

Klaus Jansen 

University of Kiel, Germany  
kj@informatik.uni-kiel.de

Arindam Khan 

Indian Institute of Science, Bangalore, India  
arindamkhan@iisc.ac.in

Malin Rau 

Univ. Grenoble Alpes, CNRS, Inria, Grenoble INP\*, LIG, Grenoble, France  
malin.rau@inria.fr

---

## Abstract

In the *Strip Packing* problem, we are given a vertical half-strip  $[0, W] \times [0, +\infty)$  and a collection of open rectangles of width at most  $W$ . Our goal is to find an axis-aligned (non-overlapping) packing of such rectangles into the strip such that the maximum height  $OPT$  spanned by the packing is as small as possible. Strip Packing generalizes classical well-studied problems such as *Makespan Minimization* on identical machines (when rectangle widths are identical) and *Bin Packing* (when rectangle heights are identical). It has applications in manufacturing, scheduling and energy consumption in smart grids among others. It is NP-hard to approximate this problem within a factor  $(3/2 - \varepsilon)$  for any constant  $\varepsilon > 0$  by a simple reduction from the *Partition* problem. The current best approximation factor for Strip Packing is  $(5/3 + \varepsilon)$  by Harren et al. [Computational Geometry '14], and it is achieved with a fairly complex algorithm and analysis.

It seems plausible that Strip Packing admits a  $(3/2 + \varepsilon)$ -approximation. We make progress in that direction by achieving such tight approximation guarantees for a special family of instances, which we call *skewed instances*. As standard in the area, for a given constant parameter  $\delta > 0$ , we call *large* the rectangles with width at least  $\delta W$  and height at least  $\delta OPT$ , and *skewed* the remaining rectangles. If all the rectangles in the input are large, then one can easily compute the optimal packing in polynomial time (since the input can contain only a constant number of rectangles). We consider the complementary case where all the rectangles are skewed. This second case retains a large part of the complexity of the original problem; in particular, it is NP-hard to approximate within a factor  $(3/2 - \varepsilon)$  and we provide an (almost) tight  $(3/2 + \varepsilon)$ -approximation algorithm.

**2012 ACM Subject Classification** Theory of computation  $\rightarrow$  Packing and covering problems

**Keywords and phrases** strip packing, approximation algorithm

**Digital Object Identifier** 10.4230/LIPIcs.APPROX/RANDOM.2020.44

**Category** APPROX



© Waldo Gálvez, Fabrizio Grandoni, Afrouz Jabal Ameli, Klaus Jansen, Arindam Khan, and Malin Rau;

licensed under Creative Commons License CC-BY

Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2020).

Editors: Jarosław Byrka and Raghu Meka; Article No. 44; pp. 44:1–44:18



Leibniz International Proceedings in Informatics

LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

**Funding** *Fabrizio Grandoni*: Partially supported by the SNSF Excellence Grant 200020B\_182865/1.

*Afrouz Jabal Ameli*: Partially supported by the SNSF Excellence Grant 200020B\_182865/1.

*Malin Rau*: Partially supported by the French research program ENERGUMEN ANR-18-CE25-0008.

## 1 Introduction

In this paper, we consider the *Strip Packing* problem, a well-studied classical rectangle packing problem (see Section 2 for a formal definition). Here, we are given a vertical half-strip of (integral) width  $W$ , plus a collection  $\mathcal{R}$  of rectangles of width at most  $W$ . Our objective is to find an axis-aligned packing of  $\mathcal{R}$  (where rectangles do not overlap) such that the maximum height spanned by the packing is as small as possible.

Strip Packing generalizes famous problems. For example, it generalizes Makespan Minimization (on identical machines) [11] when all the rectangle widths are 1 (here  $W$  would be the number of processors), and also generalizes Bin Packing [12] when all the rectangle heights are 1 (here the height  $OPT$  of the optimal solution would be the optimal number of bins). Strip Packing has several natural applications. For example, there are many manufacturing settings where rectangular pieces have to be cut out of some roll of raw material while using a rectangular piece of that roll of minimum length. Another application is the minimization of the peak energy consumption in smart-grids [28, 37]: here heights and widths model the energy consumption and duration, respectively, of a given set of jobs. For analogous reasons, it captures scenarios where a given set of jobs needs to be allocated a *consecutive* amount of a given resource (memory locations, frequencies, etc.) for a given amount of time.

Strip Packing is strongly NP-hard [17], and hence it is reasonable to consider approximation algorithms for it. A simple reduction from the *Partition* problem shows that it is not possible to obtain a  $(\frac{3}{2} - \varepsilon)$ -approximation algorithm (with polynomial running time) for any  $\varepsilon > 0$  unless  $P=NP$  (more details on this reduction are given later). The first non-trivial approximation algorithm for Strip Packing, with approximation ratio 3, was given by Baker, Coffman, and Rivest [4]. The First-Fit-Decreasing-Height algorithm (FFDH) by Coffman et al. [27] gives a 2.7-approximation. Sleator [35] gave an algorithm that generates a packing of height  $2OPT + \frac{h_{max}}{2}$ , where  $h_{max}$  is the maximum height of a rectangle in the instance, hence achieving a 2.5-approximation. Afterwards, Steinberg [36] and Schiermeyer [34] independently improved the approximation ratio to 2. Harren and van Stee [19] first broke the barrier of 2 with their 1.9396-approximation. The present best  $(\frac{5}{3} + \varepsilon)$ -approximation is due to Harren et al. [18].

The Strip Packing problem has also been studied in the pseudopolynomial setting, i.e., when  $N = n^{O(1)}$ . After a series of recent improvements [33, 1, 16, 20, 23], Jansen and Rau [22] have given a pseudopolynomial time algorithm with an almost tight  $(\frac{5}{4} + \varepsilon)$ -approximation ratio.

In terms of asymptotic approximations, the barrier of  $\frac{3}{2}$  can also be beaten. The best results in these terms are an AFPTAS presented by Kenyon and Rémila [29] which produces a solution of height  $(1 + \varepsilon)OPT + O(\frac{h_{max}}{\varepsilon^2})$ , and an APTAS which generates a solution of height  $(1 + \varepsilon)OPT + h_{max}$  by Jansen and Solis-Oba [24]. For the variant of Strip Packing *with Rotations*, where the rectangles are allowed to be rotated by 90 degrees, Jansen and van Stee [25] gave an APTAS (see also [13, 32] for related results).

### 1.1 Related Work

Strip Packing has rich connections with many other important geometric packing problems such as Two-dimensional Bin Packing (2BP) and Two-dimensional Geometric Knapsack (2GK). In 2BP, we are given a set of rectangles and unit square bins, and the goal is to pack

all the rectangles into minimum number of bins. The problem is known to be APX-hard [6] and the present best approximation ratio is 1.405 [7]. In 2GK, we are given a set of rectangles (with associated profit) and unit square knapsack, and the goal is to pack a subset of rectangles into the knapsack maximizing the total profit. This problem is strongly NP-hard even when all items are squares with unit profit [31]. The present-best approximation ratio is 1.89 due to Gálvez et al. [14] (see also [3, 26]).

Strip Packing has also been well studied for higher dimensions. The present best asymptotic approximation for 3-D Strip Packing is due to Jansen and Prädél [21] who presented a 1.5-approximation extending techniques from 2-D Bin Packing.

Another related problem is the *Independent Set of Rectangles* problem: here we are given a collection of axis-parallel rectangles embedded in the plane, and we need to find a maximum cardinality/weight subset of non-overlapping rectangles [2, 8, 9].

We refer the readers to [10, 30] for surveys on geometric packing problems.

## 1.2 Our Contribution

In this paper, we study a special case of Strip Packing, where all rectangles are *skewed*. In more detail, we say that a rectangle  $R$  is  $\delta$ -large if, for some fixed constant  $\delta > 0$ , its width is at least a  $\delta$  fraction of the width  $W$  of the strip and its height is at least a  $\delta$  fraction of the height  $OPT$  of the optimal packing; otherwise, the rectangle is  $\delta$ -skewed. We just say that a rectangle is large or skewed when  $\delta$  is clear from the context. An instance of Strip Packing is  $\delta$ -skewed if all the rectangles in the input are such.

This special case is non-trivial: in particular, the mentioned  $3/2 - \varepsilon$  hardness of approximation holds also for this special case with minor adaptations (see Section 5). We also believe that this special case is practically relevant: e.g., it captures scenarios where no job can consume a significant amount of the global resource (energy, memory space, etc.) for a significant amount of time. Our main result is as follows (see Sections 3-4).

► **Theorem 1.** *For a given constant  $\varepsilon' > 0$  and a small enough positive constant  $\delta \leq (\varepsilon')^{(1/\varepsilon')^{O(1)}}$ , there exists a polynomial-time  $(\frac{3}{2} + \varepsilon')$ -approximation algorithm for  $\delta$ -skewed Strip Packing.*

We remark that our algorithm does not need to recognize first if the instance is  $\delta$ -skewed: It always returns a feasible solution, but only if the instance satisfies the requirements, its approximation ratio is guaranteed.

Our result suggests that, in order to obtain a better (possibly  $3/2 + \varepsilon$ ) approximation for the general case of Strip Packing, one of the main obstacles is the interaction between large and skewed rectangles.

### 1.2.1 Organization

In Section 2, we introduce some useful notation and preliminary results. In Section 3, we prove the existence of a good enough packing with certain structural properties. The mentioned structure is exploited to derive an algorithm with the claimed approximation guarantee in Section 4. Section 5 contains our hardness of approximation result.

## 2 Preliminaries

A Strip Packing instance consists of a vertical strip of integral width  $W$  in the two-dimensional plane, i.e.  $[0, W] \times \mathbb{R}_{\geq 0}$ , and a set of open rectangles  $\mathcal{R}$ , where each rectangle  $R \in \mathcal{R}$  is characterized by its integral height  $h(R)$  and integral width  $w(R)$ . An embedding of  $\mathcal{R}$  is

given by specifying a bottom-left position  $(x(R), y(R))$  for each  $R \in \mathcal{R}$ . The interpretation is that  $R$  is embedded in the plane in the region  $(x(R), x(R) + w(R)) \times (y(R), y(R) + h(R))$ . An embedding is a feasible packing into the strip if the following two conditions hold: (1) each  $R \in \mathcal{R}$  is embedded inside the strip, namely  $0 \leq x(R) \leq W - w(R)$  and  $y(R) \geq 0$  and (2) rectangles do not overlap, namely, for any two  $R, S \in \mathcal{R}$ ,  $\left( (x(R), x(R) + w(R)) \times (y(R), y(R) + h(R)) \right) \cap \left( (x(S), x(S) + w(S)) \times (y(S), y(S) + h(S)) \right) = \emptyset$ . The height of a feasible packing is the maximum height spanned by any embedded rectangle, namely the maximum value of  $y(R) + h(R)$ . The goal of Strip Packing is to compute a feasible packing of minimum height  $OPT$ . W.l.o.g. we can restrict our attention to packings where the coordinates  $(x(R), y(R))$  are integral as any feasible packing can be transformed into a feasible packing satisfying this property (intuitively, by pushing rectangles to the bottom-left as much as possible while keeping feasibility).

Given a subset of rectangles  $\mathcal{S} \subseteq \mathcal{R}$ , we denote by  $w(\mathcal{S}) := \sum_{R \in \mathcal{S}} w(R)$ ,  $h(\mathcal{S}) := \sum_{R \in \mathcal{S}} h(R)$ , and  $a(\mathcal{S}) := \sum_{R \in \mathcal{S}} h(R)w(R)$  the total width, height, and area of  $\mathcal{S}$ , respectively. The operation of changing the bottom-left corner of a rectangle  $R$  in a given packing from  $(x(R), y(R))$  to  $(x(R), y(R) + a)$  will be denoted by *shifting  $R$  vertically by  $a$* . Analogously, changing the bottom-left coordinate from  $(x(R), y(R))$  to  $(x(R) + a, y(R))$  will be denoted by *shifting  $R$  horizontally by  $a$* . These operations are only allowed if the resulting packing is feasible.

A *box* of size  $w \times h$  denotes a rectangular region of width  $w$  and height  $h$ . We sometimes embed boxes into the strip similarly to rectangles. A monotone polygonal chain is a curve specified by a sequence of points  $(A_1, A_2, \dots, A_n)$  called its vertices. The curve itself consists of the line segments connecting the consecutive vertices, and we require that the x-coordinates of points  $A_i$  are non-decreasing and the segments are horizontal or vertical. We say that a rectangle  $R$  in the packing *lies above (resp. below)* one such  $P$  if for any  $x_1 \in (x(R), x(R) + w(R))$  we have that  $y(R)$  (resp.  $y(R) + h(R)$ ) is not smaller (resp. not larger) than the largest (resp., smallest) y-coordinate of  $P$  at x-coordinate  $x_1$ .

We can assume w.l.o.g. that  $W$  is lower bounded by a sufficiently large constant, in particular  $W \geq 1/\varepsilon$ . Otherwise one easily obtains a PTAS for  $\delta$ -skewed instances<sup>1</sup>.

## 2.1 Next Fit Decreasing Height

One of the most recurring tools used as a subroutine in countless results on geometric packing problems is the Next Fit Decreasing Height (NFDH) algorithm, which was originally analyzed in [27]. We will use a variant of this algorithm to pack rectangles inside a rectangular box and analyze its properties. We provide a full proof for the sake of completeness.

Suppose we are given a box  $C$  of size  $w \times h$ , and a set of rectangles  $\mathcal{R}'$  each one fitting in the box. NFDH computes in polynomial time a packing of  $\mathcal{R}'' \subseteq \mathcal{R}'$  as follows. It sorts the rectangles  $R \in \mathcal{R}'$  in non-increasing order of height  $h(R)$ , and considers rectangles in that order  $R_1, \dots, R_n$ . Then the algorithm works in rounds  $j \geq 1$ . At the beginning of round  $j$  it is given an index  $n(j)$  and a horizontal segment  $L(j)$  going from the left to the right side of  $C$ . Initially  $n(1) = 1$  and  $L(1)$  is the bottom side of  $C$ . In round  $j$  the algorithm packs a maximal set of rectangles  $R_{n(j)}, \dots, R_{n(j+1)-1}$ , with bottom side touching  $L(j)$  one next to the other from left to right (a *shelf*). The segment  $L(j+1)$  is the horizontal segment

<sup>1</sup> Choosing  $\delta$  such that  $\delta W < 1$  enforces each rectangle to have height at most  $\delta OPT$  (otherwise it would be large). A PTAS follows, e.g., from [24].

containing the top side of  $R_{n(j)}$  and ranging from the left to the right side of  $C$ . The process halts at round  $r$  when either all rectangles have been packed or  $R_{n(r+1)}$  does not fit above  $R_{n(r)}$ .

We prove the following standard result.

► **Lemma 2** ([27]). *Let  $C$  be a given box of size  $w \times h$  and  $\mathcal{R}' \subseteq \mathcal{R}$ . Assume that, for some given parameter  $\varepsilon' \in (0, 1)$ , for each  $R \in \mathcal{R}'$  one has  $w(R) \leq \varepsilon'w$  and  $h(R) \leq \varepsilon'h$ . Then NFDH is able to pack in  $C$  a subset  $\mathcal{R}'' \subseteq \mathcal{R}'$  of area at least  $a(\mathcal{R}'') \geq \min\{a(\mathcal{R}'), (1-2\varepsilon')w \cdot h\}$ . In particular, if  $a(\mathcal{R}') \leq (1-2\varepsilon')w \cdot h$ , all rectangles in  $\mathcal{R}'$  are packed.*

**Proof.** The claim trivially holds if all rectangles are packed. Thus suppose that this is not the case. Observe that  $\sum_{j=1}^{r+1} h(R_{n(j)}) > h$ , otherwise rectangle  $R_{n(r+1)}$  would fit in the next shelf above  $R_{n(r)}$ ; hence  $\sum_{j=2}^{r+1} h(R_{n(j)}) > h - h(R_{n(1)}) \geq (1 - \varepsilon')h$ . Observe also that the total width of rectangles packed in each round  $j$  is at least  $w - \varepsilon'w = (1 - \varepsilon')w$ , since  $R_{n(j+1)}$ , of width at most  $\varepsilon'w$ , does not fit to the right of  $R_{n(j+1)-1}$ . It follows that the total area of the rectangles packed in round  $j$  is at least  $(w - \varepsilon'w)h(R_{n(j+1)-1})$ , and thus

$$a(\mathcal{R}'') \geq \sum_{j=1}^r (1 - \varepsilon')w \cdot h(R_{n(j+1)-1}) \geq (1 - \varepsilon')w \sum_{j=2}^{r+1} h(R_{n(j)}) \geq (1 - \varepsilon')^2 w \cdot h \geq (1 - 2\varepsilon')w \cdot h. \blacktriangleleft$$

## 2.2 Container Packings

Similar to recent work on related problems (e.g., [14, 5]), we will exploit a *container-based* packing approach. The idea is to partition the solution into a constant number of axis-aligned rectangular regions (*containers*). The sizes (and therefore positions) of these containers can be *guessed* in polynomial time, and subsequently, rectangles are packed inside the containers in a simple way: either one next to the other from left to right (*vertical container*), or one on top of the other from bottom to top (*horizontal container*), or by means of NFDH (*area container*). We further require that the rectangles  $R$  packed into an area container of size  $w \times h$  satisfy  $w(R) \leq \varepsilon'w$  and  $h(R) \leq \varepsilon'h$  for a constant  $\varepsilon' > 0$  to be fixed later. We call this an  $\varepsilon'$ -area container.

We will make use of the following standard PTAS to pack rectangles into a constant number of containers. The basic idea is to reduce the problem to an instance of the Maximum Generalized Assignment Problem (GAP) with one bin per container, and then use a PTAS for the latter problem plus NFDH to repack rectangles in area containers. We recall that in GAP, we are given a collection of  $n$  items and a set of  $k$  (one-dimensional) bins, each one characterized by a positive size. Each item has a profit<sup>2</sup> and a positive size per bin (possibly different for different bins). Our goal is to compute a maximum profit subset of items and an assignment of them into the bins so that the total size of items packed in each bin is at most the size of the bin. GAP admits a PTAS for constant  $k$  (see e.g. Section E.2 in [15]) and the following lemma shows how to use it to pack the rectangles into a given set of containers.

► **Lemma 3.** *For any constant  $\varepsilon' > 0$ , given a set of rectangles  $\mathcal{R}$  that can be packed into a given set of  $k$  containers (each container being either vertical, horizontal or  $\varepsilon'$ -area),  $k$  constant, there is an algorithm to pack  $\mathcal{R}' \subseteq \mathcal{R}$  with  $a(\mathcal{R}') \geq (1 - 3\varepsilon')a(\mathcal{R})$  into the mentioned containers.*

<sup>2</sup> The same item might have different profit on different knapsacks; however, we do not need this extension here.

**Proof.** We let  $w(C_j) \times h(C_j)$  be the size of the  $j$ -th container  $C_j$ . We build an instance of GAP as follows. There is one item  $R$  per rectangle  $R \in \mathcal{R}$ , with profit  $a(R)$ . For each horizontal container  $C_j$ , we create a knapsack  $j$  of size  $S_j := h(C_j)$ . Furthermore, we define the size  $s(R, j)$  of rectangle  $R$  w.r.t. knapsack  $j$  as  $h(R)$  if  $h(R) \leq h(C_j)$  and  $w(R) \leq w(C_j)$ . Otherwise  $s(R, j) = +\infty$  (meaning that  $R$  does not fit in  $C_j$ ). The construction for vertical containers is symmetric. For each area container  $C_j$  we create a knapsack  $j$  of size  $S_j = a(C_j)$  and define the size  $s(R, j)$  of rectangle  $R$  w.r.t. knapsack  $j$  as  $a(R)$  if  $h(R) \leq \varepsilon' h(C_j)$  and  $w(R) \leq \varepsilon' w(C_j)$ , setting  $b(R, j) = +\infty$  otherwise (meaning that the rectangle is not small enough with respect to the dimensions of the container).

We next apply the mentioned PTAS for GAP to this instance, so as to obtain a solution  $\mathcal{R}''$  to GAP of profit at least  $a(\mathcal{R}'') \geq (1 - \varepsilon')a(\mathcal{R})$ . We build a feasible packing of  $\mathcal{R}' \subseteq \mathcal{R}''$  into the containers as follows. Let  $\mathcal{R}_j$  be the items packed into knapsack  $j$ . If  $C_j$  is vertical, we pack rectangles  $\mathcal{R}_j$  into this container bottom-most and from left to right one next to the other in any order. By definition all rectangles  $\mathcal{R}_j$  will fit. A symmetric construction works if  $C_j$  is horizontal. If  $C_j$  is area, we pack a subset  $\mathcal{R}'_j$  of  $\mathcal{R}_j$  into it using NFDH. By Lemma 2, we either have  $\mathcal{R}'_j = \mathcal{R}_j$ , or it must be the case that  $a(\mathcal{R}'_j) \geq (1 - 2\varepsilon')w(C_j)h(C_j) = (1 - 2\varepsilon')a(C_j)$ . Consider the second case. Let  $\mathcal{R}''_j = \mathcal{R}_j \setminus \mathcal{R}'_j$  be the rectangles which are not packed. Observe that  $a(\mathcal{R}_j) \leq a(C_j)$  by the feasibility of the GAP solution, hence

$$\frac{a(\mathcal{R}''_j)}{a(\mathcal{R}_j)} = 1 - \frac{a(\mathcal{R}'_j)}{a(\mathcal{R}_j)} \leq 1 - \frac{(1 - 2\varepsilon')a(C_j)}{a(\mathcal{R}_j)} \leq 2\varepsilon'.$$

Thus altogether  $a(\mathcal{R}') \geq a(\mathcal{R}'')(1 - 2\varepsilon') \geq a(\mathcal{R})(1 - 2\varepsilon')(1 - \varepsilon') \geq a(\mathcal{R})(1 - 3\varepsilon')$ .  $\blacktriangleleft$

Notice that the containers may have considerable free space inside, but the lemma just claims that the total area of the rectangles that the algorithm is not packing is negligible. Whenever this lemma is applied, we will pack the remaining rectangles into an extra rectangular box of small area and carefully argue where to place it.

### 2.3 Classification of Rectangles

From now on, we will assume that instance  $(\mathcal{R}, W)$  is  $\delta$ -skewed for some  $\delta > 0$  to be fixed later. By  $OPT$ , we denote both the optimal height and an optimal packing: the meaning will be clear from the context. We can assume that  $OPT$  is even (otherwise, multiply heights by a factor 2).

We will assume that our algorithm is given in the input a value  $OPT'$  such that  $OPT \leq OPT' \leq (1 + \varepsilon)OPT$ . This assumption can be removed as follows. We compute, say, a 2-approximation  $APX$  by Steinberg's algorithm for Strip Packing and then run our algorithm for all the (constantly many) values  $OPT' = (1 + \varepsilon)^j \frac{APX}{2(1 + \varepsilon)}$  which fit in the range  $[\frac{APX}{2(1 + \varepsilon)}, APX(1 + \varepsilon)]$ . One of these values will satisfy the claim. In order to keep the notation light, we simply use  $OPT$  to denote this value  $OPT'$ . Therefore, all the approximation factors should be scaled by a factor  $(1 + \varepsilon)$  in order to consider the true value of  $OPT$ .

Let  $\varepsilon > 0$  and assume for simplicity that  $\frac{1}{\varepsilon} \in \mathbb{N}$ . We will classify the rectangles according to their heights as follows:

- The set of *tall* rectangles  $\mathcal{T} = \{R \in \mathcal{R} : h(R) > \frac{1}{2}OPT\}$ ;
- The set of *vertical* rectangles  $\mathcal{V} = \{R \in \mathcal{R} : h(R) \in (\delta OPT, \frac{1}{2}OPT]\}$ ;
- The set of *short* rectangles  $\mathcal{S} = \{R \in \mathcal{R} : h(R) \leq \delta OPT\}$ ;

## 2.4 Linear Grouping

We need the following lemma whose proof (given in the appendix) is based on *linear grouping*, a standard technique in the area of rectangle packing. Given a subset  $\mathcal{S}$  of rectangles, let  $\mathcal{S}_{hslice}$  be the set of rectangles obtained by taking each  $R \in \mathcal{S}$  and replacing it with  $h(R)$  rectangles of height 1 and width  $w(R)$  (*horizontal slices*). We define symmetrically the set  $\mathcal{S}_{vslice}$  of *vertical slices*. Notice that any embedding of  $\mathcal{S}$  naturally induces an embedding of  $\mathcal{S}_{hslice}$  and  $\mathcal{S}_{vslice}$ .

► **Lemma 4.** *Let  $\varepsilon' > 0$  be a given constant,  $\mathcal{P}$  be a rectangular region of size  $W \times H$  and  $\mathcal{H}$  be a subset or rectangles of height at most  $\delta \cdot H$  each for some constant  $\delta \in [1, 0)$ . Suppose that  $\mathcal{H}_{hslice}$  can be packed into a set  $\mathcal{B}$  of  $K = O_{\varepsilon'}(1)$  boxes contained in  $\mathcal{P}$ . Then, for  $\delta \leq (\varepsilon'/K)^{(K/\varepsilon')^{O(1)}}$ , there exists a partition of  $\mathcal{H}$  into two sets  $\mathcal{H}^{cont}$  and  $\mathcal{H}^{disc}$  such that:*

1.  $\mathcal{H}^{cont}$  can be packed into a set of at most  $K' = O_{\varepsilon'}(1)$  horizontal and  $\varepsilon'$ -area containers, where each container is fully contained in some box in  $\mathcal{B}$ .
2.  $\mathcal{H}^{disc}$  can be packed in one horizontal container of size  $\max_{R \in \mathcal{H}}\{w(R)\} \times (\varepsilon')^2 H$  and one  $\varepsilon'$ -area container of size  $\varepsilon' W \times \varepsilon' H$ .
3. The sizes of the above containers belongs to a set that can be computed in polynomial time.

A symmetric claim holds for a subset of rectangles  $\mathcal{V}'$  of width at most  $\delta \cdot W$  such that  $\mathcal{V}'_{vslice}$  can be packed into the corresponding boxes.

## 3 Existence of a Structured Solution

In this section, we will prove our main structural result.

► **Theorem 5.** *For any given constant  $\varepsilon > 0$  and any given instance of  $\delta$ -skewed Strip Packing  $(\mathcal{R}, W)$  with  $\delta = \Omega_{\varepsilon}(1)$  small enough, there exists a feasible container packing such that the following holds:*

1. The total height of the packing is  $(\frac{3}{2} + O(\varepsilon)) OPT$ .
2. The number of containers is  $O_{\varepsilon}(1)$  and their possible sizes belong to a set that can be computed in polynomial time.
3. Given any fixed ordering of  $\mathcal{T}$  in non-increasing order of height,  $\mathcal{T}$  can be partitioned into subsequences each one fitting in precisely one vertical container.
4. The packing leaves a free rectangular region (**free box**) of size  $\varepsilon^2 W \times \frac{1}{2} OPT$ .

To achieve the above result, we proceed in three steps:

1. We describe a packing of  $\mathcal{T} \cup \mathcal{S}_{hslice}$  with height at most  $(3/2 + O(\varepsilon)) OPT$  (see Section 3.1) into  $O_{\varepsilon}(1)$  boxes. This packing leaves a free space of at least  $(1/2 + \Omega(\varepsilon)) OPT + a(\mathcal{V})$ .
2. We describe (see Section 3.2) how to pack  $\mathcal{V}_{vslice}$  within the free space of the previous packing using  $O_{\varepsilon}(1)$  extra boxes. Furthermore, we guarantee that there is a free box (not containing any rectangle) of size at least  $\Omega(\varepsilon) W \times \frac{1}{2} OPT$ . Guaranteeing the latter property is critical, and it is the main technical novelty in our approach.
3. Finally (see Section 3.3), we convert the above packing into a feasible container packing (via Lemma 4) inside the above boxes. The residual containers that do not fit into the boxes can be placed inside the free box (still leaving enough space) plus a new box of size  $W \times O(\varepsilon^2) OPT$  that can be placed on top of the previous packing.

The reason for leaving a free box will be clearer in Section 4, where we will describe our final algorithm.

### 3.1 Packing of $\mathcal{T} \cup \mathcal{S}_{hslice}$

In this section, we describe a packing of  $\mathcal{T} \cup \mathcal{S}_{hslice}$ . The proof of the following Lemma is illustrated in Figure 1.

► **Lemma 6.** *For any given constant  $\varepsilon \in (0, 1/4]$  with  $1/\varepsilon$  integral and  $\delta = \Omega_\varepsilon(1)$  sufficiently small, it is possible to pack  $\mathcal{T} \cup \mathcal{S}_{hslice}$  into the region  $\mathcal{P} = [0, W] \times [0, (\frac{3}{2} + 15\varepsilon)OPT]$  in such a way that:*

1. *Rectangles in  $\mathcal{T}$  are packed into at most  $1/\varepsilon$  vertical boxes, slices in  $\mathcal{S}_{hslice}$  are packed into at most  $1/\varepsilon + 1$  horizontal boxes, and the remaining area is partitioned into at most  $2/\varepsilon$  free boxes. Furthermore, given any fixed ordering of  $\mathcal{T}$  in non-increasing order of height, it is possible to partition  $\mathcal{T}$  into subsequences such that each subsequence fits into precisely one vertical box.*
2. *The sizes of the boxes belong to a set that can be computed in polynomial time.*
3. *The total area of the free boxes is at least  $(\frac{1}{2} + 9\varepsilon)OPT \cdot W + a(\mathcal{V})$ .*


**Proof.** Consider the embedding of  $\mathcal{T} \cup \mathcal{S}_{hslice}$  induced by the optimum solution. Let us draw the horizontal line  $y = \frac{1}{2}OPT$  and partition  $\mathcal{S}_{hslice}$  into two sets  $\mathcal{S}_{hslice}^{top}$  and  $\mathcal{S}_{hslice}^{bottom}$  corresponding to the rectangles in  $\mathcal{S}_{hslice}$  which are packed above and below the line  $y = \frac{1}{2}OPT$  respectively (notice that this line does not intersect any rectangle in  $\mathcal{S}_{hslice}$  as  $OPT$  is even by assumption). If we shift up rectangles in  $\mathcal{S}_{hslice}^{bottom}$  by  $OPT$ , we obtain a feasible packing (since the region  $[0, W] \times [OPT, \frac{3}{2}OPT]$  was empty) with final height at most  $\frac{3}{2}OPT$ . Notice that every rectangle in  $\mathcal{T}$  intersects the horizontal segment  $[0, W] \times \{\frac{1}{2}\}$ .

Now, let us shift down each rectangle  $R$  in  $\mathcal{T}$  so that its bottom coordinate becomes zero (again, the packing remains feasible). Next, we shift rectangles so that the ones in  $\mathcal{T}$  appear one next to the other in the bottom left part of the packing, in non-increasing order of height. To this aim, we proceed recursively as follows. Let  $T_1, \dots, T_q$  be the considered ordering of  $\mathcal{T}$  in non-increasing order of height. At the beginning of iteration  $i \geq 1$ , we are given a feasible packing where  $T_1, \dots, T_{i-1}$  are packed from left to right one next to the other as left as possible (and with bottom coordinate 0). We consider the region  $A_i := [w_{i-1}, W] \times [0, h(T_i)]$ , where  $w_{i-1} = \sum_{j \leq i-1} w(T_j)$ . Let  $L_i$  be the portion of  $A_i$  to the left of (the current embedding of)  $T_i$ . Note that every rectangle is either completely contained in or disjoint from  $L_i$  since  $T_i$  is the tallest rectangle contained in  $A_i$  and  $T_{i-1}$  is taller than  $T_i$ . We move  $T_i$  so that its left coordinate is  $w_{i-1}$ , and shift  $L_i$  to the right by  $w(T_i)$ , moving consistently all rectangles in  $L_i$ . Obviously the new packing satisfies the invariant for the next iteration. At the end of iteration  $q$  the packing satisfies the claim.

In the next step, we partition the area not occupied by  $\mathcal{T}$  into unit-height stripes. Notice that each rectangle in  $\mathcal{S}_{hslice}$  is fully contained in some stripe. We need (for a reason that will be clearer later) to temporarily discard, meaning that we remove them from the packing, some slices as follows. Let us say that a slice is *wide* if its width is at least  $\varepsilon W$ , and *narrow* otherwise. Consider the slices  $\mathcal{S}'_{hslice}$  in a given horizontal stripe. This set contains at most  $1/\varepsilon$  wide slices. Let  $w'$  be the total width of the remaining narrow slices and let  $w'' \leq w'$  be the largest multiple of  $\varepsilon W$ . We discard a minimal subset of narrow slices so that the remaining ones have width at most  $w''$ . We let  $\mathcal{S}_{hslice}^{disc}$  be the set of discarded slices, and  $\mathcal{S}_{hslice}^{sel}$  be the remaining (selected) slices.

Next, we push slices  $\mathcal{S}_{hslice}^{sel}$  as right as possible. Afterward, we permute the  $y$ -coordinates of slices in pairs of stripes so that stripes are sorted from top to bottom in non-increasing order of the total width of the slices contained in them. Observe that this cannot create any conflict with  $\mathcal{T}$  (hence the packing remains feasible).





■ **Figure 1** Depiction of the proof of Lemma 6. **Left:** Packing of  $\mathcal{T} \cup \mathcal{S}_{hslice}$  in the optimal solution. Light gray rectangles correspond to  $\mathcal{S}_{hslice}$ , dark gray rectangles correspond to  $\mathcal{T}$ . **Center:** By shifting  $\mathcal{S}_{hslice}^{bottom}$  to the top we can shift down the rectangles in  $\mathcal{T}$ . **Right:** We can shift now horizontally rectangles in  $\mathcal{T}$  and sort stripes. Finally one obtains  $C_{up}$  and  $C_{dw}$ .

We proceed as follows. We shift up  $\mathcal{S}_{hslice}^{sel}$  by  $11\epsilon \cdot OPT$  and construct a polygonal chain  $C_{up}$  with the following procedure. The chain starts at coordinate  $p_0^{up} = (0, (\frac{3}{2} + 11\epsilon)OPT)$ . We extend the chain to the right (possibly by a zero amount) until the chain hits some rectangle in  $\mathcal{S}_{hslice}^{sel}$ . We denote this point by  $(x_1, (\frac{3}{2} + 11\epsilon)OPT)$  and extend the chain down by  $\epsilon OPT$ , hence reaching some point  $p_1^{up} = (x_1, (\frac{3}{2} + 10\epsilon)OPT)$ . We continue from  $p_1^{up}$  in the same fashion. The procedure ends when the chain reaches the  $x$ -coordinate  $W$ . Observe that, by construction, every rectangle in  $\mathcal{S}_{hslice}^{sel}$  lies above  $C_{up}$ . Furthermore,  $C_{up}$  is defined by axis parallel segments only and at most  $1/\epsilon$  of them.

Afterward, we build symmetrically a polygonal chain  $C_{dw}$  as follows. We start from  $p_0^{dw} = (W, 0)$  and extend the chain to the left (possibly by a zero amount) until we hit some rectangle  $R$  in  $\mathcal{T}$ . Then, we extend the chain up by  $\epsilon OPT$ , hence reaching a point  $p_1^{dw} = (x_1, \epsilon OPT)$ . We continue from  $p_1^{dw}$  in the same fashion until we reach the  $x$ -coordinate  $0$ . Notice that each rectangle from  $\mathcal{T}$  lies below  $C_{dw}$ . Notice also that  $C_{dw}$  is defined by axis parallel segments only and at most  $1/\epsilon$  of them. Furthermore, it is fully below  $C_{up}$ . To see the latter, take any coordinate  $x \in (0, W)$  (which, for simplicity, is not the position of a vertex of any one of the two chains). Let  $y_{up}$  and  $y_{dw}$  be the corresponding  $y$ -coordinates in  $C_{up}$  and  $C_{dw}$ , resp. Suppose by contradiction that  $y_{dw} > y_{up}$ . Observe that by construction the segment  $x \times (y_{up}, y_{up} + \epsilon OPT)$  must hit some rectangle  $R_{up}$  in  $\mathcal{S}_{hslice}^{sel}$ . Symmetrically, the segment  $x \times (y_{dw} - \epsilon OPT, y_{dw})$  must hit some rectangle  $R_{dw}$  in  $\mathcal{T}$ . This however implies that  $R_{up}$  and  $R_{dw}$  overlapped before the shifting up of  $\mathcal{S}_{hslice}^{sel}$  by  $11\epsilon OPT$ , a contradiction<sup>3</sup>.

We claim that points  $p_i^{up}$  and  $p_i^{dw}$  have coordinates that belong to a set that can be computed in polynomial time. Recall that  $OPT$  is known to the algorithm. Notice that the  $y$  coordinates of the points  $p_i^{up}$  and  $p_i^{dw}$  are multiples of  $\epsilon OPT$ , hence they satisfy the requirement. Since we know the precise packing of  $\mathcal{T}$ , we can compute the  $x$ -coordinates of points  $p_i^{dw}$  explicitly. The  $x$ -coordinates of points  $p_i^{up}$  have value  $W$  minus the sum  $w''$  of the widths of the slices in a given strip. By the previous discarding procedure,  $w''$  is the sum of up to  $1/\epsilon$  widths of input rectangles, plus a multiple of  $\epsilon W$ . Hence we can compute the set of the possible coordinates in polynomial time.

<sup>3</sup> A shifting up by  $2\epsilon OPT$  would be sufficient to achieve a contradiction here. The extra shift by  $9\epsilon OPT$  is used to create some more free space that is needed in the following arguments.

## 44:10 A Tight $(3/2 + \varepsilon)$ Approximation for Skewed Strip Packing

Let us subdivide the area in the strip between  $C_{up}$  and  $[0, W] \times (\frac{3}{2} + 11\varepsilon)OPT$  by extending to the right the horizontal segments in  $C_{up}$ . This gives up to  $1/\varepsilon$  boxes  $\mathcal{B}_{up}$  that fully contain  $\mathcal{S}_{hslice}^{sel}$ . Symmetrically, we can subdivide the area in the strip between  $C_{dw}$  and  $[0, W]$  by extending down the vertical segments in  $C_{dw}$ . This provides up to  $1/\varepsilon$  boxes  $\mathcal{B}_{dw}$  that fully contain  $\mathcal{T}$ . Next, consider the free area between  $\mathcal{B}_{up}$  and  $\mathcal{B}_{dw}$ . By extending down the vertical sides of the boxes in  $\mathcal{B}_{up}$  until reaching  $\mathcal{B}_{dw}$  and symmetrically extending up the vertical sides of the boxes in  $\mathcal{B}_{dw}$  until reaching  $\mathcal{B}_{up}$ , we obtain a partition of the free area into up to  $2/\varepsilon$  free boxes  $\mathcal{B}_{free}$ . By the previous discussion, the possible sizes of all the mentioned boxes can be computed in polynomial time.

It remains to pack  $\mathcal{S}_{hslice}^{disc}$ . To that aim, we create a new box  $\mathcal{B}_{disc}$  of width  $W$  and height  $4\varepsilon OPT$  that we place on top of the current packing, hence increasing the total height to  $(\frac{3}{2} + 15\varepsilon)OPT$ . Notice that each  $R \in \mathcal{S}_{hslice}^{disc}$  satisfies  $w(R) \leq \varepsilon W$  and  $h(R) = 1 \leq \delta OPT$ . Hence assuming  $\varepsilon \leq 1/4$  and  $\delta \leq 4\varepsilon^2$ , Lemma 2 with  $\varepsilon' = 1/4$  guarantees that  $\mathcal{S}_{hslice}^{disc}$  can be fully packed in  $\mathcal{B}_{disc}$  via NFDH.

Properties (1) and (2) follow by construction. It remains to prove (3). Notice that by construction the area inside  $\mathcal{B}_{up}$  not occupied by  $\mathcal{S}_{hslice}^{sel}$  is at most  $\varepsilon OPT \cdot W$ . Indeed, as observed earlier, if we take any point  $(x, y)$  along  $C_{up}$ , where  $x$  does not correspond to a step of  $C_{up}$ , the segment  $x \times [y, y + \varepsilon OPT]$  hits some rectangle in  $\mathcal{S}_{hslice}^{sel}$ . Thus  $a(\mathcal{B}_{up}) \leq a(\mathcal{S}_{hslice}^{sel}) + \varepsilon OPT \cdot W$ . A symmetric argument shows that  $a(\mathcal{B}_{dw}) \leq a(\mathcal{T}) + \varepsilon OPT \cdot W$ . We can therefore conclude that

$$\begin{aligned} a(\mathcal{B}_{free}) &\geq W \cdot \left(\frac{3}{2} + 11\varepsilon\right) OPT - a(\mathcal{S}_{hslice}^{sel}) - a(\mathcal{T}) - 2\varepsilon OPT \cdot W \\ &\geq W \cdot \left(\frac{3}{2} + 9\varepsilon\right) OPT - a(\mathcal{S} \cup \mathcal{T}) \\ &\geq W \cdot \left(\frac{3}{2} + 9\varepsilon\right) OPT - OPT \cdot W + a(\mathcal{V}) = W \cdot \left(\frac{1}{2} + 9\varepsilon\right) OPT + a(\mathcal{V}). \quad \blacktriangleleft \end{aligned}$$

### 3.2 Including $\mathcal{V}_{vslice}$


In this section, we show how to incorporate  $\mathcal{V}_{vslice}$  in the packing from the previous subsection. Critically, we need to leave a free box of sufficiently large size.

► **Lemma 7.** *Consider the packing from Lemma 6 and assume  $\varepsilon$  is small enough. It is possible to pack  $\mathcal{V}_{vslice}$  inside the free boxes and furthermore define an empty rectangular region of size  $2\varepsilon^2 W \times \frac{1}{2} OPT$ .*

**Proof.** Consider the set of (at most)  $2/\varepsilon$  free boxes  $\mathcal{B}_1, \dots, \mathcal{B}_q$  sorted non-decreasingly by height. We partition them into unit-width vertical stripes  $\mathcal{S}' = \{S_1, \dots, S_k\}$  sorted in the same order, and breaking ties so that stripes of the same box appear consecutively. Recall that  $a(\mathcal{S}') \geq (\frac{1}{2} + 9\varepsilon) OPT \cdot W + a(\mathcal{V})$ .

We next place slices of  $\mathcal{V}_{vslice}$  in these stripes from bottom to top in a greedy manner. In particular, we consider rectangles  $R$  in  $\mathcal{V}_{vslice}$  in any order, and place  $R$  in the left-most stripe where it fits, as low as possible. Assume that the non-empty stripes are  $\mathcal{S}'_{used}$ . Notice that the unused space in these stripes is at most  $(w(\mathcal{S}'_{used}) - 1)(\frac{OPT}{2} - 1) + OPT \leq (\frac{w(\mathcal{S}'_{used}) + 1}{2}) OPT$ , hence

$$a(\mathcal{S}'_{used}) \leq a(\mathcal{V}) + \frac{w(\mathcal{S}'_{used}) + 1}{2} OPT.$$



■ **Figure 2** Description of the packing of  $\mathcal{V}_{vslice}$ . **Left:** Packing of  $\mathcal{T} \cup \mathcal{S}_{hslice}$  as described in Lemma 6 and the  $O_\epsilon(1)$  boxes for the free area defined by the dashed lines. **Right:** Boxes in the free area sorted by height. Even if we ignore  $\frac{1}{2}OPT$  height from each box we have enough space for  $\mathcal{V}_{vslice}$  and even to reserve space for future discarded vertical rectangles.

Let us distinguish the unused stripes  $\mathcal{S}'_{unused}$  into the ones  $\mathcal{S}^{tall}_{unused}$  of height at least  $OPT/2$ , and the remaining ones  $\mathcal{S}^{short}_{unused}$ . One has that

$$\begin{aligned}
a(\mathcal{S}^{tall}_{unused}) &\geq \left(\frac{1}{2} + 9\epsilon\right)OPT \cdot W + a(\mathcal{V}) - a(\mathcal{S}'_{used}) - a(\mathcal{S}^{short}_{unused}) \\
&\geq \left(\frac{1}{2} + 9\epsilon\right)OPT \cdot W + a(\mathcal{V}) - \left(a(\mathcal{V}) + \frac{(w(\mathcal{S}'_{used}) + 1)}{2}OPT\right) \\
&\quad - \frac{OPT}{2}w(\mathcal{S}^{short}_{unused}) \\
&= \left(\frac{1}{2} + 9\epsilon\right)OPT \cdot W - \frac{(w(\mathcal{S}'_{used}) + w(\mathcal{S}^{short}_{unused}))}{2}OPT - \frac{OPT}{2} \\
&\geq 9\epsilon OPT \cdot W - \frac{OPT}{2} \geq 8\epsilon OPT \cdot W,
\end{aligned}$$

The second last inequality follows from the fact that  $(w(\mathcal{S}'_{used}) + w(\mathcal{S}^{short}_{unused})) \leq W$ . In the last inequality, we used the assumption  $W \geq 1/\epsilon$ .

Since  $a(\mathcal{S}^{tall}_{unused}) \leq (\frac{3OPT}{2} + 11\epsilon OPT)w(\mathcal{S}^{tall}_{unused})$ , it follows that, for  $\epsilon \leq 1/22$ ,

$$w(\mathcal{S}^{tall}_{unused}) \geq \frac{8\epsilon \cdot W}{\frac{3}{2} + 11\epsilon} \geq 4\epsilon \cdot W.$$

Next, consider the set of boxes spanned by  $\mathcal{S}^{tall}_{unused}$ . All these boxes contain a free rectangular region of height  $\frac{1}{2}OPT$  induced by the bottom part of  $\mathcal{S}^{tall}_{unused}$ : let us call these regions  $F_1, \dots, F_k$ . Since the number of these regions is at most  $\frac{2}{\epsilon}$  (i.e. the total number of boxes) and their total width is at least  $4\epsilon \cdot W$ , by an averaging argument there exists one such  $F_i$  of width at least  $2\epsilon^2 W$ . ◀

### 3.3 Rounding

In this section, we show how to round the packing from Lemma 7 by means of Lemma 4, hence concluding the proof of Theorem 5.

## 44:12 A Tight $(3/2 + \varepsilon)$ Approximation for Skewed Strip Packing

**Proof of Theorem 5.** We start with the packing of  $\mathcal{T} \cup \mathcal{S}_{hslice} \cup \mathcal{V}_{vslice}$  obtained from Lemma 7. Recall that  $\mathcal{S}_{hslice}$  is packed into  $1/\varepsilon + 1$  boxes  $\mathcal{B}_S$  and  $\mathcal{V}_{vslice}$  into  $2/\varepsilon$  boxes  $\mathcal{B}_V$ . The total height of this packing is  $(\frac{3}{2} + 15\varepsilon)OPT$ , and this packing leaves a free region  $F$  of size  $2\varepsilon^2W \times \frac{1}{2}OPT$ . Provided that  $\delta$  is small enough, we can apply Lemma 4 to  $(\mathcal{S}_{hslice}, \mathcal{B}_S)$  and obtain a packing of  $\mathcal{S}$  into a set of containers fully contained in  $\mathcal{B}_S$ , plus two containers of size at most  $W \times \frac{\varepsilon^2}{2}OPT$  each. We place the latter two containers on top of the packing, hence increasing the total height by  $\varepsilon^2OPT$ . By applying the same Lemma to  $(\mathcal{V}_{vslice}, \mathcal{B}_V)$ , we obtain a packing of  $\mathcal{V}$  into a set of containers fully contained in  $\mathcal{B}_V$  plus two containers of size at most  $\frac{\varepsilon^2}{2}W \times \frac{1}{2}OPT$  each. The latter two containers can be placed inside  $F$  without further increasing the height of the packing, still leaving a free region of size  $\varepsilon^2W \times \frac{1}{2}OPT$ .

By construction, the number of used containers is  $O_\varepsilon(1)$  and their sizes belong to a set that can be computed in polynomial time.  $\blacktriangleleft$

### 4 Algorithm

In this section, we describe an algorithm based on Theorem 5 to compute our final solution.

Consider the set of containers guaranteed by Theorem 5. In polynomial time we can guess such containers by trying all possibilities. By brute force we can also compute (in polynomial time) a packing of these containers plus the free box in the strip of total height at most  $(\frac{3}{2} + O(\varepsilon))OPT$ . We guess which ones among the vertical containers contain  $\mathcal{T}$ , and pack the whole set  $\mathcal{T}$  there greedily in non-increasing order of height.

We next apply Lemma 3 with parameter  $\varepsilon' = \varepsilon^3$  to the remaining containers and to the remaining rectangles  $\mathcal{V} \cup \mathcal{S}$ . This way we can pack a set  $\mathcal{R}' \subseteq \mathcal{V} \cup \mathcal{S}$  of area at least  $a(\mathcal{V} \cup \mathcal{S})(1 - \varepsilon^3)$ . It remains to pack  $\mathcal{R}'' := (\mathcal{V} \cup \mathcal{S}) \setminus \mathcal{R}'$ ,  $a(\mathcal{R}'') \leq \varepsilon^3 a(\mathcal{V} \cup \mathcal{S}) \leq \varepsilon^3(\frac{3}{2} + O(\varepsilon))OPT \cdot W \leq 2\varepsilon^3OPT \cdot W$ . We partition  $\mathcal{R}''$  into 3 subsets and pack them as follows:


1. The rectangles  $\mathcal{V}'' \subseteq \mathcal{R}''$  of height at least  $2\varepsilon OPT$  (notice that they have height at most  $OPT/2$ ). By an area argument their total width is at most  $\frac{2\varepsilon^3OPT \cdot W}{2\varepsilon OPT} = \varepsilon^2W$ . Hence they fit in a vertical container of size  $\varepsilon^2W \times \frac{1}{2}OPT$  that can be placed in the area occupied by the free box (without increasing the height of the packing).
2. The rectangles  $\mathcal{H}'' \subseteq \mathcal{R}''$  of width at least  $\varepsilon^2W$ . By a similar area argument their total height is at most  $\frac{2\varepsilon^3OPT \cdot W}{\varepsilon^2W} = 2\varepsilon OPT$ . Hence they can be placed into a horizontal container of size  $W \times 2\varepsilon OPT$  that can be placed on top of the current packing.
3. The remaining rectangles  $\mathcal{S}'' \subseteq \mathcal{R}''$  with height at most  $2\varepsilon OPT$  and width at most  $\varepsilon^2W$ . By Lemma 2 with parameter  $\varepsilon' = \sqrt{\varepsilon}$  and for small enough  $\varepsilon$ , we can pack  $\mathcal{S}''$  by means of NFDH into an area container of size  $\varepsilon\sqrt{\varepsilon}W \times 2\sqrt{\varepsilon}OPT$  to be placed on top of the current packing.

We now have all the ingredients to prove our main theorem.

**Proof of Theorem 1.** Consider the above algorithm. Clearly it runs in polynomial time for any fixed parameter  $\varepsilon > 0$ . Furthermore, for  $\delta$  small enough, it generates a feasible packing of all rectangles of total height at most  $(\frac{3}{2} + O(\sqrt{\varepsilon}))OPT$ . Considering the initial rounding of  $OPT$  by a factor  $(1 + \varepsilon)$ , this gives a  $\frac{3}{2} + O(\sqrt{\varepsilon})$  approximation. The claim then follows by choosing  $\varepsilon$  appropriately.  $\blacktriangleleft$

### 5 Hardness of Approximation

In this section, we prove that the lower bound of  $\frac{3}{2}$  on the approximability of Strip Packing still holds in the case of  $\delta$ -skewed instances.



■ **Figure 3** Construction from Lemma 8. Light gray rectangles represent dummy rectangles and dark gray rectangles represent partition rectangles.

► **Lemma 8.** *Given  $\delta > 0$  and  $\varepsilon > 0$ , there is no polynomial-time  $(\frac{3}{2} - \varepsilon)$ -approximation for  $\delta$ -skewed Strip Packing unless  $P=NP$ .*

**Proof.** We will prove this result via a reduction from the NP-complete *Partition* problem. Recall that in *Partition* we are given a set of integers  $\mathcal{I} = \{x_1, \dots, x_n\}$  whose sum is  $p$ . Our goal is to determine whether  $\mathcal{I}$  can be partitioned into two sets  $\mathcal{I}_1$  and  $\mathcal{I}_2$  such that  $\sum_{x_i \in \mathcal{I}_1} x_i = \frac{p}{2}$ . We define our Strip Packing instance as follows: The width of the strip will be  $W = (1 + \delta/4)M$  where  $M = \frac{2p}{\delta}$ . Also, we will have  $n + \frac{4}{\delta}$  rectangles in the instance, from which  $\frac{4}{\delta}$  will have height 1 and width  $\frac{\delta}{2}M$  (*dummy rectangles*) and the remaining  $n$  rectangles will have, for each  $i = 1, \dots, n$ , height 1 and width  $x_i$  (*partition rectangles*). Notice that the instance is indeed  $\delta$ -skewed as the width of the rectangles is either  $\frac{\delta}{2}M \leq \frac{\delta}{2}W$  or at most  $p = \frac{\delta}{2}M \leq \frac{\delta}{2}W$ . Notice also that  $OPT \geq 2$  since the area of the rectangles is  $2W$ .

We will now prove that the *Partition* instance is a YES instance if and only if  $OPT = 2$ . Since all the heights in the instance are 1, as a consequence a NO instance has height at least 3, hence concluding the proof of the claim. Notice that if the *Partition* instance is a YES instance then we can pack one next to the other  $\frac{2}{\delta}$  dummy rectangles plus one side of the partition since their total width would be  $M + \frac{p}{2} = (1 + \frac{\delta}{4})M$ . We then analogously pack the rest of the rectangles on top, obtaining a packing of height 2 which is optimal as the total area of the rectangles is  $2W$  (see Figure 3). On the other hand, if the optimal height of the Strip Packing instance is 2, the subregion  $[0, W] \times [0, 2]$  in the strip must be fully occupied by rectangles. This actually implies that the horizontal segment  $[0, W] \times \{1\}$  does not intersect the interior of any rectangle in the packing: indeed otherwise the space below that rectangle could not be occupied by any other rectangle (as heights are all 1). This divides the solution into two rows of height 1 and width  $W$  which are completely filled with rectangles. The only way to divide dummy rectangles into the two rows is to have exactly  $\frac{2}{\delta}$  in each row (as the largest total width below  $W$  that they can sum up to is  $M$  and their total width is  $2M$ ), hence the remaining partition rectangles in each row have total width exactly  $\frac{p}{2}$ , forming then a feasible solution to the *Partition* instance. ◀

## References

- 1 Anna Adamaszek, Tomasz Kociumaka, Marcin Pilipczuk, and Michal Pilipczuk. Hardness of approximation for strip packing. *ACM Transactions on Computation Theory*, 9(3):14:1–14:7, 2017. doi:10.1145/3092026.

- 2 Anna Adamaszek and Andreas Wiese. Approximation schemes for maximum weight independent set of rectangles. In *54th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2013, 26-29 October, 2013, Berkeley, CA, USA*, pages 400–409. IEEE Computer Society, 2013. doi:10.1109/FOCS.2013.50.
- 3 Anna Adamaszek and Andreas Wiese. A quasi-ptas for the two-dimensional geometric knapsack problem. In Piotr Indyk, editor, *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2015, San Diego, CA, USA, January 4-6, 2015*, pages 1491–1505. SIAM, 2015. doi:10.1137/1.9781611973730.98.
- 4 Brenda S. Baker, Edward G. Coffman Jr., and Ronald L. Rivest. Orthogonal packings in two dimensions. *SIAM Journal on Computing*, 9(4):846–855, 1980. doi:10.1137/0209064.
- 5 Nikhil Bansal, Alberto Caprara, Klaus Jansen, Lars Prädél, and Maxim Sviridenko. A structural lemma in 2-dimensional packing, and its implications on approximability. In Yingfei Dong, Ding-Zhu Du, and Oscar H. Ibarra, editors, *Algorithms and Computation, 20th International Symposium, ISAAC 2009, Honolulu, Hawaii, USA, December 16-18, 2009. Proceedings*, volume 5878 of *Lecture Notes in Computer Science*, pages 77–86. Springer, 2009. doi:10.1007/978-3-642-10631-6\_10.
- 6 Nikhil Bansal, José R. Correa, Claire Kenyon, and Maxim Sviridenko. Bin packing in multiple dimensions: Inapproximability results and approximation schemes. *Mathematics of Operations Research*, 31(1):31–49, 2006. doi:10.1287/moor.1050.0168.
- 7 Nikhil Bansal and Arindam Khan. Improved approximation algorithm for two-dimensional bin packing. In Chandra Chekuri, editor, *Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2014, Portland, Oregon, USA, January 5-7, 2014*, pages 13–25. SIAM, 2014. doi:10.1137/1.9781611973402.2.
- 8 Parinya Chalermsook and Julia Chuzhoy. Maximum independent set of rectangles. In Claire Mathieu, editor, *Proceedings of the Twentieth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2009, New York, NY, USA, January 4-6, 2009*, pages 892–901. SIAM, 2009. URL: <http://dl.acm.org/citation.cfm?id=1496770.1496867>.
- 9 Timothy M. Chan and Sarel Har-Peled. Approximation algorithms for maximum independent set of pseudo-disks. *Discrete & Computational Geometry*, 48(2):373–392, 2012. doi:10.1007/s00454-012-9417-5.
- 10 Henrik I. Christensen, Arindam Khan, Sebastian Pokutta, and Prasad Tetali. Approximation and online algorithms for multidimensional bin packing: A survey. *Computer Science Review*, 24:63–79, 2017. doi:10.1016/j.cosrev.2016.12.001.
- 11 Edward Grady Coffman and John L. Bruno. *Computer and job-shop scheduling theory*. Wiley New York, 1976.
- 12 Edward G. Coffman Jr., János Csirik, Gábor Galambos, Silvano Martello, and Daniele Vigo. *Bin Packing Approximation Algorithms: Survey and Classification*, pages 455–531. Springer New York, New York, NY, 2013. doi:10.1007/978-1-4419-7997-1\_35.
- 13 Leah Epstein and Rob van Stee. This side up! *ACM Transactions on Algorithms*, 2(2):228–243, 2006. doi:10.1145/1150334.1150339.
- 14 Waldo Gálvez, Fabrizio Grandoni, Sandy Heydrich, Salvatore Ingala, Arindam Khan, and Andreas Wiese. Approximating geometric knapsack via L-packings. In Chris Umans, editor, *58th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2017, Berkeley, CA, USA, October 15-17, 2017*, pages 260–271. IEEE Computer Society, 2017. doi:10.1109/FOCS.2017.32.
- 15 Waldo Gálvez, Fabrizio Grandoni, Sandy Heydrich, Salvatore Ingala, Arindam Khan, and Andreas Wiese. Approximating geometric knapsack via L-packings. *CoRR*, abs/1711.07710, 2017. arXiv:1711.07710.
- 16 Waldo Gálvez, Fabrizio Grandoni, Salvatore Ingala, and Arindam Khan. Improved pseudo-polynomial-time approximation for strip packing. In Akash Lal, S. Akshay, Saket Saurabh, and Sandeep Sen, editors, *36th IARCS Annual Conference on Foundations of Software Technology and Theoretical Computer Science, FSTTCS 2016, December 13-15, 2016, Chennai, India*, volume 65 of *LIPICs*, pages 9:1–9:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2016. doi:10.4230/LIPICs.FSTTCS.2016.9.

- 17 M. R. Garey and David S. Johnson. "strong" np-completeness results: Motivation, examples, and implications. *Journal of the ACM*, 25(3):499–508, 1978. doi:10.1145/322077.322090.
- 18 Rolf Harren, Klaus Jansen, Lars Prädél, and Rob van Stee. A  $(5/3 + \epsilon)$ -approximation for strip packing. *Computational Geometry*, 47(2):248–267, 2014. doi:10.1016/j.comgeo.2013.08.008.
- 19 Rolf Harren and Rob van Stee. Improved absolute approximation ratios for two-dimensional packing problems. In Irit Dinur, Klaus Jansen, Joseph Naor, and José D. P. Rolim, editors, *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, 12th International Workshop, APPROX 2009, and 13th International Workshop, RANDOM 2009, Berkeley, CA, USA, August 21-23, 2009. Proceedings*, volume 5687 of *Lecture Notes in Computer Science*, pages 177–189. Springer, 2009. doi:10.1007/978-3-642-03685-9\_14.
- 20 Sören Henning, Klaus Jansen, Malin Rau, and Lars Schmarje. Complexity and inapproximability results for parallel task scheduling and strip packing. *Theory of Computing Systems*, 64(1):120–140, 2020. doi:10.1007/s00224-019-09910-6.
- 21 Klaus Jansen and Lars Prädél. A new asymptotic approximation algorithm for 3-dimensional strip packing. In Viliam Geffert, Bart Preneel, Branislav Rován, Julius Stuller, and A Min Tjoa, editors, *SOFSEM 2014: Theory and Practice of Computer Science - 40th International Conference on Current Trends in Theory and Practice of Computer Science, Nový Smokovec, Slovakia, January 26-29, 2014, Proceedings*, volume 8327 of *Lecture Notes in Computer Science*, pages 327–338. Springer, 2014. doi:10.1007/978-3-319-04298-5\_29.
- 22 Klaus Jansen and Malin Rau. Closing the gap for pseudo-polynomial strip packing. In Michael A. Bender, Ola Svensson, and Grzegorz Herman, editors, *27th Annual European Symposium on Algorithms, ESA 2019, September 9-11, 2019, Munich/Garching, Germany*, volume 144 of *LIPICs*, pages 62:1–62:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019. doi:10.4230/LIPICs.ESA.2019.62.
- 23 Klaus Jansen and Malin Rau. Improved approximation for two dimensional strip packing with polynomial bounded width. *Theoretical Computer Science*, 789:34–49, 2019. doi:10.1016/j.tcs.2019.04.002.
- 24 Klaus Jansen and Roberto Solis-Oba. New approximability results for 2-dimensional packing problems. In Ludek Kucera and Antonín Kucera, editors, *Mathematical Foundations of Computer Science 2007, 32nd International Symposium, MFCS 2007, Český Krumlov, Czech Republic, August 26-31, 2007, Proceedings*, volume 4708 of *Lecture Notes in Computer Science*, pages 103–114. Springer, 2007. doi:10.1007/978-3-540-74456-6\_11.
- 25 Klaus Jansen and Rob van Stee. On strip packing with rotations. In Harold N. Gabow and Ronald Fagin, editors, *Proceedings of the 37th Annual ACM Symposium on Theory of Computing, Baltimore, MD, USA, May 22-24, 2005*, pages 755–761. ACM, 2005. doi:10.1145/1060590.1060702.
- 26 Klaus Jansen and Guochuan Zhang. Maximizing the total profit of rectangles packed into a rectangle. *Algorithmica*, 47(3):323–342, 2007. doi:10.1007/s00453-006-0194-5.
- 27 Edward G. Coffman Jr., M. R. Garey, David S. Johnson, and Robert Endre Tarjan. Performance bounds for level-oriented two-dimensional packing algorithms. *SIAM Journal on Computing*, 9(4):808–826, 1980. doi:10.1137/0209062.
- 28 Mohammad M. Karbasioun, Gennady Shaikhet, Evangelos Kranakis, and Ioannis Lambadaris. Power strip packing of malleable demands in smart grid. In *Proceedings of IEEE International Conference on Communications, ICC 2013, Budapest, Hungary, June 9-13, 2013*, pages 4261–4265. IEEE, 2013. doi:10.1109/ICC.2013.6655233.
- 29 Claire Kenyon and Eric Rémila. A near-optimal solution to a two-dimensional cutting stock problem. *Mathematics of Operations Research*, 25(4):645–656, 2000. doi:10.1287/moor.25.4.645.12118.
- 30 Arindam Khan. *Approximation algorithms for multidimensional bin packing*. PhD thesis, Georgia Institute of Technology, Atlanta, GA, USA, 2016. URL: <http://hdl.handle.net/1853/54371>.

- 31 Joseph Y.-T. Leung, Tommy W. Tam, C. S. Wong, Gilbert H. Young, and Francis Y. L. Chin. Packing squares into a square. *Journal on Parallel and Distributed Computing*, 10(3):271–275, 1990. doi:10.1016/0743-7315(90)90019-L.
- 32 Flávio Keidi Miyazawa and Yoshiko Wakabayashi. Packing problems with orthogonal rotations. In Martin Farach-Colton, editor, *LATIN 2004: Theoretical Informatics, 6th Latin American Symposium, Buenos Aires, Argentina, April 5-8, 2004, Proceedings*, volume 2976 of *Lecture Notes in Computer Science*, pages 359–368. Springer, 2004. doi:10.1007/978-3-540-24698-5\_40.
- 33 Giorgi Nadiradze and Andreas Wiese. On approximating strip packing with a better ratio than  $3/2$ . In Robert Krauthgamer, editor, *Proceedings of the Twenty-Seventh Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2016, Arlington, VA, USA, January 10-12, 2016*, pages 1491–1510. SIAM, 2016. doi:10.1137/1.9781611974331.ch102.
- 34 Ingo Schiermeyer. Reverse-fit: A 2-optimal algorithm for packing rectangles. In Jan van Leeuwen, editor, *Algorithms - ESA '94, Second Annual European Symposium, Utrecht, The Netherlands, September 26-28, 1994, Proceedings*, volume 855 of *Lecture Notes in Computer Science*, pages 290–299. Springer, 1994. doi:10.1007/BFb0049416.
- 35 Daniel Dominic Sleator. A 2.5 times optimal algorithm for packing in two dimensions. *Information Processing Letters*, 10(1):37–40, 1980. doi:10.1016/0020-0190(80)90121-0.
- 36 A. Steinberg. A strip-packing algorithm with absolute performance bound 2. *SIAM Journal on Computing*, 26(2):401–409, 1997. doi:10.1137/S0097539793255801.
- 37 Shaojie Tang, Qiuyuan Huang, Xiang-Yang Li, and Dapeng Wu. Smoothing the energy consumption: Peak demand reduction in smart grid. In *Proceedings of the IEEE INFOCOM 2013, Turin, Italy, April 14-19, 2013*, pages 1133–1141. IEEE, 2013. doi:10.1109/INFOCOM.2013.6566904.

## A Omitted Proofs

**Proof of Lemma 4.** We prove the claim for  $\mathcal{H}$ , the case of  $\mathcal{V}$  being symmetric. For a proper parameter  $\alpha > 0$  to be fixed later, we define a rectangle  $R$  (and its horizontal slices) to be *narrow* if  $w(R) \leq \alpha^2 W$  and *wide* otherwise. We temporarily remove narrow rectangles, and pack the wide ones.

The first step in our construction is to round up the widths of the wide slices, while discarding a small-area subset of them. Let  $\beta > 0$  be a parameter to be fixed later. Let us sort the wide slices  $\mathcal{H}_{\text{hslice}}^{\text{wide}}$  in non-increasing order of width, and let us partition the obtained sequence into subsequences  $\mathcal{H}_1, \dots, \mathcal{H}_{1/\beta}$  of total height  $\beta h(\mathcal{H}_{\text{hslice}}^{\text{wide}})$  each (excluding possibly the last group that can have smaller height). For a group  $i$ , we define  $w_i^{\text{min}}$  as the minimum width in  $\mathcal{H}_i$ . For each  $i = 1, \dots, 1/\beta - 1$ , we define an injection between  $\mathcal{H}_{i+1}$  and  $\mathcal{H}_i$ . Next, we delete slices  $\mathcal{H}_1$ . Let  $\mathcal{H}_{\text{wdisc1}}$  denote the rectangles of which we removed at least one slice. Notice that all rectangles having some slice in  $\mathcal{H}_1$  have all their slices in  $\mathcal{H}_1$  excluding possibly one rectangle (which has part of its slices in  $\mathcal{H}_1$ ). Observe that  $h(\mathcal{H}_{\text{hslice}}^{\text{wide}}) \leq \frac{H}{\alpha}$  since otherwise  $a(\mathcal{H}_{\text{hslice}}^{\text{wide}}) \geq \alpha W \cdot h(\mathcal{H})$  would be too large to fit into the region of size  $W \times H$ . Hence  $h(\mathcal{H}_1) = \beta h(\mathcal{H}) \leq \frac{\beta}{\alpha} H$ . It follows that  $h(\mathcal{H}_{\text{wdisc1}}) \leq (\frac{\beta}{\alpha} + \delta)H$ . For any fixed  $\alpha$ , this quantity is at most  $\frac{(\varepsilon')^2}{2}H$  if  $\beta \leq \alpha(\varepsilon')^2/4$  and  $\delta \leq (\varepsilon')^2/4$ .

For  $i = 1, \dots, 1/\beta - 1$ , we temporarily increase the width of each  $H \in \mathcal{H}_{i+1}$  to  $w_i^{\text{min}}$ , hence getting an enlarged slice  $\bar{H}$ . Then, we move each such  $\bar{H}$  into the region that was occupied by the slice  $H' \in \mathcal{H}_i$  associated with  $H$  according to the above injection. Notice that this is possible since we removed  $\mathcal{H}_1$  and since  $w(\bar{H}) = w_i^{\text{min}} \leq w(H')$ . Let  $\bar{\mathcal{H}}$  be the final set of enlarged slices. Observe that the number of possible distinct widths in  $\bar{\mathcal{H}}$  is  $1/\beta - 1$ .



Let us focus on a specific box  $B \in \mathcal{B}$  of size  $w(B) \times h(B)$ , and let  $\mathcal{H}'_{h\text{slice}}$  be the slices contained in  $B$ . Next, we partition  $B$  into unit height stripes. We shift slices in each stripe as left as possible, and permute them so that slices in  $\overline{\mathcal{H}}$  appear to the left of each stripe. We call a *configuration*  $C$  of a stripe the sequence of (enlarged) widths  $(w_1, \dots, w_q)$  of its slices in  $\overline{\mathcal{H}}$  from left to right. Notice that there are  $1/\beta - 1$  possible enlarged widths, and each stripe can contain at most  $1/\alpha^2$  wide slices. Hence the number of possible configurations is at most  $n_{conf} = \sum_{i=0}^{1/\alpha^2} (1/\beta - 1)^i \leq 2(1/\beta - 1)^{1/\alpha^2} \leq (1/\beta)^{1/\alpha^2}$ .

We reorder the stripes in  $\mathcal{H}'_{h\text{slice}}$  vertically so that equal configurations appear consecutively from top to bottom, and stripes without narrow rectangles appear at the bottom. Suppose that the number of stripes in  $B$  with a given configuration  $C = (w_1, \dots, w_q)$  is  $h(C)$ , and  $A(C)$  is the corresponding region. We initially cover  $A(C)$  by creating  $q$  consecutive horizontal containers of height  $h(C)$  and width  $w_1, \dots, w_q$  respectively. These containers altogether cover all the wide slices in  $B$ . The width of each container belongs to a set that can be computed in polynomial time (it is the width of some input rectangle). In order to enforce the same property for their heights, we round down the height of each such container to the largest multiple  $h'(C)$  of  $\frac{\delta}{\gamma}H$  not larger than  $h(C)$ , for some parameter  $\gamma > 0$  to be fixed later. The number of these containers is  $n_{wcont} \leq Kn_{conf}$ .

We next use the obtained horizontal containers to place most of the wide rectangles. We consider the containers in non-increasing order of width and the slices of wide rectangles in the same order, breaking ties so that slices of the same rectangle appear consecutively. We also create a dummy final container of sufficient width and of height large enough to accommodate the total height of the wide slices minus the total height of the containers. Now, we place back the slices into the containers following the previous order. Notice that all slices will fit. We discard each wide rectangle whose slices are contained in 2 containers (3 is not possible) and all the wide rectangles whose slices are contained in the dummy final container. Let  $\mathcal{H}_{wdisc2}$  be the set of discarded rectangles. Their total height is

$$h(\mathcal{H}_{wdisc2}) \leq n_{wcont}\delta H + n_{wcont}\frac{\delta}{\gamma}H.$$

The above quantity is at most  $\frac{(\varepsilon')^2}{2}H$  for any choice of  $\varepsilon'$ ,  $\alpha$ ,  $\beta$ , and  $\gamma$ , provided that  $\delta \leq \frac{(\varepsilon')^2\gamma}{4n_{wcont}} \leq \frac{(\varepsilon')^2\gamma}{4K(1/\beta)^{1/\alpha^2}}$ .

So we packed all the wide rectangles into horizontal containers except for the set  $\mathcal{H}_{wdisc} = \mathcal{H}_{wdisc1} \cup \mathcal{H}_{wdisc2}$ . The latter set has, by the above discussion, height at most  $(\varepsilon')^2H$ , hence we can pack it into a container of size  $\max_{R \in \mathcal{H}_{wide}} \{w(R)\} \times (\varepsilon')^2H$ .

It remains to pack the narrow rectangles. Consider again a given box  $B$ . For each configuration  $C$ , there is some free region  $F(C)$  to the right of the containers built for  $C$  whose height is  $h'(C)$  (in particular, a multiple of  $\frac{\delta}{\gamma}H$ ) and of some width  $w(F(C))$ . We build an area container of the same height and with width equal to the largest multiple  $w'(F(C))$  of  $\frac{\alpha}{\gamma}W$  not larger than  $w(F(C))$ . We apply a similar construction to the free rectangular region  $F$  in  $B$  below all the previous containers, if any: in particular we create an area container whose width is the largest multiple of  $\frac{\alpha}{\gamma}W$  not larger than  $w(F) = w$  and whose height is the largest multiple of  $\frac{\delta}{\gamma}H$  not larger than  $h(F)$ . The total number of constructed area containers is  $n_{ncont} \leq K \cdot n_{conf}$ .

Next, we start packing the narrow rectangles in non-increasing order of height in the area containers using NFDH. Observe that these rectangles satisfy the claim of Lemma 2 with parameter  $\gamma$ . If all narrow rectangles are packed this way, we are done. Otherwise, let  $\mathcal{H}_{ncont}$  and  $\mathcal{H}_{ndisc}$  be the subset of narrow rectangles that are packed and not packed in the area containers, resp. By Lemma 2,  $a(\mathcal{H}_{ncont}) \geq (1 - 2\gamma)A_{acont}$ , where  $A_{acont}$  is the

#### 44:18 A Tight $(3/2 + \varepsilon)$ Approximation for Skewed Strip Packing

total area of the area containers. Let  $a_{free}$  be the total area in the boxes not occupied by horizontal containers. Clearly  $a_{free} \geq a(\mathcal{H}^{narrow})$  since all narrow slices did fit in a region of area not smaller than  $a_{free}$ . Due to the rounding involved in the construction, in each box there is some area which is not used by area containers nor by horizontal ones. The latter area is at most  $W \cdot \frac{\delta}{\gamma}H + H \cdot \frac{\alpha}{\gamma}W$  per container, hence at most  $\Delta \leq K \cdot (\frac{\delta+\alpha}{\gamma})WH$  in total. We can conclude that

$$a(\mathcal{H}_{ncont}) \geq (1-2\gamma)A_{acont} = (1-2\gamma)(a_{free} - \Delta) \geq (1-2\gamma)(a(\mathcal{H}^{narrow}) - K \cdot (\frac{\delta+\alpha}{\gamma})WH).$$

Thus

$$a(\mathcal{H}_{ndisc}) \leq 2\gamma \cdot a(\mathcal{H}^{narrow}) + K \cdot (\frac{\delta+\alpha}{\gamma})WH \leq (2\gamma + K \cdot (\frac{\delta+\alpha}{\gamma}))WH.$$

If we choose  $\gamma \leq (\varepsilon')^2/6$ ,  $\delta \leq \frac{(\varepsilon')^2\gamma}{6K}$  and  $\alpha \leq \frac{(\varepsilon')^2\gamma}{6K}$ , then the latter quantity is at most  $\frac{(\varepsilon')^2}{2}WH$ . Next, we create a new area container  $C_{darea}$  of size  $\varepsilon'W \times \varepsilon'H$ , and use NFDH to pack  $\mathcal{H}_{ndisc}$  in it. It is not difficult to verify that, for such values of  $\delta$  and  $\alpha$ , rectangles in  $\mathcal{H}_{ndisc}$  satisfy the conditions of Lemma 2 with parameter  $\varepsilon'$ . Thus we have

$$a(\mathcal{H}_{ndisc}) \leq \frac{1}{2}(\varepsilon')^2WH = \frac{1}{2}a(C_{darea}) \leq (1-2\varepsilon')a(C_{darea}).$$

Thus all rectangles in  $\mathcal{H}_{ndisc}$  are packed into  $C_{darea}$ .

It is possible to choose constant parameters  $\alpha$ ,  $\beta$  and  $\gamma$  such that the above conditions are all satisfied (for  $\delta$  small enough) and the total number of containers is  $O_{\varepsilon'}(1)$ . More precisely, this is true if  $\gamma = (\varepsilon')^2/6$ ,  $\alpha = (\varepsilon')^4/(36 \cdot K)$ ,  $\beta = (\varepsilon')^6/(144 \cdot K)$  and  $\delta = \frac{(\varepsilon')^2\gamma}{4K(1/\beta)^{(1/\alpha^2)}} \in \delta \leq (\varepsilon'/K)^{(K/\varepsilon')^{O(1)}}$ , leading to at most  $(K/\varepsilon')^{(K/\varepsilon')^{O(1)}}$  containers. By the above construction, the sizes of the containers belong to a set that can be computed in polynomial time. ◀