



[The VAD Journal: The journal of mechanical assisted circulation and heart failure](#)

Original Research

Improved outcomes in the treatment of post-myocardial infarction ventricular septal defect with percutaneous TandemHeart® left ventricular mechanical circulatory support

Pranav Loyalka, Angelo Nascimbene, Sriram S. Nathan, Rajko Radovancevic, Ajay S. Raman, Manish Patel, Keshava Rajagopal, Biswajit Kar, Igor D. Gregoric*

Department of Advanced Cardiopulmonary Therapies and Transplantation, University of Texas-Houston McGovern Medical School and Memorial Hermann-Texas Medical Center, Houston, TX

*Corresponding author: igor.d.gregoric@uth.tmc.edu

Citation: Loyalka, et al. Improved outcomes in the treatment of post-myocardial infarction ventricular septal defect with percutaneous TandemHeart® left ventricular mechanical circulatory support. *The VAD Journal*. 2020; 6 [Epub ahead of print].

Editor-in-Chief: Maya Guglin, University of Kentucky

Received: May 22, 2020

Accepted: July 07, 2020

Published Online: July 17, 2020

© 2020 The Author(s). This is an open access article published under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided that the original author(s) and the publication source are credited.

Funding: Not applicable

Competing interests: none

Abstract

Background

Post-myocardial infarction (MI) ventricular septal defect (VSD) is associated with 40% - 50% of peri-procedural mortalities; however, it is amenable to catheter-based therapies. We retrospectively investigated the impact of state-of-the-art bridging percutaneous left ventricular mechanical circulatory support (MCS) using the TandemHeart® (TH) ventricular assist device (VAD) on a patient with post-MI VSD.

Results

From July 2008 to March 2014, 23 patients were referred for treatment of post-MI VSD. Initially, 18/23 patients required MCS; 12 received an intra-aortic balloon pump (IABP), while 6 received initial TH support. Seven of the IABP patients later required TH support. Catheter-based device VSD closure was performed in 18 of the patients; however, three patients required conversion to conventional open cardiac surgical repair via VSD patch closure due to failure of the catheter-based approach. Five patients with TH underwent planned open cardiac surgical repair



due to an anticipated lack of suitability for catheter-based treatment. Results revealed that delayed closure after MI correlated with improved survival. Overall, 30-day and 6-month survival rates were 83% (19/23) and 70% (16/23), respectively.

Conclusions

Further, Qp/Qs ratios of <2.4 correlated with successful percutaneous VSD repair, and this assessment should be further explored as an assessment to inform clinical judgment in patients with post-MI VSD treatment.

Keywords: Ventricular septal defect, mechanical circulatory support, percutaneous repair, heart failure

Introduction

Mechanical complications of acute myocardial infarction (MI) – ventricular septal defect (VSD), mitral regurgitation (MR), and free wall rupture (FWR) carry a grave prognosis and contribute to substantial mortality and morbidity despite current therapies. However, these complications are historically uncommon, with a reported incidence of 1 to 2% in the pre-thrombolytic agent era.^{1,2} Furthermore, in the era of early myocardial reperfusion via percutaneous coronary intervention and thrombolytic agents, the incidence of mechanical complications is even lower; the incidence of VSD with thrombolysis is as low as 0.2%.^{3,4} In addition, with the advent of reperfusion strategies, the time to ventricular rupture is significantly shorter, as documented by the GUSTO1 (Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Coronary Arteries/1 Day) and SHOCK (Should We Emergently Revascularize Occluded Coronaries for Cardiogenic Shock/16 hours) trials.^{3,5}

While FWR carries the highest mortality, due to its presentation with acute cardiac tamponade and obstructive shock, VSD and MR continue to carry high mortalities and morbidities as well. Unlike post-MI papillary muscle rupture causing MR, both surgical and catheter-based treatment options exist for VSD. Broadly, two types of surgical approaches have been used for post-MI VSD. The first approach, developed by Daggett and colleagues,⁶ involves trans-infarct left ventriculotomy, infarctectomy, and closure of the VSD, which typically uses a patch prosthesis. The second and more recent approach is credited to David and colleagues,⁷⁻⁹ and it employs trans-infarct left ventriculotomy with an oversized left-sided patch prosthesis sutured to the left-sided septum and anterior left ventricular (LV) walls to “exclude” the defect and commit the infarcted septum to the right ventricular (RV) side of the patch.

With respect to surgical treatment, Arnaoutakis and colleagues, in the largest study of its kind, reported outcomes of surgical repair of post-MI VSD from the



Society of Thoracic Surgeons database.¹⁰ In this study, 2,876 patients underwent post-MI VSD repair, with a perioperative mortality of 42.9%.¹⁰ Patients undergoing treatment within 7 days of hospital admission had a 54.1% mortality rate, while the minority of patients who underwent delayed treatment had an 18.4% mortality rate. While these different survival outcomes may reflect improved hemodynamic and clinical conditions of the patients undergoing delayed VSD repair, the magnitude of the findings is striking, and suggests that, if feasible, physiological optimization both pharmacologically and via mechanical circulatory support (MCS) should be undertaken prior to VAD repair. In fact, this was proposed by Dr. T. David in his discussion of Arnaoutakis et al.¹⁰ Consistent with this hypothesis, we recently reported improved outcomes of delayed post-MI VSD surgical repair following a period of TandemHeart® (TH, LiveNova, London UK) percutaneous ventricular assist device (pVAD) support. Using this approach, we observed a 25% mortality rate at six months.¹¹

The current guidelines recommend emergent surgical repair of the VSD with concurrent myocardial revascularization via coronary artery bypass grafting, if indicated, irrespective of hemodynamic status.¹² However, the high mortality and morbidity, as well as an increasing experience with catheter-based device closure of congenital VSDs,^{13,14} have demanded the development of a comprehensive reassessment. Our proposed multi-pronged strategy for post-MI VSD management is as follows: (1) hemodynamic optimization using pharmacotherapy and MCS in the form of either intra-aortic balloon pump (IABP) counterpulsation or when needed, TH pVAD support, and (2) delayed catheter-based device and/or open cardiac surgical closure. To investigate the effectiveness of this strategy, outcomes of patients who underwent this approach of bridging MCS to delayed definitive treatment were retrospectively reviewed and analyzed.

Methods

Between July 2008 and March 2014, 23 patients were referred to our Center in a large metropolitan medical center for the management of post-MI VSDs. None of the patients had undergone emergent catheter-based or surgical VSD closure. All referred patients with post-MI VSDs were included in this study, without exclusion. The adequacy of end-organ perfusion and the presence or absence of cardiogenic shock was assessed as our Center has previously outlined in the surgical cases series by Gregoric et al.¹¹ Pharmacological optimization, comprised of LV afterload reduction and inotropic support, was initiated in all 23 patients. Patients who did not stabilize and had less severe, low-systemic cardiac output syndrome/shock underwent initial percutaneous IABP placement. In contrast, those with severe shock either initially or despite IABP support underwent percutaneous trans-atrial septal left atrial (LA) and common femoral arterial cannulation for TH pVAD support described in Gregoric et al.¹¹ This algorithm is outlined in Figure 1.

The decision to implement TH versus IAPB was exclusively based on clinical judgment. The primary step was to achieve satisfactory cardiovascular status, as assessed by systemic arterial blood pressure, Qs, central venous oximetry, and clinical and laboratory indices of end-organ perfusion. The decision to wean from




Figure 1. Strategy utilized for timing and therapeutic option in the management of post-myocardial infarction ventricular septal defects. MI (myocardial infarction); VSD (ventricular septal defect); IAPB (intra-aortic balloon pump); TH (TandemHeart™); Qp/Qs (pulmonary/systemic blood flow).

TH was attempted in the presence of adequate hemodynamics and an improved end-organ function at a pVAD flow rate of 2 L/day for two days. While on pVAD, the flow rate was adjusted to maintain mixed venous oxygen saturation >70 and mean arterial pressure >60 mmHg.

Qualitative and quantitative assessment of VSD physiology was accomplished using echocardiography and diagnostic right-sided cardiac catheterization with oximetry. Each of these modalities was used to determine pulmonary and systemic blood flow (Qp and Qs, respectively), thereby quantifying intracardiac shunt capability in all patients.


Patients who underwent successful percutaneous closure were implanted with the Amplatzer® post-infarction VSD occluder device, the Amplatzer® muscular VSD closure device, or the Atrial Septal Occluder (ASO) device (all St. Jude Medical, Inc., St. Paul, MN, USA). The implantation strategy included femoral arterial and right internal jugular venous access, with a 10Fr catheter from the systemic



venous side and a 7Fr catheter from the systemic arterial side. A Swan-Ganz catheter was navigated to the pulmonary artery to re-evaluate the degree of intracardiac shunting and was subsequently used to support a 300 cm 0.035" Storz wire (Cordis, Fremont, CA, USA) to cross the VSD from the RV. Retrograde access from the LV was obtained with a Judkins Right 7 Fr catheter (Cordis, Fremont CA, USA), and subsequently, the wire was exchanged for an endovascular snare (EN Snare Endovascular Snare System, Merit Medical, South Jordan, UT, USA). The Storz wire was then captured from the arterial access and served as a rail to deliver the Amplatzer® TorqVue 10Fr Sheath system (St. Jude Medical, St. Paul, MN, USA) in a "bodyfloss" fashion.

The Amplatzer® delivery sheath was railed across the tricuspid annulus and over the VSD in the LV. The Storz wire was externalized, the VSD occluder was loaded into the delivery system, and then it was deployed from the arterial to the venous side. Once properly positioned, intracavitary echocardiography was performed to confirm the occluder was stable, and there was no significant residual VSD.¹⁵ Subsequently, the device was released, and the delivery sheath was withdrawn (Figure 2).

Figure 2. Intracavitary echocardiogram and fluoroscopic appearance of ventricular septal defect (VSD) device across ventricular septum after deployment. Left panel: Fluoroscopic image of the TorqVue delivery system and VSD occluder successfully deployed (arrow). Note transeptal cannula for TandemHeart® support in place (*). Right panel: Intracavitary echocardiogram of VSD occluder device (arrow) in place and released.



Patients with VSDs thought *a priori* not to be amenable to catheter-based device closure, or those who underwent a failed attempt at catheter-based device closure, underwent surgical repair as outlined in Gregoric et al.¹⁶ In this subset of patients, a median sternotomy was completed, and repairs were performed using cardiopulmonary bypass (CPB) to provide total cardiac and pulmonary support with a bloodless operative field, via ascending thoracic aortic and bicaval



cannulation. The TH cannulae were maintained in place for postoperative LV MCS. Ascending thoracic aortic cross-clamping with the administration of cardioplegia (antegrade +/- retrograde) was used to render the operative field still. Trans-infarct left ventriculotomy was performed. The VSD was repaired using a woven Dacron patch sutured to a rim of healthy LV myocardium circumferentially using polypropylene suture. Depending upon intraoperative findings, either a standard or infarct exclusion patch repair was performed. After aortic cross-clamp release and initiation of inotropic and vasoactive support for the right-sided circulation, CPB was weaned and transitioned to TH pVAD support.

If the patient remained in cardiogenic shock despite TH assistance or after initial echocardiographic evaluation, defects were considered not amenable to percutaneous nor surgical closure, or in case of failed surgical closure due to size or refractory biventricular failure TH was surgically explanted and a SynCardia temporary Total Artificial Heart (TAH) (SynCardia Inc., Tucson, AZ, USA) was implanted as a bridge to transplant, as previously described.¹⁷

After a successful percutaneous or surgical VSD closure, the TH was left in place for an additional 5-7 days to facilitate myocardial recovery and ensure end-organ perfusion and then surgically explanted, as previously described.¹¹

Outcome parameters included: (1) technical success, which was defined as successful closure of the VSD without evidence of significant residual VSD or device embolization, and (2) survival at (a) 30 days, and (b) 180 days.

Eight patients included in the current manuscript were previously reported in Gregoric et al.,¹¹ as they were contemporaneous with our TH bridging and catheter-based device closure investigation.

Statistical Analysis

All statistical analyses were performed using Stata Version 12 statistical software (College Station, TX, USA). Categorical (frequency) variables were expressed as percentages or ratios. Continuous variables were expressed as means and standard deviation (SD) or medians with interquartile ranges (IQR). A comparison of Qp/Qs ratios between different types of MCS was performed with the two-sample Wilcoxon rank-sum (Mann-Whitney) test.

Results

Over the course of 5 years and 7 months, 23 patients were admitted for management of post-MI VSD. All patients in this group suffered VSD as a result of poor initial reperfusion or late presentation in the context of acute MI.

Patient baseline characteristics are reported in Table 1; briefly, this cohort includes 15 males and 8 females with a median age of 66 years (IQR 56 - 86). All patients had evidence of acute coronary syndrome a few days prior to the initial hospitalization. The date of MI and time to presentation could be identified in all patients except four due to the absence of typical signs and symptoms of angina.



Table 1. Baseline characteristics of patients in the entire group.

	N=23
Age, y	64 ± 14
Men (%)	15 (65)
Women (%)	8 (35)
EKG infarct location	
Anterior (%)	9 (39)
Inferior (%)	12 (52)
N/A (%)	2 (9)
VSD site ICE assessment	
Apical (%)	10 (42)
Basal (%)	13 (57)
Cardiogenic shock (%)	20 (88)
Median (IQR) pVAD time (days)	14 (10 – 25)
Pre-procedure pVAD time (days)	8 (5 – 10)
Post-procedure pVAD time (days)	6 (4-25)
Median (IQR) initial Qp:Qs	2.40 (1.63-3.80)

Abbreviations: VSD - ventricular septal defect, IQR - interquartile range, pVAD - percutaneous ventricular assist device, Qp - pulmonary blood flow, Qs - systemic blood flow.

Twelve patients (52%) developed an inferior infarction. Two patients (9%) had non-obstructive coronary disease as confirmed by a coronary angiogram.

Five patients (22%) were hemodynamically stabilized with pharmacological optimization (LV afterload reduction and inotropic support) alone, while 18 patients (78%) required mechanical hemodynamic support based on the primary assessment (Figure 3). Of those requiring MCS, 12 patients received IABP support, and 6 received TH support. Seven of 12 patients initially supported with IABP required a change to TH support, leaving only 5 patients on IABP.

Once hemodynamically stable, the less compromised patients (n=10; 5 on medical therapy and 5 on IABP support) underwent percutaneous VSD repair, and all survived 30 days, and 9 of 10 were alive at 180 days post-procedure.

In total, 13 patients received TH because they could not be hemodynamically stabilized either on medical therapy (n=6) or after IABP support (n=7). Of these, 8 underwent percutaneous repair initially, while 5 had a direct surgical repair. Of the 8 who underwent initial percutaneous repair, 3 failed and crossed over to surgical repair. Of the 5 TH patients who underwent percutaneous repair, 1 patient survived 30 days and was also alive at 180 days.

All 3 TH patients who underwent failed percutaneous repair and converted to surgical repair survived 180 days. Of the 5 TH patients who underwent initial surgical repair, all survived 30 days, and 3 were alive at 180 days post-procedure.



Figure 3. A total of 23 patients were included in this study. The diagram displays their interventions and outcomes. IAPB (intra-aortic balloon pump), TH (TandemHeart™), VSD (ventricular septal defect).



Based on our prior experience with surgical repair,¹¹ all patients on TH support prior to closure were continued on support after VSD closure. The median number of days on percutaneous ventricular assist device (pVAD) support was 14.0, with a median pre-procedural pVAD and post-procedural pVAD time of 6 days and 8 days, respectively.

The time-lapse between MI and defect closure ranged from 2 to 180 days, and only 4 patients underwent closure within 6 days. As a reflection of our delayed closure strategy (depicted in Figure 1), 18 patients with percutaneous closure had a procedural success rate of 78 % (14/18). A breakdown of these results includes 3 patients who were switched to successful surgical repair; 1 patient implanted with a Total Artificial Heart, as previously described,¹⁷ due to severely depressed ventricular function secondary to the extended myocardial infarction; 4 patients with percutaneous closure died within 30 days. Five patients on TH underwent directly surgical VSD without attempting percutaneous closure due to the large size and location of VSD.

The magnitude of left-to-right intracardiac shunt across the VSD correlated strongly with requirements for and types of MCS. The median Qp/Qs for this patient cohort was 2.37 (IQR 1.50-2.70). There is a significant difference in Qp/Qs ratio between the 10 patients who hemodynamically stabilized with medical therapy (n=5) or with IABP support only (n=5) and the 13 patients who required TH support to hemodynamically stabilize [median: 1.95 (IQR 1.50-2.40) vs. 2.70 (IQR 2.10 - 3.50) respectively, P=0.046].

Most patients in this study underwent percutaneous closure with the Amplatzer® post-infarction VSD occluder device.¹⁵ Three patients required Amplatzer® muscular VSD closure device and one ASO device for percutaneous closure for their septal defects.

There were no intraoperative or procedure-related deaths. The 30-day survival in all patients was 83%, with 180-day survival was 70%.

Discussion

Classical dogma regarding the pathogenesis of post-MI VSDs is that acute large trunk coronary arterial occlusion (generally the left anterior descending artery and right coronary artery, as they are responsible for interventricular septal blood supply) causes transmural septal infarction. This premise works in the setting of an isolated single coronary artery occlusive disease because robust coronary collateral circulation is commonly present in the setting of chronic multi-system coronary artery occlusive disease. After several days, large spatiotemporal gradients in the septal wall incur stress at the interfaces between infarcted and non-infarcted myocardium (due to the fact that the septum is exposed to high-stress gradients as it is situated between the LV and RV). This stress results in ventricular septal rupture and VSD formation--a dynamic perforation of the ventricular septum at the interface between the necrotic and healthy myocardium. VSD is commonly seen in the apical septum in anterior infarctions and in the basal



region in inferior infarctions.¹⁸ Although relatively rare, VSD is associated with high morbidity and mortality if managed medically or with conventional surgery.

Lundblad and colleagues describe a case series of post-MI VSD patients over a 25-year study period, and report that concomitant surgical revascularization of all stenotic vessels, beyond the occluded one, decreased mortality at 30 days; this was attributed to the improved perfusion of the ischemic border zone and the consequently reduced risk of arrhythmias.¹⁹

Another study reported improved survival for patients in whom VSD surgical repair was delayed by two weeks after infarction to allow stabilization of the defect size and sufficient maturation of the scar borders to hold sutures.²⁰ Delayed post-MI VSD closure for proper defect sizing and effective anchoring was subsequently adopted by other groups,^{21,22} and it constituted one of the foundations of our procedural strategy to surgically manage post-MI VSD.¹¹ Prior experience with patients undergoing surgical VSD repair¹¹ and prior necropsy observations¹⁸ support a 7-10 day post-MI delay period before the VSD is closed. VSDs are often observed to enlarge. This is thought to be due to the structural weakness of the infarcted septal tissue, and consequently, negatively impact the outcomes of catheter-based or open surgical closure due to dehiscence – thereby yielding residual VSD. However, the David infarct exclusion technique may be less prone to dehiscence.⁹

The Amplatzer® post-infarction VSD occluder device, when properly sized, further circumvented some of the surgical challenges. The device is a self-centering occluder that plugs the actual defect. The central part of the device fits the defect and is unaffected by the characteristics of the septal wall; its self-expanding properties allow for adjustments in the event of defect enlargement.

Currently, percutaneous closure is becoming a widespread and widely accepted method to correct post-MI VSD, and this manuscript describes our experience with percutaneous closure strategy. Our outcomes (Figure 3) are similar to the results previously reported by Landzberg and Lock,²³ Holzer et al.,²⁴ and Egbe et al.²¹ In a recent series from the Mayo Clinic, investigators observed 83% survival at 30-days post-procedure and up to 5 years of follow up.²¹

In a US registry, 30-day survival of 18 patients with percutaneous closure of post-MI VSD was 72%; however, 10 (56%) of the patients developed shock prior to defect closure, and 8 (44%) of them needed to be hemodynamically stabilized with IABP. The majority of these patients underwent the procedure at least two weeks after diagnosis.²²

Although there are still limited clinical data to support delayed VSD closure (i.e., after the acute phase), the timing of the percutaneous closure is significantly influenced by the hemodynamic status of the patient. The presence of cardiogenic shock has been repeatedly associated with a dismal survival rate. Thiele et al²⁵ have reported their experience in percutaneous post-myocardial closure in 29 patients performed within three days from presentation. Although the procedural success was 86%, the overall survival at 30 days was only 35% and was attributed to the large occurrence of left ventricular rupture, device late embolization, residual large shunt, and cardiogenic shock.



Kapur et al²⁶ used a male swine preclinical myocardial infarction model to show mechanically unloading the LV utilizing the TH reduced LV wall stress and activated signaling pathways that promoted myocardial salvage, even when coronary reperfusion was delayed. The TH provides direct left-sided cardiac unloading as opposed to venous arterial extracorporeal membrane oxygenation (VA-ECMO), which only indirectly volume-unloads the severely dysfunctional post-MI LV; thus, it was the chosen device for the post-MI VSD cases. VA-ECMO with an IABP in place reduces LV afterload, but it was theorized that the LV volume-unloading would be superior with TH pVAD support.

Our strategy of combining delayed percutaneous closure and percutaneous hemodynamic support ensured both higher short- and long-term survival rates, even in the presence of a significant fraction of patients (86%) in some degree of cardiogenic shock that required TH support, similar to previously published reports of pVAD support prior to VSD repair.^{15,16,27,28}

Due to the retrospective nature of our analysis and the relatively small patient cohort, we cannot further define specific hemodynamic and/or clinical parameters whose presence may warrant TH implementation; however, our data do support a delay in VSD closure and the use of the Qp/Qs ratio. Patients who can be hemodynamically stabilized with medical therapy or IABP support will most likely still benefit from a post-acute intervention strategy (i.e., delay VSD closure for approximately two weeks to allow tissue granulation at the VSD). A high initial Qp/Qs ratio (i.e., greater than 2.4) might be a valuable predictive tool. We included the ratio along with other clinical variables in our algorithm as a trigger to mechanical support escalation and surgical VSD repair. However, future larger studies are warranted to confirm this algorithm.

Conclusion

In conclusion, post-MI VSD is associated with extremely high mortality in patients in cardiogenic shock. The introduction of mechanical support and delayed surgical intervention in hemodynamically unstable patients may improve outcomes. Larger and longer studies should be done to provide further evidence for guidelines to advance mechanical support in the early management of post-MI VSD.

References

1. Di Summa M, Actis Dato GM, Centofanti P, Fortunato G, Patane F, Di Rosa E, et al. Ventricular Septal Rupture after a Myocardial Infarction: Clinical Features and Long Term Survival. *J Cardiovasc Surg.* 1997;38(6):589-593.
2. Reeder GS. Identification and Treatment of Complications of Myocardial Infarction. *Mayo Clinic Proc.* 1995;70(9):880-884.



3. Crenshaw BS, Granger CB, Birnbaum Y, Pieper KS, Morris DC, Kleiman NS, et al. Risk Factors, Angiographic Patterns, and Outcomes in Patients with Ventricular Septal Defect Complicating Acute Myocardial Infarction. Gusto-I (Global Utilization of Streptokinase and Tpa for Occluded Coronary Arteries) Trial Investigators. *Circulation*. 2000;101(1):27-32.
4. Yip HK, Fang CY, Tsai KT, Chang HW, Yeh KH, Fu M, et al. The Potential Impact of Primary Percutaneous Coronary Intervention on Ventricular Septal Rupture Complicating Acute Myocardial Infarction. *Chest*. 2004;125(5):1622-1628.
5. Menon V, Webb JG, Hillis LD, Sleeper LA, Abboud R, Dzavik V, et al. Outcome and Profile of Ventricular Septal Rupture with Cardiogenic Shock after Myocardial Infarction: A Report from the Shock Trial Registry. Should We Emergently Revascularize Occluded Coronaries in Cardiogenic Shock? *J Am Coll Cardiol*. 2000;36(3 Suppl A):1110-1116.
6. Daggett WM, Guyton RA, Mundth ED, Buckley MJ, McEnany MT, Gold HK, et al. Surgery for Post-Myocardial Infarct Ventricular Septal Defect. *Ann Surg*. 1977;186(3):260-271.
7. David TE. Operative Management of Postinfarction Ventricular Septal Defect. *Semin Thorac Cardiovasc Surg*. 1995;7(4):208-213.
8. Komeda M, Fremes SE, David TE. Surgical Repair of Postinfarction Ventricular Septal Defect. *Circulation*. 1990;82(5 Suppl):IV243-247.
9. David TE, Dale L, Sun Z. Postinfarction Ventricular Septal Rupture: Repair by Endocardial Patch with Infarct Exclusion. *J Thorac Cardiovasc Surg*. 1995;110(5):1315-1322.
10. Arnaoutakis GJ, Zhao Y, George TJ, Sciortino CM, McCarthy PM, Conte JV. Surgical Repair of Ventricular Septal Defect after Myocardial Infarction: Outcomes from the Society of Thoracic Surgeons National Database. *Ann Thorac Surg*. 2012;94(2):436-443; discussion 443-434.
11. Gregoric ID, Kar B, Mesar T, Nathan S, Radovancevic R, Patel M, et al. Perioperative Use of Tandemheart Percutaneous Ventricular Assist Device in Surgical Repair of Postinfarction Ventricular Septal Defect. *ASAIO*. 2014;60(5):529-532.
12. O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Jr., Chung MK, de Lemos JA, et al. 2013 Accf/Aha Guideline for the Management of St-Elevation Myocardial Infarction: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2013;127(4):e362-425.
13. Yang J, Yang L, Yu S, Liu J, Zuo J, Chen W, et al. Transcatheter Versus Surgical Closure of Perimembranous Ventricular Septal Defects in



- Children: A Randomized Controlled Trial. *J Am Coll Cardiol.* 2014;63(12):1159-1168.
14. Saurav A, Kaushik M, Mahesh Alla V, White MD, Satpathy R, Lanspa T, et al. Comparison of Percutaneous Device Closure Versus Surgical Closure of Peri-Membranous Ventricular Septal Defects: A Systematic Review and Meta-Analysis. *Catheter Cardiovasc Interv.* 2015;86(6):1048-1056.
 15. Loyalka P, Cevik C, Nathan S, Gregoric ID, Kar B. Closure of Post-Myocardial Infarction Ventricular Septal Defect with Use of Intracardiac Echocardiographic Imaging and Percutaneous Left Ventricular Assistance. *Texas Heart Institute journal / from the Texas Heart Institute of St Luke's Episcopal Hospital, Texas Children's Hospital.* 2012;39(3):454-456.
 16. Gregoric ID, Bieniarz MC, Arora H, Frazier OH, Kar B, Loyalka P. Percutaneous Ventricular Assist Device Support in a Patient with a Postinfarction Ventricular Septal Defect. *Texas Heart Institute journal / from the Texas Heart Institute of St Luke's Episcopal Hospital, Texas Children's Hospital.* 2008;35(1):46-49.
 17. Torregrossa G, Anyanwu A, Zucchetta F, Gerosa G. Syncardia: The Total Artificial Heart. *Ann Cardiothor Surg.* 2014;3(6):612-620.
 18. Edwards BS, Edwards WD, Edwards JE. Ventricular Septal Rupture Complicating Acute Myocardial Infarction: Identification of Simple and Complex Types in 53 Autopsied Hearts. *Am J Cardiol.* 1984;54(10):1201-1205.
 19. Lundblad R, Abdelnoor M, Geiran OR, Svennevig JL. Surgical Repair of Postinfarction Ventricular Septal Rupture: Risk Factors of Early and Late Death. *J Thorac Cardiovasc Surg.* 2009;137(4):862-868.
 20. Deville C, Fontan F, Chevalier JM, Madonna F, Ebner A, Besse P. Surgery of Post-Infarction Ventricular Septal Defect: Risk Factors for Hospital Death and Long-Term Results. *Euro J Cardiothor Surg.* 1991;5(4):167-174; discussion 175.
 21. Egbe AC, Poterucha JT, Rihal CS, Taggart NW, Cetta F, Cabalka AK, et al. Transcatheter Closure of Postmyocardial Infarction, Iatrogenic, and Postoperative Ventricular Septal Defects: The Mayo Clinic Experience. *Catheter Cardiovasc Interv.* 2015;86(7):1264-1270.
 22. Holzer R, Balzer D, Amin Z, Ruiz CE, Feinstein J, Bass J, et al. Transcatheter Closure of Postinfarction Ventricular Septal Defects Using the New Amplatzer Muscular Vsd Occluder: Results of a U.S. Registry. *Catheter Cardiovasc Interv.* 2004;61(2):196-201.
 23. Landzberg MJ, Lock JE. Transcatheter Management of Ventricular Septal Rupture after Myocardial Infarction. *Sem Thor Cardiovasc Surg.* 1998;10(2):128-132.



24. Holzer R, Balzer D, Cao QL, Lock K, Hijazi ZM, Amplatz Muscular Ventricular Septal Defect I. Device Closure of Muscular Ventricular Septal Defects Using the Amplatz Muscular Ventricular Septal Defect Occluder: Immediate and Mid-Term Results of a U.S. Registry. *J Am Coll Cardiol.* 2004;43(7):1257-1263.
25. Thiele H, Kaulfersch C, Daehnert I, Schoenauer M, Eitel I, Borger M, et al. Immediate Primary Transcatheter Closure of Postinfarction Ventricular Septal Defects. *Euro Heart J.* 2009;30(1):81-88.
26. Kapur NK, Paruchuri V, Urbano-Morales JA, Mackey EE, Daly GH, Qiao X, et al. Mechanically Unloading the Left Ventricle before Coronary Reperfusion Reduces Left Ventricular Wall Stress and Myocardial Infarct Size. *Circulation.* 2013;128(4):328-336.
27. Gregoric ID, Mesar T, Kar B, Nathan S, Radovancevic R, Patel M, et al. Percutaneous Ventricular Assist Device and Extracorporeal Membrane Oxygenation Support in a Patient with Postinfarction Ventricular Septal Defect and Free Wall Rupture. *Heart Surg Forum.* 2013;16(3):E150-151.
28. Kar B, Adkins LE, Civitello AB, Loyalka P, Palanichamy N, Gemmato CJ, et al. Clinical Experience with the Tandemheart Percutaneous Ventricular Assist Device. *Texas Heart Institute journal / from the Texas Heart Institute of St Luke's Episcopal Hospital, Texas Children's Hospital.* 2006;33(2):111-115.