

RESEARCH ARTICLE

Determination of Patient Learning Needs after Thyroidectomy

Zeynep Temiz¹, Didem Ozturk^{2*}, Gulay Altun Ugras³, Seher Deniz Oztekin², Emel Sengul⁴

Abstract

The purpose of this study was to determine discharge learning needs of patients undergoing thyroidectomy. The population of this descriptive study consisted of patients undergoing thyroidectomy in the Endocrine Surgery Unit of a university hospital between February and December 2013. The study included 251 patients who were discharged after thyroidectomy. Data obtained using the data collection form and the Patient Learning Needs Scale (PLNS) were analyzed by frequency, mean, standard deviation, Kruskal Wallis and student-t tests. The mean age of the patients was 47.91 ± 13.05 and 76.1% were females. The PLNS total mean score was 208.38 ± 34.91 , with the maximum score of 39.23 ± 6.80 on the subscale of treatment and complications and the minimum score of 19.45 ± 4.70 on the subscale of feelings related to condition. It was found that the PLNS total score of the patients was not influenced by age, gender, marital status ($p > 0.05$). This study demonstrated that patients had high learning needs after thyroidectomy.

Keywords: Thyroidectomy - patient learning needs - nursing - post-discharge education

Asian Pac J Cancer Prev, 17 (3), 1479-1483

Introduction

Recent advances in technology and medicine have shortened the length of hospitalization in patients undergoing surgical procedure and increased tendency to out-patient surgery. Considering patient safety and comfort, resource protection and cost-effectiveness, thyroid surgery is performed as an out-patient procedure in most centers (Terris et al., 2013; Perera, Patel and Law, 2014; Rutledge et al., 2014). In the absence of post-thyroidectomy complications, patients are discharged from hospital on the same day of surgery or hospitalized only for one day (Vrabec et al., 2003; Snyder et al., 2010). Therefore, nurses give priority to technical care and patient discharge education is poorly or inadequately provided (Tasocak, 2003; Johansson et al., 2007; Falvo, 2011).

In the lack of optimally efficacious discharge education, or of provision of appropriate information, patients experience more anxiety and problems resulting in unplanned medical visits (Moderchai, Herman, Kerzman and Irony, 2010; Dag et al., 2014). In addition, it has been reported that when patients are concerned and inadequately informed, they feel more dependent. Therefore, nurses should identify the learning needs and knowledge deficits of patients (Johansson et al., 2007; Myers and Pellino, 2009; Falvo, 2011). Throughout the discharge process, nurses are required to assess patient's condition, and provide efficient patient-centered information meeting

post-operative individual needs of patients (Sendir, Buyukyilmaz and Musovi, 2003; Johansson et al., 2007; Falvo, 2011; Galdeano et al., 2014; Krishnatreya et al., 2014). It has been reported in the literature that patients obtaining information specific to their health condition experience an accelerated healing process, show better compliance to treatment, have increased self-confidence, satisfaction, coping skills, and quality of life and decreased anxiety and unplanned medical visits, which in turn reduces health care costs and increases the quality of health care (Clark et al., 2005; McMurray et al., 2007; Piredda et al., 2008; Williams, 2008; Myers and Pellino, 2009; Fedai, Cetin and Teke, 2010).

Shortened hospital stay enhances the importance of effective use of hospitalization time in terms of discharge education (Tasocak, 2003; Johansson et al., 2007; Falvo, 2011; Sendir, Buyukyilmaz and Musovi, 2013). One of the causes of ineffective use of time has been reported in the literature to be the difference between the patients' perceptions of that which is important information and that of the nurses providing the education. In this context, the time allocated for patient education is wasted with the issues that are perceived as important by nurses and those that patients deem to be important are ignored (Bailey, 2004; Kızıl, 2008). For the appropriate identification of patients' educational needs and priorities at discharge, learning needs should be assessed using appropriate tools. The provision of discharge information specific to patients'

¹Nursing Department, Health Sciences Faculty, Artvin Coruh University, Artvin, ²Surgical Nursing Department, Florence Nightingale Nursing Faculty, Istanbul University, Istanbul, ³Nursing Department, School of Health Sciences, Mersin University, Mersin, ⁴Cerrahpasa Faculty of Medicine, Istanbul University, Istanbul, Turkey *For correspondence: didem_trk@hotmail.com

learning needs can help reduce the number of patients presenting to hospital with concern and anxiety about home care and complications. In the light of these data, this study aimed to identify discharge information needs of patients undergoing thyroidectomy and to contribute to preparation of nurses' discharge education planning on patients' needs.

Materials and Methods

Setting and Sample

This descriptive study aimed at identifying discharge learning needs of patients undergoing thyroidectomy at the Endocrine Surgery Unit of Istanbul University, Cerrahpasa Faculty of Medicine. The population of this study consisted of 251 patients undergoing thyroidectomy at the Endocrine Surgery Unit between February and December 2013, who met the study criteria.

The sample size was ensured to be as five-fold of the number of scale items (50 items) in validity and reliability studies. The inclusion criteria were the patients who had been scheduled to be discharged after thyroidectomy, who were oriented to person, place and time and had the ability to speak Turkish, and consented to participate in the study.

Instrument

The instrument developed by the researchers consisted of two sections. Section one included questions related to patients' characteristics. In section two, "The Patient Learning Needs Scale-PLNS" was used to identify patients' discharge learning needs. This scale was developed by Bubela, Galloway, McCay, McKibbin, Nagle, Pringle, Ross & Shamian in 1990, with a Cronbach's alpha coefficient of 0.95 (Bubela et al., 1990). In this study, Cronbach's alpha coefficient for the PLNS was 0.95 and that for the subscales ranged between 0.74 and 0.89. The validity and reliability study was carried out by Catal and Dicle in Turkey in 2008 (Catal and Dicle, 2008).

The PLNS is a 50-item Likert-type scale with 7 subscales. The scale items are scored on a Likert-type scale with, 1: not important 2: less important 3: fairly important 4: very important and 5: extremely important. Accordingly, patients are asked to choose the response that best describes their learning needs and priorities. Result analysis includes assessment of each subscale and total scale scores. The scores obtained from the scale range between 50 and 250 (Bubela et al., 1990).

Procedures

Data were collected by face-to-face interview technique between February 2013 and December 2013. The scale was administered to patients undergoing thyroidectomy 24-48 hours before discharge. All questions that patients had difficulty in understanding/ interpreting were explained by the researcher who accompanied the patient throughout data collection process. Patients who had difficulty reading questions or marking the responses were assisted by the researcher; the questions were read to the patients and they were given time to understand and choose the response and their responses were marked by the researcher.

Data analysis

Data were analyzed using "SPSS for Windows 21.0 (IBM Corp., Armonk, New York)". The data were analyzed statistically using frequency, percentage, mean, standard deviation. The Kruskal-Wallis and Student t test were used for the comparison of the groups. A p value of <0.05 was considered statistically significant, with a confidence interval of 95%.

Ethical Review

Prior to the study, a written consent was obtained from the Endocrine Surgery unit of Istanbul University, Cerrahpasa Faculty of Medicine and the study was approved by the Clinical Research Ethics Committee of Istanbul University Cerrahpasa Faculty of Medicine.

Before the administration of data collection tools, patients were informed about the purpose and content of the study. The patients were ensured that their privacy would be kept, their responses were to remain completely confidential and to be only used for scientific purposes by the researcher. A written and verbal consent was obtained from each patient.

Results

The mean age of the patients enrolled in this study (n:251) was 47.91 ± 13.05 . Of all patients, 76% were females, 82.5% were married, 34.7% had primary education level, and 15.9% had concerns about home care.

Table 1 presents the PLNS mean scores of the patients. The PLNS total mean score of the patients was 208.38 ± 34.91 . The highest mean score was obtained for "treatment and complications" (39.23 ± 6.80) and the lowest mean score was obtained for "feelings related to condition" (19.45 ± 4.70) (Table 1).

Table 2 presents the most important five learning needs of the patients after thyroidectomy. The patients ranked

Table 1. Patients' Learning Needs Subscales Mean Scores (n=251)

Subscales	Number of items	Mean	SD
Medications	8	34.41	5.45
Activities of living	9	37.13	6.96
Community and follow-up	6	23.80	5.01
Feelings related to condition	5	19.45	4.70
Treatment and complications	9	39.23	6.80
Enhancing quality of life	8	34.27	5.60
Skin care	5	20.06	4.13
TOTAL	50	208.38	34.91

Table 2. Rank Order of 5 Most Important Learning Needs of Patients'(n=251)

Learning Need	Mean score X \pm SD
How to care of cut/incision?	4.51 \pm 0.77
How this illness will affect my life?	4.49 \pm 0.90
How to take each medication?	4.49 \pm 0.80
What complications might occur from my illness?	4.49 \pm 0.74
What to do if I have a reaction to the medication?	4.47 \pm 0.80

Table 3. Comparison of Patients' Learning Needs Subscales Mean Scores According to Patients' Characteristics (n=251)

Patients' Learning Needs Subscales								
Characteristics	Medications	Activities of living	Community and follow-up	Feelings related to condition	Treatment and complications	Enhancing quality of life	Skin care	TOTAL
Age*								
20-35	33.28±6.53	69.20±14.92	22.48±5.77	18.26±5.44	38.33±7.19	34.02±6.69	34.15±8.18	202.17±40.54
36-50	35.18±4.20	71.69±11.30	24.45±4.42	20.17±4.29	40.24±7.19	34.76±4.87	37.31±6.40	213.62±29.60
51-65	34.06±6.28	68.18±14.63	23.49±5.29	18.91±4.84	38.64±6.47	33.83±6.10	35.37±7.90	204.72±38.42
≥66	34.45±4.62	70.58±11.24	24.58±4.54	20.41±3.83	38.70±5.12	34.16±4.58	36.91±7.13	210.29±30.29
	KW:2.302	KW:2.546	KW:4.345	KW:5.580	KW:1.618	KW:1.164	KW:5.904	KW:2.830
	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05
Gender**								
Female	34.50±5.72	70.93±13.25	23.94±5.20	19.69±4.75	39.28±6.03	34.51±5.70	36.57±7.59	209.92±35.82
Male	34.11±4.50	67.06±12.43	23.35±4.36	18.66±4.48	39.08±8.90	33.51±5.25	34.53±6.46	203.48±31.64
	t:0.484	t:1.999	t:0.805	t:1.484	t:0.203	t:1.202	t:1.876	t:1.247
	p>0.05	p<0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05
Marital status**								
Married	34.39±5.35	70.10±12.83	23.88±4.92	19.59±4.51	39.27±6.75	34.32±5.37	36.16±7.28	208.76±33.78
Single	34.50±5.95	69.56±14.64	23.40±5.45	18.75±5.49	39.09±7.11	34.04±6.65	35.70±7.85	206.56±40.21
	t:0.115	t:-0.244	t:-0.576	t:-0.958	t:-0.159	t:-0.299	t:-0.375	t:-0.379
	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05
Education level*								
Illiterate	32.90±7.54	67.85±16.84	23.33±6.12	19.95±5.24	37.57±8.62	33.14±7.17	35.66±9.52	202.85±46.64
Primary school	34.27±5.77	69.82±13.52	24.24±5.19	19.73±4.54	39.54±7.98	34.39±5.68	36.14±7.87	209.31±36.01
Secondary school								
High school and higher	34.81±3.63	72.65±8.73	24.71±3.68	20.55±4.07	39.73±3.80	34.81±4.67	37.86±5.20	214.26±24.44
	34.73±5.18	69.60±13.35	23.08±4.99	18.59±4.88	39.09±5.94	34.18±5.52	35.39±7.03	206.26±34.53
	KW:.687	KW:.682	KW:4.721	KW:6.088	KW:.150	KW:1.014	KW:3.072	KW:1.470
	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05	p>0.05
Concern status about home-care **								
Yes	35.60±3.96	38.40±6.63	24.97±4.42	20.50±4.03	40.87±4.36	35.45±4.43	20.87±3.31	216.67±28.44
No	34.18±5.66	36.89±7.01	23.58±5.09	19.25±4.79	38.92±7.14	34.05±5.78	19.91±4.26	206.81±35.85
	t:1.504	t:1.258	t:1.615	t:1.545	t:2.297	t:1.450	t:1.349	t:1.923
	p>0.05	p>0.05	p>0.05	p>0.05	p<0.05	p>0.05	p>0.05	p<0.05

*Kruskal Wallis and **Student-t tests were used

their most important information needs to be “How to care of cut/ incision?” (4.51±0.77), “How this illness will affect my life?” (4.49±0.90), “When to stop taking each medication” (4.49±0.80), “What complications might occur from my illness” (4.49±0.74) and “What to do if I have a reaction to the medication?” (4.47±0.80) (Table 2).

Table 3 presents some characteristics of patients and the comparison of the mean scores of the patients on the PLNS subscales. The PLNS total mean score of the patients aged between 20 and 35 years was 202.17±40.54, that of the patients aged between 36 and 50 years was 213.62±29.60, that of the patients aged between 51 and 65 years was 204.72±38.42 and that of those aged 66 years or older was 210.29±30.29. There was no statistically significant difference between the PLNS mean scores and age of patients (p>0.05). The PLNS mean score of female patients was 209.92±35.82 and that of male patients was 203.48±31.64. No statistically significant difference was found between the PLNS total mean scores and gender of patients (p>0.05), whereas, a statistically significant difference was noted in the “activities of living” subscale (p<0.05). The PLNS and subscales mean scores of

married patients were higher (208.76±33.78) whereas no significant difference was found between marital status and the scale and subscales mean scores (p>0.05). There was no statistically significant difference between level of education and the PLNS and subscales total mean scores (p>0.05), whereas, the patients with secondary education level had higher mean scores (214.26±24.44). It was found that the PLNS mean score of patients who had concerns about home-care at discharge was higher (216.67±28.44) and the difference between the PLNS mean score and the mean score for the subscale of “treatment and complications” was statistically significant (p<0.05) (Table 3).

Discussion

Today, in the absence of any complications, a long hospitalization is not contemplated in patients undergoing thyroidectomy and the patients are discharged in a short time. The tendency for short-stay thyroidectomy deters nurses from spending enough time with patients to provide discharge education. In this context, learning needs should

be assessed using correct tools for the correct identification of patients' discharge learning needs and priorities. This study demonstrated patients' high information needs using the PLNS scale (Table 1). Previous studies using the PLNS scale (Tasdemir et al., 2010; Demirkıran and Uzun, 2012; Erdogan, 2012; Dag et al., 2014) also reported that patients had high information needs prior to discharge, which is also consistent with our findings. On the other hand, a study of 45 surgical patients by Jacobs (2000) reported a PLNS total mean score of 106.2±48.6 and a study of cardiac catheterization patients by Carew (1999) reported a PLNS total mean score of 111.53 ±38.81, which both are lower than that found in this study (Carew, 1999; Jacobs, 2000). The higher mean score on the PLNS in this study compared to those in the latter two studies can be attributed to differences in patients' demographic characteristics, types of surgical procedures, patient education policies in the countries and institutions the studies were conducted in.

Nurses are supposed to develop new strategies to provide an effective discharge education directed at patients' educational needs. In the process of developing these strategies, it is of great importance to identify patients' priorities and primary information needs. In this study, patients ranked the most important five information needs to be "How to care of cut/ incision?", "How this illness will affect my life?", "When to stop taking each medication", "What complications might occur from my illness" and "What to do if I have a reaction to the medication?" (Table 2).

A study of neurosurgery patients by Tasdemir et al. (2010) reported that the most important five learning needs of the patients were "What the possible side effects of my treatment are", "How to manage the symptoms I have", "How this illness will affect my life" "How this illness will affect my future" and "When to stop medicine" (Tasdemir et al., 2010). A study of 212 patients undergoing hip arthroplasty by Johansson et al. (2002) reported that the most important information for the patients was "how to recognize complications" (Johansson, Hupli and Salantera, 2002).

In a study by Clark and Lan (2004), the most important learning need of patients was reported to be "what are the signs and symptoms of complications that may develop after discharge" (Clark and Lan, 2004). The results of this and similar studies suggest that nurses should focus particularly on issues related to treatment and complications and medications in the process of planning and providing discharge education to patients undergoing a surgical procedure.

It was found that patients' age, gender, and marital status had no effect on patients' learning needs, however, female patients identified the area of "activities of living" more important as a learning need and those who had concerns about home care had higher information need in the domain of "treatment and complications" (Table 3). The result of this study is consistent with those previously reported in other studies (Suhonen et al., 2005; Erdogan, 2012; Tan et al., 2013), which might have resulted from the fact that female patients perceived home care after discharge as a more important learning need as compared

to male patients. Postoperatively, patients can experience concerns about possible complications after treatment and discharge and how to manage them. After thyroidectomy, an impaired cosmetic appearance due to visible surgical incision, concerns about loss of voice or impaired voice quality (Cutcliffe and Herth, 2002), as well as, patients' and families' lack of knowledge on home care can drive patients into concerns and fears about everything that have been said and done previously (Toksal, 2005). The results of this study suggest that patients experience concerns and anxiety about post-discharge complications after thyroidectomy and these concerns may be promoted by the inability of provision of appropriate discharge education, which are also consistent with the literature.

One of the limitations of this study is that it included patients undergoing thyroidectomy in the Endocrine Surgery unit of only one university hospital. Therefore the results of this research represent only the mentioned unit. One of the other limitations is that the data were collected based on self-reporting.

In conclusion, It is important to determine patients' learning needs in terms of their priority through the use of appropriate measurements to provide patients with effective education and enhance the quality of their care. This study demonstrated that the patients, in the unit the study had been conducted, had high information needs and had greater concerns and anxiety particularly about treatment and complications. Nurses should recognize and evaluate the learning needs of their patients. These results are also important to nurses who work in practical fields.

References

- Bailey J (2004). Patient and nurses' perceptions of the cardiac patient's learning needs. The Florida State University School of Nursing. Master Thesis. USA: Florida.
- Bubela N., Galloway S, McCay E, et al (1990). The patient learning needs scale: Reliability and validity. *J Advanced Nurs*, **15**, 1181-7.
- Carew D (1999). Discharge information needs of outpatient cardiac catheterization patients. department of nursing science. master thesis. university of toronto, Canada.
- Clark JC, Lan VM (2004). Heart failure patient learning needs after hospital discharge. *Applied Nurs Res*, **17**, 150-7.
- Clark PA, Drain M, Gesell SB, et al (2005). Patient perceptions of quality in discharge instruction. *Patient Education Counseling*, **59**, 56-68.
- Cutcliffe Jr, Herth K (2002). The concept of hope in nursing 1: Its origins, background and nature. *British J Nurs*, **11**, 832-40.
- Catal E, Dicle A (2008). Hasta ogrenim gereksinimleri olcegi'nin turkiye'de gecelik ve guvenirlilik calismasi (A validity and reliability study of the patient learning needs scale in Turkey). *Dokuz Eylul Universitesi Hemsirelik Yuksekokulu Dergisi*, **1**, 19-32.
- Dag H, Donmez S, Gulec D, et al (2014). Jinekolojik operasyon geciren kadınlarnın taburculuk oncesi ogrenim gereksinimleri (Learning needs of gynelological surgery patients before discharge). *Ege Universitesi Hemsirelik Fakultesi Dergisi*, **30**, 49-59.
- Demirkıran G, Uzun O (2012). Koroner arter bypass greft ameliyatı geciren hastaların taburculuk sonrası ogrenim gereksinimleri (Post-discharge learning needs of patients who had undergone coronary artery bypass grafting surgery). *Ege Universitesi Hemsirelik Fakultesi Dergisi*, **28**, 1-12.

- Erdogan M (2012). Acık kalp ameliyati olan hastaların bilgilendirme gereksinimlerinin belirlenmesi (Determining the educational requirements of the patients with open-heart surgery). Sağlık Bilimleri Enstitüsü Hemşirelik Anabilim Dalı. Yayınlanmamış Yüksek Lisans Tezi. Baskent Üniversitesi, Türkiye: Ankara.
- Falvo DR (2011). Effective patient education: a guide to increased adherence. Sudbury, MA: Jones-Barlett Publishers.
- Fedai T, Cetin M, Teke A (2010). Tedavi sürecindeki gereksiz değişkenliklerin Pareto analizi ile değerlendirilmesi (Assessment unnecessary variations in the treatment process with Pareto analysis). *Sağlık Bilimleri Dergisi*, **19**, 184-90.
- Galdeano, LE, Furuya RK, Rodrigues MA, et al (2014). Reliability of the Cardiac Patients Learning Needs Inventory (CPLNI) for use in Portugal. *J Clin Nurs*, **23**, 1532-40.
- Jacobs V (2000). Informational needs of surgical patients following discharge. *Applied Nurs Res*, **13**, 12-8.
- Johansson K, Hupli M, Salantera S (2002). Patients' learning needs after hip arthroplasty. *J Clin Nurs*, **11**, 634-9.
- Johansson K, Salantera, S, Katajisto J (2007). Empowering orthopaedic patients through preadmission education: results from a clinical study. *Patient Education Counselling*, **66**, 84-91.
- Kızıl C (2008). Kalp yetmezliği olan hastaların eğitim gereksinimleri ve hemşirelerin bu konuda düşünceleri (The education requirement of the patient with heart failure and the nurses' thoughts about this). Sağlık Bilimleri Enstitüsü İç Hastalıkları Hemşireliği Anabilim Dalı. Yayınlanmamış Yüksek Lisans Tezi. Dokuz Eylül Üniversitesi, Türkiye: İzmir.
- McMurray A, Johnson P, Wallis M, et al (2007). General surgical patients' perspectives of the adequacy and appropriateness of discharge planning to facilitate health decision-making at home. *J Clinical Nurs*, **16**, 1602-9.
- Moderchai BB, Herman A, Kerzman H, et al (2010). Structured discharge education improves early outcome in orthopedic patients. *Int J Orthopaedic Trauma Nurs*, **14**, 66-74.
- Myers J, Pellino TA (2009). Developing new ways to address learning needs of adult abdominal organ transplant recipients. *Progress Transplant*, **19**, 160-6.
- Perera A, Patel S, Law N (2014). Thyroid surgery as a 23-hour stay procedure. *Ann Royal College Surgeons England*, **96**, 284-8.
- Piredda M, Rocci L, Gualandi R, et al (2008). Survey on learning needs and preferred sources of information to meet the needs in Italian oncology patients receiving chemotherapy. *Eur J Oncol Nurs*, **12**, 120-6.
- Rutledge J, Siegel E, Belcher R, et al (2014). Barriers to same-day discharge of patients undergoing total and completion thyroidectomy. *Otolaryngology-Head Neck Surgery*, **150**, 770-4.
- Snyder S, Hamid KS, Roberson CR, et al (2010). Outpatient thyroidectomy is safe and reasonable: experience with more than 1,000 planned outpatient procedures. *J Am College Surgeons*, **210**, 575.
- Suhonen R, Nenonen H, Laukka A, et al (2005). Patient's informational needs and information received do not correspond in hospital. *J Clinical Nursing*, **14**, 1167.
- Sendir M, Buyukyılmaz F, Musovi D (2013). Patients' discharge information needs after total hip and knee arthroplasty: A quasi-qualitative pilot study. *Rehabilitation Nurs*, **38**, 264-71.
- Tan M, Özdelikara A, Polat H (2013). Hasta öğrenim gereksinimlerinin belirlenmesi (Determination of patient learning needs). *Florence Nightingale Hemşirelik Dergisi*, **2**, 1-8.
- Tasdemir N, Guloglu S, Turan Y, et al (2010). Learning needs of neurosurgery patients. *J Neurological Sci*, **27**, 414-20.
- Tasocak G (2003). Hasta eğitimi (Patient education). İstanbul: İ.U. Basım ve Yayınevi Müdürlüğü.
- Terris DJ, Snyder S, Carneiro-Pla D, et al (2013). American thyroid association statement on outpatient thyroidectomy. *Thyroid*, **23**, 1193-202.
- Toksallı M (2005). Preoperatif dönemde hastaların hemşirelerden beklentilerinin araştırılması (Investigation of patients' expectations from nurses before the operation). Sağlık Bilimleri Enstitüsü Hemşirelik Anabilim Dalı. Yayınlanmamış Yüksek Lisans Tezi. Afyon Kocatepe Üniversitesi, Türkiye: Afyonkarahisar.
- Vrabec S, Oltmann SC, Clark N, et al (2013). A short-stay unit for thyroidectomy patients increases discharge efficiency. *J Surg Res*, **184**, 204-8.
- Williams B (2008). Supporting self-care of patients following general abdominal surgery. *J Clin Nurs*, **17**, 584-92.
- Krishnatreya M, Katakı AC, Sharma JD, et al (2014). Educational Levels and Delays in Start of Treatment for Head and Neck Cancers in North-East India. *Asian Pac J Cancer Prev*, **15**, 10867-9.