

University of Massachusetts Medical School
eScholarship@UMMS

University of Massachusetts Medical School Faculty Publications

2020-07-08


A longitudinal implementation evaluation of a physical activity program for cancer survivors: LIVESTRONG(R) at the YMCA

Jamie Faro
University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/faculty_pubs

 Part of the [Community-Based Research Commons](#), [Community Health and Preventive Medicine Commons](#), [Epidemiology Commons](#), [Health Services Administration Commons](#), [Health Services Research Commons](#), and the [Neoplasms Commons](#)

Repository Citation

Faro J, Arem H, Heston A, Hohman KH, Hodge H, Wang B, Lemon SC, Houston TK, Sadasivam RS. (2020). A longitudinal implementation evaluation of a physical activity program for cancer survivors: LIVESTRONG(R) at the YMCA. University of Massachusetts Medical School Faculty Publications. <https://doi.org/10.1186/s43058-020-00051-3>. Retrieved from https://escholarship.umassmed.edu/faculty_pubs/1794

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).


This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in University of Massachusetts Medical School Faculty Publications by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

SHORT REPORT

Open Access



A longitudinal implementation evaluation of a physical activity program for cancer survivors: LIVESTRONG® at the YMCA

Jamie M. Faro^{1*} , Hannah Arem², Ann-Hilary Heston³, Katherine H. Hohman³, Heather Hodge³, Bo Wang¹, Stephenie C. Lemon¹, Thomas K. Houston⁴ and Rajani S. Sadasivam¹

Abstract

Purpose: Increased physical activity (PA) levels in cancer survivors are associated with decreased risk of recurrence and mortality as well as additional positive health outcomes. PA interventions have shown to be efficacious, though many lack translation to and sustainability in community settings. We used dimensions of the RE-AIM framework to evaluate LIVESTRONG® at the YMCA, a nation-wide community-based PA program for cancer survivors delivered at Ys.

Methods: This was a longitudinal study design using national LIVESTRONG at the YMCA data compiled between 2010 and 2018. Data is from all YMCAs who deliver LIVESTRONG at the YMCA, submitted by Program Directors to the YMCA-USA. We assessed reach (number of participants), adoption (associations offering the program), implementation (conducting 3 fidelity checks), and organizational level maintenance (associations recently offering program). We also examined relationships between organizational characteristics (years of program existence and association area household income) and program implementation factors with member conversion rates.

Results: As of 2018, LIVESTRONG at the YMCA has reached 62,044 survivors and 245 of the 840 (29.2%) of Y associations have adopted the program. Among the adopters, 91% were aware of fidelity checks; implementation of observational (62.3%), goal setting (49.9%), and functional (64.6%) checklists varied. Most (95.1%) adopters reported offering ≥ 1 LIVESTRONG session per year (organizational-level maintenance) and a facility-level mean membership conversion percentage of $46.9 \pm 31.2\%$. Fewer years implementing the program and higher association area household income were significantly associated with a greater membership conversion rate vs their comparison. In a multiple regression model controlling for organizational characteristics, conducting the fidelity checks independently (observational, $\beta = 8.41$; goal-setting, $\beta = 9.70$; and functional, $\beta = 9.61$) and collectively ($\beta = 10.82$; 95% CI 5.90–16.80) was positively associated with higher membership conversion rates.

Conclusions: LIVESTRONG at the YMCA, in its early years, has shown promise for high reach, while adoption at more associations could be facilitated. Implementing fidelity checks along with organizational characteristics were associated with membership conversion rate. Identification of association-level strategies to increase reach, adoption, implementation, and maintenance may increase the impact of this community-based PA program.

Keywords: Cancer, Physical activity, Implementation, Evaluation, Community-based

* Correspondence: Jamie.faro@umassmed.edu

¹Department of Population and Quantitative Health Sciences, University of Massachusetts Medical School, 368 Plantation St, Worcester, MA 01605, USA
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Contributions to the literature

- LIVESTRONG at the YMCA is a nationwide physical activity program with the capacity to reach many cancer survivors, though the public health impact has yet to be examined using an implementation framework.
- Although we found the program reached many survivors, increasing adoption rates has the potential to reach a greater number of survivors. Of those reached, implementation measures and program characteristics influenced the membership conversion rate.
- This study is the first to assess implementation outcomes of a survivor community-based physical activity program and highlight the need for additional insight into implementation, organizational-level characteristics, and strategies to create more sustainable programs.

Background

In the USA, the number of cancer survivors has increased steadily as cancer death rates have decreased [1]. There were 16.9 million survivors in 2018, a number which is expected to increase to 20.3 million by 2026 [2]. Health priorities among survivors include decreasing the risk of cancer recurrence, improving the quality of life and mental health outcomes, and general health promotion [3]. Regular physical activity (PA) has been associated with lower secondary cancer recurrence and improvements in quality of life, fatigue, fitness, body composition, mood, self-esteem, and physical function [4–6]. The American College of Sports Medicine has determined the efficacy and safety of PA for survivors [7]. American Cancer Society (ACS) guidelines recommend engaging in 150 min of moderate-to-vigorous physical activity and 2 days of strength training per week [8]. However, less than < 30% of survivors are meeting guidelines [9].

As survivorship numbers grow and PA program effectiveness data increases, the need for national dissemination of programs addressing survivor needs has become evident [10]. In 2007, the YMCA of the USA (Y-USA) partnered with LIVESTRONG® to design an evidence-based 12-week exercise intervention free of cost for cancer patients and survivors, LIVESTRONG at the YMCA [10]. In brief, trained Y instructors facilitate two weekly sessions over 12 weeks to improve participants' aerobic fitness, muscle mass, strength flexibility and balance, and social support. The program has evidenced increases in cardiorespiratory fitness, PA levels, and quality of life [11, 12]. Since the initial pilot in 2008, a national infrastructure was created to increase its dissemination and implementation [10]. Since 2010, Y-USA has collected nationwide data about the program [13]. While a prior

study reported on the reach and adoption of the program as of 2015 [10], the public health field is lacking an examination of the implementation, organizational maintenance, and membership conversion rate in this evidence-based program.

The objective of this longitudinal study was to assess implementation outcomes of LIVESTRONG at the YMCA using nationwide data collected by the Y-USA between 2010 and 2018 [13]. To conduct this analysis, we used the RE-AIM (Reach, Effectiveness, Adoption, Implementation and Maintenance) framework. RE-AIM is concerned with issues related to impact in real-world settings, incorporates both individual and organizational setting level variables, and describes the population-based impact of an intervention [14]. The membership conversion rate may be one indicator of ensuring the long-term health benefits the program provides to survivors through continued physical activity; thus, a secondary aim was to determine associations between organizational characteristics and implementation measures with this RE-AIM metric. LIVESTRONG at the YMCA is the only nationwide community-based physical activity program for survivors and has been in existence for over a decade. Thus, gaining a better understanding of this large public health program that has been adopted, implemented, and maintained in real-world settings is critical to advance the field of PA and cancer survivorship. As data in this study are provided by community workers and not researchers, we are limited in some metrics and unable to examine all RE-AIM aspects. For example, effectiveness data of participant-level outcomes was not provided in this dataset but has been shown elsewhere [12, 15].

Methods

Study design and participants

This was a longitudinal study design using data collected by Y-USA about the LIVESTRONG at the YMCA program between 2010 and 2018. The Y serves 22 million people in over 800 associations and 2700 individual branches across all 50 states. Y associations may consist of single or multiple branches (up to 40) within their association and operate independently of other associations. The process of becoming a LIVESTRONG at the YMCA provider has been described elsewhere [10]. In brief, YMCAs interested in becoming providers apply to the Y-USA and complete a simple readiness assessment scored by two reviewers. If the YMCA receives a score indicating they have the capacity to become a provider, they will move on to complete a 6-month on-boarding and learning process. The Y-USA is the national resource office for all Ys and exists to serve Ys. Y-USA offers technical and administrative support throughout, though Association Program Directors are responsible for marketing, administration, oversight, and funding

within their association. This study was approved by the Institutional Review Board of the University of Massachusetts Medical School.

Data sources

Data for this study included (1) routine evaluation data collected between January 2010 and June 2018 from the Association Program Directors and (2) 2010 US Census data. Y data included organizational characteristics, overall number of participants completing the program (as well as broken down by a priori membership or decision to join after the program), and awareness and implementation of three fidelity check tools separately. Reporting of all data by Program Directors is encouraged, but not required. Program data is sent to Y-USA quarterly: January 1–March 31, April 1–June 30, July 1–September 30 and October 1–December 31. 2010 U.S. Census data were used to assess the household income of each association.

Measures

Reach and adoption

Reach has been identified as the absolute number, proportion, and representativeness of individuals who are willing to participate in an intervention [13]. While prior studies have examined representativeness on a small scale [12, 15], we did not have access to these data. To participate in LIVESTRONG at the Y, participants must (1) be aged 18 years of older, (2) have a previous cancer diagnosis, (3) receive medical clearance, and (4) be able to attend most sessions. Due to these criteria, we were unable to determine a denominator (number of eligible participants) to determine the proportion of individuals participating. However, using national data [16], we provide an estimate of the proportion of all possible cancer survivors (as of December 31, 2018) who have completed the program. Thus, reach was defined as one of the absolute number of participants completing the program and also an estimate of the proportion of cancer survivors completing the program. Adoption refers to the representativeness and proportion of given settings that adopt a program [13]. All 840 associations are eligible to apply to become a LIVESTRONG at the YMCA provider limited by the data provided, and adoption was conceptualized as the proportion of associations (out of all 840 associations) becoming approved LIVESTRONG at the YMCA providers. We calculated this rate as the number of approved LIVESTRONG at the YMCA association providers divided by the total number of Y associations.

Implementation

Implementation refers to the extent to which a program is delivered as intended [13]. While implementation

incorporates numerous components, we only assess one aspect; the extent to which implementers were aware of fidelity checklists and whether or not they used them. Due to limited data, we were unable to assess other implementation aspects, such as the consistency of delivery as intended (through results of the fidelity checks) and the time and cost of the program [17]. Fidelity checklists were designed specifically for LIVESTRONG at the YMCA by the program developers and implemented from January 2017 to June 2018. Program Directors were encouraged to use three fidelity tools conducted at least annually to ensure the program was being delivered as intended (1) *Observation Assessment Tool*—used by the Program Director to observe each Instructor conducting a session, (2) *Instructor Goal Setting*—following feedback from the observation, the Instructor and Program Director worked collaboratively to identify the Instructor's areas of strength and opportunities for improvement and document these goals, and (3) *Functional Assessments Checklist*—Program Directors observed a minimum of two Functional Assessments (baseline or 12 weeks) implemented by program Instructors. Assessing whether these checks were implemented may be one preliminary method of assessing implementation prior to examining results. Thus, implementation was conceptualized as the percentage of associations implementing fidelity checks during each reporting quarter. Program Directors reported on whether they were aware of the fidelity checklists (yes/no) and whether they completed each of the three fidelity checks (yes/no). As stated above, this assessment represents only a small section of the implementation metric and obtaining data from the actual checks themselves would provide greater insight into implementation.

Maintenance

Maintenance can refer to the extent to which a program can be sustained over time and may be measured at both the organizational and participant level [13]. Due to only organizational-level data available, we defined maintenance as the percentage of associations offering at least one 12-week session since the last full reporting year, 2017, divided by the total number of approved LIVESTRONG at the YMCA associations.

Membership conversion rate

Membership conversion rate is defined as the percentage of non-members purchasing a membership following the program cessation. Membership conversion rate (rate of non-members purchasing membership) may serve as a proxy to the lack of national-level PA data following program cessation and has been shown to predict future PA [18]. Program Directors reported the number of members and non-members who completed the

program and the number of non-members who became members following the program. We divided the total number completing the program who became members by the total number of non-members completing programs to obtain this rate.

Organizational characteristics

Y-USA maintains the number of years the association has implemented the program along with the city and zip code of the corporate branch. We categorized the duration of program implementation into low and high, by splitting associations on the median number of years ($n = 7$) offering the program. Using U.S. Census Bureau [19], median household income data was collected for each association. The sample's area household income was divided at the median (\$47,300) to classify into high- and low-household income.

Statistical analyses

Descriptive statistics were calculated. Independent t tests were used to compare membership conversion rates between organizational characteristics and implementation of fidelity checklists. Due to the variability of implementing 1, 2, and all 3 checklists, we ran all checklist models independently and cumulatively. We followed bivariate analyses with multiple linear regression analyses adjusted for organizational characteristics. These were used to determine the independent relationships between the implementation of fidelity checks and membership conversion rate. Separate models were run for all 3 fidelity variables, in addition to a 4th model run using all 3 fidelity variables. Missing data were removed from the analyses. Analyses were conducted using STATA. All statistical tests were two-tailed and considered significant at $p < 0.05$.

Results

Descriptive statistics for organizational characteristics are shown in Table 1.

Reach and adoption

Figure 1 shows the number of participants completing the program per year. In 2010, 4019 participants had completed the program, with that number increasing to 62,044 by 2018. The number of participants completing the program has steadily increased, with more rapid increases in recent reporting years. As of December 31, 2018, there were 16,806,760 US cancer survivors > 20 years of age [16]. Thus, the proportion of survivors completing the program of all US survivors was 0.004% (Table 1). Figure 2 shows the number of associations offering the program per year. The adoption rate steadily increased apart from 2014 to 2015 and was 29.2% as of June 2018.

Implementation

During 2017–2018, an estimated 91% ($n = 233$) of Y associations offering LIVESTRONG at the YMCA were aware of the fidelity checklists. Of those aware of checklists, 62.6% implemented the observational assessment checklist, 50.2% the goal-setting checklist, and 65.1% the functional assessment checklist. Of those associations aware of all checklists, 62.3% implemented the observational assessment checklist, 49.9% the goal-setting checklist, and 64.6% the functional assessment checklists, while only 40% (implemented all three checklists). Thus, of the associations aware of the checklists ($n = 223$), 144 did not implement all three checklists.

Maintenance

At the organizational level, sessions were offered during 68.2% of possible reporting quarters, while 95.1% of all

Table 1 Descriptive statistics for associations offering LIVESTRONG at the YMCA

RE-AIM element	Variable	Total N, frequency (%) or $M \pm SD$
Reach	Number and percentage of survivors completing program, N (%)	62,044 (0.004%)
Adoption	Number and percentage of Y Associations delivering program, N (%)	245 (29.2%)
Implementation	Percent aware of fidelity checklists ²	89.51%
	Percent implemented Observational checklist	62.24%
	Percent implemented Goal-setting checklist	50.19%
	Percent implemented Functional checklist	65.10%
Maintenance (organizational-level)	Percentage of associations reporting ≥ 1 sessions within last full calendar year	95.1%
<i>Membership conversion and organizational characteristics</i>		
N/A	Mean membership conversion rate per association ($M \pm SD$)	46.44 \pm 30.9
N/A	Number of years association implementing program	6.3 \pm 2.1
N/A	Mean association area household income ($M \pm SD$)	53,582.42 \pm 24,522.83

¹Partial data year

²Fidelity checklists represent data collected in 2017–2018




Fig. 1 Cumulative number of participants completing the program between the years of 2010 and 2018

on-boarded associations offered at least one session since the reporting year 2017.

Factors associated with membership conversion rates

The mean membership conversion rate was 46.4% ± 30.9, with a range from 0 to 100%. Membership conversion rates were significantly greater in associations that implemented observational, goal-setting, and functional checklists independently (see Table 2). We found similar results for those associations implementing all three checklists. In a linear regression model adjusting for organizational characteristics, implementing observational ($\beta = 8.40$, 95% CI 3.33–14.67), goal-setting ($\beta = 9.7$, 95% CI 4.85–16.63), and functional ($\beta = 9.61$, 95% CI 3.98–15.6) checklists independently and collectively ($\beta = 10.82$; 95% CI 5.90–16.80) were significantly associated with greater membership conversion rates. Associations implementing the program for < 7 years had a

significantly greater membership conversion rate than those implementing for ≥ 7 years. The membership conversion rate was significantly greater in higher household income areas as compared to lower household income areas (48.5 vs. 44.29; see Table 2).

Discussion

Our results suggest that while LIVESTRONG at the YMCA has linearly increased its participant reach and organizational adoption rates, it has room to continue to grow. Implementing community-based programs with high fidelity is challenging, and our findings suggest implementation measures of fidelity checks can be improved. While the data only shows whether checks were implemented and not actual fidelity to the program, we did find implementing checks were associated with membership conversion rates. Further, we found that area household income was associated with membership




Fig. 2 Adoption rate and number of associations trained to deliver LIVESTRONG at the YMCA between the years of 2010 and 2018

Table 2 Independent sample *t* tests comparing membership conversion rates between organizational characteristics and fidelity checklist awareness and implementation

Variable	Membership conversion rate (<i>M</i> ± <i>SD</i>)	95% <i>CI</i>	<i>p</i> value
<i>Organizational characteristics</i>			
Time implementing program (years)			< 0.001
< 7 years	49.3 ± 31.6	47.6 to 51.0	
> 7 or more years	42.8 ± 29.7	41.0 to 44.6	
Association area household income (median)*			0.001
≤ 47,300	44.3 ± 30.3	42.5 to 46.1	
> \$47,300	48.5 ± 31.4	46.7 to 50.2	
<i>Fidelity checklists**</i>			
Aware of fidelity checklists			0.832
Yes	46.6 ± 37.5	37.3 to 56.0	
No	47.5 ± 32.1	44.8 to 50.3	
Implemented observational checklist			0.002
Yes	50.6 ± 1.7	47.2 to 54.0	
No	41.6 ± 30.3	37.2 to 46.0	
Implemented goal setting checklist			< 0.001
Yes	52.5 ± 1.9	48.7 to 56.3	
No	42.2 ± 30.7	38.4 to 46.0	
Implemented functional checklist			0.001
Yes	50.6 ± 32.4	47.2 to 54.0	
No	40.8 ± 30.4	36.2 to 45.4	
Implemented all 3 checklists			< 0.001
Yes	54.2 ± 32.3	49.9 to 58.4	
No	42.8 ± 31.2	39.3 to 46.3	

P value significant at < 0.05

*Association area median household income was acquired using data from the US Census Bureau

**Fidelity checks only apply to data collected in 2017–2018

conversion rates as well. This supports the notion that inequities exist in survivor PA and potentially access to community-based programs, in low socioeconomic status areas [20].

The estimated 60,000 survivors that the program has reached is only a small fraction of the roughly 16.9 million survivors in 2018 [2]. In an attempt to provide a proportion for the number of survivors participating in the program, we found that 0.004% of US survivors completed the program. However, this value does not consider medical eligibility, location to the nearest YMCA, or other contextual factors needed to determine eligible participants. Increasing the number of participants is an important research priority. A prior examination of a subset of program participants (7%) showed the highest reported method of referral to LIVESTRONG at the YMCA was from a doctor or other healthcare professional [10]. Providers may serve on the front line to screen and refer patients to appropriate programs that fit their medical, geographic, social, and economic

preferences [21]. Recent health reforms have placed an emphasis on using electronic medical health records for surveillance [22]. Thus, integrating PA surveillance into the standard of care may provide better insight into patient characteristics, medical clearances, and referral to appropriate PA programs, such as LIVESTRONG at the YMCA.

Reach may also be increased if additional Ys adopt the program, though strategies are still needed. A previous study examining a health program in Y-affiliated sites found that adoption facilitators included organizational support, on-going financial support, matching the Ys mission and target population, novelty of the program, invitations from established partners, and program champions [23]. Barriers included limited resources and expertise, competing programs, and space and costs of offering the program. A prior examination of the Diabetes Prevention Program delivered in YMCAs found that outreach and recruitment required 2 to 20 h of staff time per week [24]. As LIVESTRONG at the YMCA is

free of cost, the YMCA staff must use some of their time to employ fundraising efforts to fund the program. Alleviating staff time and the financial burden of program costs may increase the adoption of the program and staff time to devote to outreach. The American Society of Clinical Oncology has encouraged a third-party payer system to provide coverage of services for cancer prevention and control, including those for PA [25]. Payer financial assistance may alleviate fundraising burden from Ys and provide opportunities for more Ys to adopt the program, run additional sessions, perform outreach efforts, and reach more individuals.

Program fidelity may potentially moderate the relationship between an intervention and its outcomes [26]. Less than 40% of associations were implementing all three checks which were associated with greater membership conversion rates. Fidelity is associated with an intervention's outcomes [27], and incorporating checklists is one way to measure adherence to delivering the intervention as intended. However, fidelity monitoring delivered in non-research-based settings presents several logistical concerns of self-report measures, time, and resources to complete checks while concurrently implementing the program, as well as adaptation to the local setting and drift from the intervention [28]. Implementation strategies may be needed to promote fidelity. We also found that fewer years implementing the program was associated with a higher membership conversion rate. Examining setting-specific variables affecting programs implemented over a longer time, such as funding, community saturation, change in organizational structure, adaptability of the intervention, and support from leadership [29], are warranted.

Consistent with prior data that the purchase of a fitness membership is limited to those of higher socioeconomic status [30, 31], we found that household area income was associated with membership conversion rate despite that the Y offers financial assistance to those in need. Strategies to motivate and support participants facing financial stress are needed to reduce the disparities in participation. Survivors have reported financial constraints as a barrier to exercise [32]. They also report spending 1/3 of their household income on cancer care [33]. Third-party payer systems covering survivor PA services may provide a reallocation of funds to overcome financial barriers to program attendance, including childcare, transportation, and athletic gear. Additionally, there are considerable disparities in the population being served in PA programs for survivors [34]; thus, there is a need to determine how to make even free-of-cost programs more accessible to minority survivors and those with low socioeconomic status. Providers may be able to assist in these efforts, as ACS guidelines and the Institute of Medicine recommend PA prescriptions and/or referrals be provided to survivors. However, specific recommendations on how to prescribe

or where to refer patients are not included [35]. Provider education coupled with assessing barriers to PA may aid in the PA referral process.

Several limitations should be noted. First, data is optionally self-reported from Program Directors; thus, it is unclear if an association with no report conducted sessions and our results may underestimate the outcomes of interest. Second, data is reported from Y associations rather than individual branches; therefore, it is unclear as to how individual branches perform within each association as well as a lack of branch-specific contextual factors (such as staffing, financial resources, facilities, equipment, and leadership) which may influence the capacity and performance of programs. Third, the RE-AIM metrics identified in this study are limited to the data provided. This is a strength, as measures are collected by all associations similarly, though a weakness as these measures do not fully capture all indicators of each RE-AIM aspect (such as the unknown characteristics of those not participating in the program, a denominator that captures all eligible participants for reach and the number of associations who applied to become a LIVESTRONG provider but were not approved or failed to complete the training). In lieu of long-term PA maintenance measures, our data was limited to membership conversion rates upon program cessation, which is not a construct within the RE-AIM framework. This measure also does not account for those who purchase a membership later nor assess membership use or PA behaviors in alternative settings. Fourth, our metric of household income based on census data is limited to the corporate branch within the association. Not having data at the level of the implementing branches limited our ability to understand the implementation context. Lastly, the metric of associations offering at least one session per full reporting year provides only preliminary insight into an association's organizational maintenance.

Conclusions

Applying RE-AIM to evaluate a community-based health program presents a number of complexities that are not present in traditional research-based programs [36]. We provided an examination into the implementation of this program, which will become more needed as the number of cancer survivors increases and opportunities for structured, evidence-based PA programs become critical. LIVESTRONG at the YMCA has the potential to reach many communities, successfully implement, sustain, and expand the program over the course of a decade. However, disparities in the programs' reach remain, and processes need to be integrated into the standard of care to screen and refer survivors. Future efforts should address setting-specific contextual factors to allow for the

identification of strategies and tools to enhance program implementation and maintenance. These efforts will be strengthened by studies that assess all RE-AIM measures.

Abbreviations

ACS: American Cancer Society; PA: Physical activity; RE-AIM: Reach Effectiveness Adoption Implementation Maintenance; YMCA: Young Men's Christian Association

Acknowledgements

Not applicable

Authors' contributions

JF led the conception and design of the study; was closely involved in the data cleaning, analysis, and interpretation; and wrote the manuscript. AHH, KH, and HH aided in data interpretation and contributed to the intellectual content. BW was closely involved in data cleaning, analysis, and interpretation and revised the manuscript for intellectual content. HA, SL, and TK were involved in the design of the study and data interpretation and revised the manuscript for intellectual content. RS was closely involved in the study design, data analysis, interpretation, and intellectual content of the manuscript. The authors read and approved the final manuscript.

Funding

The study was funded by a research grant from the National Cancer Institute (Grant # T32 CA172009). The funder had no role in the design, collection, analysis, and interpretation of data; in the writing of the manuscript; and in the decision to submit this manuscript for publication.

Availability of data and materials

The data that support the findings of this study are available from Y-USA but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of Y-USA.

Ethics approval and consent to participate

This study was approved by the Institutional Review Board of the University of Massachusetts Medical School.

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Population and Quantitative Health Sciences, University of Massachusetts Medical School, 368 Plantation St, Worcester, MA 01605, USA.

²Department of Epidemiology and Biostatistics, George Washington University, 950 New Hampshire Avenue NW, Room 514, Washington, DC 20052, USA. ³YMCA of the USA, 101 N Upper Wacker Dr, Chicago, IL 60606, USA. ⁴Wake Forest University School of Medicine, Medical Center Boulevard, Winston-Salem, NC 27157, USA.

Received: 18 January 2020 Accepted: 15 June 2020

Published online: 08 July 2020

References

- National Cancer Institute. Cancer Statistics 2018 [updated April 27, 2018]. Available from: <https://www.cancer.gov/about-cancer/understanding/statistics>.
- Bluthmann SM, Mariotto AB, Rowland JH. Anticipating the "Silver Tsunami": prevalence trajectories and comorbidity burden among older cancer survivors in the United States. *Cancer epidemiology, biomarkers & prevention* : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology. 2016;25(7):1029–36.
- Jarrett N, Scott I, Addington-Hall J, Amir Z, Brearley S, Hodges L, et al. Informing future research priorities into the psychological and social problems faced by cancer survivors: a rapid review and synthesis of the literature. *European Journal of Oncology Nursing*. 2013;17(5):510–20.
- Runowicz CD, Leach CR, Henry NL, Henry KS, Mackey HT, Cowens-Alvarado RL, et al. American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. *CA Cancer J Clin*. 2016;66(1):43–73.
- Courneya KS. Exercise in cancer survivors: an overview of research. *Medicine and science in sports and exercise*. 2003;35(11):1846–52.
- Schmitz KH, Campbell AM, Stuver MM, Pinto BM, Schwartz AL, Morris GS, et al. Exercise is medicine in oncology: engaging clinicians to help patients move through cancer. *CA: a cancer journal for clinicians*. 2019.
- Campbell KL, Winters-Stone KM, Wiskemann J, May AM, Schwartz AL, Courneya KS, et al. Exercise guidelines for cancer survivors: consensus statement from International Multidisciplinary Roundtable. *Medicine & Science in Sports & Exercise*. 2019;51(11):2375–90.
- Rock CL, Doyle C, Demark-Wahnefried W, Meyerhardt J, Courneya KS, Schwartz AL, et al. Nutrition and physical activity guidelines for cancer survivors. *CA: A Cancer Journal for Clinicians*. 2012;62(4):242–74.
- Ballard-Barbash R, Friedenreich CM, Courneya KS, Siddiqi SM, McTiernan A, Alfano CM. Physical activity, biomarkers, and disease outcomes in cancer survivors: a systematic review. *J Natl Cancer Inst*. 2012;104(11):815–40.
- Heston AH, Schwartz AL, Justice-Gardiner H, Hohman KH. Addressing physical activity needs of survivors by developing a community-based exercise program: LIVESTRONG(R) at the YMCA. *Clin J Oncol Nurs*. 2015;19(2):213–7.
- Tran H, Lin C, Yu F, Frederick A, Mieras M, Baccaglini L. A multicenter study on the relative effectiveness of a 12-week physical training program for adults with an oncologic diagnosis. *Supportive Care in Cancer*. 2016;24(9):3705–13.
- Irwin ML, Cartmel B, Harrigan M, Li F, Sanft T, Shockro L, et al. Effect of the LIVESTRONG at the YMCA exercise program on physical activity, fitness, quality of life, and fatigue in cancer survivors. *Cancer*. 2017;123(7):1249–58.
- Glasgow RE, Vogt TM, Boles SM. Evaluating the public health impact of health promotion interventions: the RE-AIM framework. *American Journal of Public Health*. 1999;89(9):1322–7.
- Glasgow RE, McKay HG, Piette JD, Reynolds KD. The RE-AIM framework for evaluating interventions: what can it tell us about approaches to chronic illness management? *Patient Education and Counseling*. 2001;44(2):119–27.
- Schumacher MM, McNeil P. The impact of LIVESTRONG® at the YMCA for cancer survivors. *Oncol Nurs Forum*. 2018;45(6):717–25.
- American Cancer Society. Cancer treatment & survivorship facts & figures 2019–2021 Atlanta, GA2019 [updated 2019]. Available from: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/cancer-treatment-and-survivorship-facts-and-figures/cancer-treatment-and-survivorship-facts-and-figures-2019-2021.pdf>.
- Glasgow RE, Harden SM, Gaglio B, Rabin B, Smith ML, Porter GC, et al. RE-AIM planning and evaluation framework: adapting to new science and practice with a 20-year review. *Frontiers in Public Health*. 2019;7(64).
- Ready AE, Naimark BJ, Tate R, Boreskie SL. Fitness centre membership is related to healthy behaviours. *J Sports Med Phys Fitness*. 2005;45(2):199–207.
- U.S. Census Bureau; 2015 [Available from: <http://factfinder.census.gov>].
- Kushi LH, Doyle C, McCullough M, Rock CL, Demark-Wahnefried W, Bander EV, et al. American Cancer Society guidelines on nutrition and physical activity for cancer prevention. *CA: A Cancer Journal for Clinicians*. 2012;62(1):30–67.
- Mina DS, Sabiston CM, Au D, Fong AJ, Capozzi LC, Langelier D, et al. Connecting people with cancer to physical activity and exercise programs: a pathway to create accessibility and engagement. *Current oncology (Toronto, Ont)*. 2018;25(2):149–62.
- Glasgow RE, Kaplan RM, Ockene JK, Fisher EB, Emmons KM. Patient-reported measures of psychosocial issues and health behavior should be added to electronic health records. *Health Affairs*. 2012;31(3):497–504.
- Belza B, Petrescu-Prahova M, Kohn M, Miyawaki CE, Farren L, Kline G, et al. Adoption of evidence-based health promotion programs: perspectives of Early Adopters of Enhance®Fitness in YMCA-affiliated sites. *Frontiers in Public Health*. 2015;2(164).
- Bozack A, Millstein S, Garcel JM, Kelly K, Ruberto R, Weiss L. Implementation and outcomes of the New York State YMCA diabetes prevention program: a multisite community-based translation, 2010–2012. *Preventing chronic disease*. 2014;11:E115–E.
- Bennett GG, Glasgow RE. The delivery of public health interventions via the internet: actualizing their potential. *The Annual Review of Public Health*. 2009;30(Journal Article):273–292.

26. Carroll C, Patterson M, Wood S, Booth A, Rick J, Balain S. A conceptual framework for implementation fidelity. *Implementation Science*. 2007;2(1):40.
27. Gearing RE, El-Bassel N, Ghesquiere A, Baldwin S, Gillies J, Ngeow E. Major ingredients of fidelity: a review and scientific guide to improving quality of intervention research implementation. *Clinical Psychology Review*. 2011; 31(1):79–88.
28. Heerman WJ, Schludt D, Harris D, Teeters L, Apple R, Barkin SL. Scale-out of a community-based behavioral intervention for childhood obesity: pilot implementation evaluation. *BMC Public Health*. 2018;18(1):498.
29. Durlak JA, DuPre EP. Implementation matters: a review of research on the influence of implementation on program outcomes and the factors affecting implementation. *American Journal of Community Psychology*. 2008;41(3):327.
30. Powell LM, Slater S, Chaloupka FJ, Harper D. Availability of physical activity-related facilities and neighborhood demographic and socioeconomic characteristics: a national study. *American Journal of Public Health*. 2006; 96(9):1676–80.
31. Hillsdon M, Panter J, Foster C, Jones A. Equitable access to exercise facilities. *American Journal of Preventive Medicine*. 2007;32(6):506–8.
32. Santa Mina D, Petrella A, Currie KL, Bietola K, Alibhai SMH, Trachtenberg J, et al. Enablers and barriers in delivery of a cancer exercise program: the Canadian experience. *Current oncology (Toronto, Ont)*. 2015;22(6):374–84.
33. Chino F, Peppercorn JM, Rushing C, Kamal AH, Altomare I, Samsa G, et al. Out-of-pocket costs, financial distress, and underinsurance in Cancer Care. *Out-of-Pocket Costs, Financial Distress, and Underinsurance in Cancer Care Letters*. *JAMA Oncology*. 2017;3(11):1582–4.
34. Turner RR, Steed L, Quirk H, Greasley RU, Saxton JM, Taylor SJC, et al. Interventions for promoting habitual exercise in people living with and beyond cancer. *Cochrane Database of Systematic Reviews*. 2018;9.
35. Hewitt ME. American Society of Clinical O, Ganz P, Institute of M. From cancer patient to cancer survivor: lost in transition. .
36. Shaw RB, Sweet SN, McBride CB, Adair WK, Martin Ginis KA. Operationalizing the reach, effectiveness, adoption, implementation, maintenance (RE-AIM) framework to evaluate the collective impact of autonomous community programs that promote health and well-being. *BMC Public Health*. 2019; 19(1):803.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

