




# Mathematical Modelling of Temperature Effects on the AFD Neuron of *Caenorhabditis elegans*

Zach Mobbille<sup>1,2</sup>, Rosangela Follmann<sup>1,3</sup>, Andres Vidal-Gadea<sup>4</sup>, Epaminondas Rosa Jr<sup>1,4</sup>,  
Department of Physics<sup>1</sup>, Department of Mathematics<sup>2</sup>, School of Information Technology<sup>3</sup>, School of  
Biological Sciences<sup>4</sup>, Normal, IL 61790


## Introduction

The 1-mm free-living nematode *C. elegans* displays an exceptional sensitivity to temperature fluctuations. Laser ablation studies have shown that the amphid finger-like ciliated AFD neurons in the worm are necessary and sufficient for thermotaxis [1]. We build upon a compartmental model for calcium dynamics in the chemosensitive ASER neuron of the worm [2]. Arrhenius temperature-dependent Q10 coefficients have been added, in the spirit of adapting this model to describe the temperature-sensitive dynamics of AFD. The AFD neuron is depicted schematically above and to the right. We find that the pairing of NaCl concentration fluctuations with temperature stimuli is needed for our model neuron to evoke dynamics similar to those observed experimentally.




## Linear temperature ramp below $T_C$

### Experiment [3]




### Model




$T_C = 20^\circ\text{C}$ . Gray regions in the model plot denote where  $I(t) \neq 0$ , otherwise  $I(t) = 0$

## Linear temperature ramp above $T_C$

### Experiment [4]



### Model



$T_C = 20^\circ\text{C}$ . Gray regions in the model plot denote where  $I(t) \neq 0$ , otherwise  $I(t) = 0$

## AFD Neuron Model

Variables  $c_d$ ,  $c_s$ , and  $c_a$  represent  $\text{Ca}^{2+}$  concentration in the dendrite, soma, and axon, respectively.  $y_i$  ( $i = d, s, a$ ) are inactivation variables for each compartment.

$$\frac{dc_d}{dt} = \frac{\rho}{\tau_d} [-c_d + Y_d y_d + D(W_d c_s - c_d)] + I(t)$$





$$\frac{dc_s}{dt} = \frac{\rho}{\tau_s} [-c_s + Y_s y_s + D(c_d + c_a - c_s)]$$

$$\frac{dc_a}{dt} = \frac{\rho}{\tau_a} [-c_a + Y_a y_a + D(W_a c_s - c_a)]$$

$$\frac{dy_i}{dt} = -\phi A c_i$$

$\rho = \rho_0^{(T-T_C)/10}$  and  $\phi = \phi_0^{(T-T_C)/10}$  are temperature-dependent Arrhenius coefficients for activation and inactivation, respectively.  $T$  = ambient temperature and  $T_C$  = cultivation temperature.  $I(t)$  is a time-dependent binary stimulus to the neuron, representing NaCl concentration.

## $\rho_0$ and $\phi_0$ affect sensitivity of $[\text{Ca}^{2+}]$ waveform to temperature




Dendritic responses with  $T = 30^\circ\text{C}$  &  $T_C = 20^\circ\text{C}$  for various  $\rho_0$  &  $\phi_0$ .


Top: time of dendritic  $[\text{Ca}^{2+}]$  peak for various  $\rho_0$ .  $[\text{Ca}^{2+}]$   
Bottom: time of dendritic  $[\text{Ca}^{2+}]$  peak for various  $\phi_0$ .

## Oscillating temperature ramp below $T_C$

### Experiment [3]



### Model



$T_C = 20^\circ\text{C}$ . Gray regions in the model plot denote where  $I(t) \neq 0$ , otherwise  $I(t) = 0$ .

## References and Acknowledgments

- [1] I. Mori, and Y. Ohshima. *Nature* **376** (1995).
- [2] M. Kuramochi, and M. Doi. *PLoS One* **12** (2017).
- [3] D.A. Clark et al. *J. Neurosci.* **26** (2006).
- [4] A. Yoshida et al. *Genes, Brain and Behavior* **15** (2016).
- [5] Y. Tsukada et al. *J. Neurosci.* **36** (2016).
- [6] A. Vidal-Gadea et al. *eLife* **4** (2015).

## Discussion points

- This phenomenological model can reproduce qualitative dynamics similar to those seen in experiments.
- $I(t)$  is coupled to a temperature threshold of  $T \approx T_C - 3.25^\circ\text{C}$ .
- Model neurons with  $\rho_0 > 3$  may poorly represent the AFD neuron, since it is known to detect temperature differences within a 20-second time window [5].
- $\rho_0$  modulates the  $[\text{Ca}^{2+}]$  peak time, while  $\phi_0$  modulates the rate of decay during a chemical stimulus.