

6-2020

## Efficacy of Social Cognitive Interventions on Improving Social Participation in Children with Autism Spectrum Disorder

T. Book

*Thomas Jefferson University*

A. Ching

*Thomas Jefferson University*

A. Pirilli

*Thomas Jefferson University*

J. Salem

*Thomas Jefferson University*

C. Taylor

*Thomas Jefferson University*

Follow this and additional works at: [https://jdc.jefferson.edu/student\\_papers](https://jdc.jefferson.edu/student_papers)



Part of the [Occupational Therapy Commons](#)

*See next page for additional authors*

[Let us know how access to this document benefits you](#)

---

### Recommended Citation

Book, T.; Ching, A.; Pirilli, A.; Salem, J.; Taylor, C.; Valverde, S.; Ferraro, M.; and Potvin, M. C., "Efficacy of Social Cognitive Interventions on Improving Social Participation in Children with Autism Spectrum Disorder" (2020). *Student Papers & Posters*. Paper 53.

[https://jdc.jefferson.edu/student\\_papers/53](https://jdc.jefferson.edu/student_papers/53)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Student Papers & Posters by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

**Authors**

T. Book, A. Ching, A. Pirilli, J. Salem, C. Taylor, S. Valverde, M. Ferraro, and M. C. Potvin

# Efficacy of Social Cognitive Interventions on Improving Social Participation in Children with Autism Spectrum Disorder

Book, T., Ching, A., Pirilli, A., Salem, J., Taylor, C., Valverde, S.,  
Ferraro, M., PhD, OTR/L & Potvin, M.-C., PhD, OTR/L  
June 2020

## INTRODUCTION

Autism spectrum disorder (ASD) is a developmental disability that affects approximately 1 in 54 U.S. children.<sup>1</sup> Individuals with ASD demonstrate persistent impairments in social communication and interaction, including deficits in the following: social-emotional reciprocity, nonverbal communicative behaviors, and developing, maintaining and understanding relationships.<sup>2</sup> In addition, people with ASD may demonstrate repetitive behaviors and highly restricted interests, which may increase social difficulties.<sup>2</sup> Because social impairment is a defining feature of ASD, numerous strategies have been implemented to improve social skills and thereby, social participation in this population.

The evidence for current interventions addressing social skills with this population is mixed. Some studies of sensory-based interventions have demonstrated secondary effects of improving social interaction among children with ASD.<sup>3</sup> To date, most interventions have emphasized a behavioral approach to social skill development, such as modeling and reinforcement, but limited efficacy and poor generalization has been demonstrated with this approach.<sup>4-7</sup> Crooke et al. argue that this may be in part due to the fact that the majority of treatment approaches fail to address the cognitive aspect of social interactions and relatively few studies have explored the efficacy of treatments based on social cognition.<sup>8</sup>

Cognitive-based interventions are presently used by occupational therapists working with children with developmental coordination disorder, ASD, acquired brain injury, attention deficit hyperactivity disorder, and other conditions, in order to enable fine and gross motor skill development, organizational skills, and activities of daily living completion.<sup>9-12</sup> Social cognitive training programs, developed primarily by mental

health and educational professionals to teach social skills to children with ASD, may have the potential for implementation in the field of occupational therapy to support social participation among this population. A previous systematic review examining a wide variety of occupational therapy interventions for children with ASD included a brief section on social cognitive skill training, concluding that these interventions had modest positive effects on social skill development.<sup>13</sup> To further explore this topic with up-to-date evidence, this current systematic review was conducted to examine the efficacy of social cognitive interventions to improve social participation in children with ASD.

## TERMINOLOGY

**Cognition:** the ability to acquire and use information in order to adapt to environmental demands

**Social cognitive skills:** include recognizing the difference between oneself and others, recognizing others' emotions, collaborating, sharing episodic memory, taking perspectives and experiencing theory of mind, and feeling empathy

**Social cognitive interventions:** include breaking down various cognitive components of social participation and teaching skills related to these components with increasing complexity, skill building, and repetition.<sup>14</sup>

## METHODS

An *a priori* protocol was developed prior to conducting this systematic review to increase its validity. The protocol is a step-by-step outline which includes the PICO question, search strategies for each electronic database, inclusion/

exclusion criteria, and search methodology. The protocol was developed by six collaborating reviewers and followed closely to identify, appraise, and synthesize all relevant published studies. Appendix A includes the PICO question (Table 1 ), a list of the databases searched (Table 2 ).

### **Identification of Relevant Studies:**

A systematic search of all relevant studies was conducted in February and March 2020 using the following databases: CINAHL, ERIC, PsycINFO, and PubMed. All databases were searched manually. Search restriction included quantitative group studies published in English in peer-reviewed journals. Tables 3 and 4 of the protocol provide the search terms (i.e. combination of keywords and subject headings) used to conduct the search within each electronic database. Boolean sentences used for each database are shown in Table 5 .

To be included in this systematic review, studies retrieved during the search had to meet the following criteria: (1) the population must have been children diagnosed with ASD, Asperger's syndrome, or Pervasive Developmental Disorder – Not Otherwise Specified between ages 0-21 years old; (2) the primary means of intervention must be social cognitive skill training; and (3) the outcomes of the study were based on social participation. Table 6 of the protocol provides a complete list of inclusion and exclusion criteria. In order to ensure the third criteria was met in accordance with the treatment definition, outcomes of included articles were listed and evaluated. All outcomes included in this systematic review were further categorized into three primary outcome lists, as shown in Appendix . Studies whose implementers were caregivers, parents, or teachers were excluded to ensure the highest level of intervention fidelity.

Two independent reviewers searched each database and applied the inclusion/exclusion criteria to each study retrieved during the search. Inclusion criteria was first applied to the titles and abstracts of articles. When determination of the inclusion of an article was uncertain, reviewers applied the inclusion criteria to the full text of the article. The flowchart summarizes the results of the search and application of the inclusion and exclusion criteria

( Appendix ). Each independent reviewer created a list of included articles per database, these were compared, and discrepancies were resolved through a consensus process with a third reviewer as needed. A final list of included articles across databases was created after all reviewers came to consensus.

### **Appraisal of Included Studies:**

Through the database search, 1,255 research articles were identified, 38 of which remained after exclusions were identified according to title and abstract. An in-depth review of these 38 articles identified nine articles that met the full inclusion criteria, as shown in the flowchart ( Appendix ). Following protocol, two independent reviewers appraised each article with regard to quality evidence, using predetermined criteria relevant for the study level of evidence ( Appendix ). The two reviewers then compared their independent ratings of the quality of evidence for each study. Any discrepancies were resolved and a consensus was made without the use of a third reviewer. The quality of evidence table ( Appendix ) compiles the quality of methodology rating for each included study.

The two reviewers worked independently to summarize the objective information in each study to create a description table, and again came to a consensus ( Appendix ). The consensus table of the study description includes information regarding the study design type and quality level, the data's population, statistical and clinical significance, intervention, relevant outcomes and measurements, and means/standard deviations ( Appendix ). If there was no measure of clinical significance provided in the article, the minimally detectable difference (MDD) was calculated. Practice recommendations for clinicians were generated via a modified version of the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) process.<sup>15</sup>

## **RESULTS**

Of the nine included studies, seven were quasi-experimental studies that compared results before and after the intervention without use of a control group; one of these studies used two

participant groups and the other six used only one group. The remaining two studies were of the highest level of evidence: randomized controlled trials (RCT) in which subjects were randomly assigned to the treatment and control groups and data was collected before and after the intervention.

### TERMINOLOGY

**Level of Evidence:** hierarchy of studies based on the type of study design<sup>16</sup>

**Quality of Evidence:** degree of rigor used in study methodology<sup>15</sup>

**Effect Size:** degree of difference between two interventions or the size of relationship between variables<sup>16</sup>

**Minimally detectable difference:** the degree of change that must take place to result in an actual difference<sup>16</sup>

As indicated in appendix the nine included studies consisted of level I and level III evidence.<sup>17</sup> Four of the studies provided low quality evidence and five provided moderate quality evidence. The two RCTs were of level I evidence whereas the seven quasi-experimental studies were of level III evidence.

The interventions within the studies were multi-faceted and overlapping, prohibiting their categorization by intervention type. They included guided learning, coaching, scaffolding, role-playing, and mindfulness training. Instead, the studies have been categorized by outcome, according to the basic processes of neurobiology, as described by Adolphs.<sup>18</sup> In order to achieve social participation, a person must first perceive, then process, then act upon information. The included studies primarily measured change in the following three outcomes: (1) social perception, (2) social cognition, and (3) social behavior, all of which are outlined below.

#### ***Social Perception***

Seven out of the nine studies evaluated the impact of the interventions on social perception,

which includes overall social awareness and recognition of emotion, affect, facial expression, and humor. Of these seven studies, all presented with level III evidence. The studies ranged from low to moderate quality of evidence, with three being of moderate quality and four being of low quality. Fourteen measurement data points were collected and analyzed for statistical and clinical significance. Seven were statistically significant and nine were clinically significant. Eight assessments were used to evaluate social perception measures; seven of these were valid and reliable.

#### ***Social Cognition***

All nine studies evaluated the impact of the interventions on social cognition, which included interpersonal problem solving (solutions generation), social attribution, executive function, analogical reasoning, metacognition, flexibility, and planning. Of the nine studies, two presented with level I evidence and seven presented with level III evidence. The studies ranged from low to moderate quality of evidence, with five being of moderate quality and four being of low quality. Fifty-four measurement data points were collected and analyzed for statistical and clinical significance. Thirty-six were shown to be statistically significant and 35 were shown to be clinically significant. Twelve of the 16 assessments used to evaluate social cognition measures were valid and reliable.

#### ***Social Behavior***

Eight out of nine studies evaluated the impact of the intervention on social behavior, such as socialization, maladjusted behavior, interpersonal relationships, play, and behavior regulation. Of these eight studies, two presented with level I evidence, and six presented with level III evidence. The studies ranged from low to moderate quality of evidence, with four being of moderate quality and four being of low quality. Eighteen measurement data points were collected and analyzed for statistical and clinical significance. Fourteen were shown to be statistically significant and 14 were shown to be clinically significant. Five of the six assessments used to evaluate social behavior were valid and reliable.

## SYSTEMATIC REVIEW LIMITATIONS

A large range of interventions may be considered social cognitive interventions, prompting the use of broad search terms. This strategy resulted in a high number of results in PsycNET; due to time limitations, only 500 articles were reviewed. As such, it is possible that not all the relevant evidence was found. In addition, the decision to exclude studies implemented by teachers, parents, or caregivers also limited the scope of this review.

## PRACTICE RECOMMENDATIONS

All outcomes were evaluated using a modified GRADE system, which considered the level of evidence, quality of evidence, degree of clinical significance, and benefit/cost-burden ratio for each outcome.

### **Social Perception**

#### *Weak Recommendation*

As there was a preponderance of Level III studies with positive results measuring this outcome, there is Grade C evidence supporting the use of social-cognitive interventions to improve social perception among children with ASD. The studies measuring this outcome were of low-moderate quality, and they demonstrated low clinical significance and low benefit/cost-burden ratio. Additional research with higher level evidence and higher quality design is needed.

### **Social Cognition**

#### *Weak Recommendation*

With a preponderance of Level III studies, there is grade C evidence supporting the use of social cognitive interventions to improve social cognition among children with ASD. There is a preponderance of studies measuring this outcome that meet the moderate quality criteria, however, the studies show a low degree of clinical significance and low benefit/cost-burden ratio. More rigorous research is needed to determine the estimate of effect.

### **Social Behavior**

#### *Weak Recommendation*

There is grade C evidence supporting the use of social cognitive interventions to improve social

behavior among children with ASD, as there is a preponderance of Level III studies that measure this outcome. Although the studies demonstrated a moderate effect size and moderate benefit/cost-burden ratio, the studies' design and low-moderate quality signifies that higher quality research is needed to determine the estimate of effect.

## CLINICAL IMPLICATIONS

The nine included studies in this systematic review evaluated the efficacy of social cognitive interventions in children with high-functioning autism on three outcomes. Overall, the majority of the studies indicate that there is low quality of evidence and low clinical significance to support the effectiveness of social cognitive interventions to improve social participation in children with ASD, making the potential burden and cost on families exceed the expected amount of benefits. Therefore, when considering social cognitive interventions to address social participation in children with ASD, clinicians should be aware of the limited available evidence and consider seeking alternative interventions. Further and more rigorous research should be conducted in order to determine the efficacy of social cognitive interventions on improving social participation in children with ASD.

## REFERENCES

1. Maenner, M., Shaw, K., & Baio, J. (2020). Prevalence of autism spectrum disorder among children aged 8 years. *MMWR Surveillance Summaries*, 69 (SS-4), 1-12. <https://www.cdc.gov/mmwr/volumes/69/ss/ss6904a1.htm>
  2. American Psychiatric Association (APA). (2013). *Diagnostic and statistical manual of mental disorders*. 5th ed. Arlington, VA: American Psychiatric Association.
  3. Schaaf, R. C., Benevides, T., Mailloux, Z., Faller, P., Hunt, J., Van Hooydonk, E., Freeman, R., Leiby, B., Sendekki, J., & Kelly, D. (2014). An intervention for sensory difficulties in children with autism: A randomized trial. *Journal of Autism and Developmental Disorders*, 44(7), 1493-1506. doi:10.1007/s10803-013-1983-8
-



4. Barry, T. D., Klinger, L. G., Lee, J. M., Palardy, N., Gilmore, T., & Bodin, S. D. (2003). Examining the effectiveness of an outpatient clinic-based social skills group for high-functioning children with autism. *Journal of Autism and Developmental Disorders*, 33(6), 685-701. doi:10.1023/B:JADD.0000006004.86556.e0
  5. Bellini, S., Peters, J., Benner, L., & Hopf, A. (2007). A meta-analysis of school-based social skills interventions for children with autism spectrum disorders. *Remedial and Special Education*, 28, 153-162. doi:10.1177/07419325070280030401
  6. Lee, K. Y. S., Crooke, P. J., Lui, A. L. Y., Kan, P. P. K., Mark, Y., van Hasselt, C. A., & Tong, M. C. F. (2016). The outcome of a social cognitive training for mainstream adolescents with social communication deficits in a Chinese community. *International Journal of Disability, Development and Education*, 63(2), 201-223. doi:10.1080/1034912X.2015.1065960
  7. Williams, S. K., Koenig, K., & Scahill, L. (2007). Social skills development in children with autism spectrum disorders: A review of the intervention research. *Journal of Autism and Developmental Disorders*. 37(10), 1858-1868. doi:10.1007/s10803-006-0320-x
  8. Crooke, P. J., Hendrix, R. E., & Rachman, J. Y. (2008). Brief report: Measuring the effectiveness of teaching social thinking to children with Asperger syndrome (AS) and high functioning autism (HFA). *Journal of Autism and Developmental Disorders*, 38(3), 581-591. doi:10.1007/s10803-007-0466-1
  9. Missiuna, C., DeMatteo, C., Hanna, S., Mandich, A., Law, M., Mahoney, W., & Scott, L. (2010). Exploring the use of cognitive intervention for children with acquired brain injury. *Physical & Occupational Therapy in Pediatrics*, 30(3), 205-219. doi:10.3109/01942631003761554
  10. Rodger, S., Ireland, S., & Vun, M. (2008). Can Cognitive Orientation to daily Occupational Performance (CO-OP) help children with Asperger's syndrome to master social and organisational goals? *British Journal of Occupational Therapy*, 71(1), 23-32. doi:10.1177/030802260807100105
  11. Wright, H. C., & Sugden, D. A. (1998). A school based intervention programme for children with developmental coordination disorder. *European Journal of Physical Education*, 3(1), 35-50. doi:10.1080/1740898980030104
  12. Zwicker, J. G., & Hadwin, A. F. (2009). Cognitive versus multisensory approaches to handwriting intervention: A randomized controlled trial. *OTJR: Occupation, Participation and Health*, 29(1), 40-48. doi:10.1177/153944920902900106
  13. Case-Smith, J., & Arbesman, M. (2008). Evidence-based review of interventions for autism used in or of relevance to occupational therapy. *American Journal of Occupational Therapy*, 62(4), 416-429. doi:10.5014/ajot.62.4.416
  14. Horan, W. P., Kern, R. S., Tripp, C., Helleman, G., Wynn, J. K., Bell, M., Marder, S. R., & Green, M. F. (2011). Efficacy and specificity of social cognitive skills training for outpatients with psychotic disorders. *Journal of Psychiatric Research*, 45(8), 1113-1122. doi:10.1016/j.jpsychires.2011.01.015
  15. Dijkers, M. (2013). Introducing GRADE: a systematic approach to rating evidence in systematic reviews and to guideline development. *KT Update*, 1(5), 1-9. Austin, TX: SEDL, Center of Knowledge Translation for Disability and Rehabilitation Research. [http://www.ktdrr.org/products/update/v1n5/\\_dijkers\\_grade\\_ktupdatev1n5.pdf](http://www.ktdrr.org/products/update/v1n5/_dijkers_grade_ktupdatev1n5.pdf)
-

16. Portney, L. & Watkins, M. (2009). *Foundations of clinical research: Applications to practice (3rd ed.)*. Upper Saddle River, NJ: Pearson.
17. Sackett, D.L., Rosenberg, W.M., Muir Gray, J.A., Haynes, R.B. & Richardson, W.S. (1996). Evidence-based medicine: What it is and what it isn't. *British Medical Journal*, 312, 71-72. <https://doi.org/10.1136/bmj.312.7023.71>
18. Adolphs, R. (2001). The neurobiology of social cognition. *Current opinion in neurobiology*, 11(2), 231-239. doi:10.1016/s0959-4388(00)00202-6
19. Bonete, S., Molinero, C., Mata, S., Calero, M. D., & Gómez-Pérez, M. (2016). Effectiveness of manualised interpersonal problem-solving skills intervention for children with autism spectrum disorder (ASD). *Psicothema*, 28(3), 304-310. doi:10.7334/psicothema2015.206
20. de Bruin, E. I., Blom, R., Smit, F. M. A., van Steensel, Francisca J. A., & Bögels, S. M. (2015). MYmind: Mindfulness training for youngsters with autism spectrum disorders and their parents. *Autism*, 19(8), 906-914. doi:10.1177/1362361314553279
21. Didehbani, N., Tandra, A., Kandalaft, M., Krawczyk, D., & Chapman, S. (2016). Virtual reality social cognition training for children with high-functioning autism. *Computers in Human Behaviors*, 63, 703-711. doi.org/10.1016/j.chb.2016.04.033
22. Gevers, C., Clifford, P., Mager, M., & Boer, F. (2006). Brief report: A theory-of-mind-based social-cognition training program for school-aged children with pervasive developmental disorders: An open study of its effectiveness. *Journal of Autism and Developmental Disorders*, 36(4), 567-571. doi: 10.1007/s10803-006-0095-0
23. Kenworthy, L., Anthony, L.G., & Naiman, D.Q., Cannon, L., Wills, M.C., Luong-Tran, C., Werner, M. A., Alexander, K. C., Strang, J., Bal, E., Sokoloff, J. L., & Wallace, G. L. (2014). Randomized controlled effectiveness trial of executive function intervention for children on the autism spectrum. *Journal of Child Psychology and Psychiatry*, 55(4), 374-383. doi:10.1111/jcpp.12161
24. Soorya, L., Siper, P., Beck, T., Soffes, S., Halpern, D., Gorenstein, M., Kolevzon, A., Buxbaum, J., & Wang, A.T. (2015). Randomized comparative trial of a social cognitive skills group for children with autism spectrum disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 54(3), 208-216.e1. doi:10.1016/j.jaac.2014.12.005
25. Stichter, J. P., Herzog, M., Visovsky, K., Schmidt, C., Randolph, J., Schultz, T., & Gage, N. (2010). Social Competence Intervention for youth with Asperger syndrome and high-functioning autism: An initial investigation. *Journal of Autism and Developmental Disorders*, 40(9), 1067-1079. doi:10.1007/s10803-010-0959-1
26. Stichter, J.P., O'Connor, K., Herzog, M., Lierheimer, K., & McGhee, S. (2012). Social Competence Intervention for elementary students with Aspergers syndrome and high functioning autism. *Journal of Autism and Developmental Disorders*, 42(3), 354-366. doi:10.1007/s10803-011-1249-2

#### ACKNOWLEDGMENT

We would like to thank Daniel Verbit and Theresa Edge for their assistance in developing this systematic review.



**Appendix A**  
**A *Priori* Protocol**

**Table 1A**

*PICO Question*

---

**PICO question - Does cognitive skill training improve social participation in children with ASD?**

---

<b>P - #1</b> children ages 0-21 with Autism Spectrum Disorder	<b>I - #2</b> Cognitive <b>#3</b> Intervention	<b>C -</b> n/a	<b>O -</b> Improved social participation
--	---	----------------	--

---

**Table 2A**

*List of the Databases Searched*

---

Databases Included in SR Search	Planned the Search		Conducted the Search	
	Person 1	Person 2	Person 1	Person 2
CINAHL	Carley	Amanda	Jenna	Anita
PubMed	Jenna	Anita	Steve	Tandi
PsycINFO	Steve	Tandi	Carley	Amanda
ERIC	Carley	Amanda	Tandi	Steve

---

**Table 3A**

*List of Keywords – Same for All Databases*

<b>Facet 1- ASD</b>	<b>Facet 2 - Cognitive</b>	<b>Facet 3 - Intervention</b>
- Asperger* - Autis* - “pervasive developmental disorder”	- metacognit* - cognit* - “problem solving” - coaching - “discovery learning” - “instrumental enrichment”	- training - intervention - strateg* - treatment* - therap* - habilitation - rehabilitation

**Limiters:**

CINAHL – Search keywords within abstracts; Peer review, English, Humans

PubMed – Search keywords within Title/Abstract, Sort by: Best Match, Filters: Humans; English

PsycINFO – search keywords within abstract; Filters: peer-reviewed journal articles, humans

ERIC – Search keywords within abstract; Filters: Peer-reviewed journal articles, English

**Table 4A***List of Subject Headings*

<b>Database</b>	<b>Facet 1- ASD</b>	<b>Facet 2 - Cognitive</b>	<b>Facet 3 - Intervention</b>	<b>Limiters</b>
<b>CINAHL</b>	Autistic disorder, Asperger Syndrome, Pervasive Developmental Disorder-Not Otherwise Specified	Bandura's Social Cognitive Theory	None were found	English, Peer-reviewed
<b>PubMed</b>	Child development disorders, pervasive	Cognitive remediation	None were found	English, Peer-reviewed, Journal articles, Humans; sort results by "best match"
<b>PsycINFO</b>	Autism Spectrum Disorders	Social Cognition	Intervention	Peer-reviewed Journal articles
<b>ERIC via OVID</b>	Autism, Asperger Syndrome, Pervasive Developmental Disorder	Social Cognition, Cognitive Restructuring	Intervention	English, Peer-reviewed, Journal articles

**ERIC** – You must go through OVID and select “ERIC.” Select “Multi-field search,” click the blue triangle next to “Limits” to expand that section and select limits. Search subject headings by using the dropdown “ERIC Subject Headings” and search keywords by using the dropdown “All Fields”

**PsycINFO** – Under “select databases” at the top, select PsycINFO only. To search subject headings, select “APA Thesaurus” in the main dropdown to reach “APA Thesaurus of Psychological Index Terms” page. Click on definition of desired term, and then select term to add to search. Click on “recent searches” drop down. This function provides the user to combine search terms.

**PubMed** – Under the “All Databases” dropdown menu at the top, select MeSH. If you are using the new PubMed website, there are two ways to get to MeSH. First option - click on “Advance” under the search bar. On the next page in “All Fields” dropdown menu, click on “MeSH Terms.” Second option - on the main page, scroll down to the “Explore” icon, then select “MeSH Database” below. Click on “Advance” under search bar. On the next page on the “All Fields” dropdown menu, click on “MeSH Terms.” Type and search your subject heading. Then click “Add to search builder” and “Search PubMed” on the right. For both the old and new PubMed website, search keywords by using the dropdown “All Fields.”

**CINAHL** – Use Advanced Search to find limits. Using the dropdowns, choose “MH Exact Match Subject Heading” to search subject headings and “AB Abstract” to search keywords

**Table 5A**

*Boolean Sentence for Each Database*

<b>Database Name</b>	<b>Boolean Sentence</b>
<b>CINAHL</b>	( <b>Autistic disorder</b> OR <b>Asperger Syndrome</b> OR <b>Pervasive Developmental Disorder-Not Otherwise Specified</b> OR Asperger* OR Autis* OR “pervasive developmental disorder”) AND ( <b>Bandura's Social Cognitive Theory</b> OR metacognit* OR cognit* OR “problem solving” OR coaching OR “discovery learning” OR “instrumental enrichment”) AND (training OR intervention OR strateg* OR treatment* OR therap* OR habilitation OR rehabilitation)
<b>PubMed</b>	( <b>Child development disorders, pervasive</b> OR Asperger* OR Autis* OR “pervasive developmental disorder”) AND ( <b>Cognitive remediation</b> OR metacognit* OR cognit* OR “problem solving” OR coaching OR “discovery learning” OR “instrumental enrichment”) AND (training OR intervention OR strateg* OR treatment* OR therap* OR habilitation OR rehabilitation)
<b>PsycINFO</b>	( <b>Autism Spectrum Disorders</b> OR Autis* OR Asperger* OR “pervasive developmental disorder”) AND ( <b>Social Cognition</b> OR cognit* OR metacognit* OR “problem solving” OR coaching OR “discovery learning” OR “instrumental enrichment”) AND ( <b>Intervention</b> OR training OR intervention OR strateg* OR treatment* OR therap* OR habilitation OR rehabilitation)
<b>ERIC</b>	( <b>Autism</b> OR <b>Asperger Syndrome</b> OR <b>Pervasive Developmental Disorder</b> OR Asperger* OR Autis* OR “pervasive developmental disorder”) AND ( <b>Social Cognition</b> OR <b>Cognitive Restructuring</b> OR metacognit* OR cognit* OR “problem solving” OR coaching OR “discovery learning” OR “instrumental enrichment”) AND ( <b>Intervention</b> OR training OR intervention OR strateg* OR treatment* OR therap* OR habilitation OR rehabilitation)


**\*Bolded terms = subject headings**

**Table 6A***Inclusion and Exclusion Criteria*

<b>Inclusion Criteria</b>			
<b>Population</b>	<b>Intervention and Comparison</b>	<b>Outcome</b>	<b>Other</b>
Autism Spectrum Disorder	Coaching	Social Participation, as defined by at least one of the following: Increased communication with others Increased attendance in social activities Increased prosocial behaviors, including: maintaining conversation eye contact turn taking orienting body to face peer initiating and ending conversations interpreting social cues and responding effectively maintaining socially accepted (i.e. arm's length) distance from others using effective approaches to join in activities generating or reciprocating effective facial expressions generating or reciprocating effective body language or gestures	All types of Quantitative intervention studies  Peer-reviewed
Asperger Syndrome	Scaffolding		
PPD-NOS	Priming		
All levels of Autism	Discovery Learning		
Any gender	Role playing		
Ages 0-21	Instrumental Enrichment		
	Social problem-solving		
	Cognitive orientation to (daily) occupational performance		
	Self-management		
	Social-cognitive training		
	Metacognitive strategies		
<b>Exclusion Criteria</b>			
<b>Population</b>	<b>Intervention and Comparison</b>	<b>Outcome</b>	<b>Other</b>
Caregiver/ Parent/ Teacher training	Social stories		Non-English language
Caregiver/ Parent/ Teacher implemented/mediated	Social skills training without a cognitive component		Non-human
	Video modeling		
	Cognitive Behavioral Therapy		
	Early Start Denver Model		

**Appendix B  
Flowchart**

**Figure 1**  
*Flowchart*





**Appendix C**  
***Outcome Designations***

---

**Outcomes**

---

**Social Perception**

Social awareness  
Affect Recognition  
Facial Expression  
Emotion recognition  
Perception  
Humor  
Human relatedness

---

**Social Cognition**

Interpersonal problem-solving ability  
Social cognition, abilities, attribution  
Theory of mind  
Executive function and metacognition  
Analogical reasoning  
Flexibility and planning  
Making inferences and problem-solving  
Listening skills  
First and second order beliefs  
Understanding humor

---

**Social Behavior**

Socialization  
Maladjusted behavior  
Play  
Interpersonal relationships  
Social communication  
Behavior regulation  
Social thinking

---

**Appendix D**  
**Quality and Level of Evidence Table**

Citation	Type of design	Quality Criteria										Quality Level	Evidence Level
		1	2	3	4	5	6	7	8	9	10		
Bonete et al., 2016 <sup>19</sup>	#6 Quasi-experimental (2 groups pre/post)	0	1	0	0	1	1	1	1	N/A	N/A	Moderate	Level III
de Bruin et al., 2015 <sup>20</sup>	#6 Quasi-experimental (1 group pre/post + f/u)	1	1	0	0	1	1	0	0	N/A	N/A	Moderate	Level III
Didehbani et al., 2016 <sup>21</sup>	#6 Quasi-experimental (1 group pre/post)	1	1	0	0	1	0	0	1	N/A	N/A	Moderate	Level III
Gevers et al., 2016 <sup>22</sup>	#6 Quasi-experimental (1 group pre/post)	0	1	0	0	1	0	0	0	N/A	N/A	Low	Level III
Kenworthy et al., 2014 <sup>23</sup>	#3 RCT	0	0	1	1	1	0	0	0	0	1	Moderate	Level I
Lee et al., 2016 <sup>6</sup>	#6 Quasi-experimental (1 group pre/post)	1	1	0	0	0	0	1	0	N/A	N/A	Low	Level III
Soorya et al., 2015 <sup>24</sup>	#3 RCT	1	1	1	1	0	0	0	1	0	1	Moderate	Level I
Stichter et al., 2010 <sup>25</sup>	#6 Quasi-experimental (1 group pre/post)	1	1	0	0	0	1	0	0	N/A	N/A	Low	Level III
Stichter et al., 2012 <sup>26</sup>	#6 Quasi-experimental (1 group pre/post)	1	1	0	0	0	0	0	0	N/A	N/A	Low	Level III

**Appendix E**  
**Study Description: Included Studies**

Study	Design Type, # Criteria, Quality	Population n per group Treatment/ Control	Outcome(s)	Measurement (include units)	Means and (Standard Distributions)	Statistical significance * p≤.05, **p≤.001	Clinical significance * denotes significance
Bonete et al., 2016 <sup>19</sup>	6 – Quasi 2 group pre/post 4/8 Moderate	<u>Study 1</u> n=22 Age: 7-12 Sex: 21M 1F Dx: Aspergers  <u>Study 2</u> n=15 Age: 7-12 Sex: 12M 3F Dx: Aspergers  Interpersonal Problem-Solving Skills Programme for Children  No comparison	1. interpersonal relations play/leisure time coping 2. internalized and externalized maladjusted bx 3. emotion recognition 4. causes attribution (for interpersonal problems) 5. solutions generation (for interpersonal problems) 6. interpersonal problem-solving ability (total)	1. <b>VABS - Socialisation subscale</b> 0-124; >= improved socialization ‡ 2. <b>VABS - Maladaptive Behavior subscale</b> 0-44; < = less maladaptive behavior ‡ 3. <b>ESCI : Emotion recognition subscale (ESCI-E) †</b> 4. <b>ESCI: Causes attribution subscale (ESCI-C)</b> 5. <b>ESCI: Solutions subscale (ESCI-S)</b> 6. <b>ESCI: Total (ESCI-T)</b>  ‡ Based on VABS-III; VABS-II not available † Could not find scoring information	1. <b>S1</b> Pre:128.09 (10.35) Post: 133.41 (11.35) <b>S2</b> Pre: 119.07 (11.37) Post: 126.33 (12.86) 2. <b>S1</b> Pre: 24.22 (8.22) Post: 18.00 (8.03) <b>S2</b> Pre: 29.33(12.83) Post: 24.73 (9.81) 3. <b>S1</b> Pre: 12.81 (2.70) Post: 12.95 (2.77) <b>S2</b> Pre: 13.40 (2.53) Post: 14.07 (1.87) 4. <b>S1</b> Pre: 36.55 (6.79) Post: 37.09 (6.80) <b>S2</b> Pre: 35.20 (7.61) Post: 37.40 (5.88) 5. <b>S1</b> Pre: 8.77 (3.43) Post: 9.81 (2.72) <b>S2</b> Pre: 9.07 (1.83) Post: 9.40 (1.55) 6. <b>S1</b> Pre: 58.14 (11.53) Post: 59.86 (10.27) <b>S2</b> Pre: 58.00 (8.85) Post: 60.87 (8.31)	1. <b>S1</b> p=.04* <b>S2</b> p=.01* 2. <b>S1</b> p=.002* <b>S2</b> p=.04* 3. <b>S1</b> p=.88 <b>S2</b> p=.88 4. <b>S1</b> p=.22 <b>S2</b> p=.16 5. <b>S1</b> p=.02* <b>S2</b> p=.56 6. <b>S1</b> p=.04* <b>S2</b> p=.09	1. <b>S1</b> r=.30* <b>S2</b> r=.46* 2. <b>S1</b> r=.47* <b>S2</b> r=.37* 3. <b>S1</b> r=.02 <b>S2</b> r=.03 4. <b>S1</b> r=.18* <b>S2</b> r=.26* 5. <b>S1</b> r=.34* <b>S2</b> r=.11* 6. <b>S1</b> r=.30* <b>S2</b> r=.31*
de Bruin et al., 2015 <sup>20</sup>	6 – Quasi 1 group pre/post w/ f/u 4/8 Moderate	n=23, Age: 11-23 Sex: 17 M 6F Dx: Aspergers, ASD, or PDD-NOS  <u>MyMind</u> : Mindfulness training for Youth with ASD No comparison	1. Social abilities	6a. <b>SRS Total</b> (> = poorer) 6b. <b>SRS Social Awareness</b> 6c. <b>SRS Social Cognition</b> 6d. <b>SRS Social Communication</b> 6e. <b>SRS Social Motivation</b>	6a. Post: 0.01, f/u: -0.33 6b. Post: -0.02, f/u: -0.14 6c. Post: 0.10, f/u: -0.17 6d. Post: -0.07, f/u: -0.40 6e. Post: -0.08, f/u: -0.23	6a. Post: N.G. f/u: p<.001** 6b. Post: N.G. f/u: N.G. 6c. Post: p<.10 f/u: p<.01* 6d. Post: N.G. f/u: p<.001** 6e. Post: N.G. f/u: p<.10	6a. Post: d= .01 f/u: d= -.33* 6b. Post: d= -.02 f/u: d= -.14 6c. Post: d= .10 f/u: d= -.17 6d. Post: d= -.07 f/u: d= -.40* 6e. Post: d= -.08 f/u: d= -.23*
Didehbani et al., 2016 <sup>21</sup>	6 – Quasi 1 group pre/ post 4/8 Moderate	n=30, Age: 7-16 Sex: 26M 4F Dx: n=17 ASD n=13 ADHD+ASD Other: average or higher IQ  Real-time feedback from the “coach” clinician. No comparison	1. Affect recognition (AR) 2. Social attribution/ theory of mind 3. Executive function 4. Analogical Reasoning	1a. <b>NEPSY-II Affect Recognition</b> : 1-19, >=better AR 1b. <b>Ekman60</b> : 0-60, > = better AR 2a. <b>Triangles</b> Total: 0-36, >=better social attribution 2b. <b>Intentionality</b> 3a. <b>NEPSY-II Auditory Attention (AA)</b> 3b. <b>NEPSY-II Response Test (RT)</b> 1-19, >=better executive function 4. <b>Fluid Reasoning - Analogical Reasoning Task (ART)</b> 0-24, >=better executive function	1a. Pre 8.9 (2.6), Post 10.4 (2.1) 1b. Pre 38.9 (6.6), Post 40.8 (5.8) 2a. Pre 18.5 (3.1) 19.6, Post (3.2) 2b. Pre 11.5 (3.3), Post 13.3 (3.5) 3a. Pre 7.9 (4.6) Post 8.7 (4.2) 3b. Pre 8.3 (2.6) Post 9.5 (2.9) 4. Pre 81.2 (11.9), Post 85.7 (11.1)	1a. p=.001** 1b. p=.046* 2a. p=.033* 2b. p=.016* 3a. p=.248 3b. p=.132 4. p=.016*	1a. 1.5>1.3* 1b. 1.9<3.3 2a. 1.1<1.55 2b. 1.8>1.65* 3a. 0.8<2.3 3b. 1.2<1.3 4. 4.5<5.95 (MDD)

Gevers et al., 2006 <sup>22</sup>	6 – Quasi 1 group pre/ post  2/10 Low	n=18 <b>Age:</b> 8-11 <b>Sex:</b> 13M 5F <b>Dx:</b> PDD-NOS <b>Other:</b> verbal IQ 85  Theory of Mind training  No comparison	1a. perception & imitation; pre- tense; recognition of emotions; distinction physical-mental <b>1b.</b> first order belief, understanding false belief <b>1c.</b> second order belief, understanding humor <b>2.</b> Socialization - Interpersonal relationships, Play/leisure and Social skills	<b>1. TOM Test</b> - (0-72, >=better TOM) <b>1a. TOM 1, 1b. TOM 2, 1c. TOM 3, 1d. Total TOM</b>  <b>2. VABS - Socialisation</b> <b>2a. Interpersonal relationships</b> <b>2b. Play/Leisure</b> <b>2c. Social Skills</b>	<b>1a.</b> Pre 18.4 (2.4), Post 20.9 (1.2) <b>1b.</b> Pre 27.0 (4.8), Post 32.3 (3.0) <b>1c.</b> Pre 7.7 (2.4), Post 9.5 (1.9) <b>1d.</b> Pre 52.8 (7.8), Post 62.7 (5.2)  <b>2a.</b> Pre 0.36 (0.09), Post 0.42 (0.14) <b>2b.</b> Pre 0.39 (0.10), Post 0.47 (0.10) <b>2c.</b> Pre 0.51 (0.13), Post 0.61 (0.15)	<b>1a.</b> p=.000** <b>1b.</b> p=.001** <b>1c.</b> p=.001** <b>1d.</b> p=.000**  <b>2a.</b> p=.021* <b>2b.</b> p=.013* <b>2c.</b> p=0.000**	<b>1a.</b> 2.5>1.2* <b>1b.</b> 5.3>2.4* <b>1c.</b> 1.8>1.2* <b>1d.</b> 9.9>3.9*  <b>2a.</b> 0.06>0.045* <b>2b.</b> 0.08>0.05* <b>2c.</b> 0.1>0.065*  (MDD)
Kenworthy et al., 2014 <sup>23</sup>	3 – RTC  4/10, Moderate	n T=47 (10 schools) n C=20 (4 schools) <b>Age:</b> 7-11 <b>Sex:</b> all male <b>Dx:</b> PDD, ASD  Unstuck and On Target (UOT)  Comparison: Social Skills (SS)	<b>1.</b> Problem-solving  <b>2.</b> Flexibility, Planning, & Social Appropriateness  <b>3.</b> Executive Functioning Shift, Plan/Organize  <b>4.</b> Executive Functioning Shift, Plan/Organize  <b>5.</b> ASD-related social, comm- unication, and repetitive bx  <b>6.</b> ASD-related social, comm- unication, and repetitive bx	<b>1. WASI block design</b> (> = better performance)  <b>2. Challenge Task</b> (> = more impairment) <b>2a. Flexibility, 2b. Plan, 2c. Social</b>  <b>3. BRIEF - Teacher Rated</b> (> = more impaired) <b>3a. EF Shift,</b> <b>3b. Plan/Organize</b>  <b>4. BRIEF - Parent Rated</b> <b>4a. EF Shift,</b> <b>4b. Plan/Organize</b>  <b>5. SRS - Teacher Rated</b>  <b>6. SRS - Parent Rated</b>	<b>1.</b> UOT: 3.00 (1.03), SS: -0.94 (1.11)  <b>2a.</b> UOT: -0.53 (0.07), SS: -0.15 (0.14) <b>2b.</b> UOT: -0.33 (0.07), SS: -0.22 (0.06) <b>2c.</b> UOT: 0.47 (0.16), SS: 0.26 (0.30)  <b>3a.</b> UOT: -24.00 (3.30), SS: -9.78 (3.59) <b>3b.</b> UOT: -19.14 (2.39) SS: -11.72 (3.16)  <b>4a.</b> Shift UOT: -9.56 (2.31), SS: -0.16 (2.99) <b>4b.</b> Plan/Org UOT: -5.17 (2.0), SS: 0.61 (2.90)  <b>5. SRS Teacher Rated (TR):</b> UOT: -5.4 (1.34) SS: -4.79 (2.05)  <b>6. SRS Parent Rated (PR):</b> UOT: -7.31 (1.65), SS: -4.11 (2.97)	<b>1.</b> p<.05*  <b>2a.</b> p<.05* <b>2b.</b> N.G. <b>2c.</b> N.G  <b>3a.</b> p<.01* <b>3b.</b> p<.05*  <b>4a.</b> p<.01* <b>4b.</b> p<.05*  <b>5.</b> N.G.  <b>6.</b> N.G.	<b>1.</b> d= 0.65*  <b>2a.</b> d= -0.72* <b>2b.</b> d= -0.27* <b>2c.</b> d= 0.17*  <b>3a.</b> d= -0.89* <b>3b.</b> d= -0.57*  <b>4a.</b> d= -0.66* <b>4b.</b> d= -0.45*  <b>5.</b> d = -0.08  <b>6.</b> d = -0.28*
Lee et al., 2016 <sup>6</sup>	6 – Quasi 1 group pre/post  3/8 Low	n=39 <b>Age:</b> 12 -15 yo <b>Sex:</b> 30M 9F <b>Dx:</b> 33 with ASD 6 without ASD (all had social communication impairments)  Social Thinking Training  No comparison	<b>1.</b> Social thinking - Overall  <b>2.</b> Initiation  <b>3.</b> Listening with Eyes/Brain  <b>4.</b> Abstract & Inferential Language <b>5.</b> Understanding Perspective  <b>6.</b> Gestalt Processing  <b>7.</b> Humor and Human Relatedness	<b>1.Social Thinking ILAUGH Rating Scale</b>  <b>2. Initiation Subscale</b>  <b>3. Listening with Eyes/Brain Subscale</b>  <b>4. Abstract and Inferential Language Subscale</b>  <b>5. Understanding Perspective Subscale</b>  <b>6. Gestalt Processing Subscale</b>  <b>7. Humor and Human Relatedness</b> All scores 1-5, >= greater frequency of prosocial bx)	<b>1.</b> pre m =2.74 (.56), post m= 3.14 (.48)  <b>2.</b> pre m = 2.87 (.85), post m = 3.28 (.77)  <b>3.</b> pre m= 2.70 (.69), post m = 3.21 (.63)  <b>4.</b> pre m= 2.68 (.76), post m = 3.07 (.69)  <b>5.</b> pre m = 2.72 (.66), post m = 3.15 (.57)  <b>6.</b> pre m = 2.70 (.69), post m = 3.06 (.54)  <b>7.</b> pre m = 3.14 (.67), post m = 3.24 (.62)	<b>1.</b> p<.001**  <b>2.</b> p<.001**  <b>3.</b> p<.001**  <b>4.</b> p<.001**  <b>5.</b> p<.001**  <b>6.</b> p<.001**  <b>7.</b> p=.217	<b>1.</b> d = .50*  <b>2.</b> d = .72*  <b>3.</b> d = .52*  <b>4.</b> d = .67*  <b>5.</b> d = .57*  <b>6.</b> d = .14  <b>7.</b> d = .72*

Soorya et al., 2015 <sup>24</sup>	3 – RCT 6/10 Moderate	<p><b>n</b> T=35, <b>n</b> C=34  <b>Age:</b> 8-11  <b>Sex:</b> 19M 1 F  <b>Dx:</b> ASD and a verbal IQ of 70</p> <p>Treatment (T):  Seaver-NETT (Nonverbal communication, Emotion recognition, and Theory of mind Training)</p> <p>Comparison (C):  Child-directed play “stations”. Use of reflective statements to foster communication.</p>	<p>1. Social cognition</p> <p>2. Social behavior</p>	<p><b>1a. Social Cognition Composite</b></p> <p><b>1b. DANVA2</b> (&gt; = greater ability)</p> <p><b>1c. Strange Stories Task (SST)</b>, 0-12, &gt;=more social cognition</p> <p><b>1d. Reading the Mind in the Eyes Test (RMET)</b> 5-25, (&gt; = greater ability to interpret mental states from facial cues)</p> <p><b>2a. Social Behavior Composite</b>, &lt;=better social behavior</p> <p><b>2b. Griffith Empathy Measure (GEM)</b>, 65-185, &gt;=better social behavior</p> <p><b>2c. SRS</b></p> <p><b>2d. Children’s Communication Checklist–2 (CCC)</b>, 0-18, &gt;=better social behavior</p>	<p><b>1a. Tx</b> Pre: 0.15 (0.84), Post: 0.27 (0.92)  <b>C</b> Pre: -.14 (0.78), Post: -.02 (0.76)</p> <p><b>1b. Tx</b> Pre: 84.53 (11.89), Post: 87.58 (13.51)  <b>C</b> Pre: 80.08 (12.74), Post: 81.97 (11.88)</p> <p><b>1c. Tx</b> Pre: 6.00 (3.08), Post: 6.06 (3.05)  <b>C</b> Pre: 5.17 (2.76), Post: 5.76 (2.37)</p> <p><b>1d. Tx</b> Pre: 17.03 (3.88), Post: 17.58 (4.44)  <b>C</b> Pre: 16.08 (4.6), Post: 16.03 (4.58)</p> <p><b>2a. Tx</b> Pre: 0.04 (1.00), Post: 0.34 (0.64)  <b>C</b> Pre: 0.05 (0.70), Post: -.01 (0.73)</p> <p><b>2b. Tx</b>Pre: 129.67 (25.81), Post: 132.72 (24.31)  <b>C</b> Pre: 123.62 (19.28), Post: 123.96 (20.55)</p> <p><b>2c. Tx</b> Pre: 9.34 (4.55), Post: 7.69 (3.06)  <b>C</b> Pre: 9.58 (3.02), Post: 9.22 (3.58)</p> <p><b>2d. Tx</b> Pre: 10.03 (4.78), Post: 8.88 (3.84)  <b>C</b> Pre: 10.07 (3.54), Post: 9.96 (3.46)</p>	<p>For all measures  p &lt;.05</p> <p>NETT Social Bx  p = .04</p> <p>No significant effect on the social cognition composite</p> <p>No significant interaction effect at follow-up on social behavior (p=.38) or social cognition composites (p=.79).</p>	<p><b>1a. d</b> = 0.56</p> <p><b>2a. d</b> =0.88</p> <p>Cohen’s d n.g.for other assessments</p>
Stichter et al., 2010 <sup>25</sup>	6 – Quasi 1 group pre/post  3/8 Low	<p><b>n</b>=27  <b>Age:</b> 10.83–14.75  <b>Sex:</b> all male  <b>Dx:</b> ASD, PDD-NOS, Aspergers</p> <p>Social Competence Intervention – Adolescent (SCI-A)</p> <p>No comparison</p>	<p>1. Social abilities</p> <p>2. Executive functioning, behavioral regulation, metacognition</p> <p>3. Executive functioning, making inferences, and problem-solving</p> <p>4. Facial expression recognition</p> <p>5. Facial expression and emotions recognition</p> <p>6. Theory of mind</p>	<p><b>1a. SRS Total</b></p> <p><b>1b. Social awareness (SA)</b></p> <p><b>1c. Social Cognition (SCog)</b></p> <p><b>1d. Social communication (SComm)</b></p> <p><b>1e. Social Motivation (SM)</b></p> <p><b>1f. Autistic mannerisms (AM)</b></p> <p><b>2a. BRIEF Global Executive (GE)</b></p> <p><b>2b. Behavioral Regulation (BR)</b></p> <p><b>2c. Metacognition (MC)</b></p> <p><b>3a. TOPS total</b> (&gt; = stronger abilities)</p> <p><b>3b. Making Inferences (MI)</b></p> <p><b>3c. Problem Solving (PS)</b></p> <p><b>4. DANVA</b></p> <p><b>5. RMET</b></p> <p><b>6. Faux Pas Stories (FPS):</b> (10 short narratives) (&gt; = greater accuracy of faux pas identification)</p>	<p><b>1a.</b> Pre: 107.4 (18.5), Post: 85.4 (16.7)</p> <p><b>1b.</b> Pre: 13.0 (3.0), Post: 11.2 (3.1)</p> <p><b>1c.</b> Pre: 20.1 (3.4), Post: 15.7 (3.7)</p> <p><b>1d.</b> Pre: 35.4 (7.9), Post: 29.1 (7.5)</p> <p><b>1e.</b> Pre: 16.2 (5.6), Post: 12.9 (3.4)</p> <p><b>1f.</b> Pre: 21.6 (5.6), Post: 16.4 (5.1)</p> <p><b>2a.</b> Pre: 72.9 (8.2), Post: 67.1 (8.0)</p> <p><b>2b.</b> Pre: 76.6 (8.7), Post: 68.7 (8.4)</p> <p><b>2c.</b> Pre: 69.4 (11.4), Post: 64.0 (8.9)</p> <p><b>3a.</b> Pre: 89.4 (16.8), Post: 95.8 (15.2)</p> <p><b>3b.</b> Pre: 90.0 (18.4), Post: 95.8 (16.4)</p> <p><b>3c.</b> Pre: 90.7 (18.7), Post: 96.6 (15.2)</p> <p><b>4.</b> Pre: 19.5 (2.2), Post: 20.7 (1.8)</p> <p><b>5.</b> Pre: 17.3 (3.4), Post: 18.5 (3.8)</p> <p><b>6.</b> Pre: 8.5 (1.5), Post: 9.0 (1.1)</p>	<p><b>1a.</b> p &lt;.001**</p> <p><b>1b.</b> p &lt;.01*</p> <p><b>1c.</b> p &lt;.001**</p> <p><b>1d.</b> p &lt;.001**</p> <p><b>1e.</b> p &lt;.01*</p> <p><b>1f.</b> p &lt;.001*</p> <p><b>2a.</b> p &lt;.001**</p> <p><b>2b.</b> p &lt;.001**</p> <p><b>2c.</b> p &lt;.001**</p> <p><b>3a.</b> p &lt;.001**</p> <p><b>3b.</b> p &lt;.05*</p> <p><b>3c.</b> p &lt;.05*</p> <p><b>4.</b> p &lt;.05*</p> <p><b>5.</b> p &lt;.05*</p> <p><b>6.</b> p &lt;.05*</p>	<p><b>1a.</b> SRS Total</p> <p><b>1b.</b> SA*</p> <p><b>1c.</b> SCog*</p> <p><b>1d.</b> SCom*</p> <p><b>1e.</b> SM*</p> <p><b>1f.</b> AM*</p> <p><b>2a.</b> BRIEF GE*</p> <p><b>2b.</b> BR*</p> <p><b>2c.</b> MC</p> <p><b>3a.</b> TOPS Total</p> <p><b>3b.</b> MI</p> <p><b>3c.</b> PS</p> <p><b>4.</b> DANVA*</p> <p><b>5.</b> RME</p> <p><b>6.</b> FPS</p> <p>MDD Calculated Using SD</p>

