



# CYTOTAXONOMICAL STUDIES ON THE *CALTHA PALUSTRIS* COMPLEX (RANUNCULACEAE) IN POLAND. PRELIMINARY REPORT

ELŻBIETA CIEŚLAK<sup>1</sup>, TOMASZ ILNICKI<sup>2</sup> AND MAŁGORZATA FLIS<sup>2\*</sup>

<sup>1</sup>W. Szafer Institute of Botany, Polish Academy of Sciences, ul. Lubicz 46, 31–512 Cracow, Poland

<sup>2</sup>Department of Plant Cytology and Embryology, Jagiellonian University,  
ul. Grodzka 52, 31–044 Cracow, Poland

Received February 2, 2000; revision accepted April 25, 2000

The karyology of representatives of morphologically differentiated taxa within the *Caltha palustris* complex was studied. Examination of plants from ten populations of *C. palustris* subsp. *palustris* from Poland indicated the domination of somatic chromosome numbers  $2n = 32$  and  $56$ , with  $2n = 32$  and  $56$  for var. *palustris*,  $2n = 56$  for var. *radicans* and  $2n = 56$  for var. *cornuta*. *C. palustris* subsp. *laeta* from the Tatra Mts. had chromosome number  $2n = 62$ . Mixoploidy was characteristic of the material, with a range of euploid and aneuploid chromosome numbers from  $2n = 25$  to  $2n = 94$ . Only three plants (from Mrzeżyno, Duninowo and Olkusz) uniformly had  $2n = 32$ . Detailed biometric analyses seem to point to the lack of a simple relation between karyological and morphological variability in representatives of the *C. palustris* complex occurring in Poland.

**Key words:** *Caltha palustris* L. s.l., cytological variability, chromosome numbers, morphological variability, Poland.

## INTRODUCTION


The genus *Caltha* L., with two allopatric subgenera *Caltha* and *Psychrophila* (Gay) Prantl, consists of about 40 species (Ziemmerman, 1965). The Linnaean species *Caltha palustris*, which belongs to the subgenus *Caltha*, is a circumboreal taxon, common in Poland in both the lowlands and the mountains, where it reaches up to the alpine belt. It grows in wet and marshy areas.

The great differentiation of morphological features (habit, shape of leaves and follicles) led to the division of Linnaean species *C. palustris* into independent species and/or subspecific taxa. Schott et al. (1854) distinguished six species within the complex: *C. cornuta*, *C. latifolia*, *C. laeta*, *C. intermedia*, *C. vulgaris* and *C. alpestris*. This classification was later modified several times. On the basis of follicle

polymorphism Hegi (1912) distinguished three subspecies within *C. palustris* s.l.: subsp. *genuina*, subsp. *cornuta* (Schott, Nyman et Kotschy) Beck, and subsp. *laeta* (Schott, Nyman et Kotschy) Hegi. Taking into account some extra features such as chromosome number, habitat and phytocoenoses, Ziemmermann (1965) divided the *C. palustris* complex into four subspecies, adding subsp. *minor* (Mill.) Graebner to those of Hegi (1912). In both editions of *Flora Europea* (Tutin, 1964; Akeroyd, 1993) the wide embrace of the Linnaean *C. palustris* is accepted without any infraspecific taxa distinguished, but the enormous morphological and cytological variability of the species is stressed.

The range of variability and the systematics of the *C. palustris* complex in Poland has not been clarified yet. Paczoski (1927) divided it after Schott et al. (1854) into three species (i.e. *C. cor-*

\*e-mail: flisma@grodzki.phils.uj.edu.pl



**Fig. 1.** Distribution of 74 examined samples of *Caltha palustris* s.l. in Poland. ● *C. palustris* subsp. *laeta*; ◆ *C. palustris* subsp. *palustris* var. *palustris*; ◇ *C. palustris* subsp. *palustris* var. *radicans*; ▲ *C. palustris* subsp. *palustris* var. *cornuta*. The specimens from which samples were taken for cytological tests are marked with larger symbols; the same populations are in Table 2; somatic chromosome numbers ( $2n$ ) in parentheses.

*nuta* Schott, Nyman et Kotschy; *C. laeta* Schott, Nyman et Kotschy; *C. palustris* L.). Kucowa (1985) distinguished two species within the Linnaean *C. palustris* (*C. palustris* L. s.str. and *C. laeta* Schott, Nyman et Kotschy). She also recognised two subspecies within *C. palustris* – subsp. *palustris* and subsp. *cornuta* (Schott, Nyman et Kotschy) Hegi – and two varieties within *C. laeta*, that is, var. *laeta* and var. *alpestris* (Schott, Nyman et Kotschy) Beck. Recently, Mirek et al. (1995) accepted Ziemmerman's (1965) division of the *C. palustris* complex into three subspecies: subsp. *laeta* (Schott, Nyman et Kotschy) Hegi, subsp. *palustris* and subsp. *cornuta* (Schott, Nyman et Kotschy) Hegi. The division of the complex accepted in the present paper distinguishes three varieties of *C. palustris* L. subsp. *palustris*: var. *palustris*, var. *radicans* (Forster) Beck, var. *cornuta* (Schott, Nyman et Kotschy) Borbás and *C. palustris* subsp. *laeta* (Schott, Nyman et Kotschy) Hegi.

The great morphological variability of *C. palustris* s.l. is accompanied by karyological differentiation. Different euploid and aneuploid chromosome numbers have been noted in ploidy levels ranging from  $2n = 2x = 16$  to  $2n = 11x = 88$ . The results of hitherto known karyological investigations are presented in Table 1.

The present work tests the applicability of karyological investigations in solving the taxonomic problems that arose during research on the morphological variability of this complex.

## MATERIALS AND METHODS

### MORPHOLOGY

For morphometric analysis of the *C. palustris* complex, 74 population samples of 25–30 plants each were collected from the whole area of Poland; all the main geographical regions of the country were rep-

TABLE 1. Previous chromosome reports for the *Caltha palustris* complex

Provenience of the material	Chromosome numbers 2n	Author
<b>NORTH AMERICA</b>		
Canada	32, ca 32, 56	Wcislo, 1967; Kootin-Sanwu and Woodell, 1970
U.S.A.	32, 56, 60, ca 64	Wcislo, 1967
<b>EUROPE</b>		
Austria	32, 52, 56, 57, 58, ca 58, 59, ca 59, 60, ca 60, 61, ca 61, 64	Wcislo, 1967; Smit, 1967; Reese, 1954; Kootin-Sanwu and Woodell, 1970
Czech Republic	32, 34, 35, 36, 44, 47, 55, 56, ca 56, 57, 58, 59, 61, 64	Chrtkova and Jarolimova, 1999
Denmark	56	Reese, 1954
France	32, ca 32, 48, 56	Reese, 1954; Smit, 1967; Kootin-Sanwu and Woodell, 1970
Germany	16, 28, 29, 30, 32, ca 32, 55, 56, ca 56, 57, ca 57, 58, ca 58, 59, 60, ca 60, ca 61	Engler, 1936; Reese, 1954; Wcislo, 1967; Smit, 1967; Kootin-Sanwu and Woodell, 1970; Goldblatt, 1981
Great Britain	52, 54, 55, 56, 57, 58, 60, 64, ca 64, 72, ca 80	Löve and Löve, 1948; Reese, 1954; Wcislo, 1967; Kootin-Sanwu and Woodell, 1970
Greece	32	Goldblatt, 1984
Hungary	32	Polya, 1950
Iceland	48, ca 48, 53, ca 54, 55, 56, 48–80	Löve and Löve, 1948, 1956; Reese, 1954
Italy	16, 24, 30, 32, ca 32, 33, 34, 36, 40, 42, 43, 44, 46, 48, ca 48, 50, 52, 56, ca 56, 58, 60, 61, 62, 63, 64, 72	Reese, 1954; Kootin-Sanwu and Woodell, 1970; Leoncini, 1951, 1952
Netherland	32, 56, 57	Reese, 1954; Smit, 1967
Norway	32, 56, ca 56	Reese, 1954; Kootin-Sanwu and Woodell, 1970
Poland	32, 48, 56, 58, 60	Skalińska et al., 1959; Wcislo, 1964, 1967, 1968; Falińska, 1981; Werpachowski, 1994
Portugal	32	Smit, 1967
Slovakia	32, 56	Májovský et al., 1974; Hrušovská-Osuská, 1988
Spain	32, 56	Smit, 1967
Sweden	28, 32, 48, 56, 59	Reese, 1954; Kootin-Sanwu and Woodell, 1970; Wcislo, 1967
Switzerland	60, ca 60	Reese, 1954
Former U.S.S.R.	16, 28, 32, 48, 56, ca 56, 60, 64	Wcislo, 1967; Rostovtseva, 1981; Goldblatt, 1981, 1984, 1988; Lavrenko and Serditov, 1984, 1985; Goldblatt and Johnson, 1990
Yugoslavia	32	Reese, 1954
<b>ASIA</b>		
Arctic Russia, Pamir and Altai, Kamschatka	32, 48, 56, ca 70	Kootin-Sanwu and Woodell, 1970
Japan	60, 88	Goldblatt, 1985; Goldblatt and Johnson, 1990


resented (Fig. 1). This material was analyzed biometrically according to the following five scored qualitative morphological features:

1 – margin of the top of the basal leaf (entire, serrate, dentate-crenate);

2 – margin of lobes of the basal leaf (entire, serrate, crenate, dentate-crenate);

3 – stem morphology (erect, ascending, creeping);

4 – shape of the beak (straight, weakly bent, bent, hooked);



**Fig. 2.** Scatter diagram of 74 populations of *Caltha palustris* s.l. in Poland, based on correspondence analysis versus axis 1 and axis 2, on the basis of 5 qualitative features (1–5). ● *C. palustris* subsp. *laeta*; ◆ *C. palustris* subsp. *palustris* var. *palustris*; ◇ *C. palustris* subsp. *palustris* var. *radicans*; ▲ *C. palustris* subsp. *palustris* var. *cornuta*. The specimens from which samples were taken for cytological tests are marked with larger symbols; the same numbers of populations are in Table 2; somatic chromosome numbers (2n) in parentheses.

5 – divergence of follicles (follicles straight and pressed, weakly bent, slightly divergent, distinctly bent, divergent, strongly recurved and divergent).

Nine quantitative features were also analyzed:

- 6 – angle of follicle bending;
- 7 – depth of sinus of basal leaf;
- 8 – angle of basal leaf sinus;
- 9 – ratio of follicle length (measured along the medial curve) to the shortest (straight) distance between the base of the follicle and the top of its beak;
- 10 – ratio of the difference between dorsal and abdominal edge lengths of the follicle to its length (measured along the medial curve);
- 11 – ratio of maximal width of basal leaf blade to its length (between apex and base);
- 12 – ratio of length of basal leaf blade measured along a radius angled 40° from the main axis


(connecting the base of the leaf with its apex to the length of the blade along the main axis; 13 and 14 – ratios as in 12, but with basal leaf blade measured along radii angled 80° and 140° from the main axis, respectively.

Numerical analyses were done based on the biometric data: correspondence analysis (CA) for the qualitative (1–5) features, and discriminant analysis (DA) for the quantitative features (6–14).

The aim was to establish the degree of homogeneity of the complex and the interrelations between the specific morphological groups, and indirectly to determine their variability.

#### KARYOLOGY

The specimens for karyological studies were collected from eleven natural habitats in Poland (Fig. 1) by the first author. Chromosome numbers were counted in a total of 15 plants representing all



**Fig. 3.** Scatter diagram of 74 populations of *Caltha palustris* s.l. in Poland based on discriminant analysis versus axis 1 and axis 2, on the basis of 9 quantitative features (6–14). ● *C. palustris* subsp. *laeta*; ◆ *C. palustris* subsp. *palustris* var. *palustris*; ◇ *C. palustris* subsp. *palustris* var. *radicans*; ▲ *C. palustris* subsp. *palustris* var. *cornuta*. The specimens from which samples were taken for cytological tests have been marked with larger symbols; the same numbers of populations are in Table 2; somatic chromosome numbers ( $2n$ ) in parentheses.

taxa of *C. palustris* s.l. known to occur in the country (Tab. 2).

The plants were placed in vessels with water for rooting. After several days the root tips were excised and pretreated with a saturated water solution of  $\alpha$ -bromonaphthalene for 2–4 h and fixed in 1:3 acetic alcohol. The root tips were stained in toto in acetic orcein for two weeks and squashed in 45% acetic acid. The chromosome counts were made in metaphase plates of one or two individuals from each population.

Voucher specimens are deposited in the Herbarium of the W. Szafer Institute of Botany, Polish Academy of Sciences in Cracow (KRAM).

## RESULTS

### MORPHOLOGY

On the basis of CA of 5 qualitative characters (1–5) and DA of 9 quantitative characters (6–14) (Figs. 2, 3), two main groups were distinguished within the

*C. palustris* complex. They are classified as two subspecies: subsp. *laeta* and subsp. *palustris*. Within subsp. *palustris*, apart from var. *palustris*, var. *radicans* was distinguished on the basis of the CA results (Fig. 2). The second variety has characteristically developed features as analyzed biometrically, but with procumbent stem roots at the nodes. DA provided grounds for differentiating a group of specimens classified as var. *cornuta* (Fig. 3). This variety has divergent follicles and a large angular divergence between the basal lobes. However, the range of morphological variability within this group largely overlaps that of var. *palustris*.

The lack of distinct divisions within the complex, and the occurrence of transitional forms not discussed in the present paper, points to the gradual character of variability within the *C. palustris* complex in Poland (Figs. 2, 3). The morphotypes distinguished within the *C. palustris* complex are assigned a low rank because of the lack of discontinuity in variability between these groups (Figs. 2, 3).

TABLE 2. Results of karyological studies of the *Caltha palustris* complex in Poland

Taxon	Locality (No)	N. lat.	E. long.	2n	
<i>C. palustris</i> L. subsp. <i>palustris</i> var. <i>palustris</i>	Mrzeżyno (1)	54°05'	15°17'	32	
	Duninowo (2)	54°32'	16°49'	32	
	Radnica (4)	52°05'	15°15'	32	
	Suchedniów (6)	51°03'	20°50'	56	
	Lipie (7)	51°01'	18°48'	32	
	Sandomierz (8)	50°42'	21°45'	32	
	Olkusz (9)	50°17'	19°04'	32	
	<i>C. palustris</i> L. subsp. <i>palustris</i> var. <i>radicans</i>	Frącki (3)	54°00'	23°20'	56
		<i>C. palustris</i> L. subsp. <i>palustris</i> var. <i>cornuta</i>	Zalesie-Moszczona Pańska (5)	52°27'	23°01'
Hrubieszów (10)	50°48'		23°53'	56	
<i>C. palustris</i> L. subsp. <i>laeta</i>	Dolina 5 Stawów Polskich	49°13'	20°02'	62	
	- Tatra Mts (11)				

## KARYOLOGY

Great variability in somatic chromosome number was found in the plants examined (2n from 25 to 94). The plants from Mrzeżyno, Duninowo and Olkusz, with a uniform chromosome number (2n = 4x = 32) in all root-tip metaphases, were exceptional. In most cases the chromosome number varied within individuals; different cells in their root-tips had different euploid and aneuploid chromosome numbers (mixoploidy). In each mixoploid plant, the most frequently observed chromosome number (2n = 32, 2n = 56, or 2n = 62) was treated as the main chromosome number, following Reese's rule (1954). These chromosome numbers are presented in Table 2.

In the eleven populations from Poland, somatic chromosome numbers 2n = 32, 2n = 56 and, in one case from the Tatra Mts., 2n = 62 were found (Fig. 4). More than one chromosome number, both euploid and aneuploid, were observed within one plant or even within one root: *C. palustris* subsp. *palustris* var. *radicans* (from Frącki) – 2n = 56, 64; *C. palustris* subsp. *palustris* var. *palustris* (from Radnica) – 2n = 25, 32, 60; *C. palustris* subsp. *palustris* var. *cornuta* (from Zalesie-Moszczona Pańska) – 2n = 42, 47, 56, 94; *C. palustris* subsp. *palustris* var. *palustris* (from Suchedniów) – 2n = 35, 49, 52, 56; *C. palustris* subsp. *palustris* var. *palustris* (from Lipie) – 2n = 28, 29, 32, 40, 56; *C. palustris* subsp. *palustris* var. *palustris* (from Sandomierz) – 2n = 30, 32, 35, 40, 42, 52, 53; *C. palustris* subsp. *palustris* var. *cornuta* (from Hrubieszów) – 2n = 39, 41, 48, 56, 64, 72; and *C. palustris* subsp. *laeta* (from Dolina 5 Stawów Polskich – Tatra Mts.) – 2n = 53, 57, 58, 59, 60, 61, 62.


The *C. palustris* metaphase plates show differentiation of chromosome morphology. The largest chromosome of the complement is about three times the length of the smallest one.

## DISCUSSION

None of the morphological or karyological research done so far has succeeded in dividing the *C. palustris* complex in Poland into unambiguous taxa (Wcisło, 1967; Falińska, 1974, 1976, 1981).

Our morphological studies confirmed the existence of a quite uniform group of plants from mountain habitats, classified as *C. palustris* L. subsp. *laeta* (Figs. 2, 3). The morphotype is characterized by a rounded leaf blade, a serrate or dentate-crenate basal leaf margin, and straight and pressed follicles leading to a short and straight beak. Wcisło (1967) established chromosome number 2n = 56 for this taxon from Poland. These results do not correspond exactly with Chrtkova and Jarolimova's (1999) recent data from the Czech Republic and Slovakia. They found two somatic chromosome numbers in *C. palustris* subsp. *laeta*: 2n = 32 (from hilly habitats) and 2n = 56 (from mountain habitats). Wcisło (1967) investigated several populations of subspecies from Bulgaria and determined one chromosome number, 2n = 32. In the present work, chromosome number 2n = 62 was found in this taxon in a population originating from the Tatra Mts. (Tab. 2). This number has not been reported in this subspecies before, and needs to be confirmed. This chro-

Fig. 4a–c. Metaphase plates of *Caltha palustris* s.l. (a) *C. palustris* subsp. *palustris* var. *palustris* (2n = 32; Duninowo), (b) *C. palustris* subsp. *palustris* var. *cornuta* (2n = 56; Hrubieszów), (c) *C. palustris* subsp. *laeta* (2n = 62; Tatra Mts). Bar = 10 µm.



mosome number has been reported from Italy for *C. palustris* subsp. *palustris* (Leoncini, 1952).

The data shows that morphologically subsp. *laeta* is a relatively homogenous taxon, and karyologically it is a heterogeneous one. Both  $2n = 32$  and  $2n = 56$  cytotypes have been reported for this taxon in Europe (Satzek in Skalińska et al, 1959; Wcisło, 1967; Chrtkova and Jarolimova, 1999).

*Caltha palustris* subsp. *palustris* is a much more variable taxon morphologically, as is confirmed by research from other parts of Europe (Woodell and Kootin-Sanwu, 1971; Smit, 1967). Our investigations showed the morphotype in Poland to be characterized by an erect stem, weak or distinct and divergent follicles, 30–70° angular divergence between basal lobes, and distinctly crenate to deeply dentate basal leaf margins. Biometric analysis distinguished three varieties within this subspecies: var. *palustris*, var. *cornuta* and var. *radicans*.

Varietas *palustris* is characterized by an erect stem, slightly divergent follicles, and a 40–60° angle sinus of the basal leaves. Varietas *radicans* is the most distinctive of the three. It is characterized by an ascending and creeping stem, mentioned also by Smit (1967, 1973). Less distinctive is var. *cornuta*, characterized by its strongly recurved and divergent follicles, kidney-shaped basal leaves, large-angle sinus of the basal leaves, and entire or dentate-crenate basal leaf margin. Between the varieties there are many transitional forms, whose status is not discussed here.

*Caltha palustris* subsp. *palustris* is equally variable in terms of karyology. Populations of subsp. *palustris* from Poland were characterized by the occurrence of one specific cytotype,  $2n = 56$ , except for one population with  $2n = 60$  (Wcisło, 1967). The specimens from the Czech Republic and Slovakia representing this taxon examined by Chrtkova and Jarolimova (1999) had chromosome number  $2n = 32$ , and there are individual reports of  $2n = 34$ ,  $2n = 35$  and  $2n = 36$ .

We also found cytological differentiation in the specimens of subsp. *palustris*, where numbers  $2n = 32$  and  $2n = 56$  were counted. In var. *palustris*, chromosome number  $2n = 32$  was counted in representatives of six populations, and  $2n = 56$  in one population. In var. *cornuta* and var. *radicans*  $2n = 56$  was found (Tab. 2).

Just as in subsp. *laeta*, on the basis of the data it is hard to decide on a number specific to subsp. *palustris*. As in the case of varieties distinguished morphologically, there is no specific cytotype (Tab. 2).

So far the research on *C. palustris* s.l. has shown great karyological variability within this complex. Different euploid and aneuploid chromosome numbers have been noted at ploidy levels from  $2n = 2x = 16$  to  $2n = 11x = 88$  (Tab. 1). Tetra- and heptaploid cytotypes ( $2n = 32, 56$ ) are most common in Europe, but di-, tri-, hexa-, octo- and nanoploid ( $2n = 16, 24, 48, 64, 72$ ) individuals have also been noted, together with aneuploids from  $2n = 38$  to  $2n = 70$  (Kootin-Sanwu and Woodell, 1970; Tab. 1).

Previous authors have reported different chromosome numbers within one plant (Leoncini, 1951, 1952; Reese, 1954; Kootin-Sanwu and Woodell, 1970). We also observed several numbers of chromosomes in one individual. Reese (1954) suggested that the number most often repeated can be taken as the main somatic number for plants with different chromosome numbers within their root tips.

Satzek (in Skalińska et al., 1959) and Wcisło (1967, 1968) found individuals with  $2n = 32, 56$  to be dominant in Poland, confirming observations from other parts of Europe (Smit, 1973). They also occasionally found plants with chromosome numbers  $2n = 48, 52, 58$  and  $60$ .

The results of detailed biometric analyses seem to rule out a simple relation between karyological and morphological variability in representatives of the *C. palustris* complex (also mentioned by Reese, 1954). In Poland, cytological variability is so great in the morphologically differentiated groups that taxa cannot be distinguished on the basis of their chromosome numbers. We can only state that, as in other parts of Europe (Smit, 1973), the two most common cytotypes in Poland are  $2n = 32$  and  $2n = 56$ . The difficulty in interpreting karyological differentiation within *C. palustris* s.l. is due to mixoploidy (Reese, 1954).

The great karyological variability (Tab. 1) and relatively weak differences between the groups, together with the presence of intermediate forms and the lack of genetic barriers (Wcisło, 1964, 1968), presumably means that *C. palustris* is a "young evolutionary taxon" (Hoffman, 1999).

## ACKNOWLEDGEMENTS

The authors would like to thank Professor Zbigniew Mirek, Dr. Andrzej Jankun and Dr. Andrzej Joachimiak for their valuable advice and suggestions for this paper.

This work was supported by KBN grant 6PO4C 05113.



## REFERENCES

- AKERROYD JR. 1993. *Caltha* L. In: Tutin TG, Burges NA, Chater AO, Edmondson JR, Heywood VH, Moore DM, Valentine DH, Walters S, and Webb DA [eds.], *Flora Europaea*, 81, 2nd ed. Univ. Press, Cambridge.
- CHRTKOVA A, and JAROLIMOVA V. 1999. Cytotaxonomical study of *Caltha palustris*. *Preslia* 71: 349–360.
- ENGLER A. 1936. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie*, vol. 67, 29. Verlag von Max Weg, Leipzig.
- FALIŃSKA K. 1974. Reaction of plant populations to differences in ecosystems and ecotone system. *Wiadomości Ekologiczne* 20: 356–376 (in Polish).
- FALIŃSKA K. 1976. Seasonal of *Caltha palustris* L. in forest and meadow communities. *Fragmenta Floristica et Geobotanica* 22: 529–544 (in Polish).
- FALIŃSKA K. 1981. Variability of *Caltha palustris* L. populations in garden culture. *Acta Societatis Botanicorum Poloniae* 50: 493–513.
- GOLDBLATT P. 1981. Index to plant chromosome numbers 1975–78. *Monographs in Systematic Botany* 5.
- GOLDBLATT P. 1984. Index to plant chromosome numbers 1979–81. *Monographs in Systematic Botany* 8: 324.
- GOLDBLATT P. 1985. Index to plant chromosome numbers 1982–83. *Monographs in Systematic Botany* 13: 169.
- GOLDBLATT P. 1988. Index to plant chromosome numbers 1984–85. *Monographs in Systematic Botany* 23: 186.
- GOLDBLATT P, and JOHNSON ED. 1990. Index to plant chromosome numbers 1986–87. *Monographs in Systematic Botany* 30: 163.
- HEGI G. 1912. *Illustrierte Flora vom Mittel-Europa*, vol. 3, 458. Carl Hanser Verlag, München.
- HOFFMANN MH. 1999. Biogeographical and evolutionary patterns in *Caltha* L. (Ranunculaceae). *Botanische Jahrbücher für Systematik Pflanzengeschichte und Pflanzengeographie* 121: 403–421.
- HRUŠOVSKÁ-OSUSKÁ L. 1988. Karyological study of some taxa of the flora of the northern part of Povazsky Inovec. Part I. *Acta Facultatis Rerum Naturalium Universitatis Comenianae. Ser. Botanica* 35: 70.
- KOOTIN-SANWU M, and WOODSELL SRJ. 1969. Supernumerary Chromosomes in *Caltha palustris*. *Chromosomes Today* 2: 192–196.
- KOOTIN-SANWU M, and WOODSELL SRJ. 1970. The cytology of *Caltha palustris* L. (*sensu lato*): Distribution and behaviour of the chromosome races. *Caryologia* 23: 225–239.
- KUCOWA I. 1985. *Caltha* L. In: Jasiewicz A [ed.], *Polish flora. Vascular plants. IV*, 12–15. Państwowe Wydawnictwo Naukowe, Warszawa – Kraków (in Polish).
- LAVRENKO AN, and SERDITOV NP. 1984. Karyosystematic study of the representatives of the families *Ranunculaceae* and *Paeoniaceae* in the North-East of the European part of the USSR. *Botanicheskyy Zhurnal* 70: 1346–1147.
- LAVRENKO AN, and SERDITOV NP. 1985. Chromosome numbers in some members of the *Ranunculaceae* family from Komi ASSR. *Botanicheskyy Zhurnal* 71: 1143–1144.
- LEONCINI ML. 1951. Biotipi cariologici e sistematici di *Caltha* in Italia. *Caryologia* 3: 336–350.
- LEONCINI ML. 1952. Nuove osservazioni cariologiche sul genere *Caltha*. *Caryologia* 4: 367–371.
- LOVE A, and LOVE D. 1948. Chromosome numbers of northern plant species. *University Institute of Applied Sciences Department of Agriculture, Reports, Series B* 3: 1–131. Reykjavik.
- LOVE A, and LOVE D. 1956. Cytotaxonomical conspectus of the Icelandic flora. *Acta Horti Gothoburgensis* 20: 65–290.
- MAJOVSKÝ J et al. 1974. Index of chromosome numbers of Slovakian flora. Part 4. *Acta Facultatis Rerum Naturalium Universitatis Comenianae. Ser. Botanica* 23: 5.
- MIREK Z, PIĘKOŚ-MIRKOWA H, ZAJĄC A, and ZAJĄC M. 1995. Vascular plants of Poland: a checklist. *Polish Botanical Studies Guidebook Series* 15: 1–101.
- PACZOSKI J. 1927. *Caltha* L. In: Szafer W [ed.], *Polish Flora. Vascular plants of Poland and adjacent territories. III*, 9–10. Polska Akademia Umiejętności, Kraków (in Polish).
- POLYA L. 1950. Chromosome numbers of Hungarian plants II. *Annales Universitatis Debreceniensis* 1: 46–56.
- REESE G. 1954. Euploidie, Aneuploidie und B-Chromosomen bei *Caltha palustris* L. *Planta* 44: 203–268.
- ROSTOVTSSEVA TS. 1981. Chromosome numbers of some species of the family Ranunculaceae. II. *Botanicheskyy Zhurnal* 66: 1751–1752.
- SCHOTT H, NYMAN CF, and KOTSCHY T. 1854. *Analecta Botanica*: 31–33.
- SKALIŃSKA M, CZAPIK R, and PIOTROWICZ M. 1959. Further studies in chromosome numbers of Polish Angiosperms (*Dicotyledons*). *Acta Societatis Botanicorum Poloniae* 28: 488–489.
- SMIT PG. 1967. Taxonomical and ecological studies in *Caltha palustris* L. *Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen, Ser. C, Biological and Medical Sciences* 70: 500–510.
- SMIT PG. 1973. A revision of *Caltha* (Ranunculaceae). *Blumea* 21: 119–150.
- TUTIN TG. 1964. *Caltha*. In: Tutin TG, Heywood VH, Burges NB, Valentine DH, Walters SM, Webb DA [eds.], *Flora Europaea*, 211. University Press, Cambridge.
- WCISŁO H. 1964. Experimental hybrids in the genus *Caltha* L. s.l. *Acta Biologica Cracoviensia Ser. Botanica* 7: 185–189.
- WCISŁO H. 1967. Karyological studies in *Caltha palustris* L. s.l. *Acta Biologica Cracoviensia Ser. Botanica* 10: 1–23.
- WCISŁO H. 1968. Further studies on experimental hybrids in *Caltha palustris* L. s.l. *Acta Biologica Cracoviensia Ser. Botanica* 11: 87–103.
- WERPACHOWSKI C. 1994. Effect of vegetation diversity on the population structure of marsh marigold (*Caltha palustris* L.) in the Biebrza river valley. In: Jankowska-Huflejt H and Golu-biewska E [eds.], *Conservation and management of fens*, 49–456. *Proceedings of the International Symposium on Conservation and Management of Fens*, 6–10 June 1994, 439–448. Institute for Land Reclamation and grassland Farming – Falenty, Agricultural University Faculty of Land Reclamation and Environmental Engineering, Warsaw, Warsaw-Biebrza.
- WOODSELL SRJ, and KOOTIN-SANWU M. 1971. Intraspecific variation in *Caltha palustris*. *New Phytologist* 70: 173–186.
- ZIEMMERMAN W. 1965. Familie Ranunculaceae. In: Hegi G [ed.], *Illustrierte Flora vom Mittel-Europa*, vol. 3, 78. Carl Hanser Verlag, München.