



A geometric method for infinite-dimensional chaos: Symbolic dynamics for the Kuramoto-Sivashinsky PDE on the line [☆]

Daniel Wilczak ^{*}, Piotr Zgliczyński

Faculty of Mathematics and Computer, Jagiellonian University, S. Łojasiewicza 6, 30-348 Kraków, Poland

Received 4 February 2019; revised 27 February 2020; accepted 3 June 2020

Available online 16 June 2020

Abstract

We propose a general framework for proving that a compact, infinite-dimensional map has an invariant set on which the dynamics is semiconjugated to a subshift of finite type. The method is then applied to certain Poincaré map of the Kuramoto-Sivashinsky PDE on the line with odd and periodic boundary conditions and with parameter $\nu = 0.1212$. We give a computer-assisted proof of the existence of symbolic dynamics and countable infinity of periodic orbits with arbitrary large periods.

© 2020 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

MSC: 35B30; 35B40; 65G30; 35B10; 35B45; 65N30

Keywords: Symbolic dynamics; Periodic orbits; Dissipative PDEs; Galerkin projection; Rigorous numerics; Computer-assisted proof

1. Introduction

In the study of nonlinear PDEs, there is a huge gap between what we can observe in numerical simulations and what we can prove rigorously. One possibility to overcome this problem is computer-assisted proofs. This paper is an attempt in this direction.

[☆] Research has been supported by Polish National Science Centre grant 2016/22/A/ST1/00077.

^{*} Corresponding author.

E-mail addresses: wilczak@ii.uj.edu.pl (D. Wilczak), zgliczyn@ii.uj.edu.pl, umzglicz@cyf-kr.edu.pl (P. Zgliczyński).




Fig. 1. Two approximate time-periodic orbits u^1 and u^2 .

We consider the one-dimensional Kuramoto-Sivashinsky PDE [23,33] (in the sequel we will refer to it as the KS equation), which is given by

$$u_t = -\nu u_{xxxx} - u_{xx} + (u^2)_x, \quad \nu > 0, \tag{1}$$

where $x \in \mathbb{R}$, $u(t, x) \in \mathbb{R}$ and we impose odd and periodic boundary conditions

$$u(t, x) = -u(t, -x), \quad u(t, x) = u(t, x + 2\pi). \tag{2}$$

The Kuramoto-Sivashinsky equation has been introduced by Kuramoto [23] in space dimension one for the study of front propagation in the Belousov-Zhabotinsky reactions. An extension of this equation to space dimension 2 (or more) has been given by G. Sivashinsky [33] in studying the propagation of flame front in the case of mild combustion.

The following theorem is the main result of this paper.

Theorem 1. *The system (1)–(2) with the parameter value $\nu = 0.1212$ is chaotic in the following sense. There exists a compact invariant set $\mathcal{A} \subset L^2((-\pi, \pi))$ which consists of*

1. *bounded full trajectories visiting explicitly given and disjoint vicinities U_1, U_2 of two selected periodic solutions u^1 and u^2 , respectively, with any prescribed order $(\alpha_i)_{i \in \mathbb{Z}} \in \{u^1, u^2\}^{\mathbb{Z}}$.*
2. *countable infinity of periodic orbits with arbitrary large periods. In fact, for each periodic sequence $(\alpha_i)_{i \in \mathbb{Z}} \in \{u^1, u^2\}^{\mathbb{Z}}$ there is a periodic solution of the system (1)–(2) visiting U_1 and U_2 in the order prescribed by $(\alpha_i)_{i \in \mathbb{Z}}$.*

The two special solutions u^1 and u^2 appearing in Theorem 1 are time-periodic — see Fig. 1. Profiles of initial conditions for u^1 and u^2 are shown in Fig. 2.

The proof of Theorem 1 is a mixture of topological methods and rigorous numerics. It uses neither of the special features of the Kuramoto-Sivashinsky PDE. Therefore, it should be applicable to other systems of dissipative PDEs. While there is a wide literature on computer-assisted verification of chaos for ODEs (see for example [2,15,27–29,32,36,37,39–41,50]), to the best of our knowledge, this is the first computer-assisted result of this type for PDEs.

In the topological part we provide a general framework for proving that a compact, infinite-dimensional map admits an invariant set on which the dynamics is semiconjugated to a subshift




Fig. 2. Profiles of initial conditions for time-periodic orbits u^1 and u^2 . (For interpretation of the colours in the figure(s), the reader is referred to the web version of this article.)

of finite type. The construction exploits an apparent existence of transversal heteroclinic connections between finite number of periodic points of (usually) low principal periods. These periodic orbits are then used to obtain a topological horseshoe for some higher iterate of the map. This is the same type of construction as it is used in the proof of the Smale-Birkhoff homoclinic theorem [19, Thm. 5.3.5].

In the case of the KS equation, we apply the method to certain Poincaré maps. The orbits u^1 and u^2 from Theorem 1 correspond to a fixed point and a period two point. Then, we construct symbolic dynamics along apparent heteroclinic connections between these points.

The proposed topological method is geometric, meaning that its assumptions can be expressed as a finite set of explicit inequalities. Therefore, we can use a computer to check that these inequalities are satisfied for a given map f , provided we have an algorithm that computes rigorous bounds on f on compact sets. In the case of the KS equation we have to compute rigorously bounds on the Poincaré map.

For this purpose we propose an algorithm, which allows to compute rigorous bounds on the trajectories of PDEs with periodic boundary conditions. It is applicable to a class of dissipative PDEs of the following form

$$u_t = Lu + N(u, Du, \dots, D^r u), \tag{3}$$

where $u \in \mathbb{R}^n$, $x \in \mathbb{T}^d = (\mathbb{R} \bmod 2\pi)^d$, L is a linear operator, N is a polynomial and by $D^s u$ we denote s th order derivative of u , i.e. the collection of all spatial partial derivatives of u of order s . The reason to consider polynomial and not more general functions N is technical — we need to compute the Fourier coefficients of $N(u, Du, \dots, D^r u)$. This can be achieved by taking suitable convolutions of Fourier expansions of u and its spatial partial derivatives.

We require, that the operator L is diagonal in the Fourier basis $\{e^{ikx}\}_{k \in \mathbb{Z}^d}$,

$$Le^{ikx} = -\lambda_k e^{ikx},$$

with

$$L_*|k|^p \leq \lambda_k \leq L^*|k|^p, \quad \text{for all } |k| > K \text{ and } K, L_*, L^* \geq 0, \\ p > r.$$

If the solutions are sufficiently smooth, the problem (3) can be written as an infinite ladder of ordinary differential equations for the Fourier coefficients in $u(t, x) = \sum_{k \in \mathbb{Z}^d} u_k(t) e^{ikx}$, as follows

$$\frac{du_k}{dt} = f_k(u) = -\lambda_k u_k + N_k(\{u_j\}_{j \in \mathbb{Z}^d}), \quad \text{for all } k \in \mathbb{Z}^d. \quad (4)$$

The crucial fact, which makes our approach to the rigorous integration of (4) possible is the *isolation property*, which reads:

Let

$$W = \left\{ \{u_k\}_{k \in \mathbb{Z}^d} \mid |u_k| \leq \frac{C}{|k|^s q^{|k|}} \right\}$$

where $q \geq 1$, $C > 0$, $s > 0$.

Then there exists $K > 0$, such that for $|k| > K$ there holds

$$\text{if } u \in W, \quad |u_k| = \frac{C}{|k|^s q^{|k|}}, \quad \text{then } u_k \cdot f_k(u) < 0.$$

Projection of the set W onto k th mode is a closed disc (or interval) centred at zero and of radius $r_k = \frac{C}{|k|^s q^{|k|}}$. Geometrically the isolation property means that if $|u_k| = r_k$ for some $k > K$ then the k -th component of the vector field is pointing inwards the set. As a consequence, the only way a trajectory may leave the set W forward in time is by increasing some of leading modes $|u_k|$, $k \leq K$ above the threshold r_k . However, one has to be careful with what “inwards” means in the above statement because W has empty interior as a compact subset of l_2 .

This property is used in our algorithm to obtain a priori bounds for $u_k(h)$ for small $h > 0$ and $|k| > K$, while the leading modes u_k for $|k| \leq K$ are computed using tools for rigorous integration of ODEs [3,24,30] based on the interval arithmetics [26]. Moreover, from the point of view of topological method, the isolation property is of crucial importance as it shows that on the tail we have the entry behaviour, which enable us to apply the finite dimensional tools from the dynamics.

Our algorithm for rigorous integration of (4) stems from the *method of self-consistent bounds*, which was introduced in [46] and later developed in [47–49,51]. In the present paper we propose some significant modifications to the method, discussed more in details in Section 4.

The choice of the KS equation for this study is motivated by the following facts.

- The existence of a compact global attractor, the existence of a finite-dimensional inertial manifolds for (1)–(2) are well established — see for example [4,13,12,31] and the literature cited there. We would like to emphasise, that we are not using these results in our work.
- There exist multiple numerical studies of the dynamics of the KS equation (see for example [5,20,22,21,34]), where it was shown, that the dynamics of the KS equation can be highly nontrivial for some values of parameter ν , while being well represented by relatively small number of modes.




Fig. 3. Numerically observed chaotic attractor for (1)–(2) obtained by simulation of a finite-dimensional projection of the corresponding infinite-dimensional ODE for the Fourier coefficients in $u(t, x) = \sum_{k=1}^{\infty} u_k(t) \sin(kx)$. Projection onto (u_2, u_3) plane of the intersection of the observed attractor with the Poincaré section $u_1 = 0, u_1' > 0$ is shown along with an approximate location of the two periodic points u^1, u^2 appearing in Theorem 1. The point u^1 is a fixed point for the Poincaré map and u^2 is of period two.

- There are several papers devoted to computer-assisted proofs of periodic orbits for the KS equation by Zgliczyński [51,53], Arioli and Koch [1] and by Figueras, Gameiro, de la Llave and Lessard [11,10,16].

While the choice of the odd periodic boundary conditions was motivated by earlier numerical studies of KS equation [5,22], the basic mathematical reason is the following: the equation (1) with periodic boundary conditions has the translational symmetry. This implies, that for a fixed value of ν , all periodic orbits are members of one-parameter families of periodic orbits. The restriction to the invariant subspace of odd functions breaks this symmetry, and gives a hope, that dynamically interesting objects are isolated and easier accessible for computer-assisted proofs.

Our choice of the parameter value $\nu = 0.1212$ is motivated by a numerical observation, that the Feigenbaum route to chaos through successive period doubling bifurcations [9] happens for (8) as ν decreases toward $\nu = 0.1212$ (first observed for other values of ν in [34]). For this parameter value a chaotic attractor is observed — see Fig. 3.

The content of this paper can be described as follows. In Section 2 we discuss the theorems and definitions related to covering relations and our method of obtaining infinite number of periodic points and symbolic dynamics. In Section 3 we explain the particularities of the application of topological theorems from Section 2 to the KS-equation. This ends the “geometric” part of the paper.

Our algorithm for rigorous integration of dissipative PDEs, which is necessary to compute Poincaré maps, is an independent result and it is presented in Section 4. All notions introduced there are of ‘local’ nature and there is no need to discuss them in more broad context in the introduction. These are just tools to justify the validity of our algorithm. The crucial fact which makes our algorithm to work is the isolation property, and as such it was discussed above in the introduction.

1.1. Notation

By \mathbb{C} , \mathbb{R} , \mathbb{Z} , \mathbb{N} we denote the sets of complex, real, integer and natural numbers including zero, respectively. With \mathbb{N}_+ we denote positive integers.

Let (T, ρ) be a metric space. For a set $X \subset T$, by $\text{int } X$, \overline{X} and ∂X we denote the interior, the closure and the boundary of X , respectively. If $X \subset Y \subset T$, then by $\text{int}_Y X$ and by $\partial_Y X$ we denote respectively the interior and the boundary of X with respect to the metric space (Y, ρ) . By $B(c, r) = \{x \mid \rho(c, x) < r\}$ we denote the ball of radius r centred at c . For a point $p \in T$ put $\rho(p, X) = \inf\{\rho(p, q) \mid q \in X\}$. We define $B(X, \epsilon) = \{y \mid \rho(y, X) < \epsilon\}$.

For $k \in \mathbb{N}$, $c \in \mathbb{R}^k$ and $r \geq 0$ by $B_k(c, r)$ we denote an open ball in \mathbb{R}^k of the radius r and the centre c . The norm used to define $B_k(c, r)$ will be usually the Euclidean one, but in most cases in this paper any norm can be used. By $B_k = B_k(0, 1)$ and $\overline{B}_k = \overline{B}_k(0, 1)$ we will denote the open and closed unit balls, respectively.

Let a Hilbert space $\mathcal{H} = \mathcal{X} \oplus \mathcal{Y}$ be a direct sum of two orthogonal subspaces $\mathcal{X}, \mathcal{Y} \subset \mathcal{H}$. We extend the notion of the direct sum to sets in a natural way. For $A \subset \mathcal{X}$ and $B \subset \mathcal{Y}$ we set

$$A \oplus B := \{a + b \mid a \in A, b \in B\}.$$

We will be often dealing with Galerkin projections in some space \mathcal{H} with the basis $\{e_i\}_{i \in J}$, $J \subset \mathbb{Z}^d$. We define the following projections: $\pi_k(\sum_{i \in J} u_i e_i) = u_k$ and $\pi_{\leq n}(\sum_{i \in J} u_i e_i) = \sum_{|i| \leq n} u_i e_i$ and analogously for other possible sets of indices entering into the projection. For a set $W \subset \pi_{\leq n} X$ by $\text{int}_{\leq n} W$ and $\partial_{\leq n} W$ we will denote respectively the interior and the boundary of W with respect to the set $\pi_{\leq n} X$.

2. Topological method for symbolic dynamics for maps in infinite dimension

Topological methods proved to be very useful in the context of computer-assisted study of dynamical systems. One of the most efficient in the context of studying chaotic dynamics is the *method of covering relations* introduced in [44,50] for maps with one exit (“unstable”) direction and later extended to include many exit directions in [52] known also in the literature as the method of correctly aligned windows [7,8]. In this section we extend this method to a class of maps defined on compact (possible infinite-dimensional) subsets of real normed spaces. The notion is quite similar but relies on the fixed point index for compact maps on Absolute Neighbourhood Retracts (ANRs) as defined in the book of Granas and Dugundji [17] (see also survey by Mawhin [25]). For the purpose of this article it suffices to note, that each convex and closed subset of a normed spaces is an ANR. Such an extension is motivated by the applications we keep in mind – Poincaré maps for infinite-dimensional ODEs. Another variant of this extension was discussed in [45,53].

2.1. Covering relations for maps on compact ANRs

The aim of this section is to extend the notion of covering relation to infinite-dimensional case.

Definition 2. Let X be a real normed space. A h -set, $N = (|N|, c_N, u(N))$, is an object consisting of the following data




Fig. 4. An example of an h-set in three dimensions with $u(N) = 1$ and $T_N = D_2$ – a two-dimensional closed disc. Here $N_c = \overline{B_1} \oplus D_2$.

- $|N| \subset X$ – a compact set called *the support* of N ,
- $u(N)$ – a nonnegative integer,
- c_N – a homeomorphism of X such that

$$c_N^{-1}(|N|) = \overline{B_{u(N)}} \oplus T_N =: N_c,$$

where T_N is a convex set, such that $0 \in T_N$.

The above definition generalizes the concept of *h*-sets introduced in [52] for finite-dimensional spaces, where a *h*-set is defined as the product of closed unit balls $\overline{B_u} \times \overline{B_s} \subset \mathbb{R}^{u+s}$ in a coordinate system c_N . The next definition extends the notion of covering relations from [52] to maps acting on ANRs in infinite-dimensional real normed spaces.

Definition 3. Let N and M be *h*-sets in X and Y , respectively such that $u = u(N) = u(M)$. Let $f: |N| \rightarrow Y$ be a continuous map and set $f_c := c_M^{-1} \circ f \circ c_N$. We say that N *f*-covers M , denoted by $N \xrightarrow{f} M$, if there is a linear map $L: \mathbb{R}^u \rightarrow \mathbb{R}^u$ and a continuous homotopy $H: [0, 1] \times N_c \rightarrow Y$, such that

- [CR1]: $H(0, \cdot) = f_c$,
- [CR2]: $H(1, x, y) = (L(x), 0)$, for all $(x, y) \in N_c$,
- [CR3]: $H(t, x, y) \notin M_c$, for all $(x, y) \in \partial B_u \oplus T_N$, $t \in [0, 1]$ and
- [CR4]: $H(t, x, y) \in \mathbb{R}_u \oplus T_M$ for $(x, y) \in N_c$ and $t \in [0, 1]$.

Observe that in the context of the above definition the maps f_c and H are compact, because $|N|$ is compact. This will allow us to use the fixed point index for ANRs in real normed spaces for compact maps as defined in [17].

A typical picture of a *h*-set with $u(N) = 1$ is given in Fig. 4. Fig. 5 and Fig. 6 illustrate covering relations with one and two exit directions, respectively. In Fig. 7 we illustrate the major difference between the finite-dimensional concept of covering relations from [52] and that from Definition 3. The condition

$$H(t, N_c) \cap (B_u \times \partial T_M) = \emptyset$$

from [52] is replaced by [CR4], because in general case T_M may have empty interior. Instead, we require that $H(t, N_c)$ lies in the strip $\mathbb{R}^u \oplus T_M$.

The following theorem shows the usefulness of the notion of covering relation to detect periodic orbits.




Fig. 5. An example of f -covering relation: $N \xrightarrow{f} M$. In this case, the homotopy joining $f_c(x, y)$ with a linear map $(L(x), 0)$ and satisfying [CR1]–[CR4] is simply given by $H(t, x, y) = t(L(x), 0) + (1 - t)f_c(x, y)$.




Fig. 6. An example of $N \xrightarrow{f} N$ with $N = [-1, 1]^3$, $c_N = \text{Id}$, $u(N) = 2$ and $T_N = [-1, 1]$. In this case $\pi_u f(\partial B_2 \times [-1, 1])$ covers in topologically non-trivial manner the ball (in max norm) \bar{B}_2 . Moreover $f(N) \subset \mathbb{R}^2 \times T_N$. The homotopy joining $f(x, y)$ with a linear map $(L(x), 0)$ (one can take $L(x_1, x_2) = 2(x_1, x_2)$) and satisfying [CR1]–[CR4] is simply given by $H(t, x, y) = t(L(x), 0) + (1 - t)f(x, y)$.




Fig. 7. Comparison of two definitions of covering relation. Finite-dimensional version from [52] requires that $f(N)$ does not touch $B_u \times \partial T_M$ (top panel). Condition [CR4] in Definition 3 requires, that the $f(N)$ lies in the strip $\mathbb{R}^u \times T_M$ (bottom panel).

Theorem 4. Let $X_i, i = 1, \dots, k$ be real normed spaces and let N_i be h -sets in X_i , respectively. Assume

$$N_1 \xrightarrow{f_1} N_2 \xrightarrow{f_2} \dots \xrightarrow{f_{k-1}} N_k \xrightarrow{f_k} N_1.$$

Then there exists $u_1 \in |N_1|$ such that

$$(f_i \circ \dots \circ f_1)(u_1) \in |N_{i+1}| \quad \text{for } i = 2, \dots, k - 1 \quad \text{and} \tag{5}$$

$$(f_k \circ \dots \circ f_1)(u_1) = u_1. \tag{6}$$

Proof. Put $N_{k+1} = N_1$. The set $X = X_1 \times \dots \times X_k$ with the maximum norm is a real normed space. The set

$$Y = (\mathbb{R}^u \oplus T_{N_1}) \times \dots \times (\mathbb{R}^u \oplus T_{N_k}) \subset X$$

is convex and by [17, Col. 4.4] it is an ANR. Put

$$N = (B_u \oplus T_{N_1}) \times \dots \times (B_u \oplus T_{N_k}) \subset Y.$$

Let H_i be a homotopy from the definition of covering relation $N_i \xrightarrow{f_i} N_{i+1}$. By [CR4] the range of $(f_i)_c$ and $H_i(t, \cdot)$ is in $\mathbb{R}^u \oplus T_{N_{i+1}}$ for $i = 1, \dots, k, t \in [0, 1]$. Therefore we can define a map $F: \overline{N} \rightarrow Y$ by

$$F(u_1, u_2, \dots, u_k) = ((f_k)_c(u_k), (f_1)_c(u_1), \dots, (f_{k-1})_c(u_{k-1}))$$

and a homotopy $H: [0, 1] \times \overline{N} \rightarrow Y$ by

$$H\left(t, \begin{bmatrix} (x_1, y_1) \\ (x_2, y_2) \\ \dots \\ (x_k, y_k) \end{bmatrix}\right) = \begin{bmatrix} H_k(t, x_k, y_k) \\ H_1(t, x_1, y_1) \\ \dots \\ H_{k-1}(t, x_{k-1}, y_{k-1}) \end{bmatrix}.$$

It is easy to see that for all $t \in [0, 1]$ the mapping $H(t, \cdot)$ is fixed point free on $\partial_Y N$. Indeed, if $u = ((x_1, y_1), \dots, (x_k, y_k)) \in \partial_Y \overline{N}$ then $x_i \in \partial B_u$ for some $i = 1, \dots, k$. From [CR3] we have $H_i(t, x_i, y_i) \notin \overline{B_u} \oplus T_{N_{i+1}}$ and therefore $H(t, u) \notin \overline{N}$.

Thus, the fixed point index $i(H(t, \cdot), N)$ for compact maps on ANRs [17] is well defined and does not depend on $t \in [0, 1]$.

For $t = 1$ we have

$$H\left(1, \begin{bmatrix} (x_1, y_1) \\ \dots \\ (x_k, y_k) \end{bmatrix}\right) = \begin{bmatrix} (L_k(x_k), 0) \\ (L_1(x_1), 0) \\ \dots \\ (L_{k-1}(x_{k-1}), 0) \end{bmatrix},$$

where L_i is a linear map from the covering relation $N_i \xrightarrow{f_i} N_{i+1}$. By [CR2] it is an expanding isomorphism which maps ∂B_u out of the unit ball $\overline{B_u}$. Thus, $H(1, \cdot)$ is a hyperbolic linear map and

$$|i(H(1, \cdot), N)| = 1.$$

Therefore $i(F, N) = I(H(0, \cdot), N) \neq 0$ and in consequence the mapping F has a fixed point $(\hat{u}_1, \dots, \hat{u}_k) \in N$. Now the point $u_1 = c_{N_1}(\hat{u}_1)$ satisfies (5) and (6). \square

2.2. Subshifts of finite type and symbolic dynamics

The following definitions are standard (see for example [19]). Let us fix $k > 0$ and let $(A_{ij})_{i,j=1,\dots,k}$ be $k \times k$ matrix, such that $A_{ij} \in \{0, 1\}$. We define Σ_A and Σ_A^+ by

$$\begin{aligned} \Sigma_A &= \{\alpha \in \{1, 2, \dots, k\}^{\mathbb{Z}} \mid A_{\alpha_i \alpha_{i+1}} = 1 \ \forall i \in \mathbb{Z}\}, \\ \Sigma_A^+ &= \{\alpha \in \{1, 2, \dots, k\}^{\mathbb{N}} \mid A_{\alpha_i \alpha_{i+1}} = 1 \ \forall i \in \mathbb{N}\}. \end{aligned}$$

We define a shift map σ on Σ_A and Σ_A^+ by

$$\sigma(\alpha)_i = \alpha_{i+1}, \quad \forall i \in \mathbb{Z} \ (i \in \mathbb{N}).$$

The pairs (Σ_A, σ) and (Σ_A^+, σ) are called *subshifts of finite type with transition matrix A*. Let

$$\begin{aligned} \Sigma_k &= \{1, 2, \dots, k\}^{\mathbb{Z}}, \\ \Sigma_k^+ &= \{1, 2, \dots, k\}^{\mathbb{N}}. \end{aligned}$$

We call (Σ_k, σ) and (Σ_k^+, σ) *full shifts* on k symbols.

Let $X_i, i = 1, \dots, k$ be real normed spaces and let $N_i \subset X_i$ be h -sets, such that $|N_i| \cap |N_j| = \emptyset$ for $i \neq j$. Note, we do not require that the spaces X_i are different. Let $f^\beta : |M_\beta| \rightarrow Y_\beta$ be continuous maps, for $\beta = 1, \dots, m$, where $M_\beta \in \{N_1, \dots, N_k\}$ and $Y_\beta \in \{X_1, \dots, X_k\}$. Again, we accept that two different mappings are defined on the same set $|N_i|$.

We define a *transition matrix* $(A_{ij})_{i,j=1,\dots,k}$ in the following way:

$$A_{ij} = \begin{cases} 1 & \text{if } N_i \xrightarrow{f^\beta} N_j, \text{ for some } \beta \in \{1, \dots, m\}, \\ 0 & \text{otherwise.} \end{cases}$$

Definition 5. A sequence $(x_i)_{i \in \mathbb{N}}$ is called a full trajectory with respect to family of maps (f^1, \dots, f^m) if for all $i \in \mathbb{N}$ there is $\beta_i \in \{1, \dots, m\}$ such that

$$f^{\beta_i}(x_i) = x_{i+1}.$$

A sequence $(\alpha_i)_{i \in \mathbb{N}} \in \{1, \dots, k\}^{\mathbb{N}}$ is called admissible, if there is a full trajectory $(x_i)_{i \in \mathbb{N}}$ with respect to (f^1, \dots, f^m) such that

$$x_i \in |N_{\alpha_i}|.$$

The following theorem addresses the issue of the existence of an orbit realising a non-periodic sequence of covering relations.

Theorem 6. *Every sequence of symbols $(\alpha_i)_{i \in \mathbb{N}} \in \Sigma_A^+$ is admissible. Moreover, if $(\alpha_i)_{i \in \mathbb{N}}$ is T -periodic, then the corresponding trajectory $(x_i)_{i \in \mathbb{N}}$ may be chosen to be a T -periodic sequence, too.*

Proof. From Theorem 4 every T -periodic sequence $(\alpha_i)_{i \in \mathbb{N}} \in \Sigma_A^+$ of symbols is realized by a T -periodic trajectory $(x_i)_{i \in \mathbb{N}}$.

Let us fix a sequence $(\alpha_i)_{i \in \mathbb{N}} \in \Sigma_A^+$ which is not periodic. This means, that there is a sequence $(\beta_i)_{i \in \mathbb{N}}$ such that

$$N_{\alpha_i} \xrightarrow{f^{\beta_i}} N_{\alpha_{i+1}}$$

for $i \in \mathbb{N}$. For every $T > 0$ we can construct a closed loop of covering relations

$$N_{\alpha_0} \xrightarrow{f^{\beta_0}} N_{\alpha_1} \xrightarrow{f^{\beta_1}} \dots \xrightarrow{f^{\beta_{T-1}}} N_{\alpha_T} \xrightarrow{g} N_{\alpha_0},$$

by adding an artificial covering relation (an affine map g) at the end of this sequence. For example the map g defined by $g_c(x, y) = (2x, 0)$, where $x \in \overline{B}_u$, $y \in T_N$ and the homotopy $H(\lambda, z) = g_c(z)$ is good for this purpose.

From Theorem 4 this sequence is realised by a $(T + 1)$ -periodic full trajectory $(x_i^T)_{i \in \mathbb{N}}$. Since $|N_{\alpha_0}|$ is compact, we can choose a subsequence $(x_0^{T_j})_{j \in \mathbb{N}}$ which converges to $x_0 \in |N_{\alpha_0}|$. Then for $i \in \mathbb{N}$ and $T_j > i$ we have $(f^{\beta_i} \circ \dots \circ f^{\beta_0})(x_0^{T_j}) \in |N_{\alpha_{i+1}}|$ and by the continuity of each f^{β_i} we obtain that

$$x_{i+1} := (f^{\beta_i} \circ \dots \circ f^{\beta_0})(x_0) = \lim_{j \rightarrow \infty} (f^{\beta_i} \circ \dots \circ f^{\beta_0})(x_0^{T_j}) \in |N_{\alpha_{i+1}}|.$$

The constructed sequence $(x_i)_{i \in \mathbb{N}}$ satisfies the assertion. \square

Theorem 6 is a powerful topological tool for proving symbolic dynamics and countable infinity of periodic orbits for maps. It suffices to find two different finite loops of covering relations starting from the same h -set, say N

$$\begin{aligned} N &\xrightarrow{f} M_0 \xrightarrow{f} \dots \xrightarrow{f} M_m \xrightarrow{f} N, \\ N &\xrightarrow{f} K_0 \xrightarrow{f} \dots \xrightarrow{f} K_k \xrightarrow{f} N. \end{aligned}$$

Based on them, we can construct different bi-infinite (periodic) sequences of covering relations which by Theorem 6 are realized by (periodic) trajectories of f . Another example is classical Smale horseshoe, with full transition matrix Σ_2 — see Fig. 8.




Fig. 8. Sequence of covering relations $N \xrightarrow{f} N \xrightarrow{f} M \xrightarrow{f} M \xrightarrow{f} N$ implies symbolic dynamics and countable infinity of periodic orbits for f — see Theorem 6.

3. The Kuramoto-Sivashinsky equation

Consider the equation (1) with periodic and odd boundary conditions (2) and assume that u is a classical solution to (1)–(2) given as a convergent Fourier series

$$u(t, x) = \sum_{k=1}^{\infty} -2a_k(t) \sin(kx). \tag{7}$$

The particular form of representation of u given in (7) comes from imposing on $u(t, x) = \sum_k u_k(t)e^{ikx}$ periodic and odd boundary conditions. Then we have $a_k(t) = \text{Im } u_k(t)$.

It is easy to see that (see for example [5,46]) that for sufficiently regular functions the system (1)–(2) give rise to an infinite ladder of coupled ODEs

$$\frac{da_k}{dt} = f_k(a) := k^2(1 - vk^2)a_k - k \sum_{n=1}^{k-1} a_n a_{k-n} + 2k \sum_{n=1}^{\infty} a_n a_{n+k}, \quad k \geq 1. \tag{8}$$

In order to apply topological fixed point theorems discussed in Section 2 to (8) we need to choose a topology for space of sequences $(a_k)_{k=1}^{\infty}$. We will demand that $(a_k)_{k=1}^{\infty} \in l_2$, where

$$l_2 = \left\{ (a_k)_{k=1}^{\infty} \mid \sum_{k=1}^{\infty} |a_k|^2 < \infty \right\},$$

$$\| (a_k)_{k=1}^{\infty} \|_2 = \sqrt{\sum_{k=1}^{\infty} |a_k|^2}.$$

The use of l_2 defined in terms of $(a_k)_{k=1}^{\infty}$ is quite arbitrary. More natural function spaces and norms (compare Theorem 30 in the Appendix A.1) would be the ones induced from L^2 or H^k with $k \geq 1$ on the function space of 2π -periodic functions. This, however, does not matter much, as we will consider the geometrically fast decaying sequences in l_2 , only. Geometric decay of coefficients guarantees that these sequences represent analytic functions for which replacing (1) with (8) makes sense.

For $S > 0$ and $q > 1$ we set

$$W_{q,S} = \left\{ (a_k)_{k=1}^{\infty} \mid a_k \in \mathbb{R}, |a_k| \leq Sq^{-k} \text{ for } k \geq 1 \right\},$$

$$l_{2,q} = \left\{ (a_k)_{k=1}^{\infty} \mid a_k \in \mathbb{R}, \exists S \geq 0 : |a_k| \leq Sq^{-k} \text{ for } k \geq 1 \right\} = \bigcup_{S>0} W_{q,S}.$$

Observe the set $W_{q,S} \subset l_2$ is compact. It is also compact in the topology induced by the H^k norm or L^2 norm on the space 2π -periodic functions and the convergence on $W_{q,S}$ in any of these norms and in l_2 norm is equivalent to the coordinate-wise convergence, i.e. $\lim_{j \rightarrow \infty} (a_k^j)_{k \in \mathbb{N}_+} = (c_k)_{k \in \mathbb{N}_+}$ iff for all $k \in \mathbb{N}_+$ holds $\lim_{j \rightarrow \infty} a_k^j = c_k$.

From classical results (see Theorem 30 in Appendix A.1) it follows that forward solutions of system (8) exist for all initial conditions $a \in l_2$ which defines a continuous semiflow on l_2 (a semigroup in the terminology used in [35]). In fact, we are not using this result in the sequel. For our proof we use the local semiflow restricted to some $W_{q,S}$ for suitably chosen S and q .

3.1. Symbolic dynamics in the KS equation

3.1.1. Heuristics for Galerkin projection of (8)

After an extensive numerical simulation we have found that the dynamics of Galerkin projections does not differ significantly for projections of dimension $n \geq 14$. Therefore, we set $n = 14$ and we constructed all objects that appear in our computation (like Poincaré sections, h-sets) using approximate flow generated by n -dimensional projection.

Let us consider a Poincaré section

$$\Theta = \{(a_k)_{k=1}^\infty \in \pi_{\leq n} l_2 \mid a_1 = 0 \wedge a'_1 = f_1(a) > 0\}$$

and the associated Poincaré map $P : \Theta \supset \text{dom}(P) \rightarrow \Theta$ with respect to the local flow induced by the n -dimensional Galerkin projection.

After an extensive numerical simulation we have found approximate periodic and heteroclinic points for P – see Fig. 9 and supplementary material [38]. These are

- $a^1 \in \Theta$ – an approximate fixed point for P , with one unstable direction,
- $a^2 \in \Theta$ – an approximate period-two point for P , with one unstable direction,
- $a_{1 \rightarrow 2}^i \in \Theta, i = 0, \dots, 10$ – an approximate heteroclinic chain satisfying

$$\begin{aligned} \|a^1 - a_{1 \rightarrow 2}^0\|_2 &\approx 1.676 \cdot 10^{-6}, \\ P^2(a_{1 \rightarrow 2}^i) &\approx a_{1 \rightarrow 2}^{i+1} \quad \text{for } i = 0, \dots, 9, \\ \|P^2(a_{1 \rightarrow 2}^{10}) - a^2\|_2 &\approx 3.932 \cdot 10^{-10}, \end{aligned}$$

- $a_{2 \rightarrow 1}^i \in \Theta, i = 0, \dots, 10$ – an approximate heteroclinic chain satisfying

$$\begin{aligned} \|a^2 - a_{2 \rightarrow 1}^0\|_2 &\approx 2.425 \cdot 10^{-6}, \\ P^2(a_{2 \rightarrow 1}^i) &\approx a_{2 \rightarrow 1}^{i+1} \quad \text{for } i = 0, \dots, 7 \text{ and } i = 9, \\ P^3(a_{2 \rightarrow 1}^8) &\approx a_{2 \rightarrow 1}^9, \\ \|P(a_{2 \rightarrow 1}^{10}) - a^1\|_2 &\approx 1.144 \cdot 10^{-6}. \end{aligned}$$




Fig. 9. Numerically observed heteroclinic connections for P between a^1 and a^2 in both directions.

3.1.2. Full system (8)

We consider the system (8) on $l_{2,q}$ with $q = 3/2$. In what follows we will use approximate periodic points a^1, a^2 for P and heteroclinic chains $a^i_{1 \rightarrow 2}, a^i_{2 \rightarrow 1}, i = 0, \dots, 10$ as presented in Section 3.1.1.

Analogously as in the case ODEs we can define the Poincaré map between two sections, see Appendix A.3. The notation is the same as for ODEs. We define

$$P^i_{\Pi_1 \rightarrow \Pi_2} = P^{i-1}_{\Pi_2 \rightarrow \Pi_2} \circ P_{\Pi_1 \rightarrow \Pi_2}.$$

The approximate heteroclinic loop for n -dimensional Galerkin projection was used to construct symbolic dynamics for the infinite-dimensional system (8). More precisely, we constructed

- affine Poincaré sections Θ^1 and Θ^2 , such that $a^i \in \Theta^i, i = 1, 2$,
- affine Poincaré sections $\Theta^i_{1 \rightarrow 2}$ and $\Theta^i_{2 \rightarrow 1}$, such that $a^i_{1 \rightarrow 2} \in \Theta^i_{1 \rightarrow 2}$ and $a^i_{2 \rightarrow 1} \in \Theta^i_{2 \rightarrow 1}$ for $i = 0, \dots, 10$,

- two h -sets $N^i \subset \Theta^i$, $i = 1, 2$ with $u(N^i) = 1$ (i.e. one exit direction), such that $a^i \in |N^i|$, respectively,
- two sequences of h -sets $N_{1 \rightarrow 2}^i \subset \Theta_{1 \rightarrow 2}^i$ and $N_{2 \rightarrow 1}^i \subset \Theta_{2 \rightarrow 1}^i$ for $i = 0, \dots, 10$, with $u(N_{1 \rightarrow 2}^i) = u(N_{2 \rightarrow 1}^i) = 1$, such that $a_{j \rightarrow c}^i \in |N_{j \rightarrow c}^i|$.

Explicit coordinates of the points $a^i, a_{j \rightarrow c}^i$, the Poincaré sections $\Theta^i, \Theta_{j \rightarrow c}^i$ and all h -sets listed above are given in the supplementary material [38].

Remark 7. One important parameter in our computer-assisted proof is an integer m which is by our choice set to $m = n + 1 = 15$. This is the number of explicitly stored Fourier coefficients of $(a_k)_{k=1}^\infty$ in a computer memory. The remaining coefficients, called tail, are bounded uniformly by a set represented by two real numbers S and q — see representation of h -sets in Appendix A.4.

Remark 8. The Poincaré section which contains a point $p \in \{a^1, a^2, a_{1 \rightarrow 2}^1, a_{2 \rightarrow 1}^1\}$ is chosen to be a hyperplane almost orthogonal to the flow direction of m -dimensional Galerkin projection of (8) at the point p . This means, in particular, that all Poincaré sections are defined in terms of $(a_k)_{k=1}^m$ by

$$\left\{ x \in I_{2,q} \mid \sum_{k=1}^m f_k(p)(x - p)_k = 0 \right\}.$$

We have found, that locally orthogonal sections help us in reducing overestimation when we compute rigorously Poincaré map.

Remark 9. All h -sets $N^i, N_{j \rightarrow c}^i$ are constructed to be pairwise disjoint.

Using an algorithm for rigorous integration of (8) discussed in the next sections we obtained a computer-assisted proof of the following lemma.

Lemma 10. *All the covering relations listed below are satisfied.*

$$\begin{aligned} N^1 &\xrightarrow{P^2} N^1 \xrightarrow{P^3} N_{1 \rightarrow 2}^0 \xrightarrow{P^5} N_{1 \rightarrow 2}^1 \xrightarrow{P^5} N_{1 \rightarrow 2}^2 \xrightarrow{P^5} \dots \xrightarrow{P^5} N_{1 \rightarrow 2}^{10} \xrightarrow{P^5} N^2, \\ N^2 &\xrightarrow{P^4} N^2 \xrightarrow{P^4} N_{2 \rightarrow 1}^0 \xrightarrow{P^5} N_{2 \rightarrow 1}^1 \xrightarrow{P^4} N_{2 \rightarrow 1}^2 \xrightarrow{P^5} \dots \\ &\dots \xrightarrow{P^4} N_{2 \rightarrow 1}^8 \xrightarrow{P^6} N_{2 \rightarrow 1}^9 \xrightarrow{P^5} N_{2 \rightarrow 1}^{10} \xrightarrow{P^3} N^1, \end{aligned}$$

where the starting and target sections in each of the above covering relations are determined by the h -sets appearing in the relation.

The conditions to check the covering relation with one exit direction are discussed in Appendix A.4. Some details and technical data regarding computer-assisted verification of Lemma 10 are given in Appendix A.5 and the supplementary material [38].

The subsequent remarks explain the choices made in construction of sequences of covering relations in Lemma 10.

Remark 11. The largest (in absolute value) eigenvalues of $DP(a^1)$ and $DP^2(a^2)$ are approximately -1.77 and -2.57 , respectively. Hence the expansion rate in the unstable directions along the periodic orbits is relatively mild. It turns, that using higher iterates of Poincaré map make it easier to construct h-sets and to check covering relations. As the second iterate squares both expansion and contraction factors, the stronger hyperbolicity lets the dynamics to help us in rigorous validation of covering relations for P^2 or higher iterates — there are wider margins for unavoidable overestimation when compute rigorously the image of an h-set.

Remark 12. The particular choice of the third iteration when computing $P^3(a_{2 \rightarrow 1}^8)$ comes from the fact, that it was much easier to construct coordinate system for an h-set centred at the point $P^3(a_{2 \rightarrow 1}^8)$ than at $P^2(a_{2 \rightarrow 1}^8)$. This is due to the fact, that $P^2(a_{2 \rightarrow 1}^8)$ is close to the bend in the attractor, while $a_{2 \rightarrow 1}^9$ is already very close to the fixed point a^1 .

The Poincaré maps appearing in Lemma 10 (and their iterates), despite being denoted by the same symbol P , in fact are maps between different sections, which are defined as some hyperplanes. In the definitions of sections Θ^i , $\Theta_{1 \rightarrow 2}^i$ and $\Theta_{2 \rightarrow 1}^i$ we do not specify the *crossing direction* – intersections in both directions are possible. Thus, a^1 , the fixed point for P becomes period-two point for $P_{\Theta^1 \rightarrow \Theta^1}$ and similarly a^2 is period-four point for $P_{\Theta^2 \rightarrow \Theta^2}$.

The maps P^j are compositions of j Poincaré maps between sections Θ^i , $\Theta_{1 \rightarrow 2}^i$ or $\Theta_{2 \rightarrow 1}^i$. Some of these sections are aligned so that they are almost parallel and close. For example, in $N^1 \xrightarrow{P^3} N_{1 \rightarrow 2}^0$, a trajectory starting from N^1 almost immediately cuts $\Theta_{1 \rightarrow 2}^0$ but this is not what we wanted. Then we follow the trajectory until it intersects twice $\Theta_{1 \rightarrow 2}^0$ in the vicinity of $a_{1 \rightarrow 2}^0$ — note, we allow intersections with section in both directions.

Proof of Theorem 1. Apply Theorem 6 to the transition matrix coding the covering relations from Lemma 10.

In particular, the existence of the “selected” periodic orbits u^1 and u^2 follows from the coverings $N^1 \xrightarrow{P^2} N^1$ and $N^2 \xrightarrow{P^4} N^2$. \square

4. The algorithm for rigorous bounds for solutions of (8)

The goal of this section is to describe the algorithm for rigorous integration of (4). Our algorithm stems from the *method of self-consistent bounds*, which was introduced in [46] and later developed in [47,49,48,51]. In the present paper we propose some significant modifications to the method. The new algorithm combines the High Order Enclosure method [30] with the dissipative enclosure from [51]. The other important novelty is the application of the automatic differentiation [18,14] to the infinite lader of ODEs represented by (4), therefore the use of differential inclusions as in [51] has been avoided.

Parts of our presentations will be abstract and directly applicable to system (4), while some others related to the automatic differentiation will be tailored to (8).

4.1. The setting and abstract assumptions

We will write our system as

$$a'(t) = f(a(t)) \quad (9)$$

where $f : l_2 \supset \text{dom}(f) \rightarrow l_2$. By f_k we denote the k th component of f . We assume, that the vector field (9) is of the form

$$f_k(a) = -L_k a_k + N_k(a)$$

and we make standing assumptions **C1–C5** on f listed below. We assume that there are constants

$$q > 1, \quad p > 1$$

such that

C1 there exist constants $L^* \geq L_* > 0, K \geq 0$, such that

$$L_* k^p \leq L_k \leq L^* k^p, \quad k > K,$$

C2 there exists r with

$$r < p,$$

such that for any $S > 0$ there exists $D = D(S, q)$ such that

$$|N_k(a)| \leq D k^r q^{-k} \quad \text{for } k \geq 1, a \in W_{q,S},$$

C3 for any $S > 0$ $f_k : W_{q,S} \rightarrow \mathbb{R}$ is continuous for $k = 1, 2, \dots$,

C4 for any $S > 0$ and $i, k = 1, 2, \dots$ the function $\frac{\partial N_k}{\partial a_i} : W_{q,S} \rightarrow \mathbb{R}$ is continuous and there exists $l_{\log} = l_{\log}(S, q)$, such that for all $k \in \mathbb{N}_+$ there holds

$$-L_k + \frac{1}{2} \sum_{i \in \mathbb{N}_+} \left| \sup_{a \in W_{q,S}} \frac{\partial N_k}{\partial a_i}(a) \right| + \frac{1}{2} \sum_{i \in \mathbb{N}_+} \left| \sup_{a \in W_{q,S}} \frac{\partial N_i}{\partial a_k}(a) \right| \leq l_{\log},$$

C5 for any $S > 0$, positive integer k and any partial derivative operator $D = \frac{\partial^n}{\partial a_{j_1} \dots \partial a_{j_n}}$ the function $DN : W_{q,S} \rightarrow \mathbb{R}$ is continuous.

In Section 5 we will show that conditions **C1–C5** are satisfied for the system (8).

The assumption **C2** contains two kinds of conditions; an estimate of N_k on $W_{q,S}$ and the inequality $r < p$. The estimate is satisfied for any $N(u) = P(u, Du, \dots, D^m u)$, where P is a polynomial function with the constant $r \geq m$ depending on m and the degree of P . It might turn out, however, that $r \geq p$. This happens for example for the viscous Burgers equation or the Navier-Stokes equations with periodic boundary condition in 2D or 3D. This will be avoided if instead of demanding that $|a_k| \leq S q^{-|k|}$ we will consider wider class of sets defined by $|a_k| \leq \frac{S}{q^{|k| \cdot |k|^t}}$, where $t \in \mathbb{N}_+$ — see [53,48] for more details.

The constant l_{\log} in the assumption **C4** is an upper estimate for the logarithmic norm of $\pi_{\leq n} f$ on $\pi_{\leq n} E$ for all n — see [47–49]) and Appendix A.2. This estimate will be used to obtain a uniform bound for the Lipschitz constants of the flow induced by Galerkin projections, which will allow us for a nice convergence argument in the proof Theorem 18.

The assumption **C5** is necessary in order to define the Taylor method for (9). In the next subsections we will see that **C1+C2** implies the isolation property, which plays the crucial role in our algorithm — see Remark 19.

4.2. An outline of the algorithm

The algorithm uses the following data structure to store in a computer memory a class of subsets of $l_{2,q}$

```

type  GBound
{
     $m \geq 0$       : a natural number,
     $\tilde{a} \subset \mathbb{R}^m$  : an interval vector,
     $S \geq 0$       : a real number,
     $q > 1$        : a real number.
}

```

Interval vector \tilde{a} is considered as a subset of \mathbb{R}^m , which is Cartesian product of closed intervals. An object **GBound**(m, \tilde{a}, S, q) represents a set of real sequences $(a_k)_{k>0}$ with geometric-like decay of coefficients

$$a_k \in \begin{cases} \tilde{a}_k, & 1 \leq k \leq m \\ [-S, S] \cdot q^{-k}, & k > m \end{cases}.$$

In the sequel we will use often the following decomposition $E = X_E \oplus W_E$ of a set $E = \mathbf{GBound}(m, \tilde{a}, S, q)$, where

$$X_E = \pi_{\leq m} E, \quad W_E = \pi_{> m} E.$$

Definition 13. Let $N \subset l_2$ and $h > 0$. We say that the set $E \subset l_2$ is a *rough enclosure* for N and a time h , if $N \subset E$ and for any $a_0 \in N$ any solution of (9) with initial condition $a(0) = a_0$ is defined for $t \in [0, h]$ and $a([0, h]) \subset E$.

In other words E gives a priori bounds for solutions starting from N over a time interval $[0, h]$. For a set of initial conditions $N = \mathbf{GBound}(m, \tilde{a}_N, S_N, q)$ and a time step $h > 0$ the algorithm

1. computes the rough enclosure $E = \mathbf{GBound}(m, \tilde{a}_E, S_E, q)$ for the set N and the time step h (see Section 4.4.1 and Section 4.4.2) and
2. computes a set $M = \mathbf{GBound}(m, \tilde{a}_M, S_M, q) \subset E$ such that for $a \in N$ there holds $a(h) \in M$ (see Section 4.4.3).

Step 1 may fail, which means that the algorithm could not validate the existence of solutions over the time step h for all $a \in N$. In this case we can either restart algorithm with refined data (split the initial condition N or shorten the time step h) or stop the computation. On success, in step 2 we compute a bound M for trajectories after time step $a(h)$, $a \in N$, which is usually much tighter (in the sense of inclusion) than E .

4.3. Basic existence and convergence theorems justifying the correctness of the algorithm

We work under the standing assumptions **C1–C5**.

Lemma 14. For any $S > 0$ the vector field f is continuous on $W_{q,S}$.

Proof. Let us fix $a \in W_{q,S}$. We will prove the continuity of f at a . Let us fix $\epsilon > 0$ and let $\tilde{a} \in W_{q,S}$.

We have from conditions **C1** and **C2**

$$\begin{aligned} \|f(a) - f(\tilde{a})\| &\leq \|\pi_{\leq n} f(a) - \pi_{\leq n} f(\tilde{a})\| + \|\pi_{> n} f(a)\| + \|\pi_{> n} f(\tilde{a})\| \leq \\ &\leq \|\pi_{\leq n} f(a) - \pi_{\leq n} f(\tilde{a})\| + 2 \sum_{|k|>n} Dk^r q^{-k} + 2 \sum_{|k|>n} L^* k^p q^{-k}. \end{aligned}$$

Let n be big enough to have

$$2 \sum_{|k|>n} Dk^r q^{-k} + 2 \sum_{|k|>n} L^* k^p q^{-k} < \epsilon/2.$$

From **C3** it follows that $\pi_{\leq n} f$ is continuous. Hence, there exists $\delta > 0$, such that

$$\text{if } \|a - \tilde{a}\| \leq \delta \text{ then } \|\pi_{\leq n} f(a) - \pi_{\leq n} f(\tilde{a})\| \leq \epsilon/2.$$

Gathering these estimates, we obtain

$$\text{if } \|a - \tilde{a}\| \leq \delta \text{ then } \|f(a) - f(\tilde{a})\| \leq \epsilon. \quad \square$$

Lemma 15. For $S > 0$ there holds

$$\lim_{n \rightarrow \infty} \sup_{a \in W_{q,S}} \|f(a) - f(\pi_{\leq n} a)\| = 0.$$

Proof. Let us fix $S > 0$ and $\epsilon > 0$. From Lemma 14 and the compactness of $W_{q,S}$ it follows that f is uniformly continuous on $W_{q,S}$. Therefore, there exists $\delta > 0$ such that

$$\text{if } \|a - \tilde{a}\| \leq \delta, \text{ then } \|f(a) - f(\tilde{a})\| \leq \epsilon. \tag{10}$$

Let n_0 be large enough so that $\|\pi_{> n} W_{q,S}\| < \delta$ for $n \geq n_0$. Hence, for $n \geq n_0$ it follows that

$$\|a - \pi_{\leq n} a\| = \|\pi_{> n} a\| < \delta, \quad \forall a \in W_{q,S}.$$

Combining this with (10) we obtain

$$\|f(a) - f(\pi_{\leq n} a)\| < \epsilon, \quad \forall a \in W_{q,S}.$$

This finishes the proof. \square

Definition 16. Let $N \subset W_{q,S}$ be a set of initial conditions for (9) and let us fix the time step $h > 0$. We call the set $E = \mathbf{GBound}(m, \tilde{a}_E, S_E, q) = X_E \oplus W_E$ a *First Order Enclosure* (FOE) for N over the time step h if the following conditions are satisfied:

$$N \subset E, \tag{11}$$

$$\pi_{\leq m} N + [0, h](f_1, \dots, f_m)(E) \subset \text{int}_{\leq m} X_E, \tag{12}$$

$$a_k f_k(a) < 0 \quad \text{for } a \in E, |a_k| = S_E q^{-k}, k > m. \tag{13}$$

The next lemma shows that assumptions **C1–C3** guarantee that a FOE of the form $E = \mathbf{GBound}(m, \tilde{a}_E, S_E, q)$ always exists for $N = W_{q,S}$ provided the time step h is small enough. In [53] this was done for polynomial bounds of the form $W = \left\{ a \mid |a_k| \leq \frac{S}{|k|^r} \right\}$. Later, in Theorem 18, we will prove that FOE is indeed a rough enclosure.

Lemma 17. *If $N \subset W_{q,S}$, then there exist $h > 0$ such that the set*

$$E = \mathbf{GBound}(m, \tilde{a}_E, S_E, q) := W_{q,2S}$$

is a FOE for N over the time step h for (9).

Proof. Let us set $S_E = 2S$ and

$$E = W_{q,2S}.$$

Let $D = D(E)$ be the constant from **C2**. Take a natural number

$$m > \max \left\{ K, \left(\frac{D}{L_* S_E} \right)^{\frac{1}{p-r}} \right\}.$$

The set E can be seen as $\mathbf{GBound}(m, \tilde{a}, S_E, q) = X_E \oplus W_E$, where

$$\tilde{a}_k = [-S_E q^{-k}, S_E q^k], \quad k = 1, \dots, m.$$

By the construction of E we have $N \subset E$ and therefore (11) is satisfied.

By **C1** and **C2**, for $k > m$ and $|a_k| = S_E q^{-k}$ there holds

$$\begin{aligned} a_k f_k(a) &= a_k(-L_k a_k + N_k(a)) \leq \\ &-L_k a_k^2 + |a_k N_k(a)| \leq -L_* k^p S_E^2 q^{-2k} + S_E D k^r q^{-2k} \leq \\ &S_E q^{-2k} k^r (-L_* S_E k^{p-r} + D) < \\ &S_E q^{-2k} k^r (-L_* S_E m^{p-r} + D). \end{aligned}$$

The constant m is chosen so that $-L_* S_E m^{p-r} + D < 0$ which implies $a_k f_k(a) < 0$ for $k > m$ and $|a_k| = S_E q^{-k}$. Therefore (13) is satisfied on E .

There remains to show that (12) is satisfied for sufficiently small $h > 0$. Since E is compact, by C3 the set $(f_1, \dots, f_m)(E)$ is bounded. Therefore, we can find $h > 0$ small enough such that $|hf_k(a)| < Sq^{-k}$ for $a \in E$ and $k = 1, \dots, m$. Then, for $a \in N$ and $k = 1, \dots, m$ we have

$$|a_k + [0, h]f_k(E)| < Sq^{-k} + Sq^{-k} = S_Eq^{-k}$$

and thus (12) holds true. \square

Theorem 18. *If $N \subset W_{q,S}$ and $E = \mathbf{GBound}(m, \tilde{a}_E, S_E, q) = W_{q,2S}$ is a FOE for N over the time step h , then for any $\hat{a} \in N$ there exists $a : [0, h] \rightarrow E$, a solution of (9), such that $a(0) = \hat{a}$, i.e. E is a rough enclosure for N and the time step h . This solution is unique under requirement that $a([0, h]) \subset E$.*

Moreover, for any two solutions $\hat{a}, \bar{a} : [0, h] \rightarrow E$ there holds

$$\|\hat{a}(t) - \bar{a}(t)\| \leq e^{h \log(2S,q)} \|\hat{a}(0) - \bar{a}(0)\|, \quad t \in [0, h]. \tag{14}$$

Proof. We will show that Galerkin projections of (9) with initial condition in N converge to the unique solution of the full system and we will also obtain (14) in the process.

The n th Galerkin projection of (9) can be written as follows

$$a' = \pi_{\leq n} f(a), \quad a(0) = \hat{a} \in \pi_{\leq n} l_{2,q}. \tag{15}$$

Observe that by assumptions C1, C3 and C4 the right hand side of (15) is locally Lipschitz, hence the solution to (15) is unique. Let $\varphi_n(t, a)$ be a local flow induced by (15).

In the sequel we will assume that $n \geq m$.

Step 1. We will show that $\pi_{\leq n} E$ is a priori bound for solutions of (15) with initial conditions in $\pi_{\leq n} N$, i.e. $\varphi_n(t, a)$ is defined for $t \in [0, h]$ and $a \in \pi_{\leq n} N$ and

$$\varphi_n([0, h], \pi_{\leq n} N) \subset \pi_{\leq n} E. \tag{16}$$

Indeed, since $\pi_{\leq n} E \subset E$ and E is a FOE it follows, that

$$\pi_{\leq m} N + [0, h](f_1, \dots, f_m)(\pi_{\leq n} E) \subset \text{int}_{\leq m} E, \tag{17}$$

$$a_k f_k(a) < 0 \quad \text{for } a \in \pi_{\leq n} E, \quad |a_k| = S_E q^{-k}, \quad n \geq k > m. \tag{18}$$

From (18) it follows that while $\varphi_n(t, a) \in \pi_{\leq n} E$, then for $k = m + 1, \dots, n$ the trajectory cannot reach the part of the boundary $\partial_{\leq n} E$ defined by $|a_k| = S_E q^{-k}$, because it is “repulsive” due to $\frac{d|a_k|}{dt} < 0$. The condition (17) implies that the boundary $\partial_{\leq m} E$ in the k th direction, $k = 1, \dots, m$, cannot be reached for $t \leq h$, because it is simply too far. This completes the proof of (16).

Step 2. We will show that for any $\hat{a} \in N$ the function $t \mapsto \varphi_n(t, \pi_{\leq n} \hat{a})$ converges uniformly on $[0, h]$.

From Lemma 15 it follows that

$$\delta_n = \sup_{a \in E} \|\pi_{\leq n} f(a) - \pi_{\leq n} f(\pi_{\leq n} a)\| \rightarrow 0, \quad n \rightarrow \infty.$$

Let $l = l_{\log}(2S, q)$ be a constant from condition **C4**. Using Theorem 32 (see Appendix A.2) and the Gersgorin Theorem it is easy to see that l is an upper estimate for the logarithmic norm of $\pi_{\leq n} f$ on $\pi_{\leq n} E$ for all n .

Therefore we can estimate the difference between two Galerkin projections as follows. Let us take $n_1 > n$ and let $\widehat{a}, \bar{a} \in N$.

Observe that $y(t) = \pi_{\leq n} \varphi_{n_1}(t, \pi_{\leq n_1} \bar{a})$ satisfies for $t \in [0, h]$

$$\begin{aligned} y'(t) &= \pi_{\leq n} f(\varphi_{n_1}(t, \pi_{\leq n_1} \bar{a})) \\ &= \pi_{\leq n} f(y(t)) + \pi_{\leq n} f(\varphi_{n_1}(t, \pi_{\leq n_1} \bar{a})) - \pi_{\leq n} f(\pi_{\leq n} \varphi_{n_1}(t, \pi_{\leq n_1} \bar{a})) \\ &= \pi_{\leq n} f(y(t)) + \delta(t), \quad |\delta(t)| \leq \delta_n. \end{aligned} \tag{19}$$

Therefore, from Lemma 34 (see Appendix A.2) we obtain for $t \in [0, h]$

$$\begin{aligned} \|\varphi_n(t, \pi_{\leq n} \widehat{a}) - \varphi_{n_1}(t, \pi_{\leq n} \bar{a})\| &\leq \tag{20} \\ \|\varphi_n(t, \pi_{\leq n} \widehat{a}) - \pi_{\leq n} \varphi_{n_1}(t, \pi_{\leq n} \bar{a})\| + \|\pi_{> n} \varphi_{n_1}(t, \pi_{\leq n} \bar{a})\| &\leq \\ e^{lt} \|\pi_{\leq n} \widehat{a} - \pi_{\leq n} \bar{a}\| + \delta_n \frac{e^{lt} - 1}{l} + \left(\sum_{|k|>n} S_E^2 q^{-2k} \right)^{1/2}. & \end{aligned}$$

Now if we take $\widehat{a} = \bar{a}$, then

$$\|\varphi_n(t, \pi_{\leq n} \widehat{a}) - \varphi_{n_1}(t, \pi_{\leq n} \widehat{a})\| \leq \delta_n \frac{e^{lt} - 1}{l} + \left(\sum_{|k|>n} S_E^2 q^{-2k} \right)^{1/2} \rightarrow 0,$$

when $n \rightarrow \infty$ uniformly for $(t, \widehat{a}) \in [0, h] \times N$.

Therefore we can define

$$\varphi(t, a) = \lim_{n \rightarrow \infty} \varphi_n(t, \pi_{\leq n} a).$$

Obviously $\varphi(t, a)$ is continuous on $[0, h] \times N$ as the limit of uniformly convergent sequence of continuous functions.

Step 3. We will show, that $a(t) = \varphi(t, a)$ is a solution of (9). For each $n_1 \geq n$ we have

$$\pi_{\leq n} \varphi_{n_1}(t, \pi_{\leq n_1} a) = \pi_{\leq n} a + \int_0^t \pi_{\leq n} f(\varphi_{n_1}(s, \pi_{\leq n_1} a)) ds. \tag{21}$$

From Lemma 14 and compactness of E it follows that f is uniformly continuous on E . This combined with the uniform convergence of φ_n implies that $f(\varphi_{n_1}(s, \pi_{\leq n_1} a))$ converges uniformly with $n_1 \rightarrow \infty$ to $f(\varphi(s, a))$ for $(s, a) \in [0, h] \times N$. Therefore the integral on the rhs of (21) converges and we obtain for any n

$$\pi_{\leq n} \varphi(t, a) = \pi_{\leq n} a + \int_0^t \pi_{\leq n} f(\varphi(s, a)) ds$$

which implies that

$$\varphi(t, a) = a + \int_0^t f(\varphi(s, a)) ds.$$

By differentiation we see that $t \rightarrow \varphi(t, a)$ is a solution of (9). Observe that when passing to the limit with $n, n_1 \rightarrow \infty$ in (20) we obtain that for $\widehat{a}, \bar{a} \in N$ and $t \in [0, h]$ the condition (14) is satisfied.

Step 4. It remains to show that if $a : [0, t_{\max}] \rightarrow E$ is a solution of (9) with $a(0) \in N$, then for $t \in [0, \min(t_{\max}, h)]$ holds $a(t) = \varphi(t, a(0))$.

By decreasing h if necessary without any loss of generality we can assume that $h = t_{\max}$. Observe, that for any n the function $y(t) = \pi_{\leq n} a(t)$ satisfies (19) for $t \in [0, h]$. Therefore, from Lemma 34 we obtain that for $t \in [0, h]$ there holds

$$\begin{aligned} & \|\varphi(t, a(0)) - a(t)\| \leq \\ & \|\varphi(t, a(0)) - \varphi_n(t, \pi_{\leq n} a(0))\| + \|\varphi_n(t, \pi_{\leq n} a(0)) - \pi_{\leq n} a(t)\| + \|\pi_{> n} a(t)\| \leq \\ & \|\varphi(t, a(0)) - \varphi_n(t, \pi_{\leq n} a(0))\| + \delta_n \frac{e^{lt} - 1}{l} + \left(\sum_{|k|>n} S_E^2 q^{-2k} \right)^{1/2}. \end{aligned}$$

Now passing to the limit $n \rightarrow \infty$ we obtain that $a(t) = \varphi(t, a(0))$ for $t \in [0, h]$. \square

Remark 19. Observe that (13) used to obtain (18) was of fundamental importance in the above proof, as it provides us uniform a priori bounds for high modes for all Galerkin projections. This is the isolation property mentioned in the Introduction.

Remark 20. Theorem 18 implies that for fixed $q > 1$ we have a family of continuous local semiflows defined on $W_{q,S}$ and parameterized by $S \in \mathbb{R}_+$. We can consider a ‘union’ of these semiflows to obtain a local semiflow φ on entire $l_{2,q}$. However, from the above reasoning does not follow, that φ is continuous with respect to x . This can be easily obtained from the classical existence results for KS equation recalled in Appendix A.1.

Theorem 21. For any $\bar{a} \in l_{2,q}$ there exists $a : [0, \infty) \rightarrow l_{2,q}$ a unique solution of (9) with initial condition $a(0) = \bar{a}$. The induced semiflow (semigroup) φ is continuous with respect to t and \bar{a} .

Let us emphasise, that our computer assisted proof does not depend neither on Theorem 21 nor Theorem 30. We just need a local continuous semiflow defined on $W_{q,S}$, where $S > 0$ is a computable constant, which contains finite number of sets generated by the computer program. These include selected subsets of certain Poincaré sections and enclosure of their forward trajectories unless they reach another Poincaré section.

4.4. Algorithm, more details but still on the abstract level

In the sequel we will use the following notation. For a smooth function $F: I \subset \mathbb{R} \rightarrow \mathbb{R}$ by $F^{[i]}(t)$ we denote the i th Taylor coefficient of F at t . We also write $F^{[i]}$ if $t = 0 \in I$. The same convention applies to vector valued functions $F: I \rightarrow l_2$.

In Section 5.2 we will show that the components $a_k(t)$ of the solutions to the KS equation are smooth and we give an algorithm for computation of $a_k^{[i]}$ by means of automatic differentiation. It is important to emphasise that $a_k^{[i]}$ depends on $a_k^{[j]}$ for $j = 0, \dots, i - 1$, hence it can be expressed in terms of $a_k^{[0]}$, only. Here we assume this knowledge.

4.4.1. High Order Enclosure in finite dimension

In [30] the authors propose an efficient algorithm for computation of a priori bound on the set of trajectories over a time step. It is called the High Order Enclosure method as it relies on high order Taylor expansion of the solutions. In what follows we will recall the algorithm from [6,30]. In the next section we will show, how it can be adopted to the case of infinite dimensional dissipative systems.

Consider an initial value problem for a finite dimensional ODE

$$x' = g(x), \quad x(0) \in X \subset U, \tag{22}$$

where $g \in C^{d+1}(U, \mathbb{R}^n)$, $U \subset \mathbb{R}^n$ is open and d is a natural number. Let us fix $\tilde{h} > 0$ – this is a trial time step for the numerical method used to integrate the system (22). In practice, \tilde{h} is generated by another algorithm — a time step predictor. For the reasoning given here it is irrelevant, what this value come from.

Theorem 22. [6, Theorem 3] Let $\varepsilon > 0$ be a tolerance per time step and let us set $R = [-\varepsilon, \varepsilon]^n$. Let

$$P = \sum_{i=0}^d X^{[i]}[0, \tilde{h}]^i$$

and define $E = P + R$. If

$$Z := E^{[d+1]}[0, \tilde{h}]^{d+1} \subset \text{int } R \tag{23}$$

then for all $x(0) \in X$ the solution to (22) exists for all $t \in [0, \tilde{h}]$ and $x([0, \tilde{h}]) \subset P + Z$.

The condition (23) may fail due to three reasons. First and the most common case is when the inclusion (23) is not satisfied. Given that $0 \in \text{int } R$, we can always find $h < \tilde{h}$ such that

$$Z := E^{[d+1]}[0, h]^{d+1} \subset \text{int } R$$

and the set $P + Z$ is a high order enclosure for (22) with (usually slightly) decreased time step $h < \tilde{h}$.

Second reason for failure is the situation, when the Taylor coefficient $E^{[d+1]}$ cannot be computed. This may happen for instance, when division by zero in interval arithmetics occurs or

the set E is out of the domain of g . In this case we have to decrease the trial step \tilde{h} and repeat the entire procedure (recomputation of $X^{[i]}$ required for P is not necessary). If the number of such repetitions exceeds some specified maximal value, the algorithm stops and returns **Failure**. This does not mean that the trajectories do not exist — the algorithm simply could not validate requested condition.

Third, and less common situation is when we cannot compute Taylor $X^{[i]}$ used to define P . In this case we cannot proceed and the algorithm returns **Failure**. Some higher level decisions, such as changing the order, splitting the initial condition X or just giving up have to be made.

4.4.2. High Order Enclosure in infinite dimension

The construction of the rough enclosure as FOE given in the proof of Lemma 17 was not used in our computer assisted proof, because it produces too much overestimation and we also prefer to keep the dimension m constant along the trajectory. Therefore, a bit more sophisticated routines mixing together a dissipative enclosure from [53] and high order enclosure from [6,30] are necessary.

Let us consider now an infinite dimensional dissipative PDE of the form (9). Let $N = \mathbf{GBound}(m, \tilde{a}, S, q)$ be a set of initial conditions for (9). Assume $\tilde{h} > 0$ is a trial time step given from prediction. In the algorithm given below we expect that the dynamics on modes a_k , for $k > m$ is highly dissipative. The construction of a-priori bound consists of the following steps.

1. Compute

$$P_E = \sum_{i=0}^d \pi_{\leq m} N^{[i]}[0, \tilde{h}]^i.$$

The algorithm for computation of $N^{[i]}$ for the KS equation will be given in Section 5.2.

2. Set $R = [-\varepsilon, \varepsilon]^m$ and predict an enclosure on the main modes of the form $X_E = P_E + R$.
3. Set $S_E = S$ and $h = \tilde{h}$.
4. Define the set $E = X_E \oplus W_E$, where

$$(W_E)_k = \begin{cases} 0 & k \leq m \\ [-S_E, S_E]q^{-k} & k > m \end{cases}$$

5. Check if the vector field is pointing inwards E_k for $k > m$. This is the same condition (13) as in FOE. If not satisfied, then we slightly enlarge S_E and go to step 4.
6. Check inclusion

$$Z := \pi_{\leq m} E^{[d+1]}[0, \tilde{h}]^{d+1} \subset \text{int } R$$

If the above fails, then find $h < \tilde{h}$ such that

$$Z := \pi_{\leq m} E^{[d+1]}[0, h]^{d+1} \subset \text{int } R$$

holds true.

The above algorithm may fail by the same three reasons, as in the finite dimensional case. In addition, the loop in which we enlarge the constant S_E may exceed specified maximal limit. If the algorithm does not fail, the computed set $(P_E + Z) \oplus W_E$ is a priori bound for $a([0, h])$ for $a \in N$, as guaranteed by steps 5 and 6.

4.4.3. Computation of tight enclosure for $a(h)$

Once we have a rough enclosure for $a([0, h])$, next we want to compute tight bounds for $a(h)$. Let $N = \mathbf{GBound}(m, \tilde{a}_N, S_N, q)$ be a set of initial conditions for (9). Fix $h > 0$ and assume that $E = \mathbf{GBound}(m, \tilde{a}_E, S_E, q)$ is a rough enclosure for N over the time step h . We will show, how we can bound the set $\{a(h) : a \in N\}$.

Let us fix an order of the Taylor method $d > 0$. On the main modes $k \leq m$ we will bound $a_k(h)$ by an explicit formula. For $a \in N$ by the Taylor theorem we have

$$a_k(h) \in \sum_{i=0}^d a_k^{[i]} h^i + R_k,$$

where

$$R_k = \left\{ a_k^{[d+1]} [0, h]^{d+1} \mid a \in E \right\}.$$

The bound on the tail $k > m$ will be computed using infinite set of differential inequalities. Take $D = D(E)$ from C2. By assumptions C1–C2 we have

$$a'_k(t) \leq -L_k a_k + Dk^r q^{-k} = -L_k a_k + N_k^+,$$

where $N_k^+ := Dk^r q^{-k}$. Then, for $t \in [0, h]$ and $k > m$ there holds

$$\begin{aligned} a_k(t) &\leq \frac{N_k^+}{L_k} + \left(a_k(0) - \frac{N_k^+}{L_k} \right) e^{-L_k t} \leq \\ &\frac{Dk^r q^{-k}}{L_* k^p} + \left(S_E q^{-k} - \frac{Dk^r q^{-k}}{L_* k^p} \right) e^{-L_k t} = \\ &\frac{D}{L_* k^{p-r}} q^{-k} + \left(S_E q^{-k} - \frac{D}{L_* k^{p-r}} q^{-k} \right) e^{-L_k t} = \\ &q^{-k} \left(\frac{D}{L_* k^{p-r}} + \left(S_E - \frac{D}{L_* k^{p-r}} \right) e^{-L_k t} \right) \leq \\ &q^{-k} \left(\frac{D}{L_* (m+1)^{p-r}} + S_E e^{-L_*(m+1)^s t} \right). \end{aligned}$$

In a similar way we can bound $a_k(t)$ from below. To sum up, we proved the following

Lemma 23. Let $N = \mathbf{GBound}(m, \tilde{a}_N, S_N, q)$ be a set of initial conditions for (9). Fix $h > 0$ and assume that $E = \mathbf{GBound}(m, \tilde{a}_E, S_E, q)$ is a rough enclosure for N over the time step h .

Then for $a \in N$ and $k > m$ there holds

$$|a_k(h)| \leq Sq^{-k},$$

where

$$S = \min \left\{ S_E, \frac{D}{L_*(m+1)^{p-r}} + S_E e^{-L_*(m+1)^p h} \right\}.$$

We have shown, that given an initial condition represented as **GBound** with decay $q > 1$ we can always find a rough enclosure for sufficiently small time step $h > 0$ and compute a bound on the trajectories over the time step h as **GBound** with the same geometric decay q . We use higher order time-derivatives $a_k^{[i]}$ which do not belong to $l_{2,q}$ to bound finitely many leading modes $a_k(h)$ by an explicit formula.

5. Estimates specific for the KS equation

In Section 4 we outlined an algorithm for computing rigorous enclosures on the set of trajectories in a class of vector fields satisfying assumptions **C1–C5**. We also assume that a routine that allows to compute Taylor coefficients $a_k^{[i]}$ of solutions to (9) is provided. In this section we show, that these requirements are satisfied for the Kuramoto-Sivashinsky equation (8). Analogous requirements for sets of the form $W = \left\{ |a_k| \leq \frac{C}{|k|^s} \right\}$ for a class of dissipative PDEs (including also KS equation) has been proved in [48].

5.1. Conditions C1–C5

The system (8) splits into linear and nonlinear part as follows

$$\begin{aligned} a'_k(t) &= -L_k a_k + N_k(a), \\ L_k &= k^2(1 - \nu k^2), \\ N_k(a) &= -k \sum_{n=1}^{k-1} a_n a_{k-n} + 2k \sum_{n=1}^{\infty} a_n a_{n+k}. \end{aligned}$$

C1: It is easily satisfied as L_k is a polynomial in k of degree $s = 4$.

C2: For $a \in W_{q,S}$ we have

$$\begin{aligned} |N_k(a)| &\leq k \sum_{n=1}^{k-1} S q^{-n} S q^{n-k} + 2k \sum_{n=1}^{\infty} S q^{-n} S q^{-n-k} = \\ &= k(k-1) S^2 q^{-k} + 2k S^2 q^{-k} (q^2 - 1)^{-1} \leq \\ &= 2k^2 q^{-k} S^2 \left(1 + (q^2 - 1)^{-1} \right). \end{aligned}$$

Thus, the condition **C2** is satisfied with $r = 2$ and $D = 2S^2 (1 + (q^2 - 1)^{-1})$.

C3: We have to show that f_k is continuous on l_2 . The formula for f_k splits into polynomial of degree 2, which is clearly continuous, and an infinite sum

$$a \rightarrow 2k \sum_{n=1}^{\infty} a_n a_{n+k}$$

which also is continuous as the inner product in l_2 composed with the k -shift of coordinates.

C4: Observe that

$$\frac{\partial N_k}{\partial a_i} = \begin{cases} -2ka_{k-i} + 2ka_{k+i}, & \text{if } i < k; \\ 2ka_{k+i}, & \text{otherwise.} \end{cases}$$

We estimate

$$\left| \frac{\partial N_k}{\partial u_i}(W_{q,S}) \right| \leq 4kSq^{-|k-i|}.$$

Hence

$$\begin{aligned} -L_k + \frac{1}{2} \sum_{i \in \mathbb{N}_+} \left| \sup_{a \in W_{q,S}} \frac{\partial N_k}{\partial a_i}(a) \right| + \frac{1}{2} \sum_{i \in \mathbb{N}_+} \left| \sup_{a \in W_{q,S}} \frac{\partial N_i}{\partial a_k}(a) \right| &\leq \\ -L_k + 4kS \left(\sum_{i \in \mathbb{N}_+} q^{-|i-k|} \right) &\leq \\ -L_k + 8kS \left(\sum_{i \in \mathbb{N}} q^{-i} \right) &= -L_k + \frac{8kS}{1-q}. \end{aligned}$$

Since $L_k \approx \nu|k|^4$, we see that the above expression is bounded from above and l_{\log} exists.

C5: It is obvious because N is a quadratic polynomial.

5.2. Coefficients $a_k^{[i]}$ via automatic differentiation

Let us fix an initial condition $a(0) \in l_{2,q}$. In this section prove that the Taylor coefficients $a^{[i]}$ for the solutions to (8) do exist. We propose an iterative scheme for computation bounds for $a^{[i]}$. These estimates will be also bounds for the Taylor coefficients for all Galerkin projections with sufficiently high projection dimension. The proposed scheme is a special case of the general algorithm from [43] tailored to the KS equation.

We will see that if $a(0) = a^{[0]} \in l_{2,q}$ then $a^{[i]}, i > 0$ cannot belong to $l_{2,q}$. Therefore, we will construct a strictly decreasing sequence $q = q_0 > q_1 > \dots > 1$ and represent $a^{[i]}$ as geometric bounds in l_{2,q_i} .

The nonlinear part of (8)

$$N_k(t, a) = -k \sum_{n=1}^{k-1} a_n(t)a_{k-n}(t) + 2k \sum_{n=1}^{\infty} a_n(t)a_{n+k}(t)$$

splits into a finite sum

$$E_k(t, a) = - \sum_{n=1}^{k-1} a_n(t)a_{k-n}(t) + 2 \sum_{n=1}^m a_n(t)a_{n+k}(t) \tag{24}$$

and an infinite sum

$$I_k(t, a) = \sum_{n=m+1}^{\infty} a_n(t)a_{n+k}(t).$$

Using this notation, the k th component of the vector field along the trajectory $a(t)$ reads

$$F_k(t) = k \left(k(1 - vk^2)a_k(t) + E_k(t, a(t)) + 2I_k(t, a(t)) \right).$$

The coefficients $a_k^{[i]}$ can be computed by the following iterative scheme

$$a_k^{[i+1]} = \frac{1}{i+1} F_k^{[i]}, \tag{25}$$

provided the rhs of (25) makes sense.

In what follows we will show, how we can bound $E_k^{[i]}, I_k^{[i]}$ (where the Taylor coefficients are taken with respect to time variable) and in consequence, to compute bounds on $F_k^{[i]}$.

The estimates on the infinite part $I_k^{[i]}$ are derived in the following lemma.

Lemma 24. Assume that for $j = 0, 1, \dots, i - 1$ we have already computed $a^{[j]}$ and they are represented as geometric bounds $a^{[j]} \in \mathbf{GBound}(m, \tilde{a}^{[j]}, S_j, q_j)$. Then for $k \geq 1$ there holds that

$$\left| I_k^{[i]} \right| \leq D_I q^{-k},$$

where

$$q = \min\{q_0, q_1, \dots, q_i\},$$

$$D_I = \left(\sum_{j=0}^i S_j S_{i-j} \right) \left((q^2 - 1)q^{2m} \right)^{-1}.$$

Proof. All the terms a_n and a_{n+k} which appear in the summation are bounded by geometric series. Therefore we have

$$\left| I_k^{[i]} \right| \leq \sum_{n=m+1}^{\infty} \sum_{j=0}^i \left| a_n^{[j]} a_{n+k}^{[i-j]} \right| \leq \sum_{n=m+1}^{\infty} \sum_{j=0}^i S_j q_j^{-n} S_{i-j} q_{i-j}^{-n-k}$$

$$\begin{aligned} &\leq q^{-k} \left(\sum_{j=0}^i S_j S_{i-j} \right) \sum_{n=m+1}^{\infty} q^{-2n} \\ &= q^{-k} \left(\sum_{j=0}^i S_j S_{i-j} \right) \left((q^2 - 1)q^{2m} \right)^{-1}. \quad \square \end{aligned}$$

E_k is handled by the following three lemmas.

Lemma 25. For $k = 1, \dots, 2m$ there holds that

$$\begin{aligned} E_k^{[i]} = 2 \sum_{j=0}^i \left(\sum_{n=1}^{\lfloor (k-1)/2 \rfloor} a_n^{[j]} \left(-a_{k-n}^{[i-j]} + a_{n+k}^{[i-j]} \right) + \sum_{n=\lceil k/2 \rceil}^m a_n^{[j]} a_{n+k}^{[i-j]} \right) \\ - \epsilon(k) \left(2 \sum_{j=0}^{\lfloor (i-1)/2 \rfloor} a_{k/2}^{[j]} a_{k/2}^{[i-j]} + \epsilon(i) \left(a_{k/2}^{[i/2]} \right)^2 \right), \end{aligned}$$

where $\epsilon(k) = 1$ if k is an even number and $\epsilon(k) = 0$, otherwise.

Proof. This is a direct application of the Leibniz rule to a slightly factorized form of (24). This factorization is performed in order to reduce the computational cost and the wrapping effect in evaluation of this expression in interval arithmetics. \square

Lemma 26. Assume that for $j = 0, 1, \dots, i - 1$ we have already computed $a^{[i]}$ and they are represented as geometric bounds $a^{[i]} \in \mathbf{GBound}(m, \tilde{a}^{[i]}, S_i, q_i)$. Then for $k > 2m$ there holds that

$$\left| E_k^{[i]} \right| \leq D_1 k q^{-k},$$

where

$$\begin{aligned} q &= \min\{q_0, q_1, \dots, q_i\}, \\ D_1 &= \frac{2}{2m+1} \sum_{j=0}^i \sum_{n=1}^m \left(q_{i-j}^n + q_{i-j}^{-n} \right) S_{i-j} \left| a_n^{[j]} \right| + \sum_{j=0}^i S_j S_{i-j}. \end{aligned}$$

Proof. For $k > 2m$ the formula for $E_k^{[i]}$ splits into two components $E_k^{[i]} = \Sigma_1 + \Sigma_2$, where

$$\begin{aligned} \Sigma_1 &= 2 \sum_{j=0}^i \sum_{n=1}^m a_n^{[j]} \left(-a_{k-n}^{[i-j]} + a_{n+k}^{[i-j]} \right), \\ \Sigma_2 &= - \sum_{j=0}^i \sum_{n=m+1}^{k-1-m} a_n^{[j]} a_{k-n}^{[i-j]}. \end{aligned}$$

In Σ_1 the indices $k - n$ and $n + k$ are greater than m . Therefore

$$\begin{aligned}
 |\Sigma_1| &\leq 2 \sum_{n=1}^m \sum_{j=0}^i |a_n^{[j]}| \left(S_{i-j} q_{i-j}^{n-k} + S_{i-j} q_{i-j}^{-n-k} \right) \\
 &\leq q^{-k} \left(2 \sum_{n=1}^m \sum_{j=0}^i (q_{i-j}^n + q_{i-j}^{-n}) S_{i-j} |a_n^{[j]}| \right) \\
 &\leq k q^{-k} \left(\frac{2}{2m+1} \sum_{n=1}^m \sum_{j=0}^i (q_{i-j}^n + q_{i-j}^{-n}) S_{i-j} |a_n^{[j]}| \right).
 \end{aligned}$$

In the sum Σ_2 we have $n > m$ and $k - n > m$. Therefore

$$|\Sigma_2| \leq \sum_{n=m+1}^{k-1-m} \sum_{j=0}^i S_j q_j^{-n} S_{i-j} q_{i-j}^{n-k}.$$

For $k > n$ and $i = 0, \dots, j$ there holds that

$$q_j^{-n} q_{i-j}^{n-k} = q^{-k} \left(\frac{q}{q_j} \right)^n \left(\frac{q}{q_{i-j}} \right)^{k-n} \leq q^{-k}.$$

This gives us an estimate

$$|\Sigma_2| \leq q^{-k} \sum_{n=m+1}^{k-1-m} \sum_{j=0}^i S_j S_{i-j} \leq k q^{-k} \left(\sum_{j=0}^i S_j S_{i-j} \right)$$

Gathering together bounds on Σ_1, Σ_2 we obtain the constant D_1 . \square

Lemma 27. Assume that for $j = 0, 1, \dots, i - 1$ we have already computed $a^{[j]}$ and they are represented as geometric bounds $a^{[j]} \in \mathbf{GBound}(m, \tilde{a}^{[j]}, S_i, q_i)$. Then for $m < k \leq 2m$ there holds that

$$|E_k^{[i]}| \leq D_2 k q^{-k},$$

where

$$\begin{aligned}
 q &= \min\{q_0, q_1, \dots, q_i\}, \\
 D_2 &= \max_{k=m+1, \dots, 2m} \frac{1}{k} |E_k^{[i]}| q^k.
 \end{aligned}$$

Proof. $E_k^{[i]}$ is given by an explicit finite sum. Thus we can bound D_2 in finite computation. \square

All the above considerations lead to the following estimate on the i th Taylor coefficient of F .

Lemma 28. Let $q = \min\{q_0, \dots, q_i\}$ and fix $\delta \in (1, q)$. Put

$$L = \begin{cases} (m + 1)^4 \delta^{-(m+1)} & \text{if } m > 4/\ln \delta, \\ \left(\frac{4}{e \ln \delta}\right)^4 & \text{otherwise.} \end{cases}$$

Let D_I, D_1, D_2 be constants computed as in Lemma 24, Lemma 26 and Lemma 27, respectively, and put $D = \max\{D_1, D_2\}$. Then for $k > m$ there holds that

$$\left|F_k^{[i]}\right| \leq S(q/\delta)^{-k},$$

where

$$S = L \left(|v - (m + 1)^{-2}| S_i + (m + 1)^{-3} D_I + (m + 1)^{-2} D \right).$$

Proof. $F_k^{[i]}$ is given by

$$F_k^{[i]} = k^2(1 - vk^2)a_k^{[i]} + k(E_k^{[i]} + 2I_k^{[i]}).$$

For $k > m$ we have

$$\begin{aligned} \left|F_k^{[i]}\right| &\leq k^2|1 - vk^2|S_i q_i^{-k} + kD_I q^{-k} + k^2 D q^{-k} \\ &\leq \left(k^2|1 - vk^2|S_i + kD_I + k^2 D\right) q^{-k} \\ &= \left(|v - k^{-2}|S_i + k^{-3} D_I + k^{-2} D\right) k^4 q^{-k} \\ &\leq \left(|v - (m + 1)^{-2}|S_i + (m + 1)^{-3} D_I + (m + 1)^{-2} D\right) k^4 q^{-k}. \end{aligned}$$

We would like to find a constant L such that

$$k^4 q^{-k} \leq L(q/\delta)^{-k}$$

for $k > m$. Thus L must satisfy $L \geq k^4 \delta^{-k}$ for $k > m$. The function $k \rightarrow k^4 \delta^{-k}$ attains the global maximum at $k = 4/\ln \delta$ and it is strictly decreasing to zero after reaching maximum at this point. If $m > 4/\ln \delta$ then L can be taken as $L = (m + 1)^4 \delta^{-m-1}$. Otherwise we have to take the global maximum $L = \left(\frac{4}{e \ln \delta}\right)^4$. \square

Observe that the above derivations are valid also for all n -dimensional Galerkin projections with $n \geq m$. Indeed in this situation we have

$$\pi_{\leq n} \mathbf{GBound}(m, \tilde{a}, S_i, q_i) \subset \mathbf{GBound}(m, \tilde{a}, S_i, q_i).$$

From the above Lemmas we immediately obtain the following

Remark 29. Let $a : [0, h] \rightarrow l_{2,q}$ be a solution of the KS equation (or its n -dimensional Galerkin projection with $n \geq m$), such that for $t \in [0, h]$ there holds $a(t) \in \mathbf{GBound}(m, \tilde{a}, S_0, q_0 = q)$. Then for any $i \in \mathbb{N}$ there exists $a_k^{[i]}(t)$ for $t \in [0, h]$. Moreover, it satisfies

$$|a_k^{[i]}(t)| \leq \frac{S(q/\delta)^{-k}}{i + 1}, \quad k > m$$

where S and δ are as in Lemma 28. The formulas for $k \leq m$ follow from Lemmas 24 and 25.

Appendix A

A.1. The existence and uniqueness results for Kuramoto-Sivashinsky PDE

We recall results from Section III.4.1 in [35]. There KS equation is written as

$$\frac{\partial v}{\partial t} + v \frac{\partial^4 v}{\partial x^4} + \frac{\partial^2 v}{\partial x^2} + v \frac{\partial v}{\partial x} = 0 \tag{A.1}$$

on $\Omega = [-L/2, L/2]$ subject to periodic boundary conditions

$$\frac{\partial^j v}{\partial x^j} \left(t, -\frac{L}{2} \right) = \frac{\partial^j v}{\partial x^j} \left(t, \frac{L}{2} \right), \quad j = 0, 1, 2, 3. \tag{A.2}$$

Let

$$H = \dot{L}(\Omega) = \left\{ v \in L^2(\Omega) \left| \int_{-L/2}^{L/2} v(s) ds = 0 \right. \right\},$$

$$V = H_{per}^2(\Omega) \cap H.$$

H is endowed with L^2 scalar product (denoted (\cdot, \cdot)), while V is endowed with the scalar product

$$((v, w)) = \int_{-L/2}^{L/2} \frac{\partial^2 v}{\partial x^2} \frac{\partial^2 w}{\partial x^2} dx.$$

Let A be an unbounded self-adjoint operator in H with the domain $D(A) = H_{per}^4 \cap H$ given by

$$Av = \left(\frac{\partial}{\partial x} \right)^4 v.$$

Theorem 30. For $v_0 \in H$ there exists a unique solution v of (A.1), (A.2) with $v(0) = v_0$,

$$v \in C([0, T]; H) \cap L^2(0, T; V), \quad \forall T > 0.$$

Furthermore, for $t > 0$ the function v is analytic in t with values in $D(A)$ and the mapping

$$v_0 \mapsto v(t)$$

is continuous from H into $D(A)$.

A.2. Logarithmic norms and estimates for Lipschitz constant of the flow induced by ODEs

The goal is to recall some results about the Lipschitz constant of the flow induced by ODEs based on the logarithmic norms.

Definition 31. [14, Def. I.10.4] Let Q be a square matrix; we call

$$\mu(Q) = \lim_{h>0, h \rightarrow 0} \frac{\|I + hQ\| - 1}{h}$$

the logarithmic norm of Q .

Theorem 32. [14, Th. I.10.5] The logarithmic norm is obtained by the following formulas

- for Euclidean norm

$$\mu(Q) = \text{the largest eigenvalue of } 1/2(Q + Q^T),$$

- for max norm $\|x\|_\infty = \max_k |x_k|$

$$\mu(Q) = \max_k \left(q_{kk} + \sum_{i \neq k} |q_{ki}| \right),$$

- for norm $\|x\|_1 = \sum_k |x_k|$

$$\mu(Q) = \max_i \left(q_{ii} + \sum_{k \neq i} |q_{ki}| \right).$$

Consider now the differential equation

$$x' = f(x), \quad f \in C^1(\mathbb{R}^n). \tag{A.3}$$

Let $\varphi(t, x_0)$ denote the solution of equation (A.3) with the initial condition $x(0) = x_0$. By $\|x\|$ we denote a fixed arbitrary norm in \mathbb{R}^n .

The following theorem was proved in [14, Th. I.10.6] (for a non-autonomous ODE, here we restrict ourselves to the autonomous case only and we use a different notation).

Theorem 33. Let $y : [0, T] \rightarrow \mathbb{R}^n$ be a piecewise C^1 function and $\varphi(\cdot, x_0)$ be defined for $t \in [0, T]$. Suppose that the following estimates hold:

$$\mu \left(\frac{\partial f}{\partial x}(\eta) \right) \leq l(t), \quad \text{for } \eta \in [y(t), \varphi(t, x_0)],$$

$$\left\| \frac{dy}{dt}(t) - f(y(t)) \right\| \leq \delta(t),$$

where $[y(t), \varphi(t, x_0)]$ denotes the line segment connecting $y(t)$ and $\varphi(t, x_0)$.

Then for $0 \leq t \leq T$ there holds

$$\|\varphi(t, x_0) - y(t)\| \leq e^{L(t)} \left(\|y(0) - x_0\| + \int_0^t e^{-L(s)} \delta(s) ds \right),$$

where $L(t) = \int_0^t l(s) ds$.

From the above theorem one easily derives the following.

Lemma 34. Let $y : [0, T] \rightarrow \mathbb{R}^n$ be a piecewise C^1 function and $\varphi(\cdot, x_0)$ be defined for $t \in [0, T]$. Suppose that Z is a convex set such that the following estimates hold:

$$\begin{aligned} y([0, T]), \varphi([0, T], x_0) &\in Z, \\ \mu \left(\frac{\partial f}{\partial x}(\eta) \right) &\leq l, \quad \text{for } \eta \in Z, \\ \left\| \frac{dy}{dt}(t) - f(y(t)) \right\| &\leq \delta. \end{aligned}$$

Then for $0 \leq t \leq T$ there holds

$$\|\varphi(t, x_0) - y(t)\| \leq e^{lt} \|y(0) - x_0\| + \delta \frac{e^{lt} - 1}{l}, \quad \text{if } l \neq 0.$$

For $l = 0$, there holds

$$\|\varphi(t, x_0) - y(t)\| \leq \|y(0) - x_0\| + \delta t.$$

A.3. The Poincaré transition maps between sections

Let Π_1 and Π_2 be affine hyperplanes in l_2 given by $v_i(x) + c_i = 0$ for $i = 1, 2$, where v_i the continuous linear forms. In order to define the Poincaré transition map between Π_1 and Π_2 for the local semiflow $\varphi(t, x)$ induced by (9) we need the following

- continuity of $\varphi(t, x)$ with respect to both variables,
- transversality of the trajectories $t \mapsto \varphi(t, x)$, when intersecting Π_i . For the ODE case it is enough to have $v_i(f(y)) \neq 0$ for y in the neighbourhood of intersection of $\varphi(t, x)$ with section Π_i .

In our case we will demand that analogous conditions hold in some $W_{q,S}$ close to the starting and the target sections.

We assume the standing assumptions **C1–C5** and that there exists m , such that all forms v_i are defined in terms of first m variables, i.e. $v_i(a) = v_i(\pi_{\leq m} a)$.

Definition 35. We define a *global section* as a hyperplane:

$$\Pi = \{x \in l_2 \mid v(x) + c = 0\},$$

where $v : l_2 \rightarrow \mathbb{R}$, such that $v(a) = v(\pi_{\leq m} a)$ and $c \in \mathbb{R}$.

Any convex and bounded subset $U \subset \Pi$ is called a *local section*.

Let $W_Z = \tilde{a} \oplus \pi_{> m} W_{q,S}$, where \tilde{a} is an open convex subset of $\pi_{\leq m} l_2$.

Definition 36. A local section Z is said to be *transversal* in W_Z if

$$W_Z \cap Z = Z, \quad W_Z = W_{Z,-} \cup Z \cup W_{Z,+},$$

where

$$W_{Z,-} = \{x \in W_Z \mid v(x) + c < 0\}, \quad W_{Z,+} = \{x \in W_Z \mid v(x) + c > 0\},$$

satisfying the condition

$$v(f(x)) > 0, \quad \forall x \in W_Z. \tag{A.4}$$

We will refer to (A.4) as the *transversality condition*.

We have the following easy lemma.

Lemma 37. Let Z be a local transversal section in W_Z for (9) and let $N \subset W_{q,S}$ for some $S > 0$. Assume that there exist $t_1, t_2 \in \mathbb{R}$, $t_1 < t_2$, such that the following conditions hold for all $x \in N$:

$$\varphi((t_1, t_2), x) \subset W_Z, \quad \varphi(t_1, x) \in W_{Z,-} \quad \text{and} \quad \varphi(t_2, x) \in W_{Z,+}.$$

Then, for each $x_0 \in N$, there exists unique $t_Z(x_0) \in (t_1, t_2)$, such that $\varphi(t_Z(x_0), x_0) \in Z$. Also, $t_Z : N \rightarrow [t_1, t_2]$ is continuous.

Using the above lemma we can define a map $P_{Z_1 \rightarrow Z_2} : Z_1 \rightarrow Z_2$ for two transversal local sections Z_1 and Z_2 , by

$$P_{Z_1 \rightarrow Z_2}(x) = \varphi(t_{Z_2}(x), x).$$

A.4. Representation of h -sets and computer-assisted verification of covering relations with one exit direction

Algorithms for computer-assisted verification of covering relations for finite dimensional maps with arbitrary unstable dimension are presented in [42]. Although they can be directly extended to the case of infinite dimensional systems, for self-consistency of the article we provide here a special case, when h -sets have one exit (nominally unstable) direction, i.e. $u(N) = 1$ in Definition 2.

Let us fix $q > 1$. In order to treat the system (8) rigorously on a computer we define a data structure which represents h -sets in $l_{2,q}$:

$$\begin{aligned}
 &\text{type } \mathbf{HSet} \\
 &\{ \\
 &\quad m \geq 0 \quad : \text{ a natural number,} \\
 &\quad S \geq 0 \quad : \text{ a real number,} \\
 &\quad x \in \mathbb{R}^m \quad : \text{ a vector,} \\
 &\quad A \in \mathbb{R}^{m^2} \quad : \text{ an invertible matrix.} \\
 &\}
 \end{aligned} \tag{A.5}$$

An h -set $N = \mathbf{HSet}(m_N, S_N, x_N, A_N)$ represented by the above data structure is given by

$$N = c_N \left([-1, 1]^{m_N} \oplus W \right),$$

where c_N is an invertible affine map

$$c_N \left((a_k)_{k=1}^\infty \right) = \left(x_N + A_N \cdot (a_1, \dots, a_{m_N})^T, a_{m_N+1}, a_{m_N+2}, \dots \right)$$

and

$$W = [-1, 1] \cdot \left(\underbrace{0, 0, \dots, 0}_{m_N}, S_N q^{-(m_N+1)}, S_N q^{-(m_N+2)}, \dots \right).$$

Thus, the projection of N onto m leading coordinates is a parallelepiped centred at x_N with the shape matrix A_N . The remaining coordinates are not affected by c_N and are stored as a geometrically decaying tail.

Since we assumed $u(N) = 1$ in Definition 2, the tail of N is always given by

$$\overline{T_N} = [-1, 1] \cdot \left(0, \underbrace{1, 1, \dots, 1}_{m_N-1}, S_N q^{-(m_N+1)}, S_N q^{-(m_N+2)}, \dots \right).$$

Let N_0 and N_1 be h -sets contained in $l_{2,q}$ and represented as in (A.5). Let $f : |N_0| \rightarrow l_2$ be continuous and compact map and such that $f(|N_0|) \subset W_{q,S}$, for some $S > 0$. Assume that we have a routine which for given $A \subset |N|$ computes $B \subset W_{q,S_1}$, also represented by (A.5), such that $(c_{N_1}^{-1} \circ f \circ c_{N_0})(A) \subset B$. Put

$$\begin{aligned}
 B &= \left(c_{N_1}^{-1} \circ f \circ c_{N_0} \right) ([-1, 1] \oplus \overline{T_{N_0}}), \\
 B_l &= \left(c_{N_1}^{-1} \circ f \circ c_{N_0} \right) (\{-1\} \oplus \overline{T_{N_0}}), \\
 B_r &= \left(c_{N_1}^{-1} \circ f \circ c_{N_0} \right) (\{1\} \oplus \overline{T_{N_0}}).
 \end{aligned}
 \tag{A.6}$$

Now, checking $N_0 \xrightarrow{f} N_1$ reduces to a finite set of inequalities which must be satisfied. These are:

- $\pi_i(B) \subset (-1, 1)$ for $i = 2, \dots, m$,
- $S(B) < S(N_1)$, where S is a positive constant in h -set representation (A.5),
- either $\pi_1(B_l) < -1$ and $\pi_1(B_r) > 1$ or $\pi_1(B_l) > 1$ and $\pi_1(B_r) < -1$.

The two cases in the last condition depend on whether the mapping f changes or not the orientation along the exit direction. It is easy to see that

$$\begin{aligned}
 H(t, \cdot) &= (1-t) \left(c_{N_1}^{-1} \circ f \circ c_{N_0} \right) + tL, \\
 L((a_k)_{k=1}^\infty) &= (\pm 2a_1, 0, 0 \dots),
 \end{aligned}$$

are a homotopy and a linear map required by Definition 3, where the sign in L depends on whether f preserves or not the orientation in the exit direction.

A.5. Technical data

The source code of the C++11 program that realises the computer-assisted proof of Lemma 10 is available at [38]. Below we list our choices of some parameters of the algorithms.

- All h -sets that appear in Lemma 10 are represented as a data structure (A.5) with the constant $m = 15$.
- We set $d = 4$ as the order of the Taylor method (Section 4.4.3) for rigorous integration of (8).
- High-Order Enclosure (Section 4.4.2) with $d = 4$ acts on $m = 11$ number of modes.

Verification of covering relation $N_0 \xrightarrow{f} N_1$ requires computation of three images of f — see (A.6). In Lemma 10 we verify 26 covering relations which means, that we have to check $78 = 3 \cdot 26$ inclusions.

We run the program which checks all the inequalities required for covering relations listed in Lemma 10 on a computer equipped with 64 physical cores (128 threads) Intel(R) Xeon(R) CPU E7-8867 v4 @ 2.40 GHz processors. The program finished after 40 minutes, which is the CPU time needed for the longest integration in $N_{2 \rightarrow 1}^8 \xrightarrow{P^6} N_{2 \rightarrow 1}^9$.

The algorithm for rigorous integration of the KS equation is a part of the CAPD library [3]. We tested the program with CAPD version 5.0.59 and the C++11 compiler from gcc-5.2 suite.

References

- [1] G. Arioli, H. Koch, Integration of dissipative partial differential equations: a case study, *SIAM J. Appl. Dyn. Syst.* 9 (2010) 1119–1133.

- [2] G. Arioli, P. Zgliczyński, Symbolic dynamics for the Hénon–Heiles Hamiltonian on the critical energy level, *J. Differ. Equ.* 171 (2001) 173–202.
- [3] CAPD - Computer Assisted Proofs in Dynamics, a package for rigorous numerics, <http://capd.ii.uj.edu.pl>.
- [4] P. Collet, J.-P. Eckmann, H. Epstein, J. Stubbe, Analyticity for the Kuramoto–Sivashinsky equation, *Physica D* 67 (1993) 321–326.
- [5] F. Christiansen, P. Cvitanovic, V. Putkaradze, Spatiotemporal chaos in terms of unstable recurrent patterns, *Nonlinearity* 10 (1997) 55–70.
- [6] G.F. Corliss, R. Rihm, Validating an a priori enclosure using high-order Taylor series, in: G. Alefeld, A. Frommer (Eds.), *Scientific Computing, Computer Arithmetics, and Validated Numerics*, Akademie Verlag, Berlin, 1996, pp. 228–238.
- [7] R. Easton, Isolating blocks and symbolic dynamics, *J. Differ. Equ.* 17 (1975) 96–118.
- [8] R. Easton, Homoclinic phenomena in Hamiltonian systems with several degrees of freedom, *J. Differ. Equ.* 29 (1978) 241–252.
- [9] M.J. Feigenbaum, Quantitative universality for a class of nonlinear transformations, *J. Stat. Phys.* 19 (1978) 25–52.
- [10] J. Figueras, R. de la Llave, Numerical computations and computer assisted proofs of periodic orbits of the Kuramoto–Sivashinsky equation, *SIAM J. Appl. Dyn. Syst.* 16 (2) (2017) 834–852.
- [11] J. Figueras, M. Gameiro, J.P. Lessard, R. de la Llave, A framework for the numerical computation and a posteriori verification of invariant objects of evolution equations, *SIAM J. Appl. Dyn. Syst.* 16 (2) (2017) 1070–1088.
- [12] C. Foias, B. Nicolaenko, G. Sell, R. Temam, Inertial manifolds for the Kuramoto–Sivashinsky equation and an estimate of their lowest dimension, *J. Math. Pures Appl.* 67 (1988) 197–226.
- [13] C. Foias, R. Temam, Gevrey class regularity for the solutions of the Navier–Stokes equations, *J. Funct. Anal.* 87 (2) (1989) 359–369.
- [14] E. Hairer, S.P. Nørsett, G. Wanner, *Solving Ordinary Differential Equations I, Nonstiff Problems*, Springer-Verlag, Berlin, Heidelberg, 1987.
- [15] Z. Galias, P. Zgliczyński, Computer assisted proof of chaos in the Lorenz system, *Physica D* 115 (1998) 165–188.
- [16] M. Gameiro, J.-P. Lessard, A posteriori verification of invariant objects of evolution equations: periodic orbits in the Kuramoto–Sivashinsky PDE, *SIAM J. Appl. Dyn. Syst.* 16 (1) (2017) 687–728.
- [17] A. Granas, J. Dugundji, *Fixed Point Theory*, Springer Monographs in Mathematics, 2010.
- [18] A. Griewank, *Evaluating Derivatives: Principles and Techniques of Algorithmic Differentiation*, *Frontiers in Applied Mathematics*, vol. 19, SIAM, 2000.
- [19] J. Guckenheimer, P. Holmes, *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Springer Verlag, Berlin, Heidelberg, New York, 1983.
- [20] J. Hyman, B. Nicolaenko, The Kuramoto–Sivashinsky equation: a bridge between PDEs and dynamical systems, *Physica D* 18 (1986) 113–126.
- [21] M.E. Johnson, M. Jolly, I. Kevrekidis, The Oseberg transition: visualization of the global bifurcations for the Kuramoto–Sivashinsky equation, *Int. J. Bifurc. Chaos* 11 (2001) 1–18.
- [22] M. Jolly, I. Kevrekidis, E. Titi, Approximate inertial manifolds for the Kuramoto–Sivashinsky equation: analysis and computations, *Physica D* 44 (1990) 38–60.
- [23] Y. Kuramoto, T. Suzuki, Persistent propagation of concentration waves in dissipative media far from thermal equilibrium, *Prog. Theor. Phys.* 55 (1976) 365.
- [24] R.J. Lohner, Computation of guaranteed enclosures for the solutions of ordinary initial and boundary value problems, in: J.R. Cash, I. Gladwell (Eds.), *Computational Ordinary Differential Equations*, Clarendon Press, Oxford, 1992.
- [25] J. Mawhin, Leray–Schauder degree: a half century of extensions and applications, *Topol. Methods Nonlinear Anal.* 14 (1999) 105–228.
- [26] R.E. Moore, *Methods and Applications of Interval Analysis*, SIAM, Philadelphia, 1979.
- [27] K. Mischaikow, M. Mrozek, Chaos in the Lorenz equations: a computer assisted proof, *Bull. Am. Meteorol. Soc.* 32 (1995) 66–72.
- [28] K. Mischaikow, M. Mrozek, Isolating neighborhoods and chaos, *Jpn. J. Ind. Appl. Math.* 12 (1995) 205–236.
- [29] J. Moser, *Stable and Random Motions in Dynamical Systems*, Princeton Univ. Press, 1973.
- [30] N.S. Nedialkov, K.R. Jackson, J.D. Pryce, An effective high-order interval method for validating existence and uniqueness of the solution of an IVP for an ODE, *Reliab. Comput.* 7 (6) (2001) 449–465.
- [31] B. Nicolaenko, B. Scheurer, R. Temam, Some global dynamical properties of the Kuramoto–Sivashinsky equations: nonlinear stability and attractors, *Physica D* 16 (1985) 155–183.
- [32] T. Rage, A. Neumaier, C. Schlier, Rigorous verification of chaos in a molecular model, *Phys. Rev. E* 50 (1994) 2682–2688.

- [33] G.I. Sivashinsky, Nonlinear analysis of hydrodynamical instability in laminar flames – 1. Derivation of basic equations, *Acta Astron.* 4 (11–12) (1977) 1177–1206.
- [34] Y.S. Smaylis, D.T. Papageorgiou, Predicting chaos for infinite-dimensional dynamical systems: the Kuramoto-Sivashinsky equation, a case study, *Proc. Natl. Acad. Sci. USA* 88 (24) (1991) 11129–11132.
- [35] R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, Applied Mathematical Sciences, vol. 68, Springer, 1997.
- [36] W. Tucker, Lorenz attractor exists, *C. R. Acad. Sci. Paris, Ser. I* 328 (1999) 1197–1202.
- [37] W. Tucker, A rigorous ODE solver and Smale’s 14th problem, *Found. Comput. Math.* 2 (1) (2002) 53–117.
- [38] D. Wilczak, <http://www.ii.uj.edu.pl/~{w}ilczak>, personal homepage.
- [39] D. Wilczak, Abundance of heteroclinic and homoclinic orbits for the hyperchaotic Rössler system, *Discrete Contin. Dyn. Syst., Ser. B* 11 (4) (2009) 1039–1055.
- [40] D. Wilczak, S. Serrano, R. Barrio, Coexistence and dynamical connections between hyperchaos and chaos in the 4D Rössler system: a computer-assisted proof, *SIAM J. Appl. Dyn. Syst.* 15 (1) (2016) 356–390.
- [41] D. Wilczak, P. Zgliczyński, Heteroclinic connections between periodic orbits in planar restricted circular three body problem - a computer assisted proof, *Commun. Math. Phys.* 234 (1) (2003) 37–75.
- [42] D. Wilczak, P. Zgliczyński, Topological method for symmetric periodic orbits for maps with a reversing symmetry, *Discrete Contin. Dyn. Syst., Ser. A* 17 (3) (2007) 629–652.
- [43] D. Wilczak, P. Zgliczyński, A rigorous C^1 -algorithm for integration of dissipative PDEs based on automatic differentiation and the Taylor method, in preparation.
- [44] P. Zgliczyński, Fixed point index for iterations of maps, topological horseshoe and chaos, *Topol. Methods Nonlinear Anal.* 8 (1996) 169–177.
- [45] P. Zgliczyński, Sharkovskii’s theorem for multidimensional perturbations of one-dimensional maps II, *Topol. Methods Nonlinear Anal.* 14 (1999) 169–182.
- [46] P. Zgliczyński, K. Mischaikow, Rigorous numerics for partial differential equations: the Kuramoto-Sivashinsky equation, *Found. Comput. Math.* 1 (2001) 255–288.
- [47] P. Zgliczyński, Attracting fixed points for the Kuramoto-Sivashinsky equation - a computer assisted proof, *SIAM J. Appl. Dyn. Syst.* 1 (2002) 215–235.
- [48] P. Zgliczyński, Trapping regions and an ODE-type proof of an existence and uniqueness for Navier-Stokes equations with periodic boundary conditions on the plane, *Univ. Iagel. Acta Math.* 41 (2003) 89–113.
- [49] P. Zgliczyński, On smooth dependence on initial conditions for dissipative PDEs, an ODE-type approach, *J. Differ. Equ.* 195 (2) (2003) 271–283.
- [50] P. Zgliczyński, Computer assisted proof of chaos in the Rössler equations and in the Hénon map, *Nonlinearity* 10 (1) (1997) 243–252.
- [51] P. Zgliczyński, Rigorous numerics for dissipative partial differential equations II. Periodic orbit for the Kuramoto-Sivashinsky PDE - a computer assisted proof, *Found. Comput. Math.* 4 (2004) 157–185.
- [52] P. Zgliczyński, M. Gidea, Covering relations for multidimensional dynamical systems, *J. Differ. Equ.* 202 (1) (2004) 33–58.
- [53] P. Zgliczyński, Rigorous numerics for dissipative PDEs III. An effective algorithm for rigorous integration of dissipative PDEs, *Topol. Methods Nonlinear Anal.* 36 (2010) 197–262.