

Do food trichomes occur in *Pinguicula* (Lentibulariaceae) flowers?

Krzysztof Lustofin¹, Piotr Świątek², Piotr Stolarczyk³, Vitor F. O. Miranda⁴ and Bartosz J. Płachno^{1,*}

¹Department of Plant Cytology and Embryology, Institute of Botany, Faculty of Biology, Jagiellonian University in Kraków, 9 Gronostajowa Street, 30-387 Cracow, Poland, ²Institute of Biology, Biotechnology and Environmental Protection, Faculty of Natural Sciences, University of Silesia in Katowice, 9 Bankowa Street, 40-007 Katowice, Poland, ³Unit of Botany and Plant Physiology, Institute of Plant Biology and Biotechnology, University of Agriculture in Kraków, 29 Listopada 54 Street, 31-425 Kraków, Poland and ⁴Universidade Estadual Paulista (Unesp), Faculdade de Ciências Agrárias e Veterinárias, Jaboticabal, Departamento de Biologia Aplicada à Agropecuária, São Paulo, Brazil

*For correspondence. E-mail bartosz.plachno@uj.edu.pl

Received: 8 April 2020 Returned for revision: 1 June 2020 Editorial decision: 16 June 2020 Accepted: 19 June 2020
Electronically published: 27 June 2020

- **Background and Aims** Floral food bodies (including edible trichomes) are a form of floral reward for pollinators. This type of nutritive reward has been recorded in several angiosperm families: Annonaceae, Araceae, Calycanthaceae, Eupomatiaceae, Himantandraceae, Nymphaeaceae, Orchidaceae, Pandanaceae and Winteraceae. Although these bodies are very diverse in their structure, their cells contain food material: starch grains, protein bodies or lipid droplets. In *Pinguicula* flowers, there are numerous multicellular clavate trichomes. Previous authors have proposed that these trichomes in the *Pinguicula* flower play the role of ‘futterhaare’ (‘feeding hairs’) and are eaten by pollinators. The main aim of this study was to investigate whether the floral non-glandular trichomes of *Pinguicula* contain food reserves and thus are a reward for pollinators. The trichomes from the *Pinguicula* groups, which differ in their taxonomy (species from the subgenera: *Temnoceras*, *Pinguicula* and *Isoloba*) as well as the types of their pollinators (butterflies/flies and bees/hummingbirds), were examined. Thus, it was determined whether there are any connections between the occurrence of food trichomes and phylogeny position or pollination biology. Additionally, we determined the phylogenetic history of edible trichomes and pollinator evolution in the *Pinguicula* species.
- **Methods** The species that were sampled were: *Pinguicula moctezumae*, *P. esseriana*, *P. moranensis*, *P. emarginata*, *P. rectifolia*, *P. mesophytica*, *P. hemiepiphytica*, *P. agnata*, *P. albida*, *P. ibarrae*, *P. martinezii*, *P. filifolia*, *P. gigantea*, *P. lusitanica*, *P. alpina* and *P. vulgaris*. Light microscopy, histochemistry, and scanning and transmission electron microscopy were used to address our aims with a phylogenetic perspective based on *matK*/*trnK* DNA sequences.
- **Key Results** No accumulation of protein bodies or lipid droplets was recorded in the floral non-glandular trichomes of any of the analysed species. Starch grains occurred in the cells of the trichomes of the bee-/fly-pollinated species: *P. agnata*, *P. albida*, *P. ibarrae*, *P. martinezii*, *P. filifolia* and *P. gigantea*, but not in *P. alpina* or *P. vulgaris*. Moreover, starch grains were not recorded in the cells of the trichomes of the *Pinguicula* species that have long spurs, which are pollinated by Lepidoptera (*P. moctezumae*, *P. esseriana*, *P. moranensis*, *P. emarginata* and *P. rectifolia*) or birds (*P. mesophytica* and *P. hemiepiphytica*), or in species with a small and whitish corolla that self-pollinate (*P. lusitanica*). The results on the occurrence of edible trichomes and pollinator syndromes were mapped onto a phylogenetic reconstruction of the genus.
- **Conclusion** Floral non-glandular trichomes play the role of edible trichomes in some *Pinguicula* species (*P. agnata*, *P. albida*, *P. ibarrae*, *P. martinezii*, *P. filifolia* and *P. gigantea*), which are mainly classified as bee-pollinated species that had originated from Central and South America. It seems that in the *Pinguicula* that are pollinated by other pollinator groups (Lepidoptera and hummingbirds), the non-glandular trichomes in the flowers play a role other than that of a floral reward for their pollinators. Edible trichomes are symplesiomorphic for the *Pinguicula* species, and thus do not support a monophyletic group such as a synapomorphy. Nevertheless, edible trichomes are derived and are possibly a specialization for fly and bee pollinators by acting as a food reward for these visitors.

Key words: Butterworts, carnivorous plants, floral micro-morphology, food hairs, Lentibulariaceae, trichome structure, *Pinguicula*, spur, trichomes.

INTRODUCTION

Plants offer various floral rewards for pollinators that can be divided into two groups: non-nutritive rewards (e.g. nest materials, a place of shelter, heat sources, substances for production of sexual attractants or places for mating) and nutritive

rewards (e.g. brood site, floral sweet tissue, stigmatic secretion or fatty oils) (Simpson and Neff, 1981). The most common floral nutritive rewards are nectar and pollen (Faegri and van der Pijl, 1979; Nicolson *et al.*, 2007). However, some species produce food bodies (including edible trichomes) that

are eaten by their pollinators. The cells of these structures are rich with starch grains, protein bodies or oil droplets (Young, 1986; Thien *et al.*, 2009; for orchids, see Pansarin and Maciel, 2017 and references therein). Food bodies have been recorded in several unrelated plant families: Annonaceae, Araceae, Calycanthaceae, Eupomatiaceae, Himantandraceae, Orchidaceae, Pandanaceae, Nymphaeaceae and Winteraceae (e.g. Faegri and van der Pijl, 1979; Rickson, 1979; Cox, 1982; Young, 1986; Davies *et al.*, 2002; Thien *et al.*, 2009; Endress, 2010; Pansarin and Maciel, 2017). Thus, this type of reward occurs in both evolutionarily old families via beetle pollination (Annonaceae, Calycanthaceae, Eupomatiaceae, Himantandraceae, Nymphaeaceae and Winteraceae; see Endress, 2010) as well as in the more evolutionarily derived family Orchidaceae, which now represents an evolutionary pick of diversity. Floral food bodies can be divided into two major groups: the first (which occurs, for example, in the older lineages of angiosperms, Endress, 2010) – the outgrowths (or tips) of the carpels, stamens, staminodes and tepals; and the second – the epidermal edible trichomes. These trichomes have been particularly well analysed in Orchidaceae and they were found to have evolved independently in this family about five times (genera: *Cyanaeorchis*, *Dendrobium*, *Eria*, *Maxillaria* and *Polystachya*; Pansarin and Maciel, 2017). In orchids, they are very diverse in their structure and morphology as well as in the storage of nutritive material in their cells (e.g. Davies *et al.*, 2002; Davies and Turner, 2004; Pansarin and Maciel, 2017).

Pinguicula is a monophyletic genus within the Lentibulariaceae L. family (Jobson *et al.*, 2003; Müller *et al.*, 2004; Fleischmann and Rocca, 2018) and is among the Lamiales (Schäferhoff *et al.*, 2010; Chase *et al.*, 2016) and contains about 96 species. *Pinguicula* are well known for their carnivory (e.g. Alcalá and Domínguez, 2003, 2005; Darnowski *et al.*, 2018; Heslop-Harrison, 1970; Heslop-Harrison and Heslop-Harrison, 1980; Vassilyev and Muravnik, 1988).

Pinguicula produce spurred zygomorphic flowers, which have nectar as a reward (Abrahamczyk *et al.*, 2017; Fleischmann and Rocca, 2018; Lustofin *et al.*, 2019). In *Pinguicula* flowers, there are numerous multicellular clavate trichomes at the base of the corolla – the throat; see Fig. 1A–I (Casper, 1966). Previous authors have proposed that these trichomes in the *Pinguicula* flower play the role of ‘futterhaare’ (‘feeding hairs’) and are eaten by their pollinators, or that some of them play the role of mimic pollen grains (see Fleischmann, 2016). Thus, the main aim of this study was to determine whether these trichomes of *Pinguicula* contain food reserves and thus may be a reward for potential pollinators. We selected species from the different clades, which are based on published phylogenetic proposals, within *Pinguicula* (members from three subgenera but focused on the Central American species) and also sampled species based on differences in their mating system. For this criterion, self- (i.e. a small flower with a whitish corolla) vs. outcross species (large, brightly coloured corollas, nectar guides and long spurs) were compared. Additionally, in our study, we considered the pollinator types (butterflies/fly and




FIG. 1. General morphology and micromorphology of the selected *Pinguicula* species that were examined. (A–D) General morphology and micromorphology of a *P. agnata* flower showing the entrance to the flower (En) with multicellular clavate slender trichomes (I), the throat (Th) with multicellular compact thick trichomes (II) in the front and two types of long and slender or short and compact non-glandular trichomes (III) that are located at the entrance to the spur (Sp); note the presence of a pistil (Ps) and a stamen (S) in the throat; scale bars = 2 mm, 2 mm, 1 mm and 1 mm, respectively. (E and F) Micromorphology of a *P. gigantea* flower; note the similar distribution and micromorphology of the non-glandular trichomes (I', II', III') compared with *P. agnata*; scale bars = 1 mm and 1 mm, respectively. (G and H) Micromorphology of the *P. rectifolia* throat with generative organs and many celled uniseriate slender non-glandular trichomes indicated by an acute apical cell that is located in the throat and basal part of the spur; scale bars = 1 mm and 300 μ m, respectively. (I) Micromorphology of the *P. hemiepiphytica* throat with long and slender multicellular non-glandular trichomes and a stamen; scale bar = 1 mm.

bees/hummingbirds). Fleischmann (2016) wrote that the clavate trichomes of *Pinguicula* are glandular, and therefore another task/aim was to determine whether these trichomes have the character of glands.

MATERIALS AND METHODS

Plant material

Seventeen taxa were sampled: *Pinguicula moctezumae* Zamudio & R.Z.Ortega, *P. esseriana* B.Kirchn., *P. moranensis* Kunth, *P. emarginata* Zamudio & Rzed., *P. rectifolia* Speta & F.Fuchs, *P. mesophytica* Zamudio, *P. hemiepiphytica* Zamudio & Rzed., *P. agnata* Casper, *P. albida* Wright ex Griseb., *P. ibarrae* Zamudio, *P. martinezii* Zamudio, *P. filifolia* C.Wright ex Griseb., *P. gigantea* Luhrs, *P. lusitanica* L., *P. alpina* L. and *P. vulgaris* L. [*P. vulgaris* subsp. *vulgaris* L. and *P. vulgaris*

L. subsp. *bicolor* (Wol.) Á. Löve & D. Löve]. For our study, we primarily used living material (see Table 1). However, histochemical studies were used by some authors (e.g. Hernández and Katinas, 2019) in the case of herbarium material in order to show storage material or glandular structures. Therefore, we also used herbarium material of *Pinguicula* from the Herbarium of the Institute of Botany (KRA).

Methods

The flowers were examined using light microscopy (LM), scanning electron microscopy (SEM) and transmission electron microscopy as described below. The material was fixed in a mixture of 2.5 or 5 % glutaraldehyde with 2.5 % formaldehyde in a 0.05 M cacodylate buffer (Sigma; pH 7.2) overnight or for several days, washed three times in a 0.1 M sodium cacodylate

TABLE 1. List of the *Pinguicula* species that were examined along with information regarding their infrageneric classification, the origin of the plant material and the type of pollinator for each species.

Species	Infrageneric classification	Material origin	Type of pollinator
<i>P. moctezumae</i> Zamudio & R.Z.Ortega	<i>Temnoceras</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: Mexico)	Lepidoptera (Abrahamczyk et al., 2017)
<i>P. rectifolia</i> Speta & F.Fuchs	<i>Temnoceras</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: Mexico)	Lepidoptera (flower's structure indicates that type of pollinator)
<i>P. moranensis</i> Kunth	<i>Temnoceras</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: near Santiago Juxtlahuaca, Oaxaca, Mexico 1851 m)	Lepidoptera (Villegas and Alcalá, 2018)
<i>P. emarginata</i> Zamudio & Rzed.	<i>Temnoceras</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: Mexico)	Lepidoptera (flower's structure indicates that type of pollinator)
<i>P. esseriana</i> B.Kirchn.	<i>Temnoceras</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: Mexico)	Lepidoptera (flower's structure indicates that type of pollinator)
<i>P. hemiepiphytica</i> Zamudio & Rzed	<i>Temnoceras</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: near Ixtlan de Juarez, Oaxaca, Mexico, 2209–2535 m.)	Most probably hummingbirds (Lampard et al., 2016)
<i>P. mesophytica</i> Zamudio	<i>Temnoceras</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: Cerro Miramundo, El Salvador)	Ornithophily is presumed: a watercolour showing a species of hummingbird visiting a plants of <i>Pinguicula mesophytica</i> was shown in Roccia et al. (2016)
<i>P. agnata</i> Casper	<i>Temnoceras</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: Mexico)	Diptera/Hymenoptera (flower's structure indicates that type of pollinator)
<i>P. gigantea</i> Luhrs	<i>Temnoceras</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: Mexico)	Diptera/Hymenoptera (Abrahamczyk et al., 2017)
<i>P. ibarrae</i> Zamudio	<i>Temnoceras</i>	Botanical Garden in Liberec	Diptera/Hymenoptera (flower's structure indicates that type of pollinator)
<i>P. martinezii</i> Zamudio	<i>Temnoceras</i>		Diptera/Hymenoptera (flower's structure indicates that type of pollinator)
<i>P. albida</i> Wright ex Griseb.	<i>Temnoceras</i>		Hymenoptera (Dominguez et al., 2014)
<i>P. filifolia</i> C.Wright ex Griseb.	<i>Temnoceras</i>		Hymenoptera (Dominguez et al., 2014)
<i>P. lusitanica</i> L.	<i>Isoloba</i>	Botanical Garden of Jagiellonian University in Cracow (collected from: Europa)	Diptera/Hymenoptera(?), self-pollination (Heslop-Harrison, 2004)
<i>P. alpina</i> L.	<i>Pinguicula</i>	Herbarium of Jagiellonian University in Cracow (collected from: Alps, Innsbruck, Austria; KRA 0299930)	Diptera/Hymenoptera (Molau, 1993; Nordin, 2015)
<i>P. vulgaris</i> subsp. <i>vulgaris</i> L.	<i>Pinguicula</i>	Herbarium of Jagiellonian University in Cracow (collected from: Małe Pieniny, Rezerwat Zaskalskie, Poland; KRA 71415)	Diptera/Hymenoptera (Molau, 1993)
<i>P. vulgaris</i> L. subsp. <i>bicolor</i> (Wol.) Á. Löve & D. Löve	<i>Pinguicula</i>	Herbarium of Jagiellonian University in Cracow (collected from: Dąbrowa Górnica, użytek ekologiczny 'Młaki and Pogoria I', Poland; KRA 0138573)	

buffer and post-fixed in a 1 % osmium tetroxide solution at room temperature for 1.5 h. Next, the material was treated as was previously described (Plachno *et al.*, 2017) and examined using a Hitachi H500 transmission electron microscope (Hitachi, Tokyo, Japan), which is housed at the University of Silesia in Katowice, at an accelerating voltage of 75 kV. The semi-thin sections (0.9–1.0 μm thick) that were prepared for LM were stained with aqueous methylene blue/azure II for 1–2 min (Humphrey and Pittman, 1974) and examined using Olympus BX60 and Nikon Eclipse E400 light microscopes to perform the general histology. The periodic acid–Schiff (PAS) reaction for LM (semi-thin sections) was also used to reveal the presence of insoluble polysaccharides (Wędzony, 1996), and Sudan Black B was used to detect the presence of lipids and cuticle material (Jensen, 1962).

Additionally, material that had been embedded in Technovit 7100 (Kulzer, Germany) was also examined. This material was fixed (as above), washed three times in a 0.1 M sodium cacodylate buffer, dehydrated in a graded ethanol series for 15 min at each concentration and kept overnight in absolute ethanol. Next, the samples were infiltrated for 1 h each in 3:1, 1:1 and 1:3 (v/v) mixtures of absolute ethanol and Technovit and then stored for 12 h in pure Technovit. The resin was polymerized by adding a hardener. The material was sectioned to 5 μm thickness using a rotary microtome (Microm, Adamas Instrumenten), stained with 0.1 % toluidine blue O and mounted in DPX (Sigma-Aldrich). The selected Technovit sections were stained with naphthol blue black (NBB) for total protein staining (Fisher, 1968; Mathe and Vieillescazes, 2002) or the PAS reaction was performed to visualize the starches (Wędzony, 1996).

In order to identify the main classes of the chemical compounds that are present in the trichomes, histochemical procedures with fresh or fixed flowers using Sudan III, Sudan Black B and Lugol's solution were performed in order to detect the total lipids, starch grains and proteins (Johansen, 1940), respectively.

For SEM, the flowers were fixed (as above) and later dehydrated and critical point dried using CO_2 . They were then sputter-coated with gold and examined at an accelerating voltage of 20 kV using a Hitachi S-4700 scanning electron microscope, which is housed at the Institute of Geological Sciences, Jagiellonian University in Kraków, Poland.

Phylogenetic analyses

The available *matK/trnK* DNA sequences of the *Pinguicula* species [*P. acuminata* (DQ010652.1), *P. agnata* (AF531782.1), *P. albida* (LC348432.1), *P. alpina* (AF531783.1), *P. ehlersiae* Speta & F.Fuchs (NC_023463.1), *P. elongata* Benj. (FM200224.1), *P. emarginata* (AF531785.1), *P. esseriana* (DQ010656.1), *P. filifolia* (AF531786.1), *P. gigantea* (AF531789.1), *P. gracilis* Zamudio (AF531790.1), *P. hemiepiphytica* (LC348445.1), *P. ibarrae* (LC348446.1), *P. laeana* Speta & F.Fuchs (DQ010659.1), *P. lusitanica* (DQ010661.1), *P. medusina* Zamudio & Studnička (LC348454.1), *P. moctezumae* (AF531797.1), *P. moranensis* (AF531798.1), *P. rectifolia* (AF531801.1), *P. rotundiflora* Studnička (AF531802.1), *P. sharpii* Casper & K.Kondo (AF531803.1) and *P. vulgaris* (AF531807.1)] were obtained from GenBank (NCBI) to be the ingroup. For the

outgroup, two *Genlisea* [*G. aurea* A.St.-Hil. (NC_037078.1) and *G. violacea* A.St.-Hil. (NC_037083.1)] and two *Utricularia* species [*U. foliosa* L. (KY025562.1) and *U. reniformis* A.St.-Hil. (NC_029719.2)] were used. The sequences were aligned using the online MAFFT v. 7.450 package (Katoh *et al.*, 2019). All of the gaps were treated as missing. We used three approaches to create the phylogenetic reconstructions: Bayesian inference (BI), maximum likelihood (ML) and maximum parsimony (MP). BI was determined using Mr Bayes v. 3.2.7a (Ronquist *et al.*, 2012) under the CIPRES Science Gateway v. 3.3 (Miller *et al.*, 2010). For BI, 2×10^6 generations were calculated using two runs with four chains until the standard deviation reached a value <0.01 . In each run, the trees were sampled every 100 generations at a sample frequency of 100. The first 25 % of the trees that were initially produced were discarded as burn-in. The BI was conducted using the GTR + G model and was calculated using MrModeltest v. 2.4 software (Nylander, 2004) following the Akaike information criterion (Akaike, 1973). ML was determined using the online IQ-TREE v. 1.6.12 (Nguyen *et al.*, 2015) and the obtained branch supports with the ultrafast bootstrap (10 000 replicates) (Hoang *et al.*, 2018). For the MP analyses, PAUP* v. 4.0a (build 166) program (Swofford, 2002) was used under the CIPRES Science Gateway v. 3.3 (Miller *et al.*, 2010) to obtain the bootstrap values (2000 pseudoreplicates and a heuristic search with 1000 replicates with the random addition of sequences and the branch swapping algorithm TBR). The trees that were obtained were edited using FigTree v. 1.4.3 (Rambaut, 2016). To optimize the pollinators/syndromes on the tree, we used the BI tree, and the pollinators were plotted according to published studies (listed in Table 1). The pollinator silhouettes used in Fig. 4 were designed using Freepik (<https://www.freepik.com>).

RESULTS

In our study, we observed various types of multicellular non-glandular trichomes, which differed in terms of their micromorphology (see Supplementary data Table S1). The trichome cells were highly vacuolated (Fig. 2A, B) and contained a peripheral cytoplasm with organelles such as a nucleus, mitochondria, plastids and an endoplasmic reticulum (Fig. 2C). Intranuclear paracrystalline bodies occurred in the nuclei (Fig. 2B). Staining with NBB revealed that these consisted of proteins (Fig. 2B). Some trichome cells had visible cuticular striations (Fig. 2D–F), while others had a smooth surface (Fig. 2F). The PAS reaction and Lugol's staining revealed amyloplasts with starch grains in the cells of the trichomes of the species from the subgenus *Temnoceras*: *P. agnata*, *P. albida*, *P. ibarrae*, *P. martinezii*, *P. filifolia* and *P. gigantea* (Fig. 3A–I and see 'starch' grade in Fig. 4). Starch grains were observed in these species independent of the type of trichomes (Supplementary data Table S1). Lugol's staining did not reveal any amyloplasts with starch grains in the cells of the trichomes of the species from the subgenus *Pinguicula*: *P. alpina* (Fig. 5A–C) and *P. vulgaris* (*P. vulgaris* subsp. *vulgaris* and *P. vulgaris* subsp. *bicolor*) (Fig. 5D–I) or the subgenus *Isoloba*: *P. lusitanica* (Fig. 5J–L). Moreover, this staining did not reveal any amyloplasts with starch grains in the trichome cells of species from the subgenus *Temnoceras*, which is pollinated by butterflies [*P. moctezumae*, *P. esseriana*, *P. moranensis*, *P. emarginata* and *P. rectifolia*;




FIG. 2. Structure of the non-glandular trichomes. (A) Section through the *P. albida* multicellular thick compact non-glandular trichomes that are located in the throat; note the numerous starch grains (asterisk); scale bar = 10 μ m. (B) Naphthol blue black staining of a *P. moctezumae* multicellular non-glandular trichome showing the presence of a nucleus with a paracrystalline protein inclusion (arrow); note there are no protein bodies in the cytoplasm; scale bar = 10 μ m. (C and D) Ultrastructure of a cell of a *P. agnata* non-glandular trichome; note the mitochondrion (M), nucleus (N) and prominent cuticular striations (arrowhead); scale bars = 0.7 μ m and 0.5 μ m, respectively. (E and F) Micromorphology of a *P. agnata* multicellular compact thick non-glandular trichome that is located in the front of the throat; note the cuticular striations on the surface of the apical cells (ap) and the smooth cuticle surface of the basal cell (bc); scale bars = 50 μ m and 100 μ m, respectively.

Fig. 4 ('psycho' clade Fig. 6A–G] or birds (*P. mesophytica* and *P. hemiephytica*, Fig. 7A–D; Supplementary data Table S1). Staining with NBB did not reveal any protein bodies (in either the cytoplasm or the vacuoles) in the cells of the trichomes of any of the examined species (Fig. 8A–D; Supplementary data Table S1). Staining with Sudan III did not reveal any lipid droplets in the cells of the trichomes in any of the examined species (Fig. 9A–F); however, positive staining was recorded in the cuticular striations (Fig. 9A–F).

The phylogenetic hypothesis, which was based on the *trnK/matK* sequences (Fig. 4), supports the assumption that both psychophily and ornithophily are derived for the *Pinguicula* lineages, probably from the plesiomorphic condition of myophily and/or melittophily. The ornithophily was possibly derived from the psychophily (Fig. 4). Thus, the pollination by birds has emerged at least twice as homoplasies to the *Pinguicula* species independently.




FIG. 3. PAS reaction and Lugol's staining of the *Pinguicula* species that were examined, which contain amyloplasts with starch grains (arrow, inserts) inside various types of non-glandular trichomes. (A–C) *P. agnata*; scale bars = 10 μ m, 10 μ m, 50 μ m, respectively. (D and E) *P. albida*; scale bars = 10 μ m and 50 μ m, respectively. (F) *P. ibarrae*; scale bar = 50 μ m. (G) *P. martinezii*; scale bar = 50 μ m. (H) *P. filifolia*; scale bar = 50 μ m. (I) *P. gigantea*; scale bar = 50 μ m.

DISCUSSION

We did not find the typical characters of glandular cells in the cells of the multicellular clavate trichomes. Therefore, we agree with Casper (1966, 2019) that these trichomes are non-glandular. We did show that the cells of the floral non-glandular trichomes of *P. agnata*, *P. albida*, *P. ibarrae*, *P. martinezii*, *P. filifolia* and *P. gigantea* were rich in amyloplasts that contained starch. Thus, these peculiar trichomes contain food reserves and probably function as edible trichomes. In orchids, edible trichomes (including pseudopollen-forming trichomes) are formed for a specific pollinator group, i.e. bees (Pansarin and Maciel, 2017). Thus, it is clear that in *Pinguicula* starch contained trichomes are recorded in species pollinated by bees, as showed in the 'starch' grade by the phylogenetical hypothesis (Fig. 4). Therefore, the lack of starch in the trichomes in the 'psycho' clade is a secondary loss, considering that *P. alpina*, *P. lusitanica* and *P. vulgaris* also did not present this character (Fig. 4). *Pinguicula*




FIG. 4. Phylogeny of the *Pinguicula* species based on the Bayesian inference (BI), maximum likelihood (ML) and maximum parsimony (MP) analyses of the *trnK/matK* sequences. The numbers above the branches refer to the BI posterior probability and the ML bootstrap support, respectively, and below the MP, the bootstrap support. The animal silhouettes denote the pollinator for each species. '*' indicates the homoplastic origin of the ornithophily for *P. hemiepiphytica* and *P. laeana* independently. '+' or '-' indicate the presence/absence of starch grains in the edible trichomes of the bee-/fly-pollinated species.

mesophytica is not represented in the tree but is a sister species to *P. moranensis* based on internal transcribed spacer (ITS) rDNA according to Shimai *et al.* (2007). Thus, pollination by birds is perhaps homoplastic in the *Pinguicula* species considering the known or supposed ornithophilic species (*P. hemiepiphytica*, *P. laeana* and *P. mesophytica*; Lampard *et al.*, 2016; Rocchia *et al.*, 2016).

Interestingly, not all myophilic and melittophilic species had starch in these trichomes, which enabled us to infer that these traits are not a condition for those pollination syndromes. Moreover, we did not record food reserves in the trichomes of *P. alpina* and *P. vulgaris*, which are pollinated by bees and flies (Molau, 1993; Fleischmann, 2016). Fleischmann (2016) observed various dipterans dabbling at the yellow spots on the otherwise white corolla of *P. alpina* and on the white corolla marks on the violet corolla of *P. vulgaris* and *P. leptoceras* with their proboscis. He interpreted

this behaviour as the insects trying to find nectar and pollen, and, therefore, in these species, the trichomes may guide insects to the spur. However, we do not agree with Fleischmann (2016) that they play the role of 'feeding hairs' in *P. alpina* and *P. vulgaris* because we did not find any reserve material in these trichomes. For this reason, these trichomes may play a tactile role and act as guides or they might mimic the edible trichomes of other species.

Most researchers accept that in *Pinguicula* the reward for pollinators is generally nectar because of the occurrence of a spur with glandular trichomes (Fleischmann and Rocchia, 2018; Lustofin *et al.* 2019); however, actual observations of nectar secretion and nectar analysis are rare (Zamora, 1999; Abrahamczyk *et al.* 2017; Lustofin *et al.* 2019). Although edible trichomes may act as a reward in addition to nectar, a detailed study of nectar production and secretion in *Pinguicula* is required to be absolutely certain that all *Pinguicula* species




FIG. 5. Histochemistry of the flower non-glandular trichomes from the species belonging to the *Pinguicula* and *Isoloba* subgenera; note the numerous starch grains inside the pollen grains (arrow). (A) Herbarium material of the *P. alpina* (KRA 0299930) that were examined. (B and C) Negative result of the Lugol's staining of the *P. alpina* non-glandular trichomes; scale bars = 100 μ m and 50 μ m, respectively. (D) Herbarium material of the *P. vulgaris* subsp. *vulgaris* (KRA 71415) that were examined. (E and F) Negative result of the Lugol's staining of the *P. vulgaris* subsp. *vulgaris* non-glandular trichomes; note the pollen grains (arrow) with a positive staining of the starch grains inside; scale bars = 100 μ m and 100 μ m, respectively. (G) Herbarium material of the *P. vulgaris* subsp. *bicolor* (KRA 0138573) that were examined. (H and I) Negative result of the Lugol's staining of the *P. vulgaris* subsp. *bicolor* non-glandular trichomes; note the pollen grains (arrow) with a positive staining of the starch grains inside; scale bars = 100 μ m and 100 μ m, respectively. (J–L) Negative result of the Lugol's staining of the *P. lusitanica* non-glandular trichomes; scale bars = 50 μ m, 50 μ m and 50 μ m, respectively.

produce nectar and in what quantities. In the related genera *Utricularia* (Hobbhahn *et al.*, 2006; Clivati *et al.*, 2014; Płachno *et al.*, 2017, 2018, 2019a, b) and *Genlisea* (Aranguren *et al.* 2018), the reward for pollinators is nectar. However, in some species (*U. antennifera*, *U. capilliflora*, *U. dunlopii*, *U. dunstaniae* and *U. lowriei*), the spur is significantly reduced and the corolla forms filiform appendages (Taylor, 1989; Reut and Jobson, 2010). In *U. dunlopii*, the glandular trichomes (osmophores) are densely distributed on the modified floral appendages, and therefore their scent is most probably the attractant for visiting insects (Płachno *et al.*, 2016). Although there are yellow non-glandular trichomes in the flower throats of *U. multifida* and *U. tenella*, they do not play the role of edible trichomes (Płachno *et al.*, 2019a).

In orchids, the edible trichome cells (including the pseudopollen, which is formed by the disintegration of the trichomes) contain various types of food material (see Davies, 2009 and references therein). The main food material that is found in the edible trichome of orchids in the species from the *Maxillaria* genus is protein (Davies, 2009). Starch grains were recorded in the cells of the trichomes in the species from the genera *Dendrobium* (Davies and Turner, 2004), *Cyanaeorchis* (Pansarin and Maciel, 2017), *Polystachya* (Davies *et al.*, 2002) and *Maxillaria* (Davies, 2009). Lipid droplets were recorded in the edible trichomes of *Cyanaeorchis* (Pansarin and Maciel, 2017). Thus, the edible trichomes of orchids are more diverse in the types of food material compared with *Pinguicula*.




FIG. 6. PAS reaction and Lugol's staining of various non-glandular trichomes of the *Pinguicula* species that were examined that are pollinated by Lepidoptera; note the pollen grains (arrow) with a positive staining of the starch grains inside. (A) PAS reaction of the *P. moctezumae* non-glandular trichomes that are located in the basal part of the spur; scale bar = 50 μ m. (B and C) Negative result of the Lugol's staining of the *P. moctezumae* non-glandular trichomes; scale bars = 50 μ m and 100 μ m, respectively. (D) Negative result of the Lugol's staining of the *P. esseriana* non-glandular trichomes; scale bar = 100 μ m. (E) Negative result of the Lugol's staining of the *P. moranensis* non-glandular trichomes; scale bar = 100 μ m. (F) Negative result of the Lugol's staining of the *P. emarginata* non-glandular trichomes; scale bar = 100 μ m. (G) Negative result of the Lugol's staining of the *P. rectifolia* non-glandular trichomes; scale bar = 100 μ m.




FIG. 7. PAS reaction and Lugol's staining of various non-glandular trichomes of the *Pinguicula* species that were examined that are most probably pollinated by hummingbirds. (A) PAS reaction of a *P. mesophytica* non-glandular trichome; scale bar = 50 μ m. (B) Negative result of the Lugol's staining of the *P. mesophytica* non-glandular trichomes; scale bar = 100 μ m. (C and D) Negative result of the Lugol's staining of the *P. hemiepiphytica* non-glandular trichomes; scale bars = 100 μ m and 100 μ m, respectively.




FIG. 8. Naphthol blue black (NBB) staining of various non-glandular trichomes of the selected *Pinguicula* species that were examined; note the lack of protein bodies in the cytoplasm. Nucleus (asterisk). (A) *P. agnata*; scale bar = 50 μ m. (B) *P. albida*; scale bar = 50 μ m. (C) *P. esseriana*; scale bar = 50 μ m. (D) *P. vulgaris*; scale bar = 50 μ m.




FIG. 9. Sudan III staining of various non-glandular trichomes of the selected *Pinguicula* species that were examined; note the positive staining of the cuticular striations of the non-glandular trichomes cells (arrow, insert and arrowhead) and lipids inside the pollen grains (asterisk). (A) *P. agnata*; scale bar = 100 μ m. (B) *P. rectifolia*; scale bar = 100 μ m. (C) *P. moranensis*; scale bar = 100 μ m. (D) *P. esseriana*; scale bar = 100 μ m. (E) *P. hemiepiphytica*; scale bar = 100 μ m. (F) *P. mesophytica*; scale bar = 100 μ m.

From a phylogenetic perspective, edible trichomes are symplesiomorphic for the *Pinguicula* species and are found in the species of the 'starch' grade (Fig. 4), and therefore this does

not support a monophyletic group such as a synapomorphy. However, the edible trichomes are derived and are possibly a specialization for fly and bee pollinators that act as a food reward for these visitors.

Field observations are needed to answer the question of whether insects consume ‘starch’ trichomes of *Pinguicula* flowers and thus whether these structures can be regarded as pollinators’ rewards. Checking if there is a correlation between the amount of nectar produced and the number of trichomes with starch also seems interesting.

Conclusion

Floral non-glandular trichomes play the role of edible trichomes in some *Pinguicula* species (*P. agnata*, *P. albida*, *P. ibarrae*, *P. martinezii*, *P. filifolia* and *P. gigantea*), which are primarily classified as bee-pollinated species that originated from Central and South America. It seems that in *Pinguicula* that are pollinated by other pollinator groups (Lepidoptera and hummingbirds), the non-glandular trichomes in the flowers play a role other than being a floral reward for their pollinators. However, even with a phylogenetic perspective, the gaps in knowledge are wide for several species, which does not permit a robust hypothesis. Thus, only when field studies have been undertaken can we be absolutely certain of the role of these trichomes.

SUPPLEMENTARY DATA

Supplementary data are available online at <https://academic.oup.com/aob> and consist of Table S1: micromorphology and histochemistry analyses of the food material content in various type of the *Pinguicula* flower non-glandular trichomes.

ACKNOWLEDGEMENTS

The authors would like to express their sincere and warm thanks to Dr Miroslav Studnička (director of Liberec Botanical Garden, Czech Republic) for providing some of the plant material for the study. We thank horticulturist Lucyna Kurlito for her conscientious care of the living collection of carnivorous plants that are located in the Botanical Garden of Jagiellonian University in Kraków and also thank the curator of KRA, Dr hab. Marcin Nobis for the opportunity to use the herbarium material. We cordially thank the reviewers and editors for comments, which have helped to improve the manuscript. There is no conflict of interest.

FUNDING

This research was financially supported by the Ministry of Science and Higher Education of Poland as part of the statutory activities of the Department of Plant Cytology and Embryology, Institute of Botany, Faculty of Biology, Jagiellonian University in Kraków (N18/DBS/000002). We cordially thank PIK INSTRUMENTS sp. z o.o. (<https://micro-shop.pl/english/>) for sponsoring our research.

LITERATURE CITED

- Abrahamczyk S, Kessler M, Hanley D, et al. 2017. Pollinator adaptation and the evolution of floral nectar sugar composition. *Journal of Evolutionary Biology* **30**: 112–127.
- Akaike H. 1973. Information theory and an extension of the maximum likelihood principle. In: *Second International Symposium on Information Theory* 267–281.
- Alcalá RE, Domínguez CA. 2003. Patterns of prey capture and prey availability among populations of the carnivorous plant *Pinguicula moranensis* (Lentibulariaceae) along an environmental gradient. *American Journal of Botany* **90**: 1341–1348.
- Alcalá RE, Domínguez CA. 2005. Differential selection for carnivory traits along an environmental gradient in *Pinguicula moranensis*. *Ecology* **86**: 2652–2660.
- Aranguren Y, Plachno BJ, Stpiczyńska M, Miranda VFO. 2018. Reproductive biology and pollination of the carnivorous *Genlisea violacea* (Lentibulariaceae). *Plant Biology* **20**: 591–601.
- Casper SJ. 1966. Monographie der Gattung *Pinguicula* L. *Bibliotheca Botanica* **127–128**: 1–209.
- Casper SJ. 2019. *The insectivorous genus Pinguicula (Lentibulariaceae) in the greater antilles*. Berlin: Botanischer Garten und Botanisches Museum.
- Chase MW, Christenhusz MJM, Fay MF, et al. 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* **181**: 1–20.
- Clivati D, Cordeiro GD, Plachno BJ, de Miranda VF. 2014. Reproductive biology and pollination of *Utricularia reniformis* A.St.-Hil. (Lentibulariaceae). *Plant Biology (Stuttgart, Germany)* **16**: 677–682.
- Cox PA. 1982. Vertebrate pollination and the maintenance of dioecism in *Freycinetia*. *The American Naturalist* **120**: 65–80.
- Darnowski D, Bauer U, Moran J, et al. 2018. Prey selection and specialization by carnivorous plants. In: Ellison AM, Adamec L, eds. *Carnivorous plants: physiology, ecology and evolution*. Oxford: Oxford University Press, 285–293.
- Davies KL. 2009. Morphology. In: Kull T, Arditti J, Wong SM, eds. *Orchid biology: reviews and perspectives*, X. Dordrecht: Springer, 159–184.
- Davies KL, Turner MP. 2004. Pseudopollen in *Dendrobium unicum* Seidenf. (Orchidaceae): reward or deception? *Annals of Botany* **94**: 129–132.
- Davies KL, Roberts DL, Turner MP. 2002. Pseudopollen and food-hair diversity in *Polystachya* Hook. (Orchidaceae). *Annals of Botany* **90**: 477–484.
- Domínguez Y, Silva SR, Valdés CMP, Miranda VFO. 2014. Inter- and intra-specific diversity of Cuban *Pinguicula* (Lentibulariaceae) based on morphometric analyses and its relation with geographical distribution. *Plant Ecology & Diversity* **7**: 519–531.
- Endress PK. 2010. The evolution of floral biology in basal angiosperms. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**: 411–421.
- Faegri K, Van der Pijl L. 1979. *The principles of pollination ecology*. Oxford: Pergamon Press.
- Fisher DB. 1968. Protein staining of ribboned epon sections for light microscopy. *Histochemie. Histochemistry. Histochimie* **16**: 92–96.
- Fleischmann A. 2016. *Pinguicula* flowers with pollen imitations close at night – some observations on butterwort flower biology. *Carnivorous Plant Newsletter* **45**: 84–92.
- Fleischmann A, Rocca A. 2018. Systematics and evolution of Lentibulariaceae: I. *Pinguicula*. In: Ellison AM, Adamec L, eds. *Carnivorous plants: physiology, ecology and evolution*. Oxford: Oxford University Press, 70–80.
- Hernández MP, Katinas L. 2019. Technique for the identification of osmophores in flowers of herbarium material (TIOFH). *Protoplasma* **256**: 1753–1765.
- Heslop-Harrison Y. 1970. Scanning electron microscopy of fresh leaves of *Pinguicula*. *Science* **167**: 172–174.
- Heslop-Harrison Y. 2004. *Pinguicula* L. *Journal of Ecology* **92**: 1071–1118.
- Heslop-Harrison Y, Heslop-Harrison J. 1980. Chloride ion movement and enzyme secretion from the digestive glands of *Pinguicula*. *Annals of Botany* **45**: 729–731.
- Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBboot2: improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **35**: 518–522.
- Hobbhahn N, Küchmeister H, Porembski S. 2006. Pollination biology of mass flowering terrestrial *Utricularia* species (Lentibulariaceae) in the Indian Western Ghats. *Plant Biology* **8**: 791–804.

- Humphrey CD, Pittman FE. 1974. A simple methylene blue–azure II–basic fuchsin stain for epoxy-embedded tissue sections. *Stain Technology* **49**: 9–14.
- Jensen WA. 1962. *Botanical histochemistry – principles and practice*. San Francisco: W. H. Freeman and Company.
- Jobson RW, Playford J, Cameron KM, Albert VA. 2003. Molecular phylogenetics of Lentibulariaceae inferred from plastid *rps16* intron and *trnL-F* DNA sequences: implications for character evolution and biogeography. *Systematic Botany* **28**: 157–171.
- Johansen DA. 1940. *Plant microtechnique*. New York: McGraw-Hill Book Co.
- Katoh K, Rozewicki J, Yamada KD. 2019. MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* **20**: 1160–1166.
- Lampard S, Gluch O, Robinson A, et al. 2016. *Pinguicula of Latin America*. Dorset, UK: Redfern Natural History.
- Lustofin K, Świątek P, Miranda VFO, Plachno BJ. 2019. Flower nectar trichome structure of carnivorous plants from the genus bladderworts *Pinguicula* L. (Lentibulariaceae). *Protoplasma* **257**: 245–259.
- Mathe C, Vieillescazes C. 2002. Compréhension des mécanismes de coloration des liants protéiques picturaux à l'aide du Noir Amide 10B. *L'Actualité Chimique* **7**: 11–14.
- Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *Gateway Computing Environments Workshop, GCE 2010 (New Orleans)* 1–8.
- Molau U. 1993. Reproductive ecology of the three Nordic *Pinguicula* species (Lentibulariaceae). *Nordic Journal of Botany* **13**: 149–157.
- Müller K, Borsch T, Legendre L, Porembski S, Theisen I, Barthlott W. 2004. Evolution of carnivory in Lentibulariaceae and the Lamiales. *Plant Biology (Stuttgart, Germany)* **6**: 477–490.
- Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* **32**: 268–274.
- Nicolson SW, Nepi M, Pacini E, eds. 2007. *Nectaries and nectar*. Dordrecht, The Netherlands: Springer.
- Nordin M. 2015. *Pinguicula alpina* (alpine butterwort) on the Swedish island of Gotland: pollination and reproduction. Thesis, Uppsala University (in Swedish).
- Nylander JAA. 2004. *MrModeltest v2*. Program distributed by the author.
- Pansarin ER, Maciel AA. 2017. Evolution of pollination systems involving edible trichomes in orchids. *AoB Plants* **9**: plx033.
- Plachno BJ, Stpiczyńska M, Świątek P, Davies KL. 2016. Floral micromorphology of the Australian carnivorous bladderwort *Utricularia dunlopiae*, a putative pseudocopulatory species. *Protoplasma* **253**: 1463–1473.
- Plachno BJ, Stpiczyńska M, Davies KL, Świątek P, Miranda VFO. 2017. Floral ultrastructure of two Brazilian aquatic–epiphytic bladderworts: *Utricularia cornigera* Studnička and *U. nelumbifolia* Gardner (Lentibulariaceae). *Protoplasma* **254**: 353–366.
- Plachno BJ, Stpiczyńska M, Adamec L, Miranda VFO, Świątek P. 2018. Nectar trichome structure of aquatic bladderworts from the section *Utricularia* (Lentibulariaceae) with observation of flower visitors and pollinators. *Protoplasma* **255**: 1053–1064.
- Plachno BJ, Stpiczyńska M, Świątek P, et al. 2019a. Floral micromorphology and nectar composition of the early evolutionary lineage *Utricularia* (subgenus *Polypompholyx*, Lentibulariaceae). *Protoplasma* **256**: 1531–1543.
- Plachno BJ, Stpiczyńska M, Świątek P, et al. 2019b. Floral micromorphology of the bird-pollinated carnivorous plant species *Utricularia menziesii* R.Br. (Lentibulariaceae). *Annals of Botany* **123**: 213–220.
- Rambaut A. 2016. *FigTree, version 1.4.3*. Institute of Evolutionary Biology, University of Edinburgh.
- Reut M, Jobson RW. 2010. A phylogenetic study of subgenus *Polypompholyx*: a parallel radiation of *Utricularia* (Lentibulariaceae) throughout Australasia. *Australian Systematic Botany* **23**: 152–161.
- Rickson FR. 1979. Ultrastructural development of the beetle food tissue of *Calycanthus* flowers. *American Journal of Botany* **66**: 80–86.
- Roccia A, Gluch O, Lampard S, et al. 2016. *Pinguicula of the temperate North*. Dorset, UK: Redfern Natural History.
- Ronquist F, Teslenko M, van der Mark P, et al. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**: 539–542.
- Schäferhoff B, Fleischmann A, Fischer E, et al. 2010. Towards resolving Lamiales relationships: insights from rapidly evolving chloroplast sequences. *BMC Evolutionary Biology* **10**: 352.
- Shimai H, Kondo K. 2007. Phylogenetic analysis of Mexican and Central American *Pinguicula* (Lentibulariaceae) based on internal transcribed spacer (ITS) sequence. *Chromosome Botany* **2**: 67–77.
- Simpson BB, Neff JL. 1981. Floral rewards: alternatives to pollen and nectar. *Annals of the Missouri Botanical Garden* **68**: 301–322.
- Swofford DL. 2002. *PAUP*. Phylogenetic analysis using parsimony (*and other methods)*. Version 4.0. Sunderland, MA: Sinauer Associates.
- Taylor P. 1989. The genus *Utricularia* – a taxonomic monograph. *Kew Bulletin Additional Series* **14**: 1–724.
- Thien LB, Bernhardt P, Devall MS, et al. 2009. Pollination biology of basal angiosperms (ANITA grade). *American Journal of Botany* **96**: 166–182.
- Vassilyev AE, Muravnik LE. 1988. The ultrastructure of the digestive glands in *Pinguicula vulgaris* L. (Lentibulariaceae) relative to their function. I. The changes during maturation. *Annals of Botany* **62**: 329–341.
- Villegas SG, Alcalá RE. 2018. Reproductive ecology of the carnivorous plant *Pinguicula moranensis* (Lentibulariaceae). *Plant Biology (Stuttgart, Germany)* **20**: 205–212.
- Wędzony M. 1996. *Fluorescence microscopy for botanists*. Kraków, Poland: Department of Plant Physiology Monographs 5 [in Polish].
- Young HJ. 1986. Beetle pollination of *Dieffenbachia longispatha* (Araceae). *American Journal of Botany* **73**: 931–944.
- Zamora R. 1999. Conditional outcomes of interactions: the pollinator–prey conflict of an insectivorous plant. *Ecology* **80**: 786–795.