



eCOMMONS

Loyola University Chicago
Loyola eCommons

Bioinformatics Faculty Publications

Faculty Publications and Other Works by
Department

5-7-2020

Draft Genome Sequence of *Citrobacter freundii* UFMG-H9, Isolated from Urine from a Healthy Bovine Heifer (Gyr Breed)

Silvia Giannattasio-Ferraz
Loyola University Chicago

Laura Maskeri
Loyola University Chicago

André P. Oliveira
Loyola University Chicago

Edel F. Barbosa-Stancioli
Loyola University Chicago

Catherine Putoni
Loyola University Chicago, cputonti@luc.edu

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub



Part of the [Bioinformatics Commons](#), and the [Biology Commons](#)

Recommended Citation

Giannattasio-Ferraz, Silvia; Maskeri, Laura; Oliveira, André P.; Barbosa-Stancioli, Edel F.; and Putoni, Catherine. Draft Genome Sequence of *Citrobacter freundii* UFMG-H9, Isolated from Urine from a Healthy Bovine Heifer (Gyr Breed). *Microbiology Resource Announcements*, 9, , 2020. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/MRA.00387-20>

This Article is brought to you for free and open access by the Faculty Publications and Other Works by Department at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).
© Giannattasio-Ferraz et al., 2020.



Draft Genome Sequence of *Citrobacter freundii* UFMG-H9, Isolated from Urine from a Healthy Bovine Heifer (Gyr Breed)

Silvia Giannattasio-Ferraz,^a Laura Maskeri,^b André P. Oliveira,^c Edel F. Barbosa-Stancioli,^a  Catherine Putonti^{b,d,e,f}

^aDepartamento de Microbiologia, Instituto de Ciências Biológicas, Universidade Federal Minas Gerais, Belo Horizonte, MG, Brazil

^bBioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA

^cEmpresa de Pesquisa Agropecuária de Minas Gerais–EPAMIG, Uberaba, MG, Brazil

^dDepartment of Biology, Loyola University Chicago, Chicago, Illinois, USA

^eDepartment of Computer Science, Loyola University Chicago, Chicago, Illinois, USA

^fDepartment of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA

ABSTRACT *Citrobacter freundii* is a pathogen associated with antibiotic resistance and severe infections in humans. Here, we report the draft genome sequence of *C. freundii* strain UFMG-H9, an isolate from urine from a healthy Gyr heifer.

Citrobacter freundii is a known human pathogen highly associated with the occurrence of bacteremia and antibiotic resistance (1). This species has been reported as a contaminant of wastewater (2) and food of animal origin such as milk and cheese (3). Strains of *C. freundii* have also been reported to cause urinary infections in humans (4). Here, we report the draft genome sequence of *C. freundii* strain UFMG-H9, isolated from urine from a healthy heifer belonging to a Gyr herd pure in origin, from the Agricultural Research Company of Minas Gerais State (EPAMIG) in Brazil.

Samples were collected in May 2019, and the methodology was previously approved by the Ethics Committee in Animal Experimentation of the Universidade Federal de Minas Gerais, Brazil (CEUA/UFMG; approval number 40/2019). All experiments were performed in accordance with relevant guidelines. To collect the urine, the vulva of the animal was washed with distilled water and soap, and then midstream urine was collected using a 50-ml sterile tube. The urine was kept at -20°C until processing (within 48 h). Aliquots of 2 ml were centrifuged. The supernatant was plated on lysogeny broth (LB) agar plates and incubated at 37°C overnight. Single colonies were picked, grown in LB at 37°C overnight, and then replated. This process of plating and liquid culture was repeated 3 times in order to obtain pure colonies. Following this, a single colony was inoculated in LB liquid medium and grown overnight with shaking at 37°C . A 1-ml aliquot was centrifuged, and the pellet was used for DNA extraction with the DNeasy UltraClean microbial kit (Qiagen, Hilden, Germany) according to the manufacturer's instruction. The DNA was quantified using a Qubit fluorometer. The genus and species were determined prior to whole-genome sequencing by sequencing the 16S rRNA gene sequence. Next, the DNA was sent to the Microbial Genomic Sequencing Center (MiGS) at the University of Pittsburgh for library preparation and whole-genome sequencing. Briefly, the DNA was fragmented using an Illumina tagmentation enzyme, and indices were attached using PCR. The genome sequencing was conducted using the NextSeq 550 platform. Sequencing produced 2,854,920 pairs of 150-bp reads that were trimmed using Sickle v1.33 (<https://github.com/najoshi/sickle>). The genome was assembled using SPAdes v3.13.0 with the "only-assembler" option for k values of 55, 77, 99, and 127 (5). Then, the genome coverage was calculated using BBMap v38.47 (<https://sourceforge.net/projects/bbmap/>). Finally, the genome was annotated by the NCBI

Citation Giannattasio-Ferraz S, Maskeri L, Oliveira AP, Barbosa-Stancioli EF, Putonti C. 2020. Draft genome sequence of *Citrobacter freundii* UFMG-H9, isolated from urine from a healthy bovine heifer (Gyr breed). Microbiol Resour Announc 9:e00387-20. <https://doi.org/10.1128/MRA.00387-20>.

Editor Julie C. Dunning Hotopp, University of Maryland School of Medicine

Copyright © 2020 Giannattasio-Ferraz et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catherine Putonti, cputonti@luc.edu.

Received 12 April 2020

Accepted 14 April 2020

Published 7 May 2020

Prokaryotic Genome Annotation Pipeline (PGAP) v4.11 (6). Default parameters were used for all software tools unless otherwise noted.

C. freundii strain UFMG-H9 is 4,744,381 bp long with a GC content of 50.77%. The assembly includes 20 contigs with an N_{50} value of 525,854 bp, and the genome coverage is 164×. Annotation identified 4,315 coding genes and 75 tRNAs. Further analysis of this strain can increase our understanding of the urogenital tract of this species.

Data availability. This whole-genome sequencing (WGS) project has been deposited in GenBank under accession number [JAAVMN000000000](https://ncbi.nlm.nih.gov/GenBank/entry/JAAVMN000000000). The raw reads can be found in SRA under accession number [SRR11455638](https://ncbi.nlm.nih.gov/SRA/entry/SRR11455638). This sequencing is part of BioProject accession number [PRJNA615899](https://ncbi.nlm.nih.gov/BioProject/entry/PRJNA615899).

REFERENCES

1. Liu L-H, Wang N-Y, Wu AY-J, Lin C-C, Lee C-M, Liu C-P. 2018. *Citrobacter freundii* bacteremia: risk factors of mortality and prevalence of resistance genes. *J Microbiol Immunol Infect* 51:565–572. <https://doi.org/10.1016/j.jmii.2016.08.016>.
2. Sghaier S, Abbassi MS, Pascual A, Serrano L, Díaz-De-Alba P, Said MB, Hassen B, Ibrahim C, Hassen A, López-Cerero L. 2019. Extended-spectrum β -lactamase-producing Enterobacteriaceae from animal origin and wastewater in Tunisia: first detection of O25b-B2₃-CTX-M-27-ST131 *Escherichia coli* and CTX-M-15/OXA-204-producing *Citrobacter freundii* from wastewater. *J Glob Antimicrob Resist* 17:189–194. <https://doi.org/10.1016/j.jgar.2019.01.002>.
3. Maifreni M, Frigo F, Bartolomeoli I, Innocente N, Biasutti M, Marino M. 2013. Identification of the *Enterobacteriaceae* in Montasio cheese and assessment of their amino acid decarboxylase activity. *J Dairy Res* 80: 122–127. <https://doi.org/10.1017/S002202991200074X>.
4. Jiménez-Guerra G, Borrego-Jiménez J, Gutiérrez-Soto B, Expósito-Ruiz M, Navarro-Marí JM, Gutiérrez-Fernández J. 2020. Susceptibility evolution to antibiotics of *Enterobacter cloacae*, *Morganella morganii*, *Klebsiella aerogenes* and *Citrobacter freundii* involved in urinary tract infections: an 11-year epidemiological surveillance study. *Enferm Infecc Microbiol Clin* 38:166–169. (In Spanish.) <https://doi.org/10.1016/j.eimc.2019.07.010>.
5. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
6. Tatusova T, DiCuccio M, Badredin A, Chetverin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.