


## The “Smalls” *Nephrops* Grounds (FU22) 2020 UWTV Survey Report and catch scenarios for 2021

Mikel Aristegui<sup>1</sup>, Marcin Blaszkowski<sup>1</sup>, Jennifer Doyle<sup>1</sup>, Gráinne Ryan<sup>1</sup> and Michael McAuliffe<sup>1</sup>.

<sup>1</sup> Fisheries Ecosystems Advisory Services, Marine Institute, Renville, Oranmore, Galway, Ireland.



Version 1.0



## Abstract

This report provides the main results and findings of the fifteenth annual underwater television survey on the 'Smalls grounds' ICES assessment area; Functional Unit 22. The survey was multi-disciplinary in nature collecting UWTV and other ecosystem data. A total of 40 UWTV stations were surveyed successfully (high quality image data), carried out over an isometric grid at 4.5nmi or 8.3km intervals. The precision, with a CV of 8%, was well below the upper limit of 20% recommended by SGNEPS (ICES, 2012). The 2020 abundance estimate was 33% lower than in 2019 and at 750 million is below the MSY  $B_{\text{trigger}}$  reference point (990 million). Using the 2020 estimate of abundance and updated stock data implies catch in 2021 that correspond to the F ranges in the EU multi annual plan for Western Waters are between 1238 and 1560 tonnes (assuming that discard rates and fishery selection patterns do not change from the average of 2017–2019). One species of sea pens was recorded as present at the stations surveyed: *Virgularia mirabilis*. Trawl marks were observed at 48% of the stations surveyed.

Key words: *Nephrops norvegicus*, stock assessment, geostatistics, underwater television (UWTV), benthos.

### Suggested citation:

Aristegui, M., Blaszkowski, M., Doyle, J., Ryan, G., and McAuliffe, M. 2020. The "Smalls" *Nephrops* Grounds (FU22) 2020 UWTV Survey Report and catch scenarios for 2021. Marine Institute UWTV Survey report.

## Introduction

The prawn (*Nephrops norvegicus*) are common in the Celtic Sea occurring in geographically distinct sandy/muddy areas where the sediment is suitable for them to construct their burrows (Figure 1). The *Nephrops* fishery in ICES sub-area 7 is extremely valuable with Irish landings in 2019 worth around €42 m at first sale. The Celtic Sea area (Functional Units 19-22, see Figure 1) supports a large multi-national targeted *Nephrops* fishery, mainly using otter trawls and yielding landings in the region of ~5,000 t annually over the last decade (ICES, 2020). The 2019 reported landings from the Smalls (~1639 t) were estimated to be worth in the region of €8.8 million at first sale. The Smalls ground is particularly important to the Irish demersal fleet accounting for around 13% of the fishing effort by all demersal vessels >15m between 2006 and 2009 (Gerritsen, *et al.*, 2012). The Irish demersal fleet now account for ~90% of the FU22 *Nephrops* landings (ICES, 2020). Good scientific information on stock status and exploitation rates are required to inform sustainable management of this resource.

*Nephrops* spend a great deal of time in their burrows and their emergence behaviour is influenced by several factors: time of year, light intensity, tidal strength, etc. Underwater television surveys and assessment methodologies have been developed to provide a fishery independent estimate of stock size, exploitation status and catch advice (ICES, 2009 & 2012). This is the fifteenth in a time series of UWTV surveys in the Celtic Sea FU22 "Smalls" ground carried out by the Marine Institute, Ireland.

The survey was multi-disciplinary in nature and also covered UWTV stations in FU16, FU19 and FU2021 the results of which are presented elsewhere (Aristegui *et al.*, 2020a-c).

The specific objectives of the 2020 survey are listed below:

1. To complete a survey of 40 randomised fixed isometric grid UWTV stations, with 4.5 nautical mile (nmi) spacing, on the "Smalls" *Nephrops* ground (FU22).
2. To obtain 2020 quality assured estimates of *Nephrops* burrow distribution and abundance on the "Smalls" *Nephrops* ground (FU22) and compare them with those collected in previous surveys.
3. To collect ancillary information from the UWTV footage collected at each station such as the occurrence of sea-pens, other macro benthos and fish species and trawl marks on the sea bed.
4. To sample *Nephrops* and macro benthos using a 3 metre beam trawl deployed at 10 stations once UWTV operations successfully completed.
5. To collect sediment samples for a *Nephrops* Microplastic research project.

This report details the final UWTV results of the 2020 survey and documents other data collected during the survey. Operational survey details are available in form of a survey narrative available from the scientist in charge (MA). The 2020 abundance estimate is used to generate catch scenarios for 2021 in line with procedures outlined in the stock annex for FU22 (ICES, 2017).

## Material and methods

To maintain a coefficient of variance (CV) < 20%, to achieve good spatial coverage over the ground and to generate burrow surface that reflects the underlying abundance of *Nephrops*,

a survey grid of 4.5 nm spacing has been used since 2012. The 2020 randomised isometric grid, which resulted in 40 planned stations, was generated using the “spsampl” function in the “sp” package (Pebesma & Bivand, 2005) in “R” (R Core Team, 2017). The boundary used to delineate the edge of the ground was based on information from VMS (Gerritsen & Lordan, 2011), habitat maps, and previous UWTV observations. The same boundary has been used through the time series.

The 2020 Celtic Sea survey took place on RV Celtic Voyager between the 15<sup>th</sup> July and the 21<sup>th</sup> July. The survey normally takes place in either July or August each year.

In 2020 image data were collected by a custom built camera system recording High Definition still image data at 12 frames per second with a camera angle of 75 (°). The digital images were stored on a server and were reviewed ashore through an in-house developed Image annotation R Shiny app (Aristegui, 2020). This app allows each reviewer to annotate burrows for each randomly assigned station in an efficient manner. The survey process is now paperless.

The operational protocols used were those reviewed by WKNEPHTV 2007 (ICES, 2007) and employed on other UWTV surveys in Irish waters. These protocols can be summarised as follows: At each station the UWTV sledge was deployed. Once stable on the seabed a 10 minute tow was recorded. Time referenced high definition image data was collected with a field of view or ‘FOV’ of 1.01 metre. Vessel position (DGPS) and position of sledge (using a USBL transponder) were recorded every 3 seconds. The navigational data were quality controlled using an “R” script developed by the Marine Institute (ICES, 2009) an example is shown in Figure 3. In 2020 the USBL navigational data were used to calculate distance over ground for 100% of stations.

In line with recommendations of the Workshop on *Nephrops* Burrow Counting (WKNEPS), all scientists were trained/re-familiarised using 2019 image data for training material and reference set (ICES, 2018). All counts were conducted by four trained scientists independent of each other after the survey. The numbers of *Nephrops* burrows systems (multiple burrows in close proximity which appear to be part of a single system) were counted as one. *Nephrops* activity in and out of burrows were counted and recorded for each station. Following the recommendation of SGNEPS the time for verified recounts was 7 minutes (ICES, 2009).

Presence / absence notes were also recorded on the occurrence of trawl marks, fish species and other species. Presence / absence of sea-pen species were also recorded to fulfil an OSPAR Special Request (ICES, 2011).

Finally, if there was any time during the one-minute block where counting was not possible, due to sediment clouds or other reasons, this was also estimated so that the time could be removed from the distance over ground calculations.

In 2020 the survey count data were screened to check for any unusual discrepancies using Lin’s Concordance Correlation Coefficient (CCC) with a threshold of 0.6. Lin’s CCC (Lin, 1989) measures the ability of counters to exactly reproduce each other’s counts on a scale of 1 to -1 where 1 is perfect concordance (i.e. a pairwise plot will have all points lying along the 1:1

line). A value of -1 would be generated by all points lying on the -1:1 line and a value of 0 indicates no correspondence at all. Lin's CCC quality control plots of survey count data for stations 120 to 122 is shown in Figure 4. When the count data fell below the threshold of 0.6 a third review was carried out. The paired count data that passed the Lin's CCC threshold was used in the analysis. When the paired counts did not pass the threshold an average of the three reviewers was deemed appropriate to use in the analysis.

Mean density was calculated by dividing the total number of burrow systems by the survey area observed. The USBL data were used to calculate distance over ground of the sledge. The field of view of the camera at the bottom of the screen was estimated by extrapolation at 1.01m assuming that the sledge was flat on the seabed (i.e. no sinking). Occasionally the lasers were not visible at the bottom of the screen due to sinking in very soft mud. The impact of this is a minor under estimate of densities at stations where this occurred.

From 2006 to 2014 calculation of spatial co-variance, spatial structuring, geo-statistical analysis of the mean and variance was carried out using SURFER Version 10.7.972. From 2015 the geostatistical analysis was carried out using the "RGeostats" package (Renard D., *et al*, 2015) and is available as a separate "R markdown" document. The same basic steps were carried out as in previous years; construction of experimental variogram, a model variogram produced with an exponential model, create krigged grid file using all data points as neighbours, same boundary used to estimate the domain area, mean density, total burrow abundance and calculate survey precision.

Due to time constraints beam trawling was not carried out on the "Smalls" ground this year. Sediment samples at three stations were collected for a research project to determine micro-plastics loading in *Nephrops norvegicus* and sediments.

## Results

In 2020 40 stations were completed successfully on the Smalls. A summary of the results is presented in Table 1. The density and estimated abundance decreased by around 33% in 2020. The average density and the abundance were the lowest in the time series. Figure 5 shows bubble plots of the variability between minutes and operators. These show that the burrow estimates are very consistent between minutes and counters. A combined violin and box plot of the observed burrow densities is presented in Figure 6. This shows that median and mean burrow densities are similar in most years. The inter-quartile range is between 0.2 and 0.7 in most years. However, in 2020 as in 2018 and 2016, this inter-quartile range is in the region of 0.1 to 0.4. In 2020 the mean adjusted<sup>1</sup> burrow density was 0.27 burrows/m<sup>2</sup>.

The krigged and point density data for 2006-2020 are shown in Figure 7. The krigged contours correspond well to the observed data. Highest densities are in the centre of the ground in all years. In general, densities are higher towards the south, west and central area of the ground.

---

<sup>1</sup> Note the "adjusted" density estimates in this report are adjusted by dividing by 1.3 (Table 2) to take account of edge effect over estimation of area viewed during UWTV transects (see Campbell et al 2009).

The annual survey statistics from this geo-statistical analysis are given in Table 1 and Figure 8. The 2020 estimate of 750 million burrows is below the geometric mean of the series (geomean [2006-2020]: 1184 million burrows) and is below the MSY  $B_{\text{trigger}}$  reference point of 990 million. The estimation of variance of the 2020 survey as calculated by RGeostats is low (with a CV or RSE of 8%), which is well below the SGNEPS recommendation for a CV <20% (ICES, 2012).

Figure 9 shows the standardised length frequency distributions of *Nephrops* caught using a beam trawl. Fishing operations were not carried out during 2010, 2013, 2014, 2015 and 2019 due to time constraints. For plotting purposes, the individuals <10mm caught were split evenly between males and females as it is not possible to accurately assign sex to individuals of this size. A strong cohort was apparent in the 2006 catches of around 17mm and can be tracked in catches in subsequent years. There was a shift to larger sizes in 2011 and 2012, with a shift back again to smaller sizes in 2016 to 2018.

Sea-pen presence/absence distribution across the Smalls *Nephrops* grounds is mapped in Figure 10. One sea-pen was identified from the image data in 2020, that is, *Virgularia mirabilis*. Trawl marks were noted at 48% of the stations surveyed.

The UWTV abundance data together with data from the fishery; landings, discards and removals in number, were used to calculate the harvest rate for 2019 of 8.5%. The mean weight in the landings and discards and the proportions of removal retained are also shown (Table 2).

The basis to 2021 catch scenarios are given in Table 3. The catch and landings scenarios at various different fishing mortalities were calculated in line with the stock annex of the Report of the Working Group on Celtic Seas Ecoregion (ICES, 2020) using the 2020 survey abundance (Table 3). The latest estimate of stock abundance (value from July 2020 survey, 750 million) is below the MSY  $B_{\text{trigger}}$  value (990 million). When the EU multiannual plan (MAP) for Western Waters and adjacent waters is applied (EU, 2019), catches in 2021 that correspond to the F ranges in the MAP are between 1238 and 1560 t (Table 4). This assumes that discard rates and fishery selection patterns do not change from the average of 2017–2019.

## Discussion

Since 2006 a dedicated annual UWTV survey has taken place which gives abundance estimates for this ground with high precision. The burrow abundance and mean density estimates have decreased in 2020 to the lowest level observed in the series. Fluctuations in density has also been observed in the adjacent FU20-21 and FU19 this year (Aristegui *et. al*, 2020 and Doyle *et. al*, 2020). Sudden declines followed by large increases in abundance have also been observed in other *Nephrops* stocks in the past (e.g. FU12 and FU13 in 2012-2013).

*Nephrops* in this area have been covered under the landings obligation since 2016 with several exemptions. Discard rates in weight for this FU have been around 12% in recent years. The provision of catch advice and scenarios for 2020 based on EU multiannual plan (MAP) for Western Waters assumes that discard rates and fishery selection patterns do not change from the average of 2017-2019.

The introduction of the landings obligation to *Nephrops* fisheries in 2016 should result in changes in selectivity. This is not taken into account in any of the catch advice because it is not possible to predict exactly what might happen. The main message is that any improvements in selectivity in the fishery and reductions in discards will result in increased mean weight in the catches. This will in turn reduce overall mortality on the stocks and allow for catch increases in the future.

An important objective of this UWTV survey was to collect various ancillary information. The occurrence of trawl marks on the footage is notable for two reasons. Firstly, it makes identification of *Nephrops* burrows more difficult as the trawl marks remove some signature features making accurate burrow identification more difficult. Secondly, only occupied *Nephrops* burrows will persist in heavily trawled grounds and it is assumed that each burrow is occupied by one individual *Nephrops* (ICES, 2008).

Monitoring the occurrence and frequency of sea-pens observed on this ground is important in the context of OSPAR's designations of sea-pen and burrowing megafauna communities as threatened. The sea-pen species *Virgularia mirabilis* which was seen in 2020 have been observed on previous surveys of FU22. Monitoring *Nephrops* stock and the benthic habitat is also important in the context of the MFSD indicators (e.g. sea floor integrity).

The main objectives of the survey were successfully met for the fifteenth successive year. The UWTV coverage and footage quality was excellent throughout the survey. This was mainly due to good survey planning to coincide with slack tides. The multi-disciplinary nature of the survey means that the information collected is highly relevant for a number of research and advisory applications.

The multi-disciplinary nature of the survey means that the information collected is highly relevant for a number of research and advisory applications.

## **Acknowledgments**

We would like to express our sincere thanks and gratitude to Colin McBrearty (Master) and crew of the RV. Celtic Voyager. Thanks to the onboard P&O technical staff Lukasz Pawlikowski who maintained the UWTV system throughout the survey. Thanks to Aodhán Fitzgerald, Rosemarie Butler (RVOPs) and Dave Tully (FEAS) at the Marine Institute for organising survey logistics, and also Gordon Furey and Damian Crean (P&O Maritime) for shore side support.

## **References**

- Aristegui, M. 2020. Image annotation R Shiny app. Marine Institute. <http://doi.org/d24n>
- Aristegui, M., Blaszkowski, M., Doyle, J., Hehir, I., Lynch, D., Ryan, G. and Lordan, C. 2020a. Porcupine Bank *Nephrops* Grounds (FU16) 2020 UWTV Survey Report and catch scenarios for 2021. Marine Institute UWTV Survey report.
- Aristegui, M., Doyle, J., O'Brien, S., Tully, D., McAuliffe, M., Fitzgerald, R., Fee, D., O'Connor, S., Galligan, S., Blaszkowski, M., Butler, R., and White., J. 2020b. FU19 *Nephrops*


- Grounds (FU19) 2020 UWTV Survey Report and catch scenarios for 2021. Marine Institute UWTV Survey report.
- Aristegui, M., Blaszkowski, M., Doyle, J., Fee, D., O'Connor, S. and White, J., 2020c. The Labadie, Jones and Cockburn Banks Nephrops Grounds (FU20-21) 2020 UWTV Survey Report and catch scenarios for 2021. Marine Institute UWTV Survey report.
- Campbell, N., Dobby, H., and Bailey, N. 2009. Investigating and mitigating uncertainties in the assessment of Scottish *Nephrops norvegicus* populations using simulated underwater television data. ICES Journal of Marine Science 66: 646–655. doi: 10.1093/icesjms/fsp046.
- EU. 2019. Regulation (EU) 2019/472 of the European Parliament and of the Council of 19 March 2019 establishing a multiannual plan for stocks fished in the Western Waters and adjacent waters, and for fisheries exploiting those stocks, amending Regulations (EU) 2016/1139 and (EU) 2018/973, and repealing Council Regulations (EC) No 811/2004, (EC) No 2166/2005, (EC) No 388/2006, (EC) No 509/2007 and (EC) No 1300/2008. Official Journal of the European Union, L 83: 1–17. <http://data.europa.eu/eli/reg/2019/472/oj>
- Gerritsen, H., and Lordan, C. 2011. Integrating vessel monitoring systems (VMS) data with daily catch data from logbooks to explore the spatial distribution of catch and effort at high resolution. ICES Journal of Marine Science: Journal du Conseil 68: 245–252. doi: <http://dx.doi.org/10.1093/icesjms/fsq137>.
- Gerritsen, H.D., Lordan, C., Minto, C. and Kraak, S.B.M. 2012. Spatial patterns in the retained catch composition of Irish demersal otter trawlers: High-resolution fisheries data as a management tool Fisheries Research, Volumes 129–130, October 2012, Pages 127–136. <http://dx.doi.org/10.1016/j.fishres.2012.06.019>
- ICES 2007. Report of the Workshop on the use of UWTV surveys for determining abundance in *Nephrops* stocks throughout European waters (WKNEPHTV). ICES CM: 2007/ACFM: 14 Ref: LRC, PGCCDBS.
- ICES 2009. Report of the Benchmark Workshop on *Nephrops* assessment (WKNEPH). ICES CM: 2009/ACOM:33
- ICES 2011. Report of the ICES Advisory Committee 2011. ICES Advice.2011. Book 1: Introduction, Overviews and Special Requests. Protocols for assessing the status of sea-pen and burrowing megafauna communities, section 1.5.5.3.
- ICES 2012. Report of the Study Group on *Nephrops* Surveys (SGNEPS). ICES CM 2012/SSGESST: 19. Ref: SCICOM, ACOM
- ICES 2018. Report of the Workshop of *Nephrops* Burrow Counting (WKNEPS) ICES CM: 2018/EOSG:25.
- ICES 2020. Working Group for the Celtic Seas Ecoregion (WGCSE). ICES Scientific Reports. 2:40. xx pp. <http://doi.org/10.17895/ices.pub.5978>
- Lin, L, I. 1989. A Concordance Correlation Coefficient to Evaluate Reproducibility. Biometrics 45, 255-268.
- Pebesma, E.J., R.S. Bivand, 2005. Classes and methods for spatial data in R. R News 5 (2), <https://cran.r-project.org/doc/Rnews/>




Petitgas P. and Lafont, T, 1997. EVA (Estimation VAriance). A geostatistical software on IBM-PC for structure characterization and variance computation. Version 2.

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>



Renard D., Bez N., Desassis N., Beucher H., Ors F., Laporte F. 2015. RGeostats: The Geostatistical package [version:11.0.3]. MINES ParisTech. Free download from: <http://cg.ensmp.fr/rgeostats>









**Figure 1:** FU22 Smalls grounds: *Nephrops* Functional Units (FUs) and *Nephrops* grounds (area polygons) in the greater Celtic Sea.





**Figure 2:** FU22 Smalls grounds: TV stations completed on the 2020 survey.




**Figure 3** : FU22 Smalls grounds: R - tool quality control plot for station 120 of the 2020 survey.




**Figure 4** : FU22 Smalls grounds: Lin's CCC quality control plots of count data for stations 120 to 122 of the 2020 survey.




**Figure 5** : FU22 Smalls grounds: Plots of the variability in density between minutes (top panel) and between operators (counters) (bottom panel) for each station in 2020



**Figure 6:** FU22 Smalls grounds: Violin and box plot of adjusted burrow density distributions by year from 2006 -2020. The blue line indicates the mean density over time. The horizontal black lines represent medians, white boxes the inter quartile ranges, the black vertical lines the range and the black dots are outliers.




**Figure 7:** FU22 Smalls grounds: Contour plots of the krigged density estimates by year from 2006 (top left) - 2020 (bottom left).




**Figure 8:** FU22 Smalls grounds: Time series of geo-statistical adjusted abundance estimates (in millions of burrows). The error bars indicate the 95% confidence intervals and  $B_{\text{trigger}}$  is dashed blue line.





**Figure 9:** FU22 Smalls grounds: Standardised length frequency distributions for male and female *Nephrops* caught using beam trawl during 2006 to 2018 UWTV surveys (except years 2010, 2013 – 2015, 2019 - 2020). Blue line indicates minimum conservation reference size 25 mm carapace length.



**Figure 10:** FU22 Smalls grounds: 2020 stations where *Virgularia mirabilis* (VAM) were identified. Closed circles indicated presence and open circles denotes TV stations with no sea-pen observations.

**Table 1:** FU22 Smalls grounds: Overview of geostatistical results from 2006-2020.

| Year  | Number of stations | Mean Density adjusted (burrow/m <sup>2</sup> ) | Domain Area (km <sup>2</sup> ) | Geostatistical Abundance adjusted (millions of burrows) | CV on Burrow estimate (%) |
|-------|--------------------|--|--------------------------------|---|---------------------------|
| 2006  | 100                | 0.49   | 2962                           | 1503  | 2                         |
| 2007  | 107                | 0.37   | 2955                           | 1136  | 6                         |
| 2008  | 76                 | 0.36   | 2698                           | 1114  | 6                         |
| 2009  | 67                 | 0.36   | 2824                           | 1093  | 5                         |
| 2010  | 90                 | 0.37   | 2861                           | 1141  | 4                         |
| 2011  | 107                | 0.41   | 2881                           | 1256  | 3                         |
| *2012 | 47                 | 0.49   | 2934                           | 1498  | 8                         |
| *2013 | 41                 | 0.41   | 2975                           | 1254  | 7                         |
| *2014 | 52                 | 0.53   | 2970                           | 1622  | 8                         |
| *2015 | 40**               | 0.49   | 3064                           | 1363  | 7                         |
| *2016 | 41                 | 0.31   | 3063                           | 866   | 7                         |
| *2017 | 40                 | 0.55   | 3063                           | 1600  | 5                         |
| *2018 | 42                 | 0.31   | 3063                           | 876   | 9                         |
| *2019 | 41                 | 0.40   | 3063                           | 1121  | 6                         |
| *2020 | 40                 | 0.27   | 3063                           | 750   | 8                         |

\*reduced randomised isometric grid

\*\* In 2015 seven of the stations were filled in with an estimate based on the mean density of historical stations within 2 nmi of the planned station.

**Table 2:** FU22 Smalls grounds: Inputs to catch scenarios table.

| Year | UWTV abundance estimate | 95% Confidence Interval | Landings in number | Total discards in number* | Removals in number | Harvest rate (by number) | Landings | Total discards* | Discard rate (by number) | Dead discard rate (by number) | Mean weight in landings | Mean weight in discards |
|------|-------------------------|-------------------------|--------------------|---------------------------|--------------------|--------------------------|----------|-----------------|--------------------------|-------------------------------|-------------------------|-------------------------|
|      | millions                |                         |                    | %                         |                    |                          | tonnes   |                 | %                        |                               | grammes                 |                         |
| 2003 |                         |                         | 95                 | 68                        | 146                |                          | 2065     | 720             | 41.5                     | 34.7                          | 21.7                    | 10.7                    |
| 2004 |                         |                         | 71                 | 13                        | 80                 |                          | 1828     | 202             | 15.6                     | 12.2                          | 25.9                    | 15.4                    |
| 2005 |                         |                         | 119                | 129                       | 216                |                          | 2533     | 1648            | 51.9                     | 44.7                          | 21.2                    | 12.8                    |
| 2006 | 1503                    | 70                      | 100                | 45                        | 134                | 8.9                      | 1761     | 454             | 31.1                     | 25.3                          | 17.6                    | 10.1                    |
| 2007 | 1136                    | 126                     | 165                | 181                       | 301                | 26.5                     | 2950     | 1906            | 52.3                     | 45.1                          | 17.9                    | 10.5                    |
| 2008 | 1114                    | 123                     | 144                | 26                        | 163                | 14.6                     | 3090     | 289             | 15.3                     | 12.0                          | 21.5                    | 11.1                    |
| 2009 | 1093                    | 108                     | 92                 | 33                        | 117                | 10.7                     | 2185     | 371             | 26.4                     | 21.2                          | 23.7                    | 11.3                    |
| 2010 | 1141                    | 88                      | 122                | 45                        | 155                | 13.6                     | 2714     | 636             | 26.8                     | 21.5                          | 22.3                    | 14.3                    |
| 2011 | 1256                    | 72                      | 60                 | 13                        | 70                 | 5.6                      | 1636     | 196             | 18.0                     | 14.1                          | 27.3                    | 14.9                    |
| 2012 | 1498                    | 239                     | 120                | 31                        | 144                | 9.6                      | 2618     | 347             | 20.7                     | 16.3                          | 21.8                    | 11.1                    |
| 2013 | 1254                    | 177                     | 94                 | 40                        | 124                | 9.9                      | 2257     | 497             | 30.0                     | 24.3                          | 24.1                    | 12.4                    |
| 2014 | 1622                    | 268                     | 100                | 33                        | 125                | 7.7                      | 2526     | 460             | 25.0                     | 20.0                          | 25.2                    | 13.8                    |
| 2015 | 1363                    | 180                     | 114                | 44                        | 147                | 10.8                     | 2350     | 450             | 28.0                     | 22.6                          | 20.6                    | 10.1                    |
| 2016 | 866                     | 112                     | 160                | 54                        | 200                | 23.1                     | 3329     | 519             | 25.1                     | 20.0                          | 20.8                    | 9.7                     |
| 2017 | 1600                    | 153                     | 164                | 39                        | 194                | 12.1                     | 3560     | 424             | 19.2                     | 15.2                          | 21.7                    | 10.8                    |
| 2018 | 876                     | 154                     | 98                 | 31                        | 121                | 13.8                     | 1975     | 350             | 23.7                     | 19.0                          | 20.2                    | 11.2                    |
| 2019 | 1121                    | 141                     | 81                 | 19                        | 95                 | 8.5                      | 2083     | 262             | 19.1                     | 15.1                          | 25.8                    | 13.7                    |
| 2020 | 750                     | 118                     |                    |                           |                    |                          |          |                 |                          |                               |                         |                         |

**Table 3:** The basis for the catch scenarios.

| Variable                          | Value | Notes   |
|-----------------------------------|-------|---|
| Stock abundance (2021)            | 750   | Number of individuals (million); UWTV survey 2020 |
| Mean weight in projected landings | 22.5  | Average 2017–2019 in grammes                      |
| Mean weight in projected discards | 11.8  | Average 2017–2019 in grammes                      |
| Projected discards                | 20.7  | Proportion by number; average 2017–2019           |
| Discards survival*                | 25    | Proportion by number                              |
| Projected dead discards           | 16.4  | Proportion by number; average 2017–2019           |

\*Only applied in scenarios where discarding is allowed

**Table 4:** FU22 Smalls grounds: Annual catch advice and scenarios; Discarding assumed to continue at recent average.

| Basis   | Total catch    | Dead removals | Projected landings | Projected dead discards | Projected surviving discards | Harvest rate * % | % advice change ** |
|---|----------------|---------------|--------------------|-------------------------|------------------------------|------------------|--------------------|
|   | PL + PDD + PSD | PL + PDD      | PL                 | PDD                     | PSD                          | for PL + PDD     |                    |
| ICES advice basis   |                |               |                    |                         |                              |                  |                    |
| MSY approach; F = EU MAP <sup>^</sup> : $F_{MSY} \times Stock$ Abundance 2020 / $MSY B_{trigger}$ | 1560           | 1512          | 1371               | 141                     | 47                           | 9.7              | -45                |
| MAP $F_{MSY lower} \times Stock$ Abundance 2020 / $MSY B_{trigger}$                               | 1238           | 1201          | 1088               | 112                     | 37                           | 7.7              | -45                |
| MAP $F_{MSY upper} \times Stock$ Abundance 2020 / $MSY B_{trigger}$                               | 1560           | 1512          | 1371               | 141                     | 47                           | 9.7              | -45                |
| Other options   |                |               |                    |                         |                              |                  |                    |
| F = MAP $F_{MSY}$   | 2058           | 1996          | 1809               | 187                     | 62                           | 12.8             | -27                |
| F = MAP $F_{MSY lower}$   | 1640           | 1591          | 1442               | 149                     | 50                           | 10.2             | -42                |
| F = MAP $F_{MSY upper}$ ***   | 2058           | 1996          | 1809               | 187                     | 62                           | 12.8             | -27                |
| $F_{2019}$  | 1366           | 1325          | 1201               | 124                     | 41                           | 8.5              | -52                |

<sup>^</sup> EU multiannual plan (MAP) for Western waters (EU, 2019).

\* By number.

\*\* Advice value for 2021 relative to the corresponding 2020 values (MAP advice value of 2820, 2247 and 2820 tonnes, respectively; other values are relative to  $F_{MSY}$ ).

\*\*\*  $F_{MSY upper} = F_{MSY}$  for this stock.