



Research Paper

Action of low doses of Aspirin in Inflammation and Oxidative Stress induced by $A\beta_{1-42}$ on Astrocytes in primary culture

Adrian Jorda^{1,2*}, Martin Aldasoro^{1*}, Constanza Aldasoro¹, Sol Guerra-Ojeda¹, Antonio Iradi¹, Jose M^a Vila¹, Juan Campos-Campos^{1,2}, Soraya L. Valles¹✉

1. Department of Physiology, School of Medicine, University of Valencia, Spain.
2. Faculty of Nursing and Podiatry, University of Valencia, Spain.

*These authors contributed equally to this work.

✉ Corresponding author: Dr. Soraya L. Valles. Department of Physiology. School of Medicine, University of Valencia. email: lilian.valles@uv.es Phone: 0034-963983813.© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>). See <http://ivyspring.com/terms> for full terms and conditions.

Received: 2019.10.07; Accepted: 2019.12.23; Published: 2020.03.13

Abstract

Aspirin has been used as anti-inflammatory and anti-aggregate for decades but the precise mechanism(s) of action after the presence of the toxic peptide $A\beta_{1-42}$ in cultured astrocytes remains poorly resolved. Here we use low-doses of aspirin (10^{-7} M) in astrocytes in primary culture in presence or absence of $A\beta_{1-42}$ toxic peptide. We noted an increase of cell viability and proliferation with or without $A\beta_{1-42}$ peptide presence in aspirin treated cells. In addition, a decrease in apoptosis, determined by Caspase 3 activity and the expression of Cyt c and Smac/Diablo, were detected. Also, aspirin diminished necrosis process (LDH levels), pro-inflammatory mediators (IL- β and TNF- α) and NF- κ B protein expression, increasing anti-inflammatory PPAR- γ protein expression, preventing $A\beta_{1-42}$ toxic effects. Aspirin inhibited COX-2 and iNOS without changes in COX-1 expression, increasing anti-oxidant protein (Cu/Zn-SOD and Mn-SOD) expression in presence or absence of $A\beta_{1-42}$. Taken together, our results show that aspirin, at low doses increases cell viability by decreasing inflammation and oxidative stress, preventing the deleterious effects of the $A\beta_{1-42}$ peptide on astrocytes in primary culture. The use of low doses of aspirin may be more suitable for Alzheimer's disease.

Key words: Amyloid- β ; aspirin; inflammation; oxidative stress; Alzheimer's disease

Introduction

Alzheimer's disease (AD) causes decline in memory and is the most common neurodegenerative disease implicated in the aging process [1]. The prominent features of AD include amyloid plaques, intraneuronal tangles, cell death, inflammatory changes and oxidative stress [2,3,4].

Astroglia are the most prevalent cell type in the brain [5]. These cells have roles in the brain protecting against CNS injury and repairing nervous tissue after injury [6]. Data from our laboratory demonstrated that astrocytes increase neuronal viability and mitochondrial biogenesis, protecting from oxidative stress and inflammation induced by toxic amyloid

peptide [7,8]. Also, astrocytes act in neuronal synapses, regulate the blood-brain barrier, providing nutrients to the nervous system and maintaining ion and metabolite balance, and also propagate calcium currents, release gliotransmitters, growth factors and inflammatory mediators [9,10,11]. Astroglialosis is produced by a reaction from astrocytes to inflammation, oxidative stress and cell death producing toxic products and inflammatory agents and oxidative stress mediators [3,12]. In astrocytes, complex changes and specific conflicts occur in different brain regions during the development of AD. The number of reactive astrocytes increases in

AD, phagocytizing and reducing amyloid β ($A\beta$) deposition because these cells surround amyloid plaques and secrete proinflammatory factors [13,14].

Acetylsalicylic acid (aspirin) is used frequently as a member of the nonsteroidal anti-inflammatory drugs (NSAIDs) group [15]. Aspirin induces its effects by inhibiting cyclooxygenase (COX) and suppresses prostaglandins [16,17]. Two main isoforms of COX exist, COX-1 and COX-2. COX-1 is involved in the synthesis of thromboxane A_2 (TXA_2) [18] and COX-2 in prostacyclin biosynthesis [19,20]. In epidemiological studies of AD, a high dose of aspirin produces lower prevalence of AD [21] (Nilsson *et al.* 2003). Other authors using low-dose aspirin treatment indicated promising results for aspirin [22]. In this study, we were interested in exploring the action of aspirin in both inflammatory and ROS (reactive oxygen species) events associated with Alzheimer's disease (AD) by using the amyloid β_{1-42} in astrocytes in primary culture. The accumulation and precipitation of $A\beta_{1-42}$ peptide has a neuropathological role associated with AD. However the $A\beta_{40-1}$ peptide has non-toxic effects and is used as control.

Material and Methods

Materials

This study was approved by the Bioethics Committee of the School of Medicine of the University of Valencia, and the local Government of Valencia, Spain (2016/VSC/PEA/00220). All animals (Wilson rats from Charles River laboratory) were handled according to the recommendations of the Committee and with access to food and water. Animals were sacrificed with pentobarbital by the veterinary personnel. No randomization was performed to allocate subjects in the study. Rats (6 months old) were housed alone for 21 days during pregnancy. Female rat fetuses were obtained at day 21 of pregnancy prior to delivery. The fetal cortex was obtained to create a culture of astrocytes. Every assay was performed 3, 4 or 5 times, from different mother rats. No blinding procedures for each culture were made. Female rats were excluded if they were pregnant before the study. Rats were also excluded if the delivery of rats was 21 days before pregnancy.

Dulbecco's modified Eagle's medium (DMEM) and fetal bovine serum (FBS) were obtained from Gibco (Gibco Invitrogen Corporation, Barcelona, Spain). The oligomers $A\beta$ (40-1 and 1-42), were prepared following manufacturer's instructions (Sigma-Aldrich biotechnology). Briefly, the peptides were dissolved in 100 μ M phosphate buffered saline (PBS) and preparations were heated for 24 h at 37°C

for assembly of the oligomers. Aspirin was obtained from Sigma-Aldrich biotechnology and dissolved in Krebs solution to the proper final concentration (10^{-7} M). 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) was purchased from Sigma Chemical Co. (St Louis, MO). Enzyme-linked immunosorbent assay (ELISA) kits for IL-1 β (Interleukin 1- β) and TNF- α (Tumor necrosis- α) from Pierce Biotechnology, Inc. (Rockford, USA). Western Blot Chemiluminescent Detection System (ECL) was from Amersham (Amersham Biosciences, Barcelona, Spain). Monoclonal anti-cytochrome C (Cyt c) antibody (1:500), monoclonal anti-Smac/Diablo antibody (Smac/Diablo) (1:500), monoclonal anti-nuclear factor κ B antibody (NF- κ B) (1:1000), monoclonal anti-Mn superoxide dismutase antibody (Mn-SOD) (1:500), monoclonal anti-cyclooxygenase 1 antibody (COX-1) (1:500), monoclonal anti-cyclooxygenase 2 antibody (COX-2) (1:500), monoclonal anti-inducible nitric oxide synthase antibody (iNOS) (1:500) from Santa Cruz Biotechnology (Madrid, Spain). Monoclonal anti-peroxisome proliferator-activated receptor antibody (PPAR- γ) (1:500) from Sigma Aldrich (Madrid, Spain). Polyclonal anti-Cu/Zn superoxide dismutase antibody (Cu/Zn-SOD) (1:500) from Assay Designs (Madrid, Spain). Monoclonal anti-tubulin antibody (1:1000) from Cell Signaling (Beverly, MA, USA). All other reagents were analytical or culture grade purity.

Methods

Primary Culture of Cortical Astrocytes

Cerebral cortical astrocytes were isolated from rat fetuses of 21 days gestation. Fetuses were obtained by caesarean section and decapitated. Cerebral cortices were removed and triturated 10-15 times through a Pasteur pipette. The cell suspension was filtered through nylon mesh with a pore size of 90 μ m and diluted in DMEM containing 20% fetal bovine serum (FBS) supplemented with L-glutamine (1%), HEPES (10 mM), fungizone (1%), and antibiotics (1%). Cells were plated on T75 culture flask pretreated with poli-L-lysine. Cultures were maintained in a humidified atmosphere of 5% CO $_2$ /95% air at 37°C during 20 days. After 1 week of culture, the FBS content was reduced to 10%, and the medium was changed twice a week. By immunocytochemistry, 97% of cells are GFAP positive (data not shown).

Four groups were used. Group A, control received $A\beta_{40-1}$ peptide, Group B, $A\beta_{40-1}$ peptide + aspirin, Group C, $A\beta_{1-42}$ toxic peptide and Group D, $A\beta_{1-42}$ toxic peptide + aspirin. Initially, we used $A\beta_{42-1}$ but due to its high cost, we assayed $A\beta_{40-1}$ as we have used before [7].

MTT assay

Cell viability of the cultures was determined by the MTT assay. Cells were plated in 96 well culture plate and incubated with Asp during 24 h at 10^{-11} M, 10^{-9} M, 10^{-7} M, 10^{-5} M, $A\beta_{1-42}$ 15 μ M or with $A\beta_{1-42}$ 15 μ M + 10^{-7} M Asp. After cell treatments, the medium was removed and the cortical cells were incubated with red free medium and MTT solution [0.5 mg/ml, prepared in phosphate buffer saline (PBS) solution] for 4 h at 37°C. Finally the medium was removed and formazan particles were dissolved in dimethyl sulfoxide (DMSO). Cell viability, defined as the relative amount of MTT reduction, was determined by spectrophotometry at 570 nm.

Trypan Blue Assay

Trypan blue exclusion assay was used to count the living cells and monitor cell proliferation. Astrocytes were isolated and seeded at 7×10^4 cells/35 mm dish. After 5 days of culture, cells were incubated without (control, C), with Asp (10^{-7} M), $A\beta_{1-42}$ 15 μ M or with $A\beta_{1-42}$ 15 μ M + 10^{-7} M Asp for 24 h. 1.5% trypan blue solution was applied to astrocytes cultures at room temperature for 3 min.

Lactate Dehydrogenase (LDH) Assay

To evaluate plasma membrane integrity, LDH release was determined by monitoring the leakage of the cytosolic LDH to the extracellular medium. LDH was measured spectrophotometrically at 340 nm, following the rate of conversion of reduced nicotinamide adenine dinucleotide to oxidized nicotinamide adenine dinucleotide.

Caspase 3 Activity Assay

Caspase 3 activity was measured in cytosolic fractions by using a highly sensitive colorimetric substrate, N-acetyl-Asp-Glu-Val-Asp p-nitroanilide (Ac-DEVD-pNA) following manufacturer's instructions (CalBiochem, La Jolla, CA). Enzyme activity was calculated using manufacturer's formulae, as pmol/min.

Cytokine Determination, IL-1 β and TNF α

Cells were seeded, and at time of assay, the red phenol medium was removed and replaced by PBS containing 1 mg/ml bovine serum albumin (BSA), either in the presence or absence of Asp (10^{-7} M), with $A\beta_{1-42}$ 15 μ M or $A\beta_{1-42}$ 15 μ M + 10^{-7} M Asp). IL-1 β and TNF- α concentration (pg/ml) were ascertained using ELISA kits (Pierce Biotechnology, Inc.).

Western Blot Analysis

Cultured cells were treated with lysis buffer and then mechanically degraded to release the proteins. Protein concentration was determined using modified

Lowry method. Loading buffer (0.125 M Tris-HCl, pH 6.8, 2% SDS, 0.5% (v/v) 2-mercaptoethanol, 1% bromophenolblue and 19% glycerol) was added to protein sample and heated for 5 min at 95°C. Proteins (20 μ g) were separated on SDS-PAGE gels and transferred to nitrocellulose membranes in a humid environment using a transfer buffer (25 mM Tris, 190 mM glycine, and 20% methanol). Membranes were blocked with 5% milk in TBS-T (0.05% Tween-20) and incubated with primary antibodies overnight at 4°C. Membranes were washed 3 times with wash buffer TBS-T (TBS, 0.2% Tween-20) and incubated with a secondary anti-rabbit IgG or anti-mouse IgG antibody conjugated to the enzyme horseradish peroxidase (HRP) for 1 h. Membranes were washed three times and proteins were detected using the ECL method as specified by the manufacturer. Autoradiography signals were assessed using digital image system ImageQuant LAS 4000 (GE Healthcare). Densitometry is the quantitative measurement of optical density in a photographic paper or photographic film, due to exposure to light. Concentration of protein was determined by densitometry analysis, expressed as arbitrary units relative to tubulin.

Statistical Analyses

All values are expressed as mean \pm S.D. The differences between groups were determined with unpaired Student's t-test. All statistical analyses were performed using the GraphPad Prism software (GraphPad Software Inc., San Diego, CA, USA). Statistical significance was accepted at $p \leq 0.05$.

Results

Asp and Cell Viability

The role of Asp on cell viability was studied using MTT conversion assay. Fig. 1 shows that incubation with Asp at 10^{-11} M, 10^{-9} M, and 10^{-7} M significantly increased astrocyte viability *vs* control. On the other hand, $A\beta_{1-42}$ significantly decreased cell viability (30%) compared to control cells. After incubation with $A\beta_{1-42}$ + 10^{-7} M Asp, no significant changes were detected compared to control astrocytes and contrarily, an increase in cell viability was detected compared to cells with $A\beta_{1-42}$ peptide alone.

Trypan blue exclusion assay was used to count the living cells and monitor cell proliferation. Astrocytes were isolated and seeded at 7×10^4 cells/35 mm dish. After 5 days of culture, cells were incubated without (control, C) or with Asp 10^{-7} M, $A\beta_{1-42}$ 15 μ M or $A\beta_{1-42}$ 15 μ M + Asp 10^{-7} M for 24 h. In control conditions proliferation was 0.93%, and previous incubation with Asp (10^{-7} M) increased proliferation by 9.53%. On the other hand, in presence of $A\beta_{1-42}$

proliferation decreased 12.96% and with $A\beta_{1-42}$ + Asp 10^{-7} M only decreased 5.37% (Table 1).

Table 1. Astrocytes proliferation and counting living cells

	Seeding cells	5 days of culture	After 24 h of incubation	% Proliferation
C	7	12.72±0.19	12.83±0.20	+0.93
Asp (10^{-7} M)	7	12.70±0.11	13.91±0.16*	+9.53
$A\beta$	7	12.65±0.23	11.01±0.32*	-12.96
$A\beta$ + Asp (10^{-7} M)	7	12.75±0.24	12.10±0.14*	-5.37

Astrocytes were isolated and seeded at 7×10^4 cells/35 mm dish during 5 days. At this time, cells were incubated without Asp (control, C), with Asp (10^{-7} M), Amyloid β_{1-42} (15 μ M) or Amyloid β_{1-42} (15 μ M) + Asp (10^{-7} M) for 24 h. Trypan blue exclusion was used to count the living cells and monitor cell proliferation. Data are mean \pm SD of five independent experiments (four different rats). * $p < 0.05$ vs. control.




Figure 1. Cell viability was determined by MTT assay in cells treated during 24 h. Astrocytes were incubated without Asp (control, C), with Asp at different concentrations (10^{-11} , 10^{-9} , 10^{-7} and 10^{-5} M), with $A\beta_{1-42}$ (15 μ M) or $A\beta_{1-42}$ (15 μ M) + Asp (10^{-7} M) for 24 h. Data are means \pm SD of four independent experiments (three different rats). * $p < 0.05$ vs. control. # $p < 0.05$ vs $A\beta_{1-42}$ treated cells.

LDH and Caspase 3

Incubation of the astrocytes with Asp 10^{-7} M for 24 h decreased significantly LDH values (21%) compared with control cells. With $A\beta_{1-42}$ (15 μ M) an increase of LDH release (55%) was detected compared with control cells and this data was reversed with Asp (10^{-7} M) to control values (Fig. 2A).

Incubation with Asp 10^{-7} M for 24 h, decreased Caspase 3 activity significantly (45%) compared to control cells, whereas $A\beta_{1-42}$ (15 μ M) activity was increased in presence of Asp (10^{-7} M), and prevented the toxic peptide effect (Fig 2B), indicating reduction of apoptosis when Asp is present on the culture with $A\beta_{1-42}$.

Cytochrome c and Smac/DIABLO Expression

Fig. 3A shows cytochrome c expression in astrocytes in primary culture. Asp decreased cytochrome c expression by 2.2-fold at 10^{-7} M. $A\beta_{1-42}$ increased cytochrome c expression compared to control cells. On the contrary, Asp (10^{-7} M) significantly reversed the cytochrome c increase expression induced by $A\beta_{1-42}$. Fig. 3B shows

Smac/Diablo expression in astrocytes in primary culture. Asp significantly decreased Smac/Diablo protein expression compared to control cells. Addition of $A\beta_{1-42}$ significantly increased Smac/Diablo expression (1.3-fold) compared with control cells. Furthermore, Asp 10^{-7} M decreased Smac/Diablo expression induced by $A\beta_{1-42}$ to control values. Consequently Asp addition produced a protective effect against the amyloid toxic peptide (Fig. 3).

IL-1 β and TNF- α Pro-inflammatory Cytokines

Secretion of the pro-inflammatory mediators, IL-1 β and TNF- α , were detected by ELISA. Fig 4 shows that, in astrocytes, Asp (10^{-7} M) decreased 1.6-fold IL-1 β release compared with control values (Fig 4A). Conversely, $A\beta_{1-42}$ addition increased 2.5-fold IL-1 β secretion compared with control cells. $A\beta_{1-42}$ + Asp 10^{-7} M decreased IL-1 β liberation to control values.

Fig. 4B shows that Asp (10^{-7} M) produced a significant decrease of TNF- α secretion (3.1-fold) compared to control cells. $A\beta_{1-42}$ addition increased TNF- α liberation 1.5-fold. Moreover, Asp (10^{-7} M) prevented this toxic effect produced by $A\beta_{1-42}$ peptide.

NF- κ B and PPAR- γ Expression

NF- κ B is a transcription factor that regulates positively gene expression of pro-inflammatory proteins. Fig. 5A shows that Asp decreased this protein expression about 1.25-fold compared to control astrocytes. When $A\beta_{1-42}$ was added an increase (1.5-fold) of this transcription factor was detected compared to control results. Asp (10^{-7} M) addition returned NF- κ B to control values.

PPARs family regulates negatively gene expression of pro-inflammatory proteins. Fig. 5B shows PPAR- γ expression in astrocytes in culture. Asp increased PPAR- γ expression 1.5-fold at 10^{-7} M. Addition of $A\beta_{1-42}$ decreased 1.6-fold this transcription factor compared to control results. Asp (10^{-7} M) addition returned PPAR- γ expression to control values.

COX-2 and iNOS Expression

In Fig. 6, we detected a reduction of COX-2 (panel A) expression after addition of Asp (10^{-7} M) compared with control values. The presence of the toxic peptide $A\beta_{1-42}$ increased the expression of COX-2 that was reversed by the incubation with Asp 10^{-7} M to control values. On the contrary, no changes were detected in COX-1 expression at any experimental condition (Data not shown). Fig. 6 (panel B) demonstrates a significant decrease in iNOS protein expression after Asp 10^{-7} M addition compared to

control. $A\beta_{1-42}$ produced a significant increase of iNOS expression. On the other hand, the presence of both the toxic peptide and Asp 10^{-7} M reduced significantly iNOS expression compared to $A\beta_{1-42}$.

Expression of CU/ZN-SOD and Mn-SOD Proteins

In astrocytes, Asp 10^{-7} M increased Cu/Zn-SOD (panel A) and Mn-SOD (panel B) expression compared to control cells (Fig. 7). In the presence of the $A\beta_{1-42}$ toxic peptide, a decrease of both proteins compared with control values was reversed by Asp.




Figure 2. Lactate dehydrogenase and caspase 3 activity. Astrocytes were incubated without Asp (control, C), with Asp (10^{-7} M), $A\beta_{1-42}$ (15 μ M) or $A\beta_{1-42}$ (15 μ M) + Asp (10^{-7} M) for 24 h. Panel A: Lactate dehydrogenase from supernatants of astrocytes. Panel B: Caspase 3 activity. Data are means \pm SD of four independent experiments (four different rats). * $p < 0.05$ vs. control cells. # $p < 0.05$ vs. $A\beta_{1-42}$ treated cells.




Figure 3. Cytochrome c and Smac/Diablo expression. Astrocytes were incubated without Asp (control, C), with Asp (10^{-7} M), $A\beta_{1-42}$ (15 μ M) or $A\beta_{1-42}$ (15 μ M) + Asp (10^{-7} M) for 24 h and collected to determine Cytochrome c (panel A) or Smac/Diablo (panel B) protein expression by Western-blot. A representative immunoblot is shown in the top panel. Data are means \pm SD of four independent experiments (four different rats). * $p < 0.05$ vs. control cells. # $p < 0.05$ vs. $A\beta_{1-42}$ treated cells.




Figure 4. IL-1 β and TNF- α determination. Astrocytes were incubated without Asp (control, C), with Asp (10^{-7} M), $A\beta_{1-42}$ (15 μ M) or $A\beta_{1-42}$ (15 μ M) + Asp (10^{-7} M). Cell culture supernatants were harvested and IL-1 β (Panel A) and TNF- α (Panel B) were determined by ELISA. Values are means \pm SD from four independent experiments (four different rats). * $p < 0.05$ vs. control. # $p < 0.05$ vs. $A\beta$ treated cells.




Figure 5. NF-κB and PPAR-γ expression. Astrocytes were incubated without Asp (control, C), with Asp (10^{-7} M), $A\beta_{1-42}$ ($15 \mu\text{M}$) or $A\beta_{1-42}$ ($15 \mu\text{M}$) + Asp (10^{-7} M) for 24 h and collected to determine NF-κB (panel A) or PPAR-γ (panel B) protein expression by Western-blot. A representative immunoblot is shown in the top panel. Data are means \pm SD of four independent experiments (four different rats). * $p < 0.05$ vs. control cells. # $p < 0.05$ vs. $A\beta_{1-42}$ treated cells.




Figure 6. COX-2 and iNOS expression. Astrocytes were incubated without Asp (control, C), with Asp (10^{-7} M), $A\beta_{1-42}$ ($15 \mu\text{M}$) or $A\beta_{1-42}$ ($15 \mu\text{M}$) + Asp (10^{-7} M) for 24 h and collected to determine COX-2 (panel A) or iNOS (panel B) protein expression by Western-blot. A representative immunoblot is shown in the top panel. Data are means \pm SD of four independent experiments (four different rats). * $p < 0.05$ vs. control cells. # $p < 0.05$ vs. $A\beta_{1-42}$ treated cells.




Figure 7. Cu/Zn-SOD and Mn-SOD expression. Astrocytes were incubated without Asp (control, C), with Asp (10^{-7} M), $A\beta_{1-42}$ ($15 \mu\text{M}$) or $A\beta_{1-42}$ ($15 \mu\text{M}$) + Asp (10^{-7} M) for 24 h and collected to determine Cu/Zn-SOD (panel A) or Mn-SOD (panel B) protein expression by Western-blot. A representative immunoblot is shown in the top panel. Data are means \pm SD of four independent experiments (four different rats). * $p < 0.05$ vs. control cells. # $p < 0.05$ vs. $A\beta_{1-42}$ treated cells.

Discussion

In this study we show that aspirin, at low-doses, protects from $A\beta_{1-42}$ toxic peptide actions in astrocytes in primary culture, indicating the convenience to use low doses to obtain better benefits of aspirin. The

aspirin increases cell viability and proliferation, decreases apoptosis (Caspase 3, Cyt c and Smac/Diablo) and necrosis (LDH), in the presence or absence of $A\beta_{1-42}$ peptide. Moreover, the aspirin decreases pro-inflammatory mediators (IL- β and TNF- α) and NF- κ B expression and increases anti-

inflammatory PPAR- γ protein after addition of A β ₁₋₄₂. As also inhibits COX-2 and iNOS without changes in COX-1 expression and increases anti-oxidant proteins (Cu/Zn-SOD and Mn-SOD) expression in the presence or absence of A β ₁₋₄₂.

The role of astrocytes in the brain has been reviewed [23,24,25]. It is reported that astrocytes protect neurons against A β -amyloid peptide, decreasing inflammation, and oxidative stress, and increasing cell viability and mitochondrial biogenesis [7,8,26].

Low-dose of aspirin has been reported to reduce the incidence of Alzheimer's disease [27] and also the donation of its acetyl group to molecules that suppress protein aggregation in neurodegenerative diseases, including AD [28]. This mechanism could be implicated in the astrocytic protection observed in our study. A β ₁₋₄₂ peptide induces reduction in cell viability and increases apoptosis by cytochrome c and Smac/Diablo pathways [7,29]. However, simultaneous treatment with aspirin of A β ₁₋₄₂ treated cells protects them from neurotoxicity [29]. Reactive gliosis increases glial fibrillary acidic protein (GFAP) in AD where it is highly expressed and remarkably, aspirin produces a reduction in GFAP synthesis by blocking NF- κ B in astrocytes culture [30]. It is reported that high doses aspirin can help reduce chronic AD inflammation [30]. However, our results show that aspirin, at low concentration (10^{-7} M), increases cell viability, diminishing necrosis and apoptosis. Burke *et al.* (2006) [31] reported that human daily ingestion of aspirin, 80 to 350 mg, produces plasma salicylate levels of 2.4 to 9.7 μ g/ml, corresponding to 1.3×10^{-6} to 5.4×10^{-5} M of aspirin in the culture medium [31]. According to these results, the concentrations used in our experiments correspond to doses lower than 70 mg/day, considered as low doses of aspirin [32]. However, aspirin increases apoptosis in gastric mucosal cell line in a caspase-dependent manner through Smac/Diablo pathway [33]. Our results differ due to aspirin concentration used. Redlak *et al.* (2005) [33] used a concentration of 4×10^{-2} M whereas in our study we used 10^{-7} M.

Our results indicate that low concentrations of aspirin may decrease apoptosis as opposed to high doses, protecting without increasing the adverse mechanisms observed in other paper.

After A β ₁₋₄₂ addition, we detected an increase in COX-2 without changes in COX-1 expression. Both COX proteins could lead to adverse cellular effects derived from TXA₂ and prostacyclin biosynthesis. Using LPS (bacterial lipopolysaccharide) to determine COX isoform expression in astrocytes, Font-Nieves *et al.* (2012) [34] showed a strong induction of COX-2

through an NF- κ B-dependent mechanism [34]. In chronic inflammation and in the progression of neurodegenerative diseases, NF- κ B activation plays an important role in astrocytes in primary culture [35], according to previous data obtained in our laboratory indicating that A β addition is also associated with an increase in NF- κ B activity in astrocytes [7]. Lee *et al.* (2018) [36], have indicated that downregulation of inflammatory molecules such as iNOS and different cytokines (TNF- α , IL-1 β and IL-6) could be a consequence of the NF- κ B and MAP kinases inhibition in astrocytes [36]. Furthermore, NF- κ B binding sites are present in the promoters of iNOS and COX-2 and are essential to modulate their transcription [37,38]. Results of Yao *et al.* (2014) [39] demonstrated that aspirin normalizes COX-2 and iNOS over-expression in astrocytes in part through inhibition of the NF- κ B pathway [39]. As these authors indicated, we show a reduction in the expression of iNOS, COX-2 and NF- κ B in the presence of aspirin.

Aspirin has additional targets in humans, besides the cyclooxygenases. Salicylic acid (SA), the primary metabolite of aspirin, binds human glyceraldehyde 3-phosphate dehydrogenase (GAPDH) and suppresses nuclear translocation and cell death [40]. It is known that GAPDH plays an important role in neurodegenerative diseases, including AD [41].

In neurodegenerative diseases such as AD, treatment with non-steroidal anti-inflammatory drugs (NSAIDs) has been reported [42] and COX-1 and COX-2 are the targets of those drugs. Furthermore both COX isoforms regulate PPAR γ activity through prostaglandin synthesis [43]. Also, PPAR γ has been shown to inhibit the expression of proinflammatory genes, such as iNOS [44,35] and has several inhibitory effects on inflammation, including reduction of NF- κ B transcriptional activities and promotes anti-inflammatory mediators [43]. Potent synthetic antidiabetics such as rosiglitazone are agonists for PPAR- γ [45,46] and many NSAIDs are also PPAR- γ agonists [47]. Furthermore, reduction of A β induced by indomethacin or naproxen by inhibiting BACE1 (beta-site APP-cleaving enzyme 1) activity occurs in a PPAR- γ dependent manner [48]. In our study, we detected a decrease of PPAR- γ expression after A β addition that was reversed by treatment with aspirin. It is possible that aspirin could inhibit A β synthesis through the increment of PPAR- γ .

In this study we also investigated the possible antioxidant properties of aspirin using A β ₁₋₄₂ toxic peptide as a model. We showed an increase expression of Mn-SOD and Cu/Zn-SOD proteins after aspirin addition in culture of astrocytes with and




Figure 8. Aspirin effects in astrocytes in primary culture. Asp increases cell viability, anti-inflammatory response and anti-oxidant proteins. On the other hand, Asp decreases pro-inflammatory mediators, necrosis and apoptosis.

without the toxic peptide. In agreement with our results, Dairam *et al.* (2006) [49] have demonstrated the antioxidant and neuroprotective effects of NSAIDs in an AD rat model [49]. Furthermore, aspirin-elicited lipoxin A₄ inhibited ROS synthesis induced by LPS in microglial cells [50]. Also, aspirin minimizes the effect of free radicals induced by LPS in rat dopaminergic neurons [51]. Liu *et al.* (2017) [52] indicated that aspirin at doses of 75 and 100 mg/day, similar to that used in our study, stimulates the levels of superoxide dismutase (SOD). On the other hand, production of L-NMMA in microcirculation is inhibited by aspirin, due to cyclooxygenase inhibition or SOD increase [53,54].

Also, aspirin exerts pro-oxidant effects on Mn-SOD-deficient yeast cells causing apoptosis with mitochondrial involvement [55]. Furthermore, aspirin on focal cerebral ischemia-reperfusion rats reduced MDA content [56].

Conclusion

In conclusion, our results indicate that aspirin, at low doses, prevents toxic effects induced by Aβ₁₋₄₂

peptide in astrocytes in primary culture, increasing cell viability and proliferation while decreasing apoptosis and necrosis. The key finding of our study is that aspirin at low doses prevents oxidative stress and inflammation induced by Aβ₁₋₄₂ toxic peptide. So, the administration of low doses of aspirin could be useful in AD patients (Fig. 8).

Acknowledgments

Funding sources

This work was supported in part by grant from local government of Spain: Gvcs2007-AP-001.

Ethics approval consent

All animal procedures were carried out in accordance with the European legislation on the use and care of laboratory animals (CEE 86/609). Experimental research on mice was performed with the approval of the ethics committee on animal research of the University of Valencia (Spain) and Generalitat Valenciana local government (2016/VSC/PEA/00220). All procedures were performed in

accordance with the 1964 Helsinki declaration and its later amendments.

Abbreviations

MTT: 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide; Asp: Acetylsalicylic acid; aspirin; AD: Alzheimer's disease; A β : Amyloid β ; Cu/Zn: superoxide dismutase; COX: Cyclooxygenase; Cyt c: Cytochrome c; GFAP: Glial fibrillary protein; NF- κ B: Nuclear factor κ B; PPAR- γ : Peroxisome proliferator-activated receptor.

Authors' contributions

AJ designed the study, carried out the LDH and Caspase 3 experiments, contributed to western-blot, and wrote the manuscript. MA designed the study, carried out the ELISA assays and helps to write the manuscript. CA, SGO and JCC contributed to western-blot experiments. AI contributed to statistics analysis. JV contributed to the look for references and statistics analysis. SLV conceived the study, participated in its design and coordination, and helped to draft the manuscript. All authors read and approved the final manuscript.

Competing Interests

The authors have declared that no competing interest exists.

References

- Lane CA, Hardy J, Schott JM. Alzheimer's disease. *Eur J Neurol.* 2018;25:59-70.
- Perry G, Cash AD, Smith M. Alzheimer disease and oxidative stress. *J Biomed Biotechnol.* 2002;2:120-123.
- Holtzman DM, Mandelkow E, Selkoe DJ. Alzheimer disease in 2020. *Cold Spring Harb Perspect. Med.* 2011. doi:10.1101/cshperspect.a011585
- Selkoe DJ, Hardy J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med.* 2016;8:595-608.
- Figley CR, Stroman PW. The role(s) of astrocytes and astrocyte activity in neurometabolism, neurovascular coupling, and the production of functional neuroimaging signals. *Eur J Neurosci.* 2011;33:577-588.
- Frost GR, Li YM. The role of astrocytes in amyloid production and Alzheimer's disease. *Open Biol.* 2017;7. doi: 10.1098/rsob.170228.
- Aguirre-Rueda D, Guerra-Ojeda S, Aldasoro M, Iradi A, Obrador E, Ortega A, et al. Astrocytes protect neurons from A β 1-42 peptide-induced neurotoxicity increasing TFAM and PGC-1 and decreasing PPAR- γ and SIRT-1. *Int J Med Sci.* 2015a;12:48-56.
- Aguirre-Rueda D, Guerra-Ojeda S, Aldasoro M, Iradi A, Obrador E, Mauricio MD, et al. WIN 55,212-2, agonist of cannabinoid receptors, prevents amyloid β 1-42 effects on astrocytes in primary culture. *PLoS One.* 2015b. doi: 10.1371/journal.pone.0122843.
- Nedergaard M, Ransom B, Goldman SA. New roles for astrocytes: redefining the functional architecture of the brain. *Trends Neurosci.* 2003;26:523-530.
- Malarkey EB, Pappas V. Mechanisms of glutamate release from astrocytes. *Neurochem Int.* 2018;52:142-154.
- Verkhatsky A, Olabarria M, Noristani HN, Yeh CY, Rodriguez JJ. Astrocytes in Alzheimer's disease. *Neurotherapeutics.* 2010;7:399-412.
- Giunta B, Fernandez F, Nikolic WC, Obrego E, Rrapo E, Town T, et al. Inflammation as a prodrome to Alzheimer's disease. *J Neuroinflammation.* 2008;5:51-65.
- Hou L, Liu Y, Wang X, Ma H, He J, Zhang Y, et al. The effects of amyloid- β 42 oligomer on the proliferation and activation of astrocytes in vitro. *Vitr Cell Dev Biol Anim.* 2011;47:573-580.
- Farina C, Aloisi F, Meinl E. Astrocytes are active players in cerebral innate immunity. *Trends Immunol.* 2017;28:138-145.
- Green GA. Understanding NSAIDs: from aspirin to COX-2. *Clin Cornerstone.* 2001;3:50-60.
- Vane JR, Botting R. The mechanism of action of aspirin. *Thromb Res.* 2003;110:255-258.

- Ertmer C, Rehberg S, Westphal M. Cyclooxygenase-2 inhibition and increased arterial vasoconstriction to vasopressin: what is the link? *Crit Care Med.* 2008;36:353-354.
- Caughey GE, Cleland LG, Penglis PS, Gamble JR, James MJ. Roles of cyclooxygenase (COX)-1 and COX-2 in prostanoid production by human endothelial cells: selective up-regulation of prostacyclin synthesis by COX-2. *J Immunol.* 2001;167:2831-2838.
- Aldasoro M, Mauricio MD, Serna E, Cortina B, Segarra G, Medina P, et al. Effects of aspirin, nimesulide, and SC-560 on vasopressin-induced contraction of human gastroepiploic artery and saphenous vein. *Crit Care Med.* 2008;36:193-197.
- McAdam BF, Catella-Lawson F, Mardini IA, Kapoor S, Lawson JA, FitzGerald GA. Systemic biosynthesis of prostacyclin by cyclooxygenase (COX)-2: the human pharmacology of a selective inhibitor of COX-2. *Proc Natl Acad Sci USA.* 1999;96:272-277. Erratum in: *Proc Natl Acad Sci USA.* 1999;96:5890.
- Nilsson SE, Johansson B, Takkinen S, Berg S, Zarit S, McClearn G. Does aspirin protect against Alzheimer's dementia? A study in a Swedish population-based sample aged or 80 years. *Eur J Clin Pharmacol.* 2003;59:313-319.
- Kern S, Skoog I, Ostling S, Kern J, Björjesson-Hanson A. Does low-dose acetylsalicylic acid prevent cognitive decline in women with high cardiovascular risk? A 5-year follow-up of a non-demented population-based cohort of Swedish elderly women. *BMJ Open.* 2012. doi:10.1136/bmjopen-2012-001288.
- Heneka MT, Carson MJ, El Khoury J, Landreth GE, Brosseron F, Feinstein DL, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol.* 2015;14:388-405.
- Nirzhor SSR, Khan RI, Neelotpol S. The Biology of Glial Cells and Their Complex Roles in Alzheimer's Disease: New Opportunities in Therapy. *Biomolecules.* 2018;10. doi: 10.3390/biom8030093.
- Jorda A, Cauli O, Santonja JM, Aldasoro M, Aldasoro C, Obrador E, et al. Changes in Chemokines and Chemokine Receptors Expression in a Mouse Model of Alzheimer's Disease. *Int J Biol Sci.* 2019;15:453-463.
- Yao Y, Huang JZ, Chen Y, Hu HJ, Tang X, Li X. Effects and mechanism of amyloid β 1-42 on mitochondria in astrocytes. *Mol Med Rep.* 2018;17:6997-7004.
- Pomponi MF, Gambassi G, Pomponi M, Masullo C. Alzheimer's Diseases fatty acids we eat may be linked to a specific protection via low-dose aspirin. *Aging Dis.* 2010;1:37-59.
- Ayyadevara S, Balasubramaniam M, Kakraba S, Alla R, Mehta JL, Shmookler-Reis RJ. Aspirin-Mediated Acetylation Protects Against Multiple Neurodegenerative Pathologies by Impeding Protein Aggregation. *Antioxid Redox Signal.* 2017;27:1383-1396.
- Parmar HS, Houdek Z, Pesta M, Vaclava C, Dvorak P, Hatina J. Protective Effect of Aspirin against Oligomeric A β 42 Induced Mitochondrial Alterations and Neurotoxicity in Differentiated EC P19 Neuronal Cells. *Curr Alzheimer Res.* 2017;14:810-819.
- Bae MK, Kim SR, Lee HJ, Wee HJ, Yoo MA, Ock Oh S, et al. Aspirin-induced blockade of NF- κ B activity restrains up-regulation of glial fibrillary acidic protein in human astroglial cells. *Biochim Biophys Acta.* 2006;1763:282-289.
- Burke A, Smyth E, Fitzgerald G, Goodman and Gilman's the Pharmacological Basis of Therapeutics, 11th ed. McGraw-Hill, New York. 2006.
- He J, Whelton PK, Vu B, Klag MJ. Aspirin and risk of hemorrhagic stroke: a meta-analysis of randomized controlled trials. *JAMA.* 1998;280:1930-1935.
- Redlak MJ, Power JJ, Miller TA. Role of mitochondria in aspirin-induced apoptosis in human gastric epithelial cells. *Am J Physiol Gastrointest Liver Physiol.* 2005;289:G731-8.
- Font-Nieves M, Sans-Fons MG, Gorina R, Bonfill-Teixidor E, Salas-Pédomo A, Márquez-Kisinousky L, et al. Induction of COX-2 enzyme and down-regulation of COX-1 expression by lipopolysaccharide (LPS) control prostaglandin E2 production in astrocytes. *J Biol Chem.* 2012;287:6454-6468.
- Newcombe EA, Camats-Perna J, Silva ML, Valmas N, Huat TJ, Medeiros R. Inflammation: the link between comorbidities, genetics, and Alzheimer's disease. *J Neuroinflammation.* 2018;24:276.
- Lee JE, Park JS, Lee YY, Kim DY, Kang JL, Kim HS. Anti-inflammatory and anti-oxidant mechanisms of an MMP-8 inhibitor in lipoteichoic acid-stimulated rat primary astrocytes: involvement of NF- κ B, Nrf2, and PPAR- γ signaling pathways. *J Neuroinflammation.* 2018;23:326.
- Chen CH, Sheu MT, Chen TF, Wang YC, Hou WC, Liu DZ, et al. Suppression of endotoxin-induced proinflammatory responses by citrus pectin through blocking LPS signaling pathways. *Biochem Pharmacol.* 2006;72:1001-1009.
- Wu KK. Control of cyclooxygenase-2 transcriptional activation by pro-inflammatory mediators, Prostaglandins Leukot Essent Fatty Acids. 2005;72:89-93.
- Yao C, Yang D, Wan Z, Wang Z, Liu R, Wu Y, et al. Aspirin-triggered lipoxin A4 attenuates lipopolysaccharide induced inflammatory response in primary astrocytes. *Int Immunopharmacol.* 2014;18:85-89.
- Choi HW, Tian M, Manohar M, Harraz MM, Park SW, Schroeder FC, et al. Human GAPDH Is a Target of Aspirin's Primary Metabolite Salicylic Acid and Its Derivatives. *PLoS One.* 2015. doi:10.1371/journal.pone.0143447.
- El Kadmiri N, El Khachibi M, Slassi I, El Moutawakil B, Nadifi S, Soukri A. Assessment of GAPDH expression by quantitative real time PCR in blood of Moroccan AD cases. *J Clin Neurosci.* 2017;40:24-26.

42. Lleo A, Galea E, Sastre M. Molecular targets of non-steroidal anti-inflammatory drugs in neurodegenerative diseases. *Cell Mol Life Sci.* 2007;64:1403-1408.
43. Wahli W, Michalik L. PPARs at the crossroads of lipid signaling and inflammation. *Trends Endocrinol Metab.* 2012;23:351-363.
44. Iglesias J, Morales L, Barreto GE. Metabolic and Inflammatory Adaptation of Reactive Astrocytes: Role of PPARs. *Mol Neurobiol.* 2017;54:2518-2538.
45. Li AC, Glass CK. PPAR- and LXR-dependent pathways controlling lipid metabolism and the development of atherosclerosis. *J Lipid Res.* 2004;45:2161-2173.
46. Sastre M, Klockgether T, Heneka MT. Contribution of inflammatory processes to Alzheimer's disease: molecular mechanisms. *Int J Dev Neurosci.* 2006;24:167-176.
47. Lehmann JM, Lenhard JM, Oliver BB, Ringold GM, Kliewer SA. Peroxisome proliferator-activated receptors alpha and gamma are activated by indomethacin and other non-steroidal anti-inflammatory drugs. *J Biol Chem.* 1997;272:3406-3410.
48. Sastre M, Dewachter I, Landreth GE, Willson TM, Klockgether T, van Leuven F, et al. Nonsteroidal anti-inflammatory drugs and peroxisome proliferator-activated receptor-gamma agonists modulate immunostimulated processing of amyloid precursor protein through regulation of beta-secretase. *J Neurosci.* 2003;23:9796-97804.
49. Dairam A, Chetty P, Daya S. Non-steroidal anti-inflammatory agents, tolmetin and sulindac, attenuate oxidative stress in rat brain homogenate and reduce quinolinic acid-induced neurodegeneration in rat hippocampal neurons. *Metab Brain Dis.* 2006;21:221-233.
50. Wu Y, Zhai H, Wang Y, Li L, Wu J, Wang F, et al. Aspirin-triggered lipoxin A₄ attenuates lipopolysaccharide-induced intracellular ROS in BV2 microglia cells by inhibiting the function of NADPH oxidase. *Neurochem Res.* 2012;37:1690-1696.
51. Wang F, Zhai Z, Huang L, Li H, Xu Y, Qiao X, et al. Aspirin protects dopaminergic neurons against lipopolysaccharide-induced neurotoxicity in primary midbrain cultures. *J Mol Neurosci.* 2012;46:153-161.
52. Liu F, Yang H, Li G, Zou K, Chen Y. Effect of a small dose of aspirin on quantitative test of 24-h urinary protein in patients with hypertension in pregnancy. *Exp Ther Med.* 2017;13:37-40.
53. Rosenblum WI, Nishimura H, Nelson GH. L-NMMA in brain microcirculation of mice is inhibited by blockade of cyclooxygenase and by superoxide dismutase. *Am J Physiol.* 1992;262:H1343-1349.
54. Grosser N, Schröder H. Aspirin protects endothelial cells from oxidant damage via the nitric oxide-cGMP pathway. *Arterioscler Thromb Vasc Biol.* 2003;23:1345-1351.
55. Farrugia G, Bannister WH, Vassallo N, Balzan R. Aspirin-induced apoptosis of yeast cells is associated with mitochondrial superoxide radical accumulation and NAD(P)H oxidation. *FEMS Yeast Res.* 2013;13:755-768.
56. Qiu LY, Yu J, Zhou Y, Chen CH. Protective effects and mechanism of action of aspirin on focal cerebral ischemia-reperfusion in rats. *Yao Xue Xue Bao.* 2003;38:561-564.