



The Open Medicinal Chemistry Journal

Content list available at: <https://openmedicinalchemistryjournal.com>



RESEARCH ARTICLE

Dose-Dependent Tissue Distribution of K117, a Bis-pyridinium Aldoxime, in Rats

Huba Kalász^{1,*}, Gellért Karvaly², Kamil Musilek³, Kamil Kuca³, Jung Young-Sik⁴, Barbara Malawska⁵, Ernest A. Adeghate⁶, Syed M. Nurulain⁷, Judit Szepesy¹, Tibor Zelles¹ and Kornélia Tekes⁸

¹Department of Pharmacology and Pharmacotherapy, Semmelweis University, 1089 Budapest, Orczy út 2-4, Hungary

²Department of Laboratory Medicine, Semmelweis University, 1089 Budapest, Orczy út 2-4, Hungary

³Department of Chemistry, University of Hradec Kralove, 500 03 Hradec Kralove, Rokitanskeho 62, Czech Republic

⁴Medicinal Science Division, Korea Research Institute of Chemical Technology, Daejeon, Republic of Korea

⁵Department of Physicochemical Drug Analysis, Jagiellonian University, 30688 Krakow, ul. Medyczna 9, Poland

⁶Department of Anatomy, College of Medicine, United Arab Emirates University, 17666, Al Ain, UAE

⁷Department of Biosciences, COMSATS Institute of Information Technology, Islamabad, Pakistan

⁸Department of Pharmacodynamics, Semmelweis University, 1089 Budapest, Orczy út 2-4, Hungary

Abstract:

Background:

Bis-pyridinium aldoximes are reactivators of the paraoxon-inhibited butyrylcholinesterase enzyme. Paraoxon is the active product of parathion, a widely used insecticide.

Objective:

The objective of this study is to examine the dose-dependent distribution of K117, a bis-pyridinium aldoxime in rat tissues.

Materials and Methods:

White male Wistar rats were intramuscularly injected with various doses of K117; the animals were sacrificed 30 minutes after injections. The dose-dependent body distribution of K117 was determined using reversed-phase HPLC.

Results:

Dose-dependent distribution of K117 in body tissues was linear in the serum and other body tissues throughout the whole range of the concentrations studied. However, the of distribution was not observed in the brain and cerebrospinal fluid, especially with high doses.

Conclusion:

The body distribution of K117 significantly depends on doses used, the p-value is: 500 nmol, i.m., when applied in the range of 100 to 10,000 nmol.

Keywords: K117, Dose-dependence, Pharmacokinetics, Wistar rats, Intramuscular injections, Linear Pattern.

Article History

Received: November 02, 2018

Revised: January 27, 2019

Accepted: February 06, 2019


1. INTRODUCTION

There are two types of cholinesterase enzymes in humans and animals. The acetylcholinesterase enzyme (AcChE, EC 3.1.1.7) is located in many tissues (red blood cell membranes,


the central nervous system, peripheral organs, cholinergic and non-cholinergic fibers *etc.*) and functions postsynaptically with an extremely high catalytic activity at nerve synapses to terminate the effect of acetylcholine by its hydrolyzation.

The butyrylcholinesterase enzyme (BuChE, EC 3.1.1.8), also known as pseudocholinesterase or plasma esterase, serves as a backup for AcChE [1]. It is responsible for the quick

* Address correspondence to this author at the Department of Pharmacology and Pharmacotherapy, Semmelweis University, 1089 Budapest, Orczy út 2-4, Hungary; Tel: +36302747986; Fax: +3612104412; E-mail: drkalasz@gmail.com



K117



K127

Fig. (1). Chemical structure of K117 and that of K127.

inactivation through the hydrolysis of different endogenous and exogenous esters in the blood plasma. The potential poisons of AcChE are scavenged by BuChE, and the human body has BuChE (approximately 680 nanomoles) about tenfold as much as AcChE [2]. BuChE also plays an essential role in metabolizing cocaine, heroin, mivacurium, succinylcholine and procaine. BuChE-deficiency results in increased sensitivity to succinylcholine (a widely used depolarizing neuromuscular blocking agent in clinical practice). It is generally accepted that BuChE has an essential role in the inactivation of toxic compounds including organophosphorous esters. Measuring BuChE serves as a biomarker for organophosphate exposure [3]. Either AcChE- or BuChE-deficiency is an indicator of a special depression that is common in pesticide handlers [4].

Reactivation of AcChE- or BuChE-deficiency caused by insecticide poisoning is crucial to the survival of a poisoned patient. Karasova *et al.* [5] reported that both K117 and K127 were among the best reactivators of BuChE, when some currently available and several newly synthesized pyridinium aldoximes were studied on rats intoxicated by tabun. The reactivation efficiency of K117, and also that of HI-6, obidoxime, triedoxime, K127, K206, K250, K251, K269, K347, K628, were experimentally determined in plasma and the brain by Kovarik *et al.* [6]. K117 showed similar effectivity as HI-6 in peripheral tissues. Jun *et al.* [7] compared the reactivation potency of several pyridinium aldoximes *in vitro* on paraoxon-inhibited human AcChE and BuChE. Two of the reactivators (obidoxime and trimedoxime) worked well on inhibited AcChE, giving a reactivation rate over 75% when used in a concentration of 100 μ M concentration. However, none of the classical pyridinium aldoximes (pralidoxime, methoxime, obidoxime, trimedoxime, HI-6) produced the reactivation rate of BuChE over 10%, even when they were used in a concentration of 100 μ M. Kuca *et al.* [5, 7, 8] found

that K117 and atropine co-doses work efficiently *in vitro* on rats intoxicated by tabun.

Sakurada *et al.* [9, 10] were the first to detect and measure that pralidoxime penetrated through the blood-brain barrier. Similar determinations and statements were made by Kalász *et al.* [11, 12], who expanded the analysis of pyridinium aldoximes on the cerebrospinal fluid.

This paper presents a dose-dependent tissue penetration of the K117 from the site of its intramuscular application to various target organs.

2. MATERIALS AND METHODS

2.1. Chemicals and Solvents

All solvents and chemicals were bought from commercial sources in the best possible quality. Pyridinium aldoximes (K117 and K127) were supplied by the Department of Chemistry, University of Hradec Kralove, Czech Republic. The chemical structures of these two compounds are given in Fig. (1).

2.2. Animals and Animal Treatment

White male Wistar rats weighing 180-199 grams were obtained from Toxicoop (Budapest, Hungary). Two animals were treated intramuscularly (i.m.) with an adequate dose of a freshly prepared aqueous solution of K117 (0.1, 0.3, 1.0, 3.0 and 10.0 μ mol for each pair of rats). The rats were sacrificed 30 minutes after treatment, keeping the ethical regulation of Semmelweis University. Body fluids (serum, cerebrospinal fluid) were taken and certain organs/tissues (brain, eyes, lungs, testes, liver, kidneys and inner ear) were dissected and treated with perchloric acid, homogenized and centrifuged. HPLC determinations of K117 were done using K127 as an internal standard (Figs. 2 and 3), as detailed in our previous publication [13].




Fig. (2). Calibration curve of K117 determination. $R^2 > 0.99$.




Fig. (3). Representative chromatograms of K117 and the internal standard K127. (A) rat serum (340 ng/mL K117), (B) rat kidney homogenate (3430 ng/mL K117), (C) calibration sample (500 ng/mL K117), (D) blank serum.

Table 1. Dose-dependence of tissue and body fluid concentrations of K 117 injected intramuscularly to rats.

Sample (tissue or body fluid)	Concentration ($\mu\text{g/ml} \pm \text{SD}$) Determined by HPLC-UV	Tissue Concentration (ng/ml)	Tissue concentration related to the serum concentration (%)	Concentration, related to the dose of 1 μmol
Dose: 0.1 μmol				
Serum	0.011 \pm 0.005	110	100.0	0.567
Brain	0.011 \pm 0.005	55	50.00	3.235
CSF	0.010 \pm 0.000	50	45.45	5.880
Eyes	0.010 \pm 0.002	50	45.45	0.990
Lung	0.055 \pm 0.004	275	250.0	2.350
Testis	0.006 \pm 0.004	30	27.27	5.450
Liver	0.043 \pm 0.005	215	195.5	39.09
Kidney	0.340 \pm 0.012	1700	1545	42.50
Inner ear	0.079 \pm 0.005	395	359.0	3.361
Dose: 0.3 μmol				
Serum	0.022 \pm 0.008	220	100.0	0.378
Brain	0.015 \pm 0.004	75	34.09	1.470
CSF	0.012 \pm 0.003	60	27.27	2.352
Eyes	0.014 \pm 0.005	70	31.82	0.460
Lung	0.051 \pm 0.002	255	115.9	0.720
Testis	0.009 \pm 0.004	45	20.45	2.727
Liver	0.077 \pm 0.027	385	175.0	3.208
Kidney,	0.768 \pm 0.009	3840	1745	0.920
Inner ear	0.088 \pm 0.010	440	200.0	0.838
Dose: 1.0 μmol				
Serum	0.194 \pm 0.003	1940	100.0	1.00
Brain	0.034 \pm 0.003	170	8.762	1.00
CSF	0.017 \pm 0.011	85	4.381	1.00
Eyes	0.101 \pm 0.013	505	26.03	1.00
Lung	0.234 \pm 0.006	1170	60.30	1.00
Testis	0.011 \pm 0.004	55	2.835	1.00
Liver	0.080 \pm 0.010	400	20.61	1.00
Kidney,	2.781 \pm 0.026	13905	716.7	1.00
Inner ear	0.235 \pm 0.006	1175	60.56	1.00
Dose: 3.0 μmol				
Serum	0.576 \pm 0.013	5760	100.0	0.989
Brain	0.083 \pm 0.006	415	7.204	0.813
CSF	0.023 \pm 0.016	115	1.996	0.451
Eyes	0.222 \pm 0.005	1111	19.28	0.733
Lung	0.601 \pm 0.013	3005	52.17	0.856
Testis	0.093 \pm 0.009	465	8.072	2.818
Liver	0.194 \pm 0.020	970	16.84	0.808
Kidney	4.698 \pm 0.159	23490	407.8	0.563
Inner ear	0.442 \pm 0.064	2210	38.36	0.626
Dose: 10.0 μmol				
Serum	2.289 \pm 0.121	22890	100.0	1.179
Brain	0.360 \pm 0.004	1800	7.863	1.058
CSF	0.075 \pm 0.004	375	1.638	0.441
Eyes	0.763 \pm 0.035	3815	16.66	0.755
Lung	1.788 \pm 0.033	8940	39.05	1.770
Testis	0.484 \pm 0.003	2420	10.57	4.400
Liver	0.153 \pm 0.002	765	3.342	0.190
Kidney	11.219 \pm 0.099	56095	245.06	0.403
Inner ear	1.246 \pm 0.047	6230	27.21	0.530

3. RESULTS

Dose-dependence of K117 levels in various body compartments of rats is given in Table 1.

Table 1 shows very high levels of K117 in the lungs, liver, kidney and Inner ear compared to those in serum. These relatively high levels continuously decrease with time in the lungs and inner ear, while they continue to increase in the liver and kidney.

4. DISCUSSION

During their *in vitro* experiments, Jun *et al.* [7] compared AcChE and BuChE reactivating process of several K-compounds against paraoxon-inhibition, using a concentration as high as 100 μM , as the overall (*in vivo*) toxicity does not limit the dose used. However, studying the dose-dependence in *in vivo* experiments injections up to 100 μM could be applied, as higher doses of K117 were toxic [6]. Horn *et al.* [14] experimentally proved that K117 fulfils one of the basic requirements of an adequate antidote, it does not even influence the enzyme activity of BuChE in excess (1,000 μM). Thereby, K117 belongs to the group of pyridinium aldoximes that can be potentially used (K27, K48, K74, K75, K99, K127, K203, *etc.*) in medical practice. The BuChE reactivation power of K117 is preferable in paraoxon-inhibited BuChE, while its (*in vitro*) activity on tabun-inhibited enzyme is not significant.

It is the circulating blood that supplies K117 from the site of i.m. injection to each organ, tissue and cell of rats. However, special barriers of the organism (*e.g.* blood-brain barrier, blood-testis barrier *etc.*) can either totally or partially hinder the transfer of K117 to special organs such as the central nervous system and the organs of reproduction. Each biological barrier has its own characteristics. When brain concentrations are compared to the serum levels of K117 no proportionality can be observed. About 50% of K117 could penetrate into the brain when a dose of 0.1 μmol was given. However, this ratio decreased to 9% when 1 μmol and to 7% when 10 μmol K117 were applied, respectively. This dynamic function of the blood-brain-barrier was even more expressed for CSF; at a dose of 0.1 μmol . The relative concentration of K117 in the CSF compared to that in the serum, was 45%, however, this ratio decreased to 5% and 1.6% using doses of 1 μmol and 10 μmol , respectively. Lorke *et al.* [15] also demonstrated dynamic changes in blood-brain barrier and blood-CSF barrier functions. The relatively high proportion of K-117 in the kidneys, 30 minutes following intramuscular administration, compared to serum and the liver *versus* serum, indicates the essential role of the kidneys in the removal of K117 from the organism. It also indicates that K117 is hydrophilic and, therefore, is excreted from the body *via* the kidney.

The inner ear has a multi-compartmental structure (peri-, endolymph) with several different barrier systems (*e.g.* blood-endolymph, blood-perilymph, CSF-perilymph) [16, 17], which makes the explanation of the relatively high K117 concentrations in the inner ear difficult. Experimental data suggest that a much slower elimination from the perilymph [18, 19] is presumably a major factor in the development of substance accumulation in the inner ear.

CONCLUSION

The analysis of tissue concentrations of K117 levels relative to serum concentrations is a useful method to determine the special characteristics of the penetration of this compound into critically important organs and tissues. The validated reversed-phase HPLC bioanalytical method developed was sensitive and selective enough for detailed pharmacokinetic measurements.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

All animal experiments were carried out according to the ethical requirements of Semmelweis University (permission number of Governmental Office of Pest County, Hungary, PE/EA/385-5/2018.)

HUMAN AND ANIMAL RIGHTS

No human were used in the study. All the procedures followed were in accordance with the standards set forth in the eighth edition of "Guide for the Care and Use of Laboratory Animals" (grants.nih.gov/grants/olaw/guide-for-the-care-and-use-of-laboratory-animals_prepub.pdf published by the National Academy of Sciences, The National Academies Press, Washington, D.C.).

CONSENT FOR PUBLICATION

Not applicable.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

This project is based on a Corsorcium Agreement between Korea Research Institute (Daejeon, Korea), Jagiellonian University (Krakow, Poland), University of Hradec Kralove (Hradec Kralove, Czech Republic) and Semmelweis University (Budapest, Hungary).

The authors express their thanks to Mr. János Horváth and Ms. Bogi Szalacsi for their advice. Technical assistance was provided by Ms. Györgyike Guth, Ms. Krisztina Kecskés, Mr. Zoltán Szimrök and Mr. András Keglevich.

This project was financially supported by the grant NN126968 of National Research, Development and Innovation Office (NRDIO, Budapest, Hungary) this research was supported by Ministry of Education, Youth and Sports of the Czech Republic (No. 8F17004) via V4-Korea Joint Research Program and also by Kalász Teaching and Research Co. (Budapest, Hungary). Grant No. K128875 of NRDIO served as partial source of salary to JSz.

REFERENCES

- [1] Nachon, F.; Brazzolotto, X.; Trovaslet, M.; Masson, P. Progress in the development of enzyme-based nerve agent bioscavengers. *Chem. Biol. Interact.*, **2013**, *206*(3), 536-544. [<http://dx.doi.org/10.1016/j.cbi.2013.06.012>] [PMID: 23811386]
- [2] Masson, P.; Lockridge, O. Butyrylcholinesterase for protection from organophosphorus poisons: catalytic complexities and hysteretic

- behavior. *Arch. Biochem. Biophys.*, **2010**, *494*(2), 107-120. [http://dx.doi.org/10.1016/j.abb.2009.12.005] [PMID: 20004171]
- [3] Lockridge, O. Review of human butyrylcholinesterase structure, function, genetic variants, history of use in the clinic, and potential therapeutic uses. *Pharmacol. Ther.*, **2015**, *148*, 34-46. [http://dx.doi.org/10.1016/j.pharmthera.2014.11.011] [PMID: 25448037]
- [4] Strelitz, J.; Engel, L.S.; Keifer, M.C. Blood acetylcholinesterase and butyrylcholinesterase as biomarkers of cholinesterase depression among pesticide handlers. *Occup. Environ. Med.*, **2014**, *71*(12), 842-847. [http://dx.doi.org/10.1136/oemed-2014-102315] [PMID: 25189163]
- [5] Karasova, J.Z.; Kassa, J.; Jung, Y.S.; Musilek, K.; Pohanka, M.; Kuca, K. Effect of several new and currently available oxime cholinesterase reactivators on tabun-intoxicated rats. *Int. J. Mol. Sci.*, **2008**, *9*(11), 2243-2252. [http://dx.doi.org/10.3390/ijms9112243] [PMID: 19330072]
- [6] Kovarik, Z.; Katalinić, M.; Sinko, G.; Binder, J.; Holas, O.; Jung, Y.S.; Musilova, L.; Jun, D.; Kuca, K. Pseudo-catalytic scavenging: searching for a suitable reactivator of phosphorylated butyrylcholinesterase. *Chem. Biol. Interact.*, **2010**, *187*(1-3), 167-171. [http://dx.doi.org/10.1016/j.cbi.2010.02.023] [PMID: 20206154]
- [7] Jun, D.; Musilova, L.; Kuca, K.; Kassa, J.; Bajgar, J. Potency of several oximes to reactivate human acetylcholinesterase and butyrylcholinesterase inhibited by paraoxon *in vitro*. *Chem. Biol. Interact.*, **2008**, *175*(1-3), 421-424. [http://dx.doi.org/10.1016/j.cbi.2008.05.004] [PMID: 18617161]
- [8] Kuca, K.; Jun, D.; Junova, L.; Musilek, K.; Hrabnova, M.; da Silva, J.A.V.; Ramalho, T.C.; Valko, M.; Wu, Q.; Nepovimova, E.; França, T.C.C. Synthesis, Biological evaluation, and docking studies of novel bisquaternary aldoxime reactivators on acetylcholinesterase and butyrylcholinesterase inhibited by paraoxon. *Molecules*, **2018**, *23*(5), E1103. [http://dx.doi.org/10.3390/molecules23051103] [PMID: 29735900]
- [9] Sakurada, K.; Matsubara, K.; Shimizu, K.; Shiono, H.; Seto, Y.; Tsuge, K.; Yoshino, M.; Sakai, I.; Mukoyama, H.; Takatori, T. Pralidoxime iodide (2-pAM) penetrates across the blood-brain barrier. *Neurochem. Res.*, **2003**, *28*(9), 1401-1407. [http://dx.doi.org/10.1023/A:1024960819430] [PMID: 12938863]
- [10] Okuno, S.; Sakurada, K.; Ohta, H.; Ikegaya, H.; Kazui, Y.; Akutsu, T.; Takatori, T.; Iwadata, K. Blood-brain barrier penetration of novel pyridinealdoxime methiodide (PAM)-type oximes examined by brain microdialysis with LC-MS/MS. *Toxicol. Appl. Pharmacol.*, **2008**, *227*(1), 8-15. [http://dx.doi.org/10.1016/j.taap.2007.09.021] [PMID: 17964625]
- [11] Kalász, H.; Szöko, E.; Tábi, T.; Petroianu, G.A.; Lorke, D.E.; Omar, A.; Alaffi, S.; Jasem, A.; Tekes, K. Analysis of pralidoxime in serum, brain and CSF of rats. *Med. Chem.*, **2009**, *5*(3), 237-241. [http://dx.doi.org/10.2174/157340609788185882] [PMID: 19442213]
- [12] Kalász, H.; Szegi, P.; Jánoki, G.; Balogh, L.; Pöstényi, Z.; Musilek, K.; Petroianu, G.A.; Siddiq, A.; Tekes, K. Study on medicinal chemistry of K203 in wistar rats and beagle dogs. *Curr. Med. Chem.*, **2013**, *20*(16), 2137-2144. [http://dx.doi.org/10.2174/0929867311320160006] [PMID: 23531217]
- [13] Tekes, K. Distribution of K117, a bispyridinium aldoxime with regenerating activity on diminished butyrylcholinesterase enzyme activity *in preparation*;
- [14] Horn, G.; Wille, T.; Musilek, K.; Kuca, K.; Thiermann, H.; Worek, F. Reactivation kinetics of 31 structurally different bispyridinium oximes with organophosphate-inhibited human butyrylcholinesterase. *Arch. Toxicol.*, **2015**, *89*(3), 405-414. [http://dx.doi.org/10.1007/s00204-014-1288-5] [PMID: 24912784]
- [15] Lorke, D.E.; Kalász, H.; Petroianu, G.A.; Tekes, K. Entry of oximes into the brain: A review. *Curr. Med. Chem.*, **2008**, *15*(8), 743-753. [http://dx.doi.org/10.2174/092986708783955563] [PMID: 18393843]
- [16] Juhn, S.K. Barrier systems in the inner ear. *Acta Otolaryngol. Suppl.*, **1988**, *458*, 79-83. [http://dx.doi.org/10.3109/00016488809125107] [PMID: 3245438]
- [17] Sun, W.; Wang, W. Advances in research on labyrinth membranous barriers. *J. Otol.*, **2015**, *10*(3), 99-104. [http://dx.doi.org/10.1016/j.joto.2015.11.003] [PMID: 29937790]
- [18] Tran Ba Huy, P.; Bernard, P.; Schacht, J. Kinetics of gentamicin uptake and release in the rat. Comparison of inner ear tissues and fluids with other organs. *J. Clin. Invest.*, **1986**, *77*(5), 1492-1500. [http://dx.doi.org/10.1172/JCI112463] [PMID: 3700652]
- [19] Chen, Z.; Duan, M.; Lee, H.; Ruan, R.; Ulfendahl, M. Pharmacokinetics of caroverine in the inner ear and its effects on cochlear function after systemic and local administrations in Guinea pigs. *Audiol. Neurotol.*, **2003**, *8*(1), 49-56. [http://dx.doi.org/10.1159/000067893] [PMID: 12566692]