# Coupling of Arctic ozone and stratospheric dynamics and its influence on surface climate:

## The role of CFC concentrations

Abstract ID: EGU2020-9118


Marina Friedel<sup>1</sup>, Gabriel Chiodo<sup>1,2</sup>, Stefan Muthers<sup>3</sup>, Julien Anet<sup>4</sup>, Andrea Stenke<sup>1</sup>, Thomas Peter<sup>1</sup>

<sup>1</sup> ETH Zürich, Institute for Atmospheric and Climate Science, Zürich, Switzerland

- <sup>2</sup> Applied Physics and Applied Math, Columbia University, New York (NY), USA
- <sup>3</sup>German Meteorological Service, Freiburg, Germany
- <sup>4</sup>ZHAW School of Engineering, Winterthur, Switzerland

## Coupling of Arctic ozone and stratospheric dynamics

**ETH**zürich





## Motivation & Method

#### Why?

Understanding ozone - circulation - climate coupling could result in better seasonal predictions and long-term climate projections!


#### How?

- 2 Chemistry Climate Models (SOCOL & WACCM)
- Contrasting runs with (CHEM) and without (NOCHEM) interactive chemistry (interactive vs. prescribed O<sub>3</sub>)
- Contrasting preindustrial vs. year-2000 time slice simulations (200 years each)
- Contrasting Northern Annular Mode (NAM) indices for SSW composites



#### **ETH** zürich

## Preliminary Results - WACCM



WACCM **preindustrial** runs:

- $\rightarrow$  Stronger downward propagation in **NOCHEM**
- → Longer lasting signal in lower stratosphere in CHEM

WACCM **year-2000**:

-3.8

-2.8

-1.9

0.9

0.0

-0.9

-1.9

-2.8

-3.8

NAM index

- $\rightarrow$  Stronger downward propagation in CHEM
- → Longer lasting signal in lower stratosphere in CHEM





Ozone feedbacks seem to enhance the downward propagation of Sudden Stratospheric Warming signals in recent times (CHEM vs. NOCHEM year 2000).

Ozone feedbacks seem to extend and intensify the sudden stratospheric warming signature in the lower stratosphere (CHEM vs. NOCHEM).

CFC and/or GHG concentrations might influence the sign of the ozonecirculation feedback (preindustrial vs. year-2000). Internal variability might also play a role.



### References



Polvani, L.M. and D.W. Waugh, 2004: Upward Wave Activity Flux as a Precursor to Extreme Stratospheric Events and Subsequent Anomalous Surface Weather Regimes, *J. Climate*, **17**, 3548–3554, https://doi.org/10.1175/1520-0442(2004)017<3548:UWAFAA>2.0.CO;2

Muthers, S., Anet, J. G., Stenke, A., Raible, C. C., Rozanov, E., Brönnimann, S., Peter, T., Arfeuille, F. X., Shapiro, A. I., Beer, J., Steinhilber, F., Brugnara, Y., and Schmutz, W.: The coupled atmosphere–chemistry–ocean model SOCOL-MPIOM, *Geosci. Model Dev.*, **7**, 2157–2179, https://doi.org/10.5194/gmd-7-2157-2014, 2014.

Butler, A. H., Sjoberg, J. P., Seidel, D. J., and Rosenlof, K. H.: A sudden stratospheric warming compendium, *Earth Syst. Sci. Data*, **9**, 63–76, https://doi.org/10.5194/essd-9-63-2017, 2017.

