



INDEX OF TEXAS ARCHAEOLOGY

Open Access Gray Literature from the Lone Star State

Volume 2010

Article 31

2010

Recalibrated Chronological Framework for Texas Archaeology- Geoarchaeology

Ken L. Lawrence
SWCA, klawrence@swca.com

Follow this and additional works at: <https://scholarworks.sfasu.edu/ita>

 Part of the [American Material Culture Commons](#), [Archaeological Anthropology Commons](#), [Environmental Studies Commons](#), [Other American Studies Commons](#), [Other Arts and Humanities Commons](#), [Other History of Art, Architecture, and Archaeology Commons](#), and the [United States History Commons](#)

Tell us how this article helped you.

Cite this Record

Lawrence, Ken L. (2010) "Recalibrated Chronological Framework for Texas Archaeology-Geoarchaeology," *Index of Texas Archaeology: Open Access Gray Literature from the Lone Star State*: Vol. 2010, Article 31.

<https://doi.org/10.21112/ita.2010.1.31>

ISSN: 2475-9333

Available at: <https://scholarworks.sfasu.edu/ita/vol2010/iss1/31>

This Article is brought to you for free and open access by the Center for Regional Heritage Research at SFA ScholarWorks. It has been accepted for inclusion in Index of Texas Archaeology: Open Access Gray Literature from the Lone Star State by an authorized editor of SFA ScholarWorks. For more information, please contact cdsscholarworks@sfasu.edu.

Recalibrated Chronological Framework for Texas Archaeology-Geoarchaeology

Licensing Statement

FAIR USE AND AUTHOR'S PERMISSION STATEMENT

Fair Use

This work is protected by the Copyright Laws of the United States (Public Law 94-553, section 107). Consistent with fair use as defined in the Copyright Laws, brief quotations from this material are allowed with proper acknowledgment. Use of this material for financial gain without the author's express written permission is not allowed.

Duplication Permission

As the copyright holder of this work I, Ken Lawrence, refuse permission to copy in excess of the "Fair Use" exemption without written permission.

A RECALIBRATED CHRONOLOGICAL FRAMEWORK FOR TEXAS
ARCHAEOLOGY–GEOARCHAEOLOGY

Thesis

Presented to the Graduate Council of
Texas State University-San Marcos
in Partial Fulfillment
of the Requirements

for the Degree

Master of ARTS

by

Ken Lawrence, B. A.

San Marcos, Texas
December 2010


A RECALIBRATED CHRONOLOGICAL FRAMEWORK FOR TEXAS

ARCHAEOLOGY-GEOARCHAEOLOGY


Committee Members Approved:


C. Britt Bousman, Chair


Stephen L. Black


Charles D. Frederick

Approved:


J. Michael Willoughby
Dean of the Graduate College

COPYRIGHT

By

Ken L. Lawrence

2010

FAIR USE AND AUTHOR'S PERMISSION STATEMENT

Fair Use

This work is protected by the Copyright Laws of the United States (Public Law 94-553, section 107). Consistent with fair use as defined in the Copyright Laws, brief quotations from this material are allowed with proper acknowledgment. Use of this material for financial gain without the author's express written permission is not allowed.

Duplication Permission

As the copyright holder of this work I, Ken Lawrence, refuse permission to copy in excess of the "Fair Use" exemption without written permission.

DEDICATION

Those who have come before whose research and insights we benefit greatly from and to those who will follow that will continue the work and correct our mistakes. This thesis is also dedicated to Freda Gibson (my first real teacher), my wife Susan (who has been my biggest supporter), and my study buddy Baxter (who we miss terribly).

ACKNOWLEDGEMENTS

As with any large endeavor, the assistance and influence of numerous people are needed. Such is the case with this thesis, which owes an enormous debt of gratitude to a multitude of individuals. Foremost, I am particularly grateful and consider myself fortunate to have the guidance of my esteemed committee, Dr. Britt Bousman, Dr. Steve Black, and Dr. Charles Frederick. If it were not for the knowledge, supervision, and prodding of my advisor Dr. Bousman, this thesis would not have been attempted. The insight, considerable knowledge of Texas prehistory, and significant encouragement of Dr. Steve Black were greatly appreciated. Further, the extensive knowledge of Dr. Charles Frederick regarding geoarchaeology and Texas drainages served as an inspiration before and during the research of the thesis. I hope I never quit learning from, and being inspired by, each of these committee members.

Deep appreciation is owed to a host of individuals who have directly or indirectly contributed or motivated me through the years. I am grateful to each of the following individuals who knowingly or unknowingly inspired and/or helped me. Dr. Jim Abbott, Michael Aiuvalasit, Dr. Michael Blum, Dr. Karl Butzer, Chris Caran, Dr. Reid Ferring, Victoria (Tory) Flores, Mary Gibson, Paul Heinrich, Dr. Vance Holliday, Dr. John Hoopes, Dr. Don Johnson, Chris Lintz, Dr. Brad Logan, Dr. Ray Mauldin, Dr. Rolfe Mandel, Charles Mear, Holly Meier, Dr. Lee Nordt, Tiffany Osburn, Elton Prewitt (and everyone at SHUMLA), Dr. Alan Skinner, Dr. Alston Thoms, Dr. Steve Tomka, and Melissa Winans.

I am also eternally grateful to a number of my colleagues and close friends who have been extremely helpful, provided significant support, and have greatly assisted me over the years. These individuals include everyone at SWCA as well as Jimmy Barrera, Stacey Buchanan, Shawn and Betty Cantrell, Steve and Carole Carpenter, Josh Haefner, Dr. Brett Houk, Dr. Jon Lohse, Kevin A. Miller, Lindsey Naslenas, Eric Oksanen, Chris Ringstaff, Dr. Matt Seddon, Kevin Stone, Suzanne Trussell, Ernest Wingate, and David Yelacic.

I would be nowhere without the love and support of my family. To my grandmother (Freda) I owe my appreciation for writing, which ultimately set me on my current path. To my parents (Gary and Judy) I owe my stubbornness to accomplish what I set my sights upon, to keep aiming for higher goals, and the ability to not take it all too seriously. To my brother (Eric) of whom I am extremely proud, I owe my competitive spirit and the appreciation of friends. Finally, to my best friend and wife (Susan) I owe everything. I am grateful for her advice, criticism, encouragement, and patience. I can never repay her for love and support particularly over the course of graduate school.

I am eternally grateful to all of these individuals for their encouragement and assistance and am a far better person for knowing each of them.

This manuscript was submitted on November 1, 2010.

Table of Contents

DEDICATION	v
ACKNOWLEDGEMENTS.....	vi
Table of Contents	viii
List of Tables	xi
List of Figures	xii
Abstract.....	xiv
CHAPTER 1	1
Introduction-Setting.....	1
Geology.....	4
CHAPTER 2	8
Methods and Principles.....	8
Radiocarbon Research	8
Bayesian Statistics	12
Methods.....	13
CHAPTER 3	18
Synthesis of Geoarchaeological Investigations Nueces River Basin	18
Previous Investigations	20
Geomorphic/Alluvial History	23
Upper- Dry Frio River (Woodrow Heard site).....	23
Lower- Choke Canyon.....	26
Calibration Results.....	29
CHAPTER 4	40
Recalibrated Geoarchaeological Framework within the Guadalupe-San Antonio River Basins	40
Previous Investigations	42
Guadalupe River Basin	45
San Antonio River Basin.....	46
Geomorphic/Alluvial History.....	48
Upper Extent San Antonio River Basin	48
Middle Extent San Antonio River Basin.....	53
Lower Extent San Antonio-Nueces Coastal Basin	56
Calibration Results.....	59
Upper Extent.....	59
Middle Extent	66
Lower Extent	69
CHAPTER 5	71
Recalibrated Geoarchaeological Framework within the Colorado River Basin of	

Texas.....	71
Previous Investigations	73
Geomorphic/Alluvial History.....	78
Upper Extent.....	78
Lower Extent	86
Calibration Results.....	91
Upper Extent.....	92
Lower Extent	95
CHAPTER 6.....	102
Recalibrated Geoarchaeological Framework within the Brazos River Basin of Texas.....	102
Previous Investigations	102
Upper Extent.....	107
Middle Extent	112
Lower Extent	122
Calibration Results.....	127
Upper Extent.....	127
Middle Extent	131
Lower Extent	135
CHAPTER 7	138
Recalibrated Geoarchaeological Framework with the Trinity River Basin.....	138
Previous Investigations	140
Geomorphic/Alluvial History.....	144
Upper Extent Trinity River Basin.....	144
Calibration Results.....	149
CHAPTER 8	158
Patterns and Correlations across Texas River Basins and Region	158
Intra-Basinal Patterns.....	159
Nueces River Basin.....	159
Inter-Basinal and Regional Patterns	164
Causal Factors	170
Climate	171
Eustasy	175
Correlations	181
Synchronous Event I.....	181
Synchronous Event II.....	184
Synchronous Event III	185
Synchronous Event IV	187
Summary	189
CHAPTER 9	191
Conclusions and Future Research	191
Interpretations.....	192
Future Research	197
Glossary	200

APPENDIX I-Nueces River Basin OxCal Results.....	202
APPENDIX II-San Antonio River Basin OxCal Results	211
APPENDIX III-Colorado River Basin OxCal Results.....	217
APPENDIX IV-Brazos River Basin OxCal Results	221
APPENDIX V: Trinity River Basin OxCal Results	234
APPENDIX VI-Categorized Bibliography	239
REFERENCES	251
VITA	283

List of Tables

Table 3.1	Previous Geoarchaeological Investigations in the Nueces River Basin.	21
Table 3.2	Radiocarbon data from Nueces River Basin.	30
Table 4.1	Previous Investigations in the San Antonio-Guadalupe River Basins.	43
Table 4.2	Radiocarbon data within the San Antonio River Basin.	60
Table 5.1	Previous Investigations within the Colorado River Basin.	74
Table 5.2	Radiocarbon data within Colorado River Basin.	93
Table 6.1	Previous Investigations within Brazos River Basin.	105
Table 6.2	Radiocarbon data from Brazos River Basin.	116
Table 7.1	Previous Investigations within Trinity River Basin.	141
Table 7.2	Radiocarbon data from Trinity River Basin.	150

List of Figures

Figure 1.1	Overview of Texas River Basins with Examined Study Areas.....	1
Figure 1.2	Overview of Natural Regions of Texas:	5
Figure 3.1	Nueces River Basin:	18
Figure 3.2	Initial Depositional History Nueces River Basin.	23
Figure 3.4	Calibration Plot of Woodrow Heard radiocarbon assays;.....	35
Figure 3.5	Calibrated Depositional History Nueces River Basin.....	36
Figure 4.1	Overview of Guadalupe-San Antonio River Basins:.....	40
Figure 4.2	Initial Depositional History San Antonio River Basin.	48
Figure 4.3	Idealized Profile of Jonas Terrace site (41ME29)	50
Figure 4.4	Cross-section at Richard Beene site on Medina River.....	54
Figure 4.5	Profile of Swan Lake deposits	57
Figure 4.6	Calibrated Depositional History San Antonio River Basin.	62
Figure 4.7	Calibration Plot of Jonas Terrace site (41ME29) radiocarbon assays.	63
Figure 4.8	Select Calibration Plots of the Richard Beene site radiocarbon assays;.....	67
Figure 5.1	Overview of Colorado River Basin:.....	71
Figure 5.2	Initial Depositional History Colorado River Basin.	78
Figure 5.3a	Profile of 41TG307 on Concho River.....	81
Figure 5.3b	Profile of 41TG309 on Concho River.....	82
Figure 5.4a	Idealized Cross-section of Colorado River Basin at the Eagle Lake locality	87
Figure 5.4b	Idealized Cross-section of Columbus Bend Members 1–3 at the Austin, West Point, and Columbus locations.....	88
Figure 5.5	Calibrated Depositional History Colorado River Basin.....	96
Figure 5.6	Select Calibration Plot of Lower Extent of Colorado River Basin.....	99
Figure 6.1	Overview of Brazos River Basin:	102
Figure 6.2	Initial Depositional History Brazos River Basin.	107
Figure 6.3	Idealized Profile of Lubbock Lake site	109
Figure 6.4	Idealized Cross-section of Cowhouse Creek (adapted from Nordt 1992: Figure 29).....	114
Figure 6.5	Idealized Cross-section of Brazos River at A&M study area.....	123
Figure 6.6	Calibrated Depositional History Brazos River Basin.....	129
Figure 6.7	Select Calibration Plot of Cowhouse Creek assays	133
Figure 7.1	Overview of Trinity River Basin:	138
Figure 7.2	Stratigraphic Columns from Aubrey Clovis site.....	144
Figure 7.3	Initial Depositional History Trinity River Basin.....	147
Figure 7.4	Calibrated Depositional History Trinity River Basin.....	154
Figure 7.5	Select Calibration Plot of Upper Trinity River Basin assays	155
Figure 8.1	Examples of a Hiatus and Chronometric Gap.	160
Figure 8.2	Patterns along Tributary Drainages in Study.	161
Figure 8.3	Patterns along Trunk Portions of Texas Drainages in Study.	165
Figure 8.4	Regional Climatic Patterns.....	168
Figure 8.5	Eustatic Patterns.	178

Figure 8.6	Calibrated cultural chronology contrasted with Synchronous Event I-IV	183
Figure 8.7	Radiocarbon dates from Central Texas burned rock middens.	188
Figure 9.1	Radiocarbon Frequency at Fort Hood Equated with Cultural Activity... ..	195

ABSTRACT

A RECALIBRATED CHRONOLOGICAL FRAMEWORK FOR TEXAS

ARCHAEOLOGY–GEOARCHAEOLOGY

By

Ken Lawrence, B. A.

Texas State University-San Marcos

December 2010

SUPERVISING PROFESSOR: C. Britt Bousman

Radiocarbon assays from select archaeological-geoarchaeological research projects within Texas river basins were compiled and recalibrated using the same calibration curve (i.e., INTCAL09). Chronometric data from investigations within the Nueces, San Antonio, Colorado, Brazos, and Trinity River basins were uniformly calibrated to construct a consistent chronological framework. Once calibrated, the analogous chronometric data were then used to compare drainage basins, paleoenvironmental data, and cultural chronologies across Texas and the region. These comparisons revealed four periods (Synchronous Events I–IV) in the Holocene that occurred simultaneously within all of the examined drainage basins. Synchronous Event I dating to 8,750–8,250 cal yr BP (~6800–6300 BC), Synchronous Event II dating to 7,000–6,250 cal yr BP (~5050–4300 BC), and Synchronous Event III 5,250–5,000 cal yr BP (~3300–3050 BC) are apparent periods of instability. While Synchronous Event IV occurs at 1,000–750 cal yr BP (~AD 950–1200) represents a period of stability. These events may be attributed to previously identified widespread climatic changes and seemingly coincide with several transitions in the archaeological record.

CHAPTER 1

Introduction-Setting

This study regards the construction of a recalibrated radiocarbon baseline from select archaeological-geoarchaeological research projects in Texas. The compilation of the chronometric data was the result of an extensive archival review that focused on previous research from sites containing deep, intact alluvial stratigraphy, which encompassed the Late Pleistocene–Holocene. Subsequently, select radiocarbon assays from previous investigations were compiled and recalibrated using the same calibration curve (i.e., INTCAL09). The uniform calibration of the radiocarbon assays provided a consistent chronological framework that can be used to compare drainage basins, paleoenvironmental data, and cultural chronologies across Texas.

There are several interrelated research objectives for the current study. The primary objective is to recalibrate radiocarbon data from previous archaeological-geoarchaeological investigations within select Texas drainage basins (Figure 1.1). The recalibration of these data will provide a chronological baseline for the comparison of Texas drainage basins and cultural components contained therein. Further, this chronological baseline will enable the evaluation of the depositional histories of Texas drainage basins for characterizing the integrity of each basin at various times. This study also compares the depositional histories of the selected drainages within a basin to each other (i.e., intra-basinal) and with other drainages in Texas and the region (i.e., inter-




Figure 1.1 Overview of Texas River Basins with Examined Study Areas.

1) Woodrow Heard, 2) Choke Canyon Reservoir, 3) Jonas Terrace, 4) Richard Beene, 5) Copano Bay, 6) San Angelo, 7) Lower Extent Colorado River, 8) Lubbock Lake, 9) Fort Hood, 10) A&M-College Station, and 11) Upper Trinity River Basin.

basinal) in an effort to discern similarities. Observed patterns within this study are then correlated with external mechanisms (e.g., climate and eustatic effects), which may have contributed to these occurrences. Finally, the depositional patterns of the basins are compared with the archaeological record in Texas to demonstrate a correlation.

This chapter includes a review of the geology of the selected Texas drainage basins. Chapter 2 reviews of the Methods and Background for this study. Subsequent to this is a discussion of the selected drainage basins of this study, beginning with the Nueces River Basin (Chapter 3), the Guadalupe-San Antonio River basin (Chapter 4), the Colorado River basin (Chapter 5), the Brazos River basin (Chapter 6), and concludes with the Trinity River basin (Chapter 7). Each drainage basin chapter briefly reviews previous investigations conducted within the basin.

The radiocarbon datasets selected for recalibration because they satisfied three primary criteria (assays of charcoal, assays 'corrected' for isotopic fractionation, and those assays in good stratigraphic context). Additionally, selected datasets have been utilized by other researchers for characterizing the depositional and/or paleoenvironmental history of these study areas. These data from widely accepted studies were recalibrated in part to demonstrate the implications for recalibrating chronometric data. There are numerous previous investigations with chronometric data scattered within drainage basins that were not selected for recalibration. Those datasets were omitted because either they 1) did not meet the predefined criteria, 2) had been recently calibrated, 3) had a sparse chronometric dataset, or 4) a combination of these factors. Several of the recalibrated datasets within this study do suffer from some of these concerns, but were included out of necessity.

Chapter 8 consolidates and reviews the results from all of the basins and discusses the identification of geomorphic patterns and possible correlations derived from the calibrated chronological baseline. Chapter 9 summarizes the conclusions from the recalibrated basins and interpretations that developed during its course, as well as reviewing several study areas and topics for future research and investigation.

Geology

With some exceptions, the surface geology of the selected drainage basins in Texas typically cross progressively younger geological deposits as they trend toward their terminus at the Gulf of Mexico. The geology of the High Plains region, which contains the headwaters of the Colorado and Brazos River basins include erosionally softer sedimentary Tertiary and Quaternary deposits before descending into significantly older Lower Cretaceous (Colorado River basin) or Permian deposits (Brazos River basin) in North-central Texas (Spearing 1991). The sedimentary Permian-aged deposits crossed by the Brazos River basin and portions of the Colorado River basin include soft beds of shale, sandstone, and gypsum. The Trinity River basin joins these basins as they cross lower and upper Cretaceous-aged deposits of limestone, sandstone, marl, and shale (Ferring 2001; Spearing 1991; Williams 2004). The Colorado and Brazos River basins enter into the Edwards Plateau region while the Trinity River basin exits the Cretaceous-age surface geology and enters Paleocene deposits of fine grained mixed clastic including clay, silt, and sand (USGS 2010). Within the Edwards Plateau, the Brazos and Colorado River basins are joined by the headwaters of the Guadalupe-San Antonio and Nueces River basins. The southern and eastern edges of the Edwards Plateau are indicated by the Balcones escarpment formed by a fault zone (Spearing 1991). Uplift along the Balcones

Fault has subsequently caused the edges of the Edwards Plateau to erode away along waterways that form flat-topped hills with stair-stepped tiers clearly demarcating its separation from the Coastal Plains. Below the Cretaceous age geology of the Plateau, the Colorado, Brazos, Guadalupe-San Antonio, and Nueces River basins enter the broad Tertiary and Quaternary age deposits of the Coastal Plains. The nearly level Coastal Plains consist of progressively younger beds from the Tertiary and Quaternary composed of sandstone, siltstone, and mudstone, while the drainages throughout the basin largely contain Holocene age deposits of clay, silt, sand, and gravel.

The implications of the respective geology for each of the basins regard the development of their channels. Notably, Ferring (1994:150, 2001) argues that the underlying bedrock lithology influences the Trinity River basin. Specifically, the resilience of the surface geology to erosion affects the evolution of the upper basin landforms and consequently the vegetation. The limestone bedrock in the western portion of the Trinity River basin erodes into calcareous soil supporting a prairie environment while bedrock composed of noncarbonate deposits (e.g., sandstone and shale) erode into non-calcareous soils that supports a mixed forest environment (Ferring 1994:150). A corollary of this is the vegetation of a region influences the rate of erosion and sediment budget for the drainage (Figure 1.2). Ferring (1994:150) proposes that areas with non-calcareous soils are more susceptible to erosion with the implication being possibly poor conditions for the preservation of cultural deposits. Similarly, the drainage basins that cross the Edwards Plateau, in particular the Colorado River basin, cross limestone of varying resistance, thin marls, shale, and chalk (Abbott 1994:359–360).




Figure 1.2 Overview of Natural Regions of Texas: 1) Piney Woods, 2) Oak Woods & Prairies, 3) Blackland Prairie, 4) Gulf Coast Prairies & Marshes, 5) Coastal Sand Plain, 6) South Texas Brush Country, 7) Edwards Plateau, 8) Llano Uplift, 9) Rolling Plains, 10) High Plains, 11) Trans Pecos, and 12) Marine Environment (adapted from TPWD 2009).

Once the drainage basins enter the Coastal Plain, the deposits become progressively younger and composed of sandstone, siltstone, and mudstone while the drainages throughout the basins largely contains localized deposits of Holocene age clay, silt, sand, and gravel. Simply put, as the drainage basins transition from the more resistant limestone to the less resilient bedrock lithology of the nearly level Coastal Plains, the drainages convert from smaller bedrock incised channels with low-moderate sinuosity to larger floodplain channels with high sinuosity (Thoms and Mandel 2007). Thus, the lower drainage basins have higher potential for well-preserved, stratigraphically isolated cultural deposits. Consequently, the quantity and integrity of alluvial deposits within a drainage largely depends upon the portion of a basin (e.g., upper, middle, or lower extent) within which it is located.

CHAPTER 2

Methods and Principles

Radiocarbon Research

Previous researchers have expertly and thoroughly addressed the history and application of radiocarbon analysis (e.g., Libby 1955; Lowe and Walker 1997; Taylor 1987, 1997, 2009). Briefly however, few discoveries have had such a profound impact on archaeology as that of the process of radiocarbon (^{14}C) dating (Ramsey 2008:249). Since radiocarbon's discovery by Willard Libby in the late 1940s, researchers could now scarcely imagine conducting archaeological research without it (Bowman 1990; Huntley 1985; Libby 1955; Taylor 1987, 1997, 2009). The introduction of radiocarbon analyses provided absolute dates for archaeological deposits in contrast to the gross approximation of relative dating from methods like stratigraphy. A chronological framework could now be constructed with a foundation accepted by almost all researchers. Further, the temporal information derived from radiocarbon analysis allowed comparisons to other sites, regions, or countries. Whereas, the relative dating supplied by stratigraphic provenience was only applicable to an immediate area or region. The ramifications of the process on the field of archaeology were substantial and widespread.

There are two analysis methods for the measurement of ^{14}C , the original conventional method and the more recent Accelerator Mass Spectrometry (AMS) (Bowman 1990:12; Lowe and Walker 1997:241). At its simplest, the conventional method consists of counting the remaining electrons of a weighted sample to determine

the ^{14}C rate of emission (Bowman 1990:12). By contrast, the AMS method 'accelerates' particles in a sample in order to compare the proportion of ^{14}C atoms to the atoms identified in the ^{13}C and ^{12}C in the sample (Bowman 1990:12; Lowe and Walker 1997:241). Both methods effectively provide a similar output, which (most importantly) can be converted to a chronological measurement. In this study, a predominance of the selected samples utilized conventional methods and a distinction is made when AMS methods were used.

The principle behind radiocarbon dating is that one of the isotopes of carbon (^{14}C), a naturally occurring element in all living organisms, has a prolonged rate of decay that can be measured (Banning 2000; Bowman 1990; Christen 1994; Hedman 2007; Libby 1955). Libby determined that once an organism dies its remains stop accumulating ^{14}C and the radioactive decay of ^{14}C could be determined, which he calculated had a half-life of 5568 ± 30 years (Banning 2000; Bowman 1990; Libby 1955). Although, subsequent research has refined the ^{14}C half-life to be 5730 ± 30 years, laboratories continue to use Libby's calculation (e.g., 5568 ± 30 years) to prevent confusion (Banning 2000; Bowman 1990; Christen 1994; Libby 1955; Lowe and Walker 1997; Stuiver and Polach 1977). One of Libby's assumptions was that a constant amount of ^{14}C has entered the atmosphere throughout time and that this exchange has been evenly distributed (Christen 1994; Taylor 1987, 1997, 2009). However, researchers have since determined that the amount of ^{14}C has extensively fluctuated globally and throughout time. To account for these atmospheric ^{14}C variations (sometimes called the de Vries effect) among other key assumptions of Libby's determined to be incorrect, a suite of operating parameters were adopted (Stuiver and Polach 1977). Some of the parameters

include researchers using the aforementioned 5568 ± 30 year half-life, using A. D. 1950 as the beginning point of before present (BP), and a recognition of isotopic fractionation effects whereby ^{14}C samples must be 'normalized' to a baseline of $\delta^{13}\text{C}$ value of -25.0‰ (Hua 2009; Mook and Waterbolk 1985; Stuiver and Polach 1977; Taylor 1997:67–68). In short, a system of calibration must be implemented to identify the ^{14}C discrepancies in order to use them in a chronological application (Ramsey 2008:260).

Another common error with the ^{14}C dating process is the misinterpretation of the results by researchers (Banning 2000). One often-occurring mistake is equating of ^{14}C result with a calendar age by archaeological researchers. The ^{14}C result is not a calendar date, but rather, a ratio of isotopes (Bartlein et al. 1995; Blockley et al. 2007; McCormac and Baillie 1993; Lowe and Walker 1997:243; Mock and Bartlein 1995; Ramsey 2008, 2009:337; Stuiver and Suess 1966). When this disparity occurs, the researcher often rejects, misuses, and/or incorrectly reports the results, which may lead to a misinterpretation that cascades throughout their research. Considering that the use and reporting of ^{14}C results (calibrated and conventional) differ throughout North American and European journals, it is small wonder that these errors and misconceptions occur (Taylor 1997:68–69).

The incorporation of dendrochronology (tree-ring dating) with radiocarbon analysis has been the most effective calibration method, which has been an instrument of calibration has only occurred since the late 1950s (Taylor 1997). One method of this process uses the known age provided by tree rings and then processes them through radiocarbon analysis typically dating the tree rings in decadal or bidecadal year increments (Lowe and Walker 1997; Nash 1999; Taylor 1997; Walker 2005). The result

of such analyses from around the world has been the refinement and extension of the calibration curve (Klein et al. 1982). Although researchers have identified these atmospheric variations, they have also determined that the extent of the discrepancies is more widespread and occasionally more pronounced than previously recognized.

This ^{14}C calibration curve is continually being supplemented and refined from collected data and since 1981 published in various journals (e.g., *Radiocarbon*) and on the internet (Banning 2000:268). One particular group, the IntCal Working Group (IWG), comprised of international scientists from a variety of disciplines, is developing an internationally agreed-upon calibration curve (Blackwell and Buck 2008:227). This international curve (IntCal) is frequently updated and presented for the use of researchers (e.g., Reimer et al. 2009). Recent calibration curves have been produced in 1998 (IntCal98), 2004 (IntCal04), and most recently in 2009 (IntCal09). Of primary relevance, with the introduction of each of these calibration curves, the radiocarbon datasets that had been previously calibrated using earlier calibration curves (e.g., IntCal98) are not fully comparable with the most recent calibration curve. Consequently, the ^{14}C data that had been previously calibrated needs 'adjustment', sometimes significantly to be correlated with more recent ^{14}C data. The most dramatic alterations to ^{14}C data regard those assays that date to roughly 7,000 ^{14}C BP or older, where calibration data previously has been more sparse.

Due to these abundant deviations of the ^{14}C fluctuations, numerous techniques have been developed for the calibration, interpretation, and presentation of the results (e.g., Acabado 2009; Blackwell et al. 2006; McCormac et al. 1993; Michael and Klein 1979; Ralph et al. 1973; Steier et al. 2001; Talma and Vogel 1993; Taylor et al. 1996;

Wohlfarth 1996). One process includes calibration software (e.g., OxCal, Bcal, and Calib), which interprets the conventional ^{14}C data with a probability distribution (Banning 2000; Ramsey 1995, 1998, 2001, 2008, 2009). In essence, the calibration software plots the ^{14}C dating results onto the calibration curve and then characterizes the probability of the interpreted outcome. There are more than a half dozen software packages, but the one used for this study is OxCal Version 4.1.6, which can incorporate Bayesian statistics into the characterization of the dating results.

Bayesian Statistics

Broadly defined, Bayesian statistics enables a researcher to incorporate data from the calibration curve with new data (e.g., ^{14}C results) as well as accounting for prior information (e.g., stratigraphy) to suggest the most probable outcome (Bayliss and Ramsey 2004; Blackwell and Buck 2008; Buck et al. 1991; Buck et al. 1992; Buck et al. 1994; Buck et al. 1996, Buck et al. 2004; Buck 2004; Christen 1994; Heaton et al. 2009; Ramsey 2009; Sharon 2001). The technique uses Baye's theorem that expresses the uncertainty of an event or set of parameters occurring before and subsequent to the results of an analysis (Buck 2004; Buck et al. 1996; Christen 1994). Put another way, the Bayesian analysis of ^{14}C calibration data considers all possibilities of the outcome (prior) with the measured data (likelihood), and then determines the probability of those results occurring (Ramsey 2009). A prior probability is inferred from relative dates (e.g., stratigraphy) and then compared with the likelihood probability that is interpreted from the absolute dates (e.g., ^{14}C dates), which culminates in the determination of the posterior probability.

When comparing multiple events simultaneously, the large number of individual

combinations that are considered for each outcome (iteration) can become infinite. To aid in these analyses by calibration software, a sampling algorithm such as the Metropolis-Hastings or equivalent is typically used (Ramsey 2009). Metropolis-Hastings is an algorithm of the Markov chain Monte Carlo (MCMC) method, which randomly examines each event across a defined distribution gradually increasing the confidence of the result (Breyer 2009; Heaton et al. 2009; Ramsey 2009). Further, the implementation of MCMC allows for the inclusion of the uncertainty of multiple factors that can allow for the comparison of points as well as their deviations on the curve (Buck and Blackwell 2004:1101; Everitt 2002; Upton and Cook 2006).

In summary, the analysis and interpretation of ^{14}C results requires calibration due to extensive atmospheric ^{14}C variations that have been recognized. The calibration of the ^{14}C dates, whether individually or in multiple sets, are plotted on an internationally agreed upon calibration curve (currently IntCal09), which is continually updated and periodically published. Due to these refinements, the results of previous ^{14}C analyses, even though calibrated, require adjustment. In addition, numerous software packages are used for the calibration of the ^{14}C results that use some form of Bayesian statistics to characterize the results most accurately. For the calibration and comparison of multiple events, the Bayesian analysis commonly utilizes MCMC sampling algorithms (e.g., Gibbs Sampler and Metropolis-Hastings) for the most appropriate outcome (Ramsey 2009).

Methods

Several additional concepts and definitions warrant discussion and clarification. A brief review of relevant components pertaining to this study follows and, when

appropriate, a more detailed discussion is provided in later chapters. There were three main criteria used for selecting the radiocarbon assays for this recalibration study.

Namely, 1) assays of charcoal were given priority over other dated materials; 2) samples that have been previously 'corrected' for isotopic fractionation, and finally; 3) datasets composed of samples in good stratigraphic context.

Previous investigators have noted that bulk humate (both sediment and soil) tends to date inconsistently, sometimes drastically older (approximately 1,000–1,500 years) than comparable charcoal samples (Abbott 1994:375; Fowler et al. 1986; Grimm et al. 2009; Mandel et al. 2007:50; Matthews 1985; Martin and Johnson 1994; Nordt 1992:9–10; Wang et al. 1996). One reason proposed for this phenomenon is attributed to mean residence time (MRT) of the soil sample. Simply put, the MRT of the bulk humate is a weighted average of the organic components within the sample (Lowe and Walker 1997: 247–248; Schaetzl and Anderson 2005). Thus, any radiocarbon analyses of bulk humate samples may encompass a suite of organic matter that could provide an imprecise measurement beyond use. For this study, bulk humate samples from previous investigations have been avoided as much as possible. Only when charcoal samples were not available for an important allostratigraphic unit (e.g., Jackson alluvium in Fort Hood) were bulk humate samples used. Although, these data from bulk humate samples are included in summary table they have not been included in recalibration exercise unless essential.

Radiocarbon years (^{14}C) are not calendar years, but rather a measurement of remaining ^{14}C isotopes (McCormac and Baillie 1993; Lowe and Walker 1997:243; Ramsey 2009:337; Stuiver and Suess 1966; Taylor 1997:68). Thus, the ^{14}C years must be

converted to a calendrical format (i.e., calibrated) (Mook and Waterbolk 1985:20; Van der Plicht and Hogg 2006:238). Regarding the issue of isotopic fractionation (i.e., $\delta^{13}\text{C}$ value of -25.0‰), all samples selected for this study have been adjusted for fractionation by the initial investigators and represent the 'corrected' value. Similarly, several assays reviewed and calibrated within this study were derived from marine shell. Researchers have long recognized a discrepancy between radiocarbon dates from terrestrial samples and shell from a marine environment. This difference (reservoir effect) differs by as much as 400 radiocarbon years from their terrestrial equivalent and must be corrected (Stuiver and Braziunas 1993). Those shell assays reviewed from previous investigations for this study were not corrected or undetermined to have been corrected for this reservoir effect. Likewise, the current study did not correct for the reservoir effect when calibrating shell assays. This study has also adopted the nomenclature of Nordt (1992) and the journal *American Antiquity* for distinguishing between uncalibrated radiocarbon and calibrated radiocarbon results. Specifically, in this study uncalibrated radiocarbon years are reported as ' ^{14}C yr BP' while calibrated radiocarbon samples are indicated as '**cal yr BP**'.

Regarding the third criteria of context, previous investigations that had conducted extensive radiocarbon analyses of depositional stratigraphy (e.g., allostratigraphic units) were almost exclusively selected. The term allostratigraphic unit used herein follows the definition indicated by the North American Commission on Stratigraphic Nomenclature (NACSN). Briefly, this refers to a mappable body of sedimentary rock bounded by a discontinuity (NACSN 2005:1578). In this study the use of allostratigraphic unit is appropriate because it provides a recognizable system for characterizing fluvial deposits

of previous investigations reviewed here (Jacobson et al. 2003:36–37). However, the nomenclature of the initial researchers has been adopted when available. The purpose of selecting datasets in this context is to provide additional information (e.g., the prior) for Bayesian analyses and provide for the previously mentioned phase model and sequential setting to get the most probable statistical outcome. Further, the recalibration and characterization of allostratigraphic units of one study area ideally can be contrasted with the results of other allostratigraphic units in other basins. The objective of this ^{14}C recalibration is to uniformly calibrate and present the results to produce a consistent chronological framework. Once ^{14}C results are on the same baseline, they can be used by researchers for regional comparisons and refinement of cultural chronologies. Additionally, the depositional history of select Texas drainage basins as well as the chronology of cultural activities can be reevaluated.

For the calibration and comparison of multiple events from previous investigations, the current study used the OxCal v4.1.6 program, which utilizes the MCMC Metropolis-Hastings sampling algorithm, a collaborative component of Bayesian analyses (Ramsey 2009). For specific OxCal v4.1.6 operations, the calibration curve utilized IntCal09 and the analyses were primarily conducted using default settings. However, output was set at both 68.2% (1σ) and 95.4% (2σ) and rounded to the nearest decade for a minimum of 30,000 iterations, but frequently went over 3 million iterations. Due to the nature of the samples selected for recalibration (i.e., multiple samples) a calibration model was utilized. Specifically, these analyses focused on previous investigations with radiocarbon samples collected from recognized horizons in a stratigraphic setting. Using these criteria, analyses were conducted under the assumption

that horizons at lower elevations will be older than shallower samples (i.e., Law of Superposition). When recalibrating the suites of radiocarbon assays for this study they were grouped in the stratigraphic horizon identified (when available) by the previous investigator. Consequently, the analyses were conducted in OxCal using the phase model under the Sequential setting. This setting allows for sets of radiocarbon samples to be grouped in a particular sequence (i.e., stratigraphic) and contrasted both within the group and against other groups (Ramsey 2009). Further, the Sequential setting operates under the assumption that another group (e.g., horizon) cannot be temporally contiguous or overlapping (Ramsey 2009). Ultimately, the intent of using this suite of techniques is to derive as much information as possible from previous research, compare the results with an equivalent metric, and, to substantiate or update previous interpretations when necessary.

CHAPTER 3

Synthesis of Geoarchaeological Investigations Nueces River Basin

The Nueces River basin of Texas is relatively small basin (16,800 square miles), which composes a significant portion of South-Central Texas (Durbin 1999; Weddle 2010). With the exception of its headwaters, the Nueces River basin is south of the Edwards Plateau (Figure 3.1). It crosses the South Texas Plains then the Gulf Prairies and Marshes before entering Nueces Bay in San Patricio and Nueces Counties. Upon exiting the Plateau, waterways cross the coastward-sloping Coastal Plains and become gradually more sinuous. Some of the prominent tributary drainages contributing to the roughly 315-mile-long (500 km) Nueces River include the West Nueces, Sabinal, Frio, Dry Frio, Atascosa, and Leona rivers as well as Indian, Seco, Hondo, Verde, San Miguel, and Hackberry creeks (NRA 2010; Weddle 2010).

Three drainages form the main trunks of this basin with the other tributaries dendritically draining into these waterways (Figure 3.1). The primary trunk, the Nueces River, drains the western portions of the basin beginning in Edwards County and crossing Uvalde, Zavala, Dimmit, La Salle, McMullen, Live Oak, Nueces, and San Patricio Counties. Within Live Oak County, the Nueces River converges with the other two main basin trunks, the Frio and Atascosa rivers. The Frio River, begins at a spring in Real County where it trends southward, is joined by the Sabinal River in Uvalde County, and meets its confluence with the Nueces River 250 miles (400 km) from its source




Figure 3.1 Nueces River Basin: 1) Woodrow Heard, 2) Choke Canyon Reservoir, and 3) Paine 1991 study area.

(Texas Parks and Wildlife Department 2009). The third main waterway is Atascosa River, a seasonal drainage, which begins as two minor branches (North and West Prongs) in Bexar and Medina Counties, respectively. The Atascosa River runs south-southeast along the eastern margins of the basin prior to converging with the Frio River in Live Oak County.

Previous Investigations

Archaeological investigations, particularly in the Edwards Plateau region, have been conducted within the Nueces River basin since the 1930s (Sayles 1935). However, only a select few have conducted a serious examination of the geomorphic history within the basin (Table 3.1). The investigations that incorporated geology and archaeology occurred relatively early with Mear's (1953) master thesis work along the Sabinal River. Since that time, roughly a dozen geoarchaeological investigations have been carried out, but most of these were typically assessing site integrity or had a similarly narrow focus (e.g., Brown et al. 1982; Scott and Fox 1982; Taylor and Highley 1995).

One investigation of particular relevance occurred in the early 1980s at the northern extent of the Nueces River Basin occurred at the Woodrow Heard site (41UV88) on the Dry Frio River in northern Uvalde County (Figure 3.1). Although not extensive, a component of this research involved a deliberate geoarchaeological investigation of the Dry Frio River valley around the site and included a geomorphic assessment with a series of radiocarbon analyses (Decker et al. 2000).

One of the larger, more comprehensive archaeological investigations in South Texas was undertaken in the 1970s at Choke Canyon Reservoir situated on the lower reaches of the Frio River in Live Oak and McMullen Counties (Figure 3.1).

Table 3.1 Previous Geoarchaeological Investigations in the Nueces River Basin

Drainage Extent	Drainage	Resource	Project-Site(s)	Geoarchaeologist or Researcher	Chronometric Data*
Lower	Nueces River	Johnson 1933, Price 1933, Weeks 1933, 1945	Coastal region	E. Johnson, W. Price, and A. Weeks	Relative
Upper	Sabinal River	Mear 1953, 1990, 1995, 1998	Sabinal River Valley; thesis, articles	Charles Mear	Relative; some uncorrected 14c
Upper	Nueces River	Gustavson 1978	Nueces River; article	Thomas Gustavson	Relative
Lower	Frio River	Brown et al. 1982; Hall et al. 1982; Scott and Fox 1982	Choke Canyon Reservoir; 41LK31, 41LK32, 41LK67, and 41LK202; CRM report	Russell Bunker (vol. 5)	Relative; sparse 14c
Lower	Hackberry Creek-Atascosa/Frio River	Taylor and Highley 1995	Loma Sandia (41LK28); TxDOT Report	Vance T. Holliday	Relative; Uncorrected, uncalibrated 14c
Lower	Nueces River	Baskin and Cornish 1989; Cornish and Baskin 1995	Lower Nueces River sedimentation; article	F. G. Cornish and J. A. Baskin	Very sparse 14c
Lower	Nueces River	Ricklis 1988, 1993; Ricklis and Blum 1997; Ricklis and Cox 1998	McKinzie Site (41NU221); 41NU266, 41NU268, White's Point sites; article	Robert Ricklis	14c shell; Relative
Lower	Nueces River	Durbin 1999	Late quaternary geomorphic change to lower Nueces River; Dissertation	James M. Durbin	TL; OSL; Relative
Upper	Dry Frio River	Decker, Black, and Gustavson 2000	Woodrow Heard (41UV88); TxDOT Report	Thomas Gustavson	Relative; calibrated 14c
Lower	Nueces River	Simms 2005	Late quaternary coastal shelf evolution of Nueces River; Dissertation	Alexander Simms	Corrected, calibrated 14c

indicates study selected for calibration

* TL=thermoluminescence, OSL=optically stimulated luminescence

The investigations, utilizing a broad spectrum of multi-disciplinary approaches consisted of survey, testing, and data recovery of a number of archaeological sites (Hall 2010). Despite significant research contributions, the chronometric data for these multi-year investigations was severely limited.

In the late 1980s, research was conducted regarding sedimentation of the Nueces River during the Late Quaternary (Figure 3.1). These investigations, conducted on the lower extent of the Nueces River, identified four terraces (Angelita, Fort Lipantitla, Bluntzer, and Corpus Christi) in addition to the modern floodplain (Baskin and Cornish 1989; Cornish and Baskin 1995:193). These researchers identified several allostratigraphic units that they correlated to the Holocene. These units include the Cayamon Creek Allomember 1 (CCA-1) associated with the Terminal Pleistocene/Early Holocene, Cayamon Creek Allomember 2 (CCA-2) Middle to Late Holocene, and the Cayamon Creek Allomember 3 (CCA-3) Recent (Cornish and Baskin 1995).

Beginning in the mid 1980s, investigations were conducted at the McKinzie Site (41NU221) along the Nueces River overlooking Nueces Bay (Figure 3.1). This research was compared to results from 22 other archaeological sites in similar settings at Baffin Bay, Copano Bay, Guadalupe Bay, and Lavaca Bay (Ricklis 1988, 2004; Ricklis and Blum 1997; Ricklis and Cox 1998). Situated in upland settings, the sites all had stratified shell middens that provided 80 radiocarbon assays (Ricklis and Blum 1997).

Also in the lower extent of the basin, a more recent geomorphic investigation along the Nueces River at the Gulf of Mexico was the doctoral research by Durbin (1999). The research examined the responses of the Nueces River to changes in the climate and fluctuations in sea level using new research and previous investigations

(Durbin 1999). Encompassing the last 120,000 years, a component of this research investigated the proposal that rising sea levels instigated valley aggradation while conversely dropping sea levels caused valley incision (Durbin 1999).

Geomorphic/Alluvial History

The result of these geoarchaeological investigations is a set of Late Quaternary stratigraphic records within the Nueces River basin (Figure 3.2). These are discussed geographically, beginning at the upper (northern) extent of the basin and then downstream to the lower extent of the basin to the Nueces Bay.

Upper- Dry Frio River (Woodrow Heard site)

The investigations at the Woodrow Heard site (41UV88) provided chronometric data for the Dry Frio River and the basin (Figure 3.1). This research provided thirty radiocarbon samples, predominantly composed of charcoal (n=23) and fourteen of these samples, including two derived from humate, were used to establish a geomorphic chronology of the valley (Gustavson 2000:114–123).

The geoarchaeological analysis at Woodrow Heard identified two stratigraphic units (Units I and II) composing the Dry Frio River terrace (Decker et al. 2000:114–117). The base of Unit I was not observed during investigations, but the observed stratigraphy consisted of a series of fining upward deposits beginning with gravels (Figure 3.3). The chronometric data from Unit I indicated deposition prior to 8,000 ¹⁴C yr BP (Decker et al. 2000:117). Subsequent to this, but prior to roughly 6,400 ¹⁴C yr BP, the drainage migrated laterally and began to deposit Unit II (Decker et al. 2000:117). In addition, a disconformity separates the two stratigraphic units, suggesting a period of erosion. Unit II consists of four internal deposits (Units IIa–IId) from oldest to youngest, respectively




Figure 3.2 Initial Depositional History Nueces River Basin.




Figure 3.3 Profile of Woodrow Heard Stratigraphy (adapted from Decker et al. 2000: Figure 54).

(Figure 3.2). Each of these units was recognized by slight differences in parent material. The radiocarbon data indicate that Unit IIa was deposited between 6,400–6,000 ^{14}C yr BP, Unit IIb was deposited between 5,650–4,710 ^{14}C yr BP, and Unit IIc was deposited sometime after 3,270 ^{14}C yr BP (Decker et al. 2000:117–124). No chronometric data was available for Unit IId.

Lower- Choke Canyon

The Choke Canyon project in Live Oak and McMullen Counties was one of the largest projects conducted within the basin. Unfortunately, the chronometric data was severely limited. During Phase I investigations at the Possum Hollow Site (41LK201), over 70 radiocarbon samples were collected, but only seven were processed (Highley 1986). Similarly, on the Gates-Rowell Site (41LK31/32), only three radiocarbon samples were measured (Scott and Fox 1982:34). In contrast, the Phase II investigations of Choke Canyon Reservoir fared better chronometrically. Forty-three (MASCA corrected) radiocarbon samples (wood charcoal) collected from seven prehistoric sites were submitted for analyses (Hall et al. 1986).

Several notable results from the Choke Canyon project stand out. The first is that there were several gaps in the radiocarbon dates. The most prominent gaps occurred between 5,780–4,790 ^{14}C yr BP, 4,610–4,130 ^{14}C yr BP, 3,810–3,360 ^{14}C yr BP, and 1,800–1,520 ^{14}C yr BP (Hall et al. 1986:585–588). The researchers attributed the gaps to possible sampling bias, preservation, human settlement pattern discontinuities, or a combination of these factors (Hall et al. 1986:587). Regarding the chronological assessment of the Frio River valley stratigraphy, the researchers identified that this was an area needing future research (Hall et al. 1986:590). Even though the Choke Canyon

investigations comprised the largest group of radiocarbon assays in Southern Texas and a geomorphic study was undertaken (i.e., Bunker 1982), radiocarbon dating of the depositional history of the Dry Frio River was not a research focus.

Bunker (1982) did recognize four terraces (1–4) in the Frio River Valley encompassing modern to Pleistocene in age (Figure 3.2). Other, older, terraces were observed in the valley, but these were characterized as discontinuous and isolated (Bunker 1982:499). No chronometric data was indicated for Terrace 4, but Bunker (1982:501) infers that it is Pleistocene in age. The third terrace (Terrace 3), containing most of the Choke Canyon Reservoir archaeological sites, began construction sometime before 5,330 ^{14}C yr BP and was characterized by extensive lateral migration eroding older deposits and bearing a coarse bedload until roughly 2,280 ^{14}C yr BP (Bunker 1982:514–515). Sometime after 2,280 ^{14}C yr BP, the Frio River straightened, incised and began construction of the second terrace. Terrace 2 was described as having a decrease in overbank flooding episodes with more stability (Bunker 1982:515). No chronometric data were available for determining the end of Terrace 2 construction and beginning of Terrace 1. However, Bunker (1982:511) suggests that sometime after 2,000 ^{14}C yr BP, the Frio River incised isolating Terrace 2 and forming the modern floodplain (Terrace 1).

Near the coast, Durbin (1999) identified Pleistocene Deweyville allostratigraphic units and Holocene Post-Deweyville (PD) allostratigraphic units (Figure 3.2). The Pleistocene Deweyville units consisted of High Deweyville (HD), Middle Deweyville (MD), and Low Deweyville (LD) and were dated using Thermoluminescence (TL) and Optically Stimulated Luminescence (OSL) methods. The more recent Post-Deweyville

(PD) unit was recognized as Terminal Pleistocene/Early Holocene (PD1), Middle to Late Holocene (PD2), and Recent (PD3) (Durbin 1995:119–131). Durbin (1999:124) did correlate PD1 to the Cayamon Creek Allomember 1 (CCA-1) identified by Cornish and Baskin (1995). Similarly, units PD2 and PD3 correlate with CCA-2 and CCA-3, respectively. Unfortunately, the chronometric data for the Holocene PD units was limited to two OSL dates from Durbin's study (1999:118–124) and a single radiocarbon sample from another investigation by Cornish and Baskin (1995).

Durbin (1999) concluded that climate and sea level (eustatic) fluctuations affect valley deposits. However, sea level affects were not as pronounced, possibly to less than 40–100 km from the coast, as previous researchers had proposed (Durbin 1999:149–150; Etheridge et al. 1998). Further, the Pleistocene Deweyville units were deposited under cooler and moister conditions than the Holocene when sea levels were lower. Consequently, the Deweyville units had elevated flow regimes producing larger meanders and coarser bed loads with each unit incising into previous, units resulting in stair-stepped terraces (Durbin 1999:180–183). In contrast, the Terminal Pleistocene to Holocene PD units were constructed under the transition to progressively more arid and warmer climates and rising sea levels. These conditions decreased the flow regime and increased the amounts of sediment contribution through erosion causing gradual vertical aggradation with each unit covering the preceding one (Durbin 1999:180–183). Thus, the stratigraphy of the PD units is well-defined allostratigraphic units (Durbin 1999:182–183). Unfortunately, due to the paucity of chronometric data for the PD units, the chronology of these units is poorly defined.

Near to Durbin's research were the investigations conducted in Nueces Bay by Ricklis (2004) examining shell middens on upland sites overlooking coastal estuaries. Among several important observations, this work is helpful to correlate sea level rise with prehistoric occupation along the coast (Ricklis 2004; Ricklis and Blum 1997; Ricklis and Cox 1998). Two occupation hiatuses were observed between 6,800–5,900 ^{14}C yr BP and 4,200–3,000 ^{14}C yr BP and correspond to higher sea levels (Ricklis 2004:175–177; Ricklis and Blum 1997:299–300).

Calibration Results

For a variety of reasons, only the suite of assays from the Woodrow Heard site at the northern extent of the basin met this study's calibration criteria (e.g., stratigraphic control and $\delta^{13}\text{C}$ corrected). Although the chronometric data from Ricklis' (2004) investigations do not meet the criteria for this study, his dates have been recalibrated due to its broad implications for the Nueces River basin and the coast in general. Despite the limited dataset, the calibration of select radiocarbon samples from the Woodrow Heard site and Ricklis' research proved beneficial and informative (Table 3.2). This is particularly true of the Woodrow Heard assays. Specifically, the stratigraphic history of the Dry Frio River may, in actuality, be more reflective of basin changes affected by climate.

The calibration of Unit I at the Woodrow Heard site revealed that this horizon was constructed prior to 9,480 cal yr BP and ended after 8,810 cal yr BP. Following the construction of Unit I an apparent period of lateral migration southward occurred between 8,810–7,420 cal yr BP when the oldest date of Unit IIa occurs. Beginning prior to 7,420 cal yr BP, Unit IIa was deposited until after 6,880 cal yr BP.

Table 3.2 Radiocarbon Data from Nueces River Basin

Locality	Sample#	¹³ C Adjusted 14c yr BP	Dev +/-	Depth (m)	$\delta^{13}C$	Material sampled	Initial 68.2%	Initial 95.4%	IntCal09 68.2%	IntCal09 95.4%
Upper Extent of Nueces River basin										
Dry Frio River Woodrow Heard-Geological Unit I										
Area A; Near Feat 14, Level 10	CAMS-14500	8400	50		-26.1	Indeterminate wood	8430-8430		9480-9310	9510-9280
Area D; Feature 75	Beta-112979	8330	50		-25.8	Indeterminate	8360-8260		9430-9280	9470-9140
Area A; S6/W2, Level 12	CAMS-14501	8250	60		-26.9	Indeterminate wood	8280-8160		9380-9120	9410-9030
Area A; S6/W2, Level 8	CAMS-9057	8030**	70		-26.5	Soto/yucca leaf base	9000-7960		9030-8790	9130-8660
Area A; S6/W0 Level 10	CAMS-14496	8010	50		-25.2	Live Oak wood type	8060-7960		9010-8810	9030-8720
Dry Frio River Woodrow Heard-Geological Unit Iia										
Area D; Feature 72	Beta-112976	6430	40		-25.2	Indeterminate	6480-6380		7420-7310	7430-7270
Area D; Feature 74	Beta-112978	6110	60		-25.7	Indeterminate	6160-6040		7160-6910	7170-6850
Area D; Feature 73	Beta-112977	6090	50		-25.5	Indeterminate	6130-6030		7150-6890	7160-6850
Area D; Feature 71	Beta-112974	6080	50		-25.9	Live Oak wood	6110-6010		7150-6880	7160-6800
Dry Frio River Woodrow Heard-Geological Unit Iib										
Area D; between Features 37 and 73	Beta-112981	5650	50		-22.6	Humate	5740-5640		6480-6320	6530-6300
Area A; near Feature 18	CAMS-9054	4710**	60		*	Juniper wood	4770-4650		5580-5320	5590-5310
Area D; Feature 49	CAMS-9063	4670**	60		-22.1	Soto/yucca leaf base	4730-4610		5470-5310	5590-5300
Area C; S74/W2 Level 9	CAMS-9060	4520**	50		*	Live Oak wood	4570-4470		5310-5080	5440-4990
Dry Frio River Woodrow Heard-Geological Unit Iic										
Area D; 60cm above Feature 37	Beta-112980	3270	50		-23.8	Humate	3340-3240		3570-3440	3630-3390
Middle Extent Nueces River basin										

Table 3.2 Radiocarbon Data from Nueces River Basin

Frio River Choke Canyon Radiocarbon Phase II assays-Horizon 4										
41LK51 N998 E982	TX-4690	6360	90	98.4-98.3	wood	7380-7060		7420-7160		7440-7000
41LK51 N998 E982	TX-4691	5860	80	98.4-98.3	wood	6400		6780-6560		6890-6480
41LK201 N500 E996	TX-4673	2520	70	97.75-97.65	wood	2670-2610		2760-2610		2780-2500
41LK201 N500 E996-997	TX-4672	2710	60	97.85-97.75	wood	2790-2770		2860-2750		2950-2740
Frio River Choke Canyon Radiocarbon Phase II assays-Horizon 3										
41LK201 N491 E1043	TX-4665	2450	60	97.65-97.55	wood	2430		2450-2350		2560-2330
41MC296 N1023 E986-987	TX-4669	2090	70	98.90-98.80	wood	1920-1900		2150-1950		2310-1890
41MC296 N1023 E986-987	TX-4681	1860	60	98.90-98.80	wood	1790		1870-1720		1930-1620
41MC296 N1023 E986-987	TX-4679	2020	70	99.00-98.90	wood	1920-1900		2060-1880		2160-1820
41MC296 N1023 E986-987	TX-4680	1440	80	99.00-98.80	wood	1360		1520-1300		1550-1260
Frio River Choke Canyon Radiocarbon Phase II assays-Horizon 2										
41MC29 N997 E1013	TX-4688	2320	70	99.30-99.20	wood	2360				
41MC29 N997 E1013	TX-4689	2060	80	99.30-99.20	wood	2010-1940				
41MC296 N1022-1024 E985-987	TX-4687	1110	60	99.20-99.10	wood	1040		1040-930		1150-790
41MC296 N1022-1024 E985-987	TX-4685	780	60	99.30-99.20	wood	750-730		770-670		910-650
41MC296 N1025-1027 E997-999	TX-4686	750	70	99.30-99.20	wood	720		770-650		910-560
Frio River Choke Canyon Radiocarbon Phase II assays-Horizon 1										
41MC296 N1024 E989	TX-4684	320	50	99.50-99.40	wood	440-360		460-340		490-310
41MC296 N1025-1027 E997-999	TX-4683	290	50	99.50-99.40	wood	430-340		460-350		480-300
41MC296 N1022-1023 E985-987-989	TX-4682	450	60	99.40-99.30	wood	520		490-330		520-310
41MC296 N1023-1024 E985-987	TX-4677	430	80	Surf-99.40	wood	520		480-330		510-310
41MC296 N1024 E985	TX-4678	330	60	Surf-99.40	wood	430		470-330		490-310

Table 3.2 Radiocarbon Data from Nueces River Basin

41LK201 N497-499-500 E997-998	TX-4667	360	50	99.15-99.05		wood	480-450	470-330	500-310
41LK201 N507-508 E1013	TX4668	320	60	99.20-99.10		wood	440-360	460-340	490-310
Lower Extent Nueces River basin									
Ricklis and Blum 1997 Shell midden Sea Level Dataset									
41NU281	Beta-80018	6600	60			oyster	7525-7390	7560-7430	7580-7420
41SP153	TX-7024	6550	120			scallop	7509-7299	7570-7320	7660-7250
41NU266	Beta-80015	6500	60			oyster	7395-7290	7470-7320	7560-7280
41NU266	Beta-53647	6430	50			oyster	7381-7237	7420-7320	7430-7260
41NU266	Beta-80009	6390	60			oyster	7371-7216	7420-7260	7430-7170
41NU266	Beta-80014	6380	50			oyster	7360-7215	7420-7260	7430-7170
41NU266	Beta-80007	6270	70			oyster	7220-7035	7280-7020	7420-6980
41SP153	TX-7082	6210	90			oyster	7189-7010	7250-7000	7320-6880
41SP153	Beta-57043	6110	80			oyster	7167-6857	7160-6890	7250-6750
41SP136	TX-7302	6070	110			oyster	7159-6798	7160-6780	7250-6670
41NU266	Beta-53073	5980	70			oyster	6888-6737	6910-6730	7000-6660
41NU281	Beta-80017	5050	50			Rangia flex.	5895-5730	5900-5740	5910-5660
41SP153	TX-7308	4990	70			oyster	5888-5650	5890-5640	5900-5600
41JK24	Beta-57912	4940	70			oyster	5736-5638	5740-5600	5900-5580
41NU266	Beta-53072	4900	70			oyster	5716-5589	5720-5580	5720-5580
41NU221	TX-5264	5030	90			Rangia flex.	5919-5654	5900-5660	5890-5470
41NU221	TX-5263	4850	90			Rangia flex.	5724-5474	5710-5710	5860-5320
41NU221	TX-5265	4810	90			Rangia flex.	5647-5336	5650-5330	5730-5320
41NU184	TX-5303	4790	70			Rangia flex.	5633-5336	5600-5330	5650-5320
41SP156	TX-7081	4750	90			oyster	5592-5325	5590-5320	5660-5300
41NU281	Beta-80019	4750	50			Rangia flex.	5581-5332	5590-5330	5590-5320
41SP153	TX-7083	4500	70			oyster	5298-4991	5300-5040	5440-4880
41SP15	TX-6963	4430	70			scallop	5257-4875	5280-4870	5290-4860
41NU266	Beta-53198	4430	70			Human bone	5243-4873	5280-4870	5290-4860
41SP148	TX-7307	4080	270			Charcoal	4986-4229	4960-4150	5440-3860
41NU266	Beta-80006	4380	60			Rangia flex.	5036-4863	5040-4860	5280-4840
41NU267	Beta-80013	4370	50			oyster	4986-4862	5030-4860	5270-4830
41SP153	TX-7309	4080	270			Charcoal	4962-4229	4960-4150	5440-3860
41NU268	Beta-80009	4260	70			Rangia flex.	4865-4636	4960-4640	5040-4570

Table 3.2 Radiocarbon Data from Nueces River Basin

41SP156	TX-6881	4210	70			scallop	4859-4614	4850-4620	4950-4520
41SP1534	TX-7310	4190	90			scallop	4861-4568	4850-4580	4960-4440
41SP120	TX-7311	3970	80			quahog	4533-4353	4530-4290	4810-4150
41SP177	Beta-47105	2840	60			oyster	3156-2873	3070-2860	3160-2790
41NU266	Beta-80005	2990	60			oyster	3258-3069	3320-3070	3350-2990
41CL2	UGA-6152	2611	89			oyster	2837-2611	2850-2510	2920-2360
41AS3	TX-5664	2610	60			quahog	2773-2740	2840-2540	2860-2490
41CL3	Beta-77687	2610	60			oyster	2764-2727	2840-2540	2860-2490
41NU267	Beta-80012	2580	50			oyster	2752-2716	2770-2540	2790-2480
41JK35	Beta-57915	2479	70			Rangia cun.	2736-2357	2720-2460	2730-2360
41CL3	Beta-77686	2450	60			oyster	2713-2355	2700-2360	2720-2350
41SP136	TX-7303	2230	60			oyster	2340-2155	2330-2150	2350-2060
41CL3	Beta-77685	2160	60			oyster	2303-2051	2310-2060	2330-2000
41CL74	UGA-6151	2006	86			Rangia cun.	2060-1873	2110-1870	2300-1730
41NU268	Beta-80008	1910	60			Rangia cun.	1891-1752	1930-1740	2300-1730
41CL2	UGA-6151***	1756	88			oyster	1818-1559	1810-1560	1890-1410
41CL3	Beta-77684	1810	60			oyster	1816-1626	1830-1630	1710-1410
41SP153	TX-7084	1760	70			oyster	1816-1749	1780-1560	1870-1520
41JK24	Beta-57911	1720	60			oyster	1812-1551	1700-1550	1820-1510
41SP43	TX-5891	1850	70			quahog	1873-1711	1880-1710	1950-1600
41SP43	TX-5893	1660	70			quahog	1689-1515	1700-1420	1730-1390
41SP43	TX-6062	1659	60			quahog	1682-1514	1690-1420	1710-1410
41NU266	Beta-80016	1670	50			Rangia cun.	1611-1522	1690-1520	1710-1410
41SP43	TX-5892	1580	70			scallop	1546-1395	1540-1390	1690-1320
41CL3	Beta-77683	1640	60			oyster	1569-1418	1610-1410	1700-1400
41SP149	TX-7304	1440	70			Rangia cun.	1407-1296	1400-1290	1520-1190
41SP120	TX-7313	1370	60			scallop	1338-1270	1350-1180	1390-1170
41SP120	TX-7312	1180	70			whelk	1217-974	1180-980	1270-960
41SP43	TX-523	1220	50			scallop	1235-1067	1240-1070	1280-1010
41SP43	TX-521	1220	50			scallop	1235-1067	1240-1070	1280-1010
41SP43	TX-522	1110	40			scallop	1062-970	1060-970	1170-930
41SP120	TX-6924	1160	50			oyster	1161-996	1170-980	1240-960
41SP120	TX-6926	1030	70			quahog	990-919	1060-800	1170-770
41SP120	TX-6919	1020	70			scallop	982-915	1060-790	1080-740
41SP120	TX-6925	980	70			whelk	960-793	960-790	1060-730

Table 3.2 Radiocarbon Data from Nueces River Basin

41SP120	TX-7305	910	60		scallop	926-741	920-770	940-690
41SP120	TX-7306	910	70		quahog	928-738	920-760	940-690
41RF21	TX-6127	760	130		Bison bone	790-576	900-560	930-530
41RF21	TX-6125	790	70		Bison bone	768-675	780-660	910-560
*standard value of -25.0 assumed by previous investigators								
**radiocarbon date not listed on original table								
***duplicate sample number in original text								

The Dry Frio River again migrated southward between 6,880–6,480 cal yr BP when the oldest Unit IIb sample is indicated. Of note, this date is derived from a humate sample (Beta-112981) and may not accurately reflect the age of the genesis of Unit IIb (e.g., Grimm et al. 2009; Matthews 1985). Another reason for doubting the accuracy is comparing this date to the other Unit IIb results where the humate sample is distinctly out of sync with the others (Figure 3.4). Regardless, Unit IIb definitely was being deposited by 5,580 cal yr BP and continued until sometime after 5,080 cal yr BP when the drainage again migrated southward. The lateral migration appears to have occurred between 5,080–3,570 cal yr BP when the oldest date for Unit IIc is indicated. Unfortunately, only one sample (Beta-112980) is available for Unit IIc, which is a humate sample and none for the overlying Unit IId.

Comparing the initial Woodrow Heard radiocarbon calibration results to the recalibration of this study demonstrates some significant differences. The most apparent changes are exhibited in the older assays of the site, particularly in Unit I (Figure 3.5). The recalibration results push the ages of Unit I and Unit IIa back about 1,000 years.

To a lesser degree the age of Unit IIb has been pushed back about 750 years older than the initial calibration (Figure 3.5). Although there was a shift in the recalibration of Unit IIc, the magnitude of this shift is marginal.

The observed change from the initial Woodrow Heard calibrations to the current study is expected. Simply put, the majority of recent improvements to the radiocarbon calibration curve are applied to the older end of the curve where calibration data (i.e., dendrochronology) are sparser. Thus, most adjustments of a recalibration will typically




Figure 3.4 Calibration Plot of Woodrow Heard radiocarbon assays; arrow indicates anomalous assay.




Figure 3.5 Calibrated Depositional History Nueces River Basin.

be exhibited in the older assays of a study. Such was the case of the recalibration of the Woodrow Heard radiocarbon assays.

A final observation regarding the recalibration of the Woodrow Heard assays is the use of the MCMC analysis. The implementation of MCMC is most useful when radiocarbon results in stratigraphic context overlap temporally. Due to the sizable chronological gaps between each of the stratigraphically defined Woodrow Heard geological units, the application of MCMC did not measurably refine the recalibration results. Therefore, the MCMC analysis of the Woodrow Heard assays was not informative.

As previously mentioned, the recalibration of Ricklis' dataset was also conducted for this study. These assays were reexamined in order to determine if the two occupational hiatuses identified by Ricklis would be altered using the most recent calibration data. The cultural hiatuses, identified by gaps in the radiocarbon results, were recognized to have occurred between 6,800–5,900 ^{14}C yr BP and 4,200–3,000 ^{14}C yr BP (Ricklis 2004; Ricklis and Blum 1997; Ricklis and Cox 1998). These chronometric data were composed of a combination of oyster, scallop, *Rangia flex.*, Quahog, and wood charcoal from 23 coastal archaeological sites (Ricklis and Blum 1997; Table I). Interestingly, the results of the recalibration of these data are very similar to the initial calibration (Table 3.2). Although there are some slight variations between the two calibrations, there are no appreciable differences. Therefore, the timing of the previously identified occupation hiatuses and corollary rapid sea level transgressions still appears to have occurred at 6,800–5,900 ^{14}C yr BP and 4,200–3,000 ^{14}C yr BP.

The results of these recalibrated Nueces River basin study areas are examined further, contrasted with other recalibrated analyses, and correlated with extrinsic factors in Chapter 8.

CHAPTER 4

Recalibrated Geoarchaeological Framework within the Guadalupe-San Antonio River Basins

The Guadalupe and San Antonio River basins encompass a significant portion of South-Central Texas (Figure 4.1). Both basins begin in the Edwards Plateau and extend from the southern margins of the Plateau southeastward across the coastal plain before emptying into the Gulf of Mexico. The roughly 252 mile (405 km) Guadalupe River and 180 mile (290 km) San Antonio River drain a combined 10,250 mile² (26,545 km²) basin area (Donecker 2010; Smyrl 2010). The Edwards Plateau contains a complex system of interrelated aquifers, springs, and rivers. The most prominent of these is the Edwards Aquifer, a large subterranean reservoir underlying the Hill Country in which water filters through porous Lower Cretaceous limestone directly above resistant pre-Cretaceous formations (Edwards Aquifer 2009). As such, this groundwater discharge provides excellent water sources supplying springs, creeks, and rivers. In addition to the Guadalupe River, the most prominent contributors in the basin include the Comal and San Marcos rivers followed by the Blanco River, Coleta Creek, and Sandies Creek. About seven miles north of its terminus at the coastal San Antonio Bay near Tivoli, Texas, the San Antonio River empties into the Guadalupe River.

For the San Antonio River, some of the prominent drainages include the Medina River as well as Cibolo, Leon, Salado, San Pedro, Marcelinas, Culebra, Ecleto, and Medio creeks. Three drainages compose the main trunks of the San Antonio River basin




Figure 4.1 Overview of Guadalupe-San Antonio River Basins: 1) Jonas Terrace site, 2) Richard Beene site, 3) Copano Bay study area, 4) Gatlin site, 5) San Marcos study area, and 6) McNeill Ranch site.

with the other tributaries dendritically spread around and draining into one of the three trunk waterways.

The primary trunk is the San Antonio River that runs roughly down the middle of the basin through Bexar, Wilson, Karnes, Goliad, and Refugio Counties. Within Refugio County the San Antonio River converges and drains into the Guadalupe River, which in turn empties into the San Antonio Bay and then the Gulf of Mexico. The second trunk is the Medina River, a perennial waterway that begins at a spring in the Edwards Plateau county of Bandera where it trends southeastward for about 116 miles along the southern margins of the basin before draining into the San Antonio River. The third main waterway is Cibolo Creek, a seasonal drainage, which begins at a spring in Kendall County and runs south-southeast roughly 96 miles along the northern margins of the basin prior to converging with the San Antonio River in Wilson County.

Previous Investigations

Within the Guadalupe River Basin, the earliest notable investigations with a geoarchaeological component is at Berger Bluff (41GD30) occurring in the 1970s (Brown 2006). Within the San Antonio River basin, the earliest noteworthy geoarchaeological investigation occurred in the 1980s (Table 4.1). Since that time about a dozen significant geoarchaeological investigations have been carried out in the Guadalupe and San Antonio River Basins. The following review of the Guadalupe and San Antonio River basins encompasses some of the more prominent investigations associated with geomorphic examinations, beginning in the upper reaches of each basin and continuing downstream to the gulf.

Table 4.1 Previous Geoarchaeological Investigations in the Guadalupe-San Antonio River Basins

Basin Extent	Drainage	Resource	Guadalupe River Basin		Project-Site(s)	Geoarchaeologist or Researcher	Chronometric Data
			Resource	Project-Site(s)			
Lower	Coletto Creek	Brown 2006		Berger Bluff Site (41GD30) dissertation	K. Brown	14c; relative	
Upper	Upland	Toomey 1993; Toomey et al. 1993		Hall's cave; articles and dissertation	Rickard Toomey	14c	
Middle	San Marcos River	Ringstaff 2000		41HY165 Thesis	C. Ringstaff	limited 14c; relative	
Middle	Blanco-San Marcos Rivers	Schroeder and Oksanen 2002		Armstrong Site (41CW54)	E. Schroeder	14c; relative	
Upper	Upland	Cooke et al. 2003; Cooke 2005, Cooke et al. 2007		Hall's cave Edwards Plateau articles and dissertation	Mary Cooke	Relative, AMS	
Lower	Guadalupe River	Aluvalasit 2006, 2007		McNeill-Gonzales Site (41VT141) thesis and article	M. Aluvalasit	OSL; Relative	
Upper	Guadalupe River	Houk et al. 2008		Gatlin site (41KR621) CRM report	J. Abbott and C. Frederick	14c; relative	
Middle	San Marcos River	Oksanen 2008		Icehouse site (41HY161) thesis	E. Oksanen	14c (AMS); relative	
Middle	San Marcos River	Nickels and Bousman 2010		San Marcos Springs site (41HY160) CRM report	L. Nordt and B. Bousman	14c (AMS); relative	
San Antonio-Medina River Basin							
Middle	Salado Creek	Black and McGraw 1985		Panther Springs site (41BX228) CRM report	Glen L. Evans	14c; relative	
Lower	Aransas River; Copano Bay	Paine 1991; Prewitt and Paine 1987		Swan Lake site (41AS16) dissertation and article	Jeffery Paine	14c (mostly shell); relative; temporally diagnostic artifacts	
Upper	South Fork San Geronimo Creek	Johnson 1995; Johnson and Goode 1994		Jonas Terrace Site (41ME29) CRM report	Leroy Johnson	14c; relative	
Middle	Leon Creek	Tennis and Hard 1995		Archaeological Survey Upper Leon Creek Terraces (41BX47); CRM report	L. Nordt	Relative	
Middle	Leon Creek	Tennis 1996		Upper Leon Creek Terraces (41BX47) CRM report	L. Nordt	uncorrected 14c; relative	

Table 4.1 Previous Geoarchaeological Investigations in the Guadalupe-San Antonio River Basins

Middle	Medio Creek	Houk and Nickels 1997	Lackland Airforce Base (41BX1070, 41BX1076, 41BX1088, 41BX1090, 41BX1091, 41BX1102, 41BX1103, and 41BX1114) CRM report	L. Nordt	14c; relative
Middle	Salado and Panther Springs Creeks	Black, Karbula, Frederick, and Mear 1998	Wurzbach Parkway Number-6 Site (41BX947, 41BX948, 41BX996, 41BX1062, and 41BX1603) CRM report	C. Frederick and C. Mear	14c; relative
Middle	San Antonio River	Houk, Miller, Meadows, and Ringstaff 1999	Brackenridge Park 41BX323 CRM report	L. Nordt	
Middle	Cibolo Creek	Hudler 2000	Dissertation	D. Hudler	no radiocarbon
Middle	Culebra Creek	Nickels, Bousman, Leach, and Cargill 2001	Culebra Creek Site (41BX126) CRM report	L. Nordt and C. Frederick	14c; relative
Middle	San Antonio River	Thoms 2001	41BX1239 CRM report	S. Christopher Caran, David Kuehn, and Rolfe Mandel	Relative
Middle	Medina River	Nordt, Boutton, Jacob, and Mandel 2002	Richard Beene article (41BX831)	L. Nordt, T. Boutton, J. Jacob, and R. Mandel	14c; relative
Middle	Salado and Panther Springs Creeks	Weston 2003	Walker Ranch 41BX1271	Russell Greaves	Relative
Middle	Leon Creek	Collins, Hudler, and Black 2003	Pavo Real (41BX52) CRM report	Charles Johnson and Michael Collins	14c; relative
Middle	San Antonio River	Osburn and Kuehn 2006	Blue Wings SAWS 41BX1623 CRM report	David Kuehn	Relative
Middle	Medina River	Thoms and Mandel 2007	Richard Beene Report (41BX831)	R. Mandel, J. Jacob, and L. Nordt	14c
Middle	San Antonio River	Osburn, Frederick, and Ward 2007	Missions Reach 41BX254, 41BX256, 41BX1621, and 41BX1628 CRM report	C. Frederick	14c; relative
Middle	San Antonio River	Lawrence, Carpenter, Bousman, Miller, Bement, and Lowe 2007	41BX1239 CRM report	B. Bousman	14c; Relative

indicates study selected for calibration

* TL=thermoluminescence, OSL=optically stimulated luminescence, relative (stratigraphy-temporally diagnostic artifacts)

Guadalupe River Basin

In the 2000s, archaeological excavations were conducted by SWCA at the Gatlin site (41KR621) located in the upper extent of the Guadalupe River in eastern Kerr County (Houk et al. 2008). This stratified archaeological site contained cultural activities from the Archaic to Late Prehistoric. One aspect of the site investigations of particular relevance was to characterize the Guadalupe River deposits at different locations (Abbott 2008; Frederick 2008). Specifically, the Gatlin site's location in the Edwards Plateau was contrasted with the deposition of the river off the Edwards Plateau along the margins of the Balcones Escarpment (Abbott 2008).

Although not situated in a drainage setting, Hall's Cave is significant to this study because of the extensive paleoenvironmental investigations and chronometric analyses. Hall's Cave is located at the northern extent of the Guadalupe Basin and the deposits within the Central Texas cave have been a subject of several informative studies (Cooke et al. 2003; Cooke 2005; Cooke et al. 2007; Toomey 1993).

In the late 1990s through the 2000s, a series of investigations were conducted at several archaeological sites (e.g., 41HY160, 41HY161 and 41HY165) along the San Marcos River in San Marcos, Texas. These investigations were conducted by Texas State University Archaeological Field Schools and the Center for Archaeological Studies (CAS) associated with the Aquarena Center at the confluence of Sink Creek and the San Marcos River (Nickels and Bousman 2010; Oksanen 2008; Ringstaff 2000). A component of these investigations included a geoarchaeological examination of the sites with some chronometric analyses. The research identified stratified deposits extending back to the Late Pleistocene and over 11,000 years of cultural activity.

Further downstream the San Marcos River near its confluence with the Blanco River investigations were conducted in the late 1990s at the Armstrong Site (41CW54). The Armstrong Site is situated on a relict channel of the San Marcos River and was investigated by Paul Price and Associates (Schroeder and Oksanen 2002). Work at this stratified site included a geomorphic assessment of the site's stratigraphy coupled with chronometric analyses.

At the lower extent of the Guadalupe Basin in Victoria County, investigations were conducted at the McNeill-Gonzales site (41VT141) in the early 2000s. This site is located in the Coastal Plains and situated on a terrace of the Guadalupe River with deposits extending into the Late Pleistocene (Aiuvalasit 2006, 2007). A primary component of the research consisted of a geoarchaeological analysis of the site to characterize and date the site and drainage stratigraphy.

In 1979, multi-year archaeological investigations began at the Berger Bluff site (41GD30) by the Center for Archaeological Research (CAR). This work was associated with the construction of the Coletto Creek Reservoir. This site is about 9.5 miles (15.3 km) west of Victoria and situated on a high bluff overlooking Coletto Creek a tributary of the Guadalupe River (Figure 4.1). These investigations focused on the geomorphology of the Coletto Creek valley, which included a robust chronometric sampling strategy in order to characterize the geochronology of the site and drainage (Brown 2006).

San Antonio River Basin

In the mid 1980s and early 1990s, archaeological excavations were conducted at the Jonas Terrace site (41ME29) in northeastern Medina County (Johnson 1995; Johnson and Goode 1994). The site is situated on the South Fork of the San Geronimo Creek, a

tributary of the Medina River and contains deposits dating back to the early Holocene. One of the many research avenues undertaken at the site was to date the stratigraphy of the terrace and site as well as reexamine the climates of the Holocene of the eastern Edwards Plateau. Of particular relevance, the researchers compiled a dataset of chronometric data from previous Central Texas investigations and calibrated them to a comparable scale.

In 1991 and 1995, archaeological investigations were carried out for the Applewhite Reservoir project located on the Medina River, a tributary of the San Antonio River (Thoms and Mandel 2007). This project included the excavation at the Richard Beene Site (41BX831), which is located southwest of San Antonio situated on an alluvial terrace (Applewhite Terrace) of the Medina River (Figure 4.1). A significant component of these investigations focused on the geomorphology of the Medina River valley, including a robust chronometric sampling strategy to characterize the geochronology of the site and drainage (Mandel et al. 2007). As such, these investigations at this unprecedented site containing over 7 m of Pleistocene-Holocene alluvial deposits were the first serious and most extensive consideration of Late Quaternary depositional history in the San Antonio River basin.

In the late 1980s, investigations were conducted along the coast at the Swan Site (41AS16) and the Copano Bay area (Paine 1991; Prewitt and Paine 1987). The site is located on the adjacent Aransas River, technically outside of the San Antonio River basin, but the implications of the research are germane to this examination. Of particular relevance, Paine (1991) examined the valley fill near the site to determine changes in sea level and climate over the last 100,000 years. Although this research primarily used

offshore bore samples for radiocarbon analyses, several assays were conducted from the Swan Site coupled with temporally diagnostic artifacts.

Geomorphic/Alluvial History

The depositional history of the San Antonio River basin is composed of a representative selection of several of the aforementioned investigations (Figure 4.2). These selections each have a reasonably extensive, stratified dataset that embodies the various extents (i.e., upper, middle, and lower) of the basins. These interpreted stratigraphic records are arranged geographically beginning at the upper (northern) extent of the basin upstream of the Balcones Escarpment. The middle extent includes the drainage sections from the Balcones Escarpment to the Gulf Prairies and Marshes region. The lower extent of the basin covers the Gulf Prairies and Marshes to the Gulf of Mexico (see Figure 1.2). The selections include the Jonas Terrace site (41ME29) for the upper basin, Richard Beene site (41BX831) for the middle, and the Copano Bay area with the Swan Lake site (41AS16) for the lower extent (Figure 4.2).

Upper Extent San Antonio River Basin

The research at Jonas Terrace was not the most geomorphically focused, but did comprehensively evaluate the site's stratigraphy, enabling a depositional comparison. One focus of research regarded levels of phosphate by horizon with the inference that low amounts of phosphate were implied with fast aggradation while high phosphate quantities inferred slow aggradation (Johnson 1995:29–30). Similarly, the presence and quantities




Figure 4.2 Initial Depositional History San Antonio River Basin.

of various snail species (e.g., *Rabdotus mooreanus* and *Oligyra orbiculata*) were interpreted to correspond with various environments. The researchers identified six strata (i.e., Strata 1–6) at the site all containing cultural materials (Johnson 1995:30–32). The following stratigraphic description was primarily derived from the researcher's description of Unit 23 (Figure 4.3).

The lowermost of the site deposits is Stratum 6, which rested unconformably on bedrock and characterized as alluvial sediments with fluctuating amounts of phosphate and a dominant presence of *Oligyra orbiculata* (Johnson 1995:30–32). Based on these data, Johnson (1995:30–31) interpreted Stratum 6 as having slow aggradation with vegetation cover that was deposited prior to ~5,280 cal yr BP (3330 BC) and ending sometime after ~4,770 cal yr BP (2820 BC) with an erosive event. The overlying Stratum 5 is composed of a thin horizon of colluvial deposits exhibiting low phosphate amounts and an increase in *Rabdotus mooreanus* (i.e., Prairie Rabdotus) snail species. The researchers infer a relatively quick aggradation with a more open vegetation cover for Stratum 5. This horizon has one radiocarbon date indicating an age of 4,400–4,230 cal yr BP (2450–2280 BC). The deposition of the overlying Stratum 4 occurred sometime prior to ~3,460–3,260 cal yr BP (1510–1310 BC) and continued at least until ~2,350 cal yr BP (400 BC). The *terminus post quem* for Stratum 4 is tentative in that the researchers recovered an assay (Beta-62339) from atop a burned rock midden upon the surface of Stratum 4, which dates to a more recent ~1,170 cal yr BP (AD 780). In contrast, the previously mentioned date of ~2,350 yr BP (400 BC) is derived from within Stratum 4 and is seemingly more reflective of the horizon's *terminus post quem* (Figure 4.3). Regardless, Stratum 4 is characterized as containing high phosphate amounts with a




Figure 4.3 Idealized Profile of Jonas Terrace site (41ME29), strata are numbered along right side of profile while sample numbers (1–12) are within profile (adapted from Johnson and Goode 1994: Figure 3).

continued dominance of *Rabdotus mooreanus* and a notable abundance of cultural materials (Johnson 1995:31). These data imply a slower aggradation for Stratum 4 with a continuation of more open vegetation cover.

No chronometric data are available for Stratum 3; this horizon is described as exhibiting a decrease in phosphates, a replacement of *Rabdotus mooreanus* with *Oligyra orbiculata* snail species, and a noticeable decrease in cultural materials. Johnson (1995:31) interprets these data as reflecting a return of tree cover and the fast aggradation of Stratum 3.

The overlying alluvial horizon Stratum 2 began aggrading prior to 1,870 cal BP (AD 80), which notably precedes the aforementioned troublesome assay (Beta-62339) of Stratum 4. Stratum 2 is described as similar to that of the underlying Stratum 3 with low amounts of phosphate and cultural materials, but with a general drop in snail quantities (Johnson 1995:31). This horizon also may have had a relatively fast aggradation that appears to have ended sometime after 1,280 cal BP (AD 670).

The surface layer Stratum 1 is characterized as a partially disturbed horizon of colluvial-alluvial deposits with a slight increase in *Rabdotus mooreanus* and cultural materials. Stratum 1 has one radiocarbon date indicating an age of 1,060–920 cal yr BP (AD 890–1030).

The researchers propose that the environment during the time of Stratum 6 was cooler and moister than present until the approximate terminus of the horizon. The climate steadily became more arid and warmer (i.e., xeric) until peaking around 3,850 cal yr BP (1900 BC) sometime prior to the deposition of Stratum 4. From this apex, the climate became cooler and moister reaching comparable levels of today around the

beginning of Stratum 3 (Johnson 1995). It bears repeating that the Stratum 4 /Stratum 3 boundary is tentative considering Stratum 3 has no chronometric data while the *terminus post quem* for Stratum 4 is the problematic assay (Beta-62339). Nevertheless, the environment is interpreted to continue to cool reaching a relative nadir around 1,950 cal yr BP (AD 0) and returning to conditions similar to today near the terminus of Stratum 2. During the deposition of Stratum 1, the climate is interpreted to have become increasingly warmer and arid reaching a peak around 750 cal yr BP (AD 1200) before becoming cooler and more mesic.

Middle Extent San Antonio River Basin

One of the most intensive geoarchaeological investigations was that conducted at the Richard Beene site in southern Bexar County (Figure 4.1). The site is situated on the right bank of the Medina River and is located about 1.2 miles (2 km) upstream from the drainage's confluence with Leon Creek. The researchers identified five terrace landforms within the Medina River valley at this location that from oldest to youngest consist of the Walsh Terrace (T₄), the Leona Terrace (T₃), the Applewhite Terrace (T₂), the Miller Terrace (T₁), and the modern floodplain (Mandel et al. 2007). Similarly, seven depositional units (Units A1–A7) were recognized primarily related to the Applewhite Terrace (T₂), which contains the Richard Beene site and was the focus of the investigations (Figure 4.4).

Mandel and others (2007:35) interpreted Unit A1 as a coarse-grained depositional unit underlying the Applewhite Terrace, which has an unknown beginning but ceased aggrading before 33,000 ¹⁴C yr BP, when Unit A2 is dated. The age of Unit A2 is based upon chronometric data indicating a beginning around 33,000 ¹⁴C yr BP and continuing

until 20,000 ^{14}C yr BP. Next, Unit A3 extends from the Late Pleistocene into the early Holocene starting around 20,000 ^{14}C yr BP and terminated deposition about 8,600 ^{14}C yr BP. Near the upper boundary of Unit A3, deposits are characterized as cumulic in which soil pedogenesis (Perez Paleosol) formed while alluvium was gradually added. This paleosol in Unit A3 contains the first evidence of cultural activities at the site, which are interpreted to reflect Early Archaic (Angostura) occupations. Also, the upper boundary of the Perez Paleosol exhibits truncation suggesting a discontinuous surface and erosion. Subsequent to this period of erosion, Unit A4 begins deposition containing another paleosol (Elm Creek paleosol). Unit A4 and the Elm Creek Paleosol within contains a few artifacts and extends from 8,600–7,000 ^{14}C yr BP before terminating. The Elm Creek Paleosol is capped with the depositional horizon Unit A5 (Medina Horizon).

The researchers indicate that Unit A5 received the most intensive stratigraphic analyses at the site and contains cultural materials from the Early and Middle Archaic occupations (Mandel et al. 2007:50–52). Among other observations, this unit is noted to have an increase in sand deposits that continued into the following Unit A6 (Leon Creek Horizon) and that may represent an increase in fluvial energy during this time. Further, the upper boundary of Unit A5 contains at least two buried soils that have welded together and have been designated the Medina Pedocomplex. The numerous chronometric assays within the Medina Pedocomplex date the span of Unit A5 to 7,000–4,400 ^{14}C yr BP before being overlain by Unit A6.

The following stratigraphic Unit A6 is noted to have a buried soil (Leon Creek Paleosol), which exhibits evidence of two types of development (Mandel et al. 2007:52–53). Specifically, pedogenesis seems to have started during an extended period of surface





Figure 4.4 Cross-section at Richard Beene site on Medina River (adapted from Thoms and Mandel 2007: Figure 3.3).

stability in Unit A6 and again when aggradation renewed and cumulative development occurred. Although not overtly stated by the researchers, the period of stability may indicate a temporary change in environment during the interpreted span (i.e., 4,100–2,800 ^{14}C yr BP) of Unit A6.

The final stratigraphic unit studied on the Applewhite Terrace is Unit A7. This depositional unit is characterized as encompassing 2,800 ^{14}C yr BP to the present and exhibits evidence of a decrease in fluvial deposits, particularly from 1,200–400 ^{14}C yr BP. The researchers notably correlate the paucity of Late Prehistoric occupation features at the Richard Beene site to the slower deposition in Unit A7 whereby, they argue, created a palimpsest.

Lower Extent San Antonio-Nueces Coastal Basin

For the lower extent of the San Antonio Basin, research in the Copano Bay area was selected for review. Of note, the Swan site (41AS16) is located on the Aransas River, which is situated outside of the San Antonio River basin bounded on the opposite side by the Nueces River Basin (Figure 4.1). Regardless, the site and most importantly the geomorphic investigations in the Copano Bay area are adjacent to the San Antonio Basin and relevant to this study (Figure 4.5). Specifically, Paine (1991) used a variety of datasets (e.g., sea cores, trench profiles, archaeological investigations) to examine sea levels influenced by changes in the climate. Extending back over the last 100,000 years, the research dated these changes using previous research, new radiocarbon data, and temporally diagnostic artifacts (Paine 1991; Prewitt and Paine 1987).

Focusing on the Late Pleistocene-Holocene, Paine (1991) recognized two phases over the last 18,000 years. Broadly defined, the period from 18,000 to 5,000 years ago

interpreted to be a transgressive phase of sea level rise followed by a relative sea level stillstand encompassing the last 5,000 years Paine (1991:57). Within these phases there are 'pulses' that represent transgressive sequences that alter from a dominance of fluvial, deltaic, or estuarine deposits (Paine 1991:60–61). During the Holocene, Paine (1991:61–64) recognizes three transgressive pulses of rising sea level occurring at 10,000–9,000 years ago, 7,500–6,000 years ago, and 5,000–4,000 years ago. Of note, mean sea levels (MSL) at these times were below modern levels as much as 27.5 m.

The first transgressive pulse (10,000–9,000 years ago) is interpreted to be a transition from fluvial (i.e., stream) to marine (i.e., sea) influenced deposition signifying a rapid rise in sea level. Paine (1991:61–64) indicates this pulse is followed by a transition from marine back to stream deposition at roughly 9,000–7,000 years ago suggesting a stillstand or possible drop in sea level. Subsequently, the second transgressive pulse represented by a transition from stream to marine deposition occurred around 7,500–6,000 years ago signifying another rise in sea level. The second transgressive pulse is followed by a transition from marine to stream deposition at roughly 6,000–5,000 years ago suggesting another stillstand. Around 5,000–4,000 years ago, the third transgressive pulse is suggested by a transition from stream to marine deposition interpreted to be a slow sea level rise. Interestingly, Paine (1991:170–171) interprets that sea levels at this time rose above present day levels by as much as 0.9 m beginning as early as 5,300 years ago and lasting until roughly 2,600 years ago. This third transgressive pulse ushers in a sea level stillstand, which covers the last 5,000 years.




Figure 4.5 Profile of Swan Lake deposits (adapted from Paine 1991: Figure 42).

Calibration Results

The radiocarbon datasets for the previously reviewed Jonas Terrace site (41ME29), Richard Beene site (41BX831), and the Copano Bay area (e.g., 41AS16) were recalibrated (Table 4.2 and Figure 4.6). Beginning at the Jonas Terrace site on San Geronimo Creek of the San Antonio basin, the results are presented from this point in the upper limits of the basin followed by the Richard Beene results downstream and finally the chronometric data at Copano Bay area at the coast.

Upper Extent

Thirteen radiocarbon assays were selected from the Jonas Terrace assemblage, all derived from charcoal (Johnson 1995:Table 1). Generally, the recalibration of the assays did not dramatically alter the initial results (Johnson 1995). The most beneficial result of the current study was to increase the precision of the previous results. Beginning with Stratum 6 the oldest identified horizon, the recalibration indicates that deposition began sometime prior to 5,140 cal yr BP and continued subsequent to 4,680 cal yr BP. Sometime prior to 4,380 cal yr BP, the deposition of Stratum 6 ended and the overlying Stratum 5 began.

Stratum 5 is represented by one radiocarbon assay (i.e., Beta-62347) that suggests a terminated around 4,180 cal yr BP. In contrast, the overlying Stratum 4 is dated by six radiocarbon assays indicating deposition began prior to 3,420 cal yr BP and termination after 2,380 cal yr BP. As previously discussed, the *terminus post quem* for Stratum 4 is uncertain in that one of the six assays appears to be anomalous (Beta-62339), which is particularly apparent when examining the calibration plot for this dataset (Figure 4.7).

Table 4.2 Radiocarbon Data San Antonio-Medina River Basins

Locality	Sample#	¹³ c Adjusted ¹⁴ c yr BP	Dev +/-	Depth (m)	δ ¹³ c	Material sampled	Initial 68.2%	Initial 95.4%	IntCal09 68.2%	IntCal09 95.4%
Upper Extent San Antonio River Basin										
San Geronimo Creek (Jonas Terrace) Straturn 6-Johnson 1995 Table 1										
Date 1; Straturn 6	Beta-62340	4480	80		-25.8	Charcoal, buried stump	5280-5100, 5090-4990		5140-4890	5280-4850
Date 2; Straturn 6, Burned rock midden 25I	Beta-62343/CAM S-6503	4370	90		-26.9	Charcoal; AMS	5050-4840		5040-4860	5240-4740
Date 3; Straturn 6, Burned rock midden area 25F	Beta-62341/CAM S-6501	4180	60		-27	Charcoal; AMS	4770-4610		4840-4680	4870-4590
San Geronimo Creek (Jonas Terrace) Straturn 5-Johnson 1995 Table 1										
Date 4; Straturn 5, Burned rock midden area 25D	Beta-62347/CAM S-6506	3870	60		-27.7	Charcoal; AMS	4400-4230		4380-4180	4430-4100
San Geronimo Creek (Jonas Terrace) Straturn 4-Johnson 1995 Table 1										
Date 5; Straturn 4, Soil Column III, Burned rock midden area	Beta-62348/CAM S-6507	3140	80	0.4-0.5	-26.8	Charcoal; AMS, live oak wood	3460-3260		3420-3210	3520-3080
Date 6; Straturn 4, Soil Column IV, Burned rock midden area	Beta-62349/CAM S-6508	2600	70	0.3-0.4	-26	Charcoal; AMS, live oak wood	2780-2710, 2590-2500		2780-2540	2850-2430
Date 7; Straturn 4, South Block 3F	Beta-62338	2570	60		-25.8	Charcoal; AMS	2760-2710, 2630-2500		2740-2520	2780-2410
Date 8; Straturn 4, South Block 24H	Beta-62342/CAM S-6502	2400	70		-26.3	Charcoal; AMS	2710-2640, 2490-2340		2650-2370	2710-2310
Date 9; Straturn 4, South Block 23G	Beta-62346/CAM S-6505	2420	60		-28.3	Charcoal; AMS	2700-2640, 2490-2350		2660-2380	2710-2350
Date 10; Straturn 4, Soil Column IV, Burned rock midden area	Beta-62339/ETH-10478	1295	55	0-0.1	-26.5	Charcoal; AMS	1280-1170		1480-1330	1520-1320
San Geronimo Creek (Jonas Terrace) Straturn 2-Johnson 1995 Table 1										
Date 11; Straturn 2, Feat. 10, North Block 18C or 18D	Beta-11250	1830	110		-26.7	Charcoal	1870-1610		1380-1300	1450-1290

Table 4.2 Radiocarbon Data San Antonio-Medina River Basins

Date	Beta-AMS	1430	60	-26.4	Charcoal; AMS	1360-1280	1360-1300	1410-1270
Date 12; Stratum 2, North Block 27C	Beta-26345/CAM S-6504	1430	60	-26.4	Charcoal; AMS	1360-1280	1360-1300	1410-1270
San Geronimo Creek (Jonas Terrace) Stratum 1-Johnson 1995 Table 1								
Date 13; Stratum 1, Feat. 1, South Block 87	Beta-75905/CAM S-15805	1060	70	-26.3	Charcoal	1100-920	1060-920	1170-810
Middle Extent of San Antonio River Basin								
Medina River-Richard Beene-Somerset Horizon Unit A2								
BHT 54; Somerset paleosol?	Beta-47528	32850	530	n/a	charcoal burned surface; AMS		37970-36650	38740-36450
Medina River-Richard Beene-Perez Horizon Unit A3								
Block S BHT 39; Upper Soil 7	Beta-47526	12745	190	n/a	charcoal, Feature 95; AMS		14270-15290-15900	15890-14120
Block N; Perez paleosol	Beta-80974	8810	60	-24	charcoal		10070-9730-9620	10150-9620
Block T; Perez paleosol	Beta-47527	8805	75	-25	charcoal isolated frags; AMS		10070-9710-10150	10160-9590
Block T; Perez paleosol	Beta-80687	8640	60	-26	charcoal, Feature 106		9700-9550-9890	9850-9530
Medina River-Richard Beene-Elm Creek Horizon Unit A4								
Block K; Elm ck paleosol?	Beta-44386	8080	130	-26	charcoal, Feature 64?		8600-8920-8630-9400	9120-8540
Block M; Elm ck paleosol?	Beta-78656	7910	60	-25.5	charcoal, Feature 80		8590-8890-8890	8960-8560
Block M; Elm ck paleosol?	Beta-78657	7740	50	-25.4	charcoal, Feature 80		8410-8590-8590	8610-8420
Block O; Elm ck paleosol?	Beta-47529	7645	70	n/a	charcoal, Feature 109; AMS		8220-8550-8410-8590	8590-8370
Medina River-Richard Beene-Lower Medina Horizon Unit A5								
Block G	Beta-47530	7000	70	n/a	charcoal, Feature 43		7680-7940-7940	7950-7680
Block G	Beta-47523	6985	65	n/a	charcoal, tree burn		7590-7920-7920	7940-7680
Block G	Beta-47525	6930	65	n/a	charcoal, Feature 44		7620-7930-7930	7920-7640
Block G	Beta-47524	6900	70	n/a	charcoal, AMS Feature 30		7590-7920-7920	7910-7610

Table 4.2 Radiocarbon Data San Antonio-Medina River Basins

Block G	AA-20400	6700	110	6.37	n/a	charcoal, AMS	7340-7790	7660-7480	7780-7400
Medina River-Richard Beene-Upper Medina Horizon Unit A5									
Lower Block A; top of Medina pedocomplex?	Beta-38700	4570	70	2.6	-26.3	charcoal, tree burn	4980-5470	5400-5090	5470-5000
Block U; Medina pedocomplex?	AA-20402	4510	110	3.59	n/a	charcoal	4860-5470	5330-5030	5500-4900
Block U; Medina pedocomplex?	GX-21746	4430	55	3.43	n/a	charcoal	4870-5290	5240-4970	5290-4880
Block U; Medina pedocomplex?	AA-20401	4380	100	3.43	n/a	isolated charcoal, AMS	4660-5310	5250-4940	5310-4860
Medina River- Richard Beene Leon Creek horizon Unit A6									
Block A; near base of Unit A6	Beta-43330	4135	70	2.5	-24.5	charcoal, tree burn	4450-4830	4720-4510	4810-4430
Leon ck paleosol	Beta-36702	3090	70	1.3-4	n/a	charcoal, Feature 1	3080-3470	3390-3220	3460-3100
Lower Extent									
Paine 1991; Table 5 Copano Bay-Egery Island MSA-1									
MSA-1:217, Egery Island	TX-6061	3950	100	2.17		Humate; floodplain	4529-4279	4500-4180	4700-4020
MSA-1:157, Egery Island	TX-6060	2950	60	1.57		humate; clay dune	3215-3004	3230-3020	3320-2950
MSA-1:133, Egery Island	TX-6059	2380	70	1.33		humate; clay dune	2701-2343	2670-2400	2710-2340
MSA-1:82, Egery Island	TX-6058	2250	70	0.82		humate; clay dune	2345-2152	2320-2150	2360-2060
Paine 1991; Table 5 Copano Bay-Egery Island MSA-5 Location									
MSA-5:33, Egery Island	TX-6103	4630	60	0.33		Chione shell	5451-5295	5466-5300	5581-5064
MSA-5:33, Egery Island	TX-6102	4580	90	0.33		mixed shell	5300-5055	5450-5050	5580-4970
*standard value of -25.0 assumed by previous investigators									
**radiocarbon date not listed on original table									
***duplicate sample number in original text									




Figure 4.6 Calibrated Depositional History San Antonio River Basin.




Figure 4.7 Calibration Plot of Jonas Terrace site (41ME29) radiocarbon assays.

Although the sample is derived from charcoal, its provenance is questionable (i.e., atop a burned rock midden upon the surface of Stratum 4). Consequently, for the purposes of this study the sample Beta-62339 has been rejected.

The next chronometric data is within Stratum 2 exhibits a beginning prior to 1,380 cal yr BP and termination subsequent to 1,300 cal yr BP. These data infer an approximate 1,000 year gap between Stratum 4 and Stratum 2, which includes the undated Stratum 3. The final horizon Stratum 1 has one radiocarbon assay that indicates a beginning prior to 1,060 cal yr BP and continues to the present.

A comparison of the initial and current calibrations does not exhibit any striking differences. The most apparent distinction regards the terminus of Stratum 4, but this is a result of rejecting sample Beta-62339 rather than adjustments from a more recent calibration curve. The omission of this sample effectively broadens the possible temporal range for Stratum 3 whereas the initial results implied a very brief Stratum 3. Similarly, the use of the MCMC analyses on this dataset did not markedly refine the results. However, it was instrumental in pointing out the anomalous radiocarbon sample.

Considering these recalibration results, Johnson's (1995) interpretations stand without any notable adjustments, particularly the environmental reconstruction. One exception may regard interpretations associated with Stratum 3. Specifically, the horizon was interpreted to be concurrent with a return of arboreal cover and rapid alluvial deposition, presumably a short-lived depositional horizon. If correct, the adjustment in timing to Stratum 3 in combination with a return in tree pollen correlates with a previously identified spike in arboreal pollen in Central Texas. Bousman (1998:212) interpreted a jump in arboreal canopy to have occurred around 2,000 ¹⁴C yr BP. Similar

to the low arboreal canopy interpreted for Strata 4 and 2 at Jonas Terrace, Bousman (1998:Figure 7) identified that the spike in arboreal canopy was preceded and followed by low arboreal pollen counts. Further, a comparison of the environment associated with each of the Jonas Terrace strata appear to correlate with that proposed by Bousman (1998:Figure 7). The sole exception is the spike of arboreal pollen around 3,500 ^{14}C yr BP, which is not identified at Jonas Terrace. However, this exception seemingly falls within the gap between Strata 5 and 4 (Figure 4.7). Although not conclusive, the data seems to reflect that the proposed environments are comparable particularly when considering the recalibration data.

Middle Extent

Nineteen radiocarbon assays were selected from the Richard Beene assemblage all derived from charcoal (Mandel et al. 2007: Table 3.4). The calibration of select radiocarbon samples from Richard Beene proved beneficial and, to varying degrees, the results generally pushed back the age of the previously reported assays.

Beginning with Unit A3 (Perez Horizon) at the Richard Beene site, the calibration revealed that this horizon was constructed prior to 15,290 cal yr BP and ended subsequent to 9,550 cal yr BP. Using other lines of evidence (e.g., soil carbon) in conjunction with charcoal, Mandel and others (2007:35–48) indicates that Unit A3 possibly began forming around 20,000 ^{14}C yr BP and ceased aggrading about 8,600 ^{14}C yr BP (Mandel et al. 2007:39–46). Further, this depositional unit contained cultural materials interpreted to represent Early Archaic occupations. The presence of the Perez Paleosol, which caps this horizon suggests that this depositional unit ended with a period of stability and seemingly followed by a period of erosion as evidenced by the

disconformity separating it from the overlying Unit A4 (Mandel et al. 2007:Figure 3.10). This erosive event terminated prior to 8,920 cal yr BP as suggested by the oldest assay in Unit A4.

Unit A4 (Elm Creek Horizon) appears to have been relatively brief ending construction sometime after 8,410 cal yr BP, but also has a paleosol (Elm Creek Paleosol) suggestive of a period of prolonged stability (Mandel et al. 2007:48–49). Prior to 7,910 cal yr BP, Unit A5 (Medina Horizon) began deposition capping the underlying Unit A3 (Table 4.2). As previously mentioned, Mandel and others (2007:51) noted several buried soils welded together within this depositional unit identified as the Medina Pedocomplex. Four radiocarbon assays date the Medina Pedocomplex (i.e., Upper Medina Horizon) collected from the top of the horizon while five assays date the lower portions of the horizon (i.e., Lower Medina). The Medina Pedocomplex assays indicate that pedogenesis likely began around 5,400 cal yr BP continuing until sometime after 4,940 cal yr BP as indicated by the latest Unit A5 assay. Although the Medina Horizon extends from 7,910–4,940 cal yr BP, there is an apparent hiatus of about 2,000 years separating the Lower and Upper Medina portions (Figure 4.8). However, this gap may be attributed to differing sample elevations; the assays were collected from about 4 m vertical difference between the upper and lower sample sets.

Subsequent to Unit A5, the overlying Unit A6 (Leon Creek) began aggrading prior to 4,720 cal yr BP (Table 4.2). The Unit A6 Leon Creek Horizon extends from 4,720–3,220 cal yr BP and is capped by the (Leon Creek Paleosol), which again suggests a period of stability prior to the deposition of the Unit A7 (Modern Horizon) mantle. No chronometric assays are available for the final unit the Modern Horizon only Late




Figure 4.8 Select Calibration Plots of the Richard Beene site radiocarbon assays; arrow indicates hiatus period between Lower and Upper Medina Horizon.

Prehistoric artifacts. Thus, the researchers interpreted Unit A7 to extend from the end of Unit A6 to the present (Mandel et al. 2007:53).

Overall, several interesting findings were determined by contrasting the results of the initial calibration with those of the current recalibration study. First, the differences between the results of the two calibrations were not as pronounced as is typical (Table 4.5). The recalibration of the Richard Beene radiocarbon data trended very close to the initial results and occasionally skewed younger. The most evident adjustments involve the Unit A4 Elm Creek Horizon, which the calibration has shortened by roughly 700 years. To a lesser extent, the *terminus post quem* for both Unit A5 and Unit A6 horizons were identified to have occurred earlier than the initial calibration. The majority of these refinements are attributed to the implementation of the MCMC analysis of the recalibration results. The performance of the MCMC application appears to have been beneficial in that the largest refinements to the Richard Beene results were due to this statistical analysis.

Lower Extent

There are several issues with the radiocarbon dataset for the Copano Bay area. One issue regards the provenience information for the samples, only four of the ten radiocarbon assays could be placed into a stratigraphic context (Figure 4.5). Further, none of the assemblage is charcoal, but rather composed of soil humate and shell (Paine 1991:Table 5). With these limitations in mind, the samples were recalibrated due to the implications of their results. Specifically, Paine (1991:170–171) had interpreted these samples to represent distinct depositional events particularly in relation to the previously mentioned third transgressive pulse when sea level rose above present day levels by as

much as 0.9 m. Therefore, the radiocarbon assays from the Copano Bay area were selectively chosen to determine how the recalibration of the assays would alter the initial interpretation.

Six radiocarbon assays were selected from the Copano Bay area assemblage all from Egery Island (Table 4.3). Four of the samples were in a stratified context (MSA-1) while the remaining two are indicated by Paine (1991:134) to date the sea level highstand. Of note, only the four stratified radiocarbon assays were subjected to the phase model and Bayesian analyses of OxCal (Figure 4.4). The remaining two ‘unstratified’ samples (i.e., TX-6102 and TX-6103) were each calibrated individually.

The results of the two ‘unstratified’ samples exhibit some parity. They both are from shell samples that indicate the third transgressive highstand began sometime prior to 5,466 cal yr BP and continued beyond 5,050 cal yr BP (Table 4.2). In contrast, the oldest stratified assay (TX-6061) is from a buried soil and is indicated to date to 4,500–4,180 cal yr BP. This horizon is associated with alluvial floodplain deposits, which Paine (1991:134) implies represents a drop in sea level (i.e., regression) had begun. A disconformity separates this horizon with the overlying clay horizon. This incision or erosional event occurred sometime between 4,180–3,230 cal yr BP indicated by sample (TX-6060) that dates a period of clay dune growth. This horizon signifying clay dune growth is one of four strata observed by Paine (1991:Figure 42). Each stratum of clay dune growth is separated by a disconformity with the fourth stratum composing the modern surface. The three periods of clay dune growth are calibrated to be 3,230–3,020 cal yr BP, 2,670–2,400 cal yr BP, and 2,320–2,150 cal yr BP.

CHAPTER 5

Recalibrated Geoarchaeological Framework within the Colorado River Basin of Texas

The Colorado River is the largest drainage contained entirely within Texas, which extends about 600 miles (965 km) and throughout its course drops in elevation about 3,400 feet from its headwaters in Dawson County to its terminus at the Gulf of Mexico at Matagorda, Texas south of Bay City (Comer and Kleiner 2010). The drainage trends almost exclusively southeast as it winds through the Southern High Plains into the Edwards Plateau where it runs through a bedrock confined valley before exiting the Balcones escarpment onto the relatively level Coastal plain (Blum 1992; Blum and Valastro 1994; Comer and Kleiner 2010). The Colorado River basin encompasses about 110, 000 km² (42,475 square miles) with approximately 92 percent of the drainage network portion situated north of the Balcones Escarpment. Using a drainage basin division recognized by Blum (1992:18), the Colorado River basin is divided into two parts consisting of an upper and lower extent demarcated at the Balcones Escarpment (Figure 5.1). Along its course, the Colorado River crosses a diverse assemblage of physiographic settings beginning in the Southern High Plains and drops into rolling prairies of the North Central Plains where it trends east-south eastward before turning southward to wind through series of canyons between the Central Texas Uplift (i.e., Llano Uplift) and the Edwards Plateau. The drainage abruptly emerges out of the Edwards Plateau at the Balcones Escarpment and crosses a narrow band of Blackland




Figure 5.1 Overview of Colorado River Basin: 1) O. H. Ivie Reservoir, 2) San Angelo study area, and 3) Lower Colorado River study area.

Prairie before entering the interior Coastal Plain, and finally the Coastal Prairies and Marshes.

In the Upper Extent, the basin begins in a series of intermittent ephemeral draws that gradually converge to form the Colorado River. The principle tributaries in the Upper Extent include the Pedernales, Llano, San Saba, and Concho Rivers while the primary tributaries of the Lower Extent include Onion, Big Sandy, Sandy, Cedar, Alum, Pin Oak, and Caney Creeks. Similar to the adjacent Guadalupe River, the Colorado River as it exits the steeper Edwards Plateau and enters the relatively flat Coastal Plain, they convert from a smaller bedrock incised channel with low-moderate sinuosity to a larger floodplain with an increase in sinuosity.

Blum (1992:57–59) recognizes three components (i.e., gathering, transport, and deposition) of the basin in order to characterize the hydrology of the Colorado River basin. The gathering component encompasses the entire upper extent of the Colorado River basin and, as the name implies, collects the sediment from the network. The transport component moves the sediment from the Plateau downstream and extends across the interior Coastal Plain from the Balcones Escarpment downstream to roughly Columbus, Texas. Finally, the deposition component deposits the drainage's materials and roughly extends from Columbus, Texas to the coast.

Previous Investigations

The Colorado River basin is second only to the Brazos River basin for previous investigations (Table 5.1). Investigations have been conducted along the Colorado River basin for a little over 100 years, but the predominance of this research has been in the Lower Extent of the basin (Blum 1992; Blum and Valastro 1994). Since the

Table 5.1 Previous Geoaarchaeological Investigations in the Colorado River Basin

Drainage Extent		Drainage	Resource	Project-Site(s)	Geoaarchaeologist or Researcher	Chronometric Data
Upper	Clear Creek	Fiore 1976	Thesis; Burnet County	R. Fiore	Relative	
Upper	Colorado River	Lintz, Treirweiller, Oglesby, O'Neill, Doering, and McFaul 1991	Mitchell County Reservoir	W. Doering and M. McFaul	14c	
Upper	Colorado and Concho Rivers	Blum and Valastro, Jr. 1992	O. H. Ivie Reservoir research article	M. Blum and S. Valastro	uncorrected 14c; relative	
Upper	Colorado and Concho Rivers	Lintz, Treirweiller, Oglesby, Blum, O'Neill, Holloway, Scott-Cummings, and Scurlock 1993	O. H. Ivie Reservoir	M. Blum and C. Lintz	uncorrected 14c; relative	
Upper	North Concho River	Quigg and Peck 1995	Rush Site (41TG346)	J. Quigg and J. Peck	14c	
Upper	Concho River	Quigg, Peck, Lintz, Treece, Frederick, Clem, Ellis, Schubert, and Abbott 1996	41TG307 and 41TG309	C. Frederick and J. Abbott	14c; relative	
Upper	Middle Concho River	Mauldin and Nickels 2001; Nordt and Bousman 2001	Twin Buttes Reservoir Project (41TG378, 41TG379, and 41TG410)	L. Nordt and B. Bousman	14c; relative	
Upper	Pedernales River	Blum 1987	Thesis	M. Blum	14c data, relative	
Upper	Pedernales River	Blum and Valastro, Jr. 1989	Annals of American Geographers article	M. Blum	14c data, relative	
Middle	Pedernales River	Blum, Toomey III, and Valastro, Jr. 1994	Article focused on paleoclimate change reflected in Texas drainages (Concho River, Pedernales River, Sabinel River, Cowhouse Creek, and Town Creek)	M. Blum, R. Toomey III, and S. Valastro Jr.	uncorrected 14c; relative	

Table 5.1 Previous Geoarchaeological Investigations in the Colorado River Basin

Lower	Barton Creek, Onion Creek, Shoal Creek, and Bull Creek	Urbanec 1963		Austin area-Thesis	D. Urbanec	Relative			
Lower	Colorado River	Weber 1968		Thesis	G. Weber	Relative			
Lower	Colorado River	Tinkler 1971			K. Tinkler	Relative?			
Lower	Colorado River	Wallis 1976		Thesis	J. C. Wallis	Relative			
Lower	Colorado River	Mandel 1980		Thesis	R. Mandel	No 14c data; relative			
Lower	Colorado River	Caran and Baker 1986			S. C. Caran and V. Baker	Relative?			
Lower	Colorado River	Frederick 1987		Austin Mastodon Site-Thesis	C.Frederick	14c; relative			
Lower	Middle Onion Creek	Ricklis and Collins 1994		Barton Site 41HY202 and Mustang Branch 41HY209	J. Abbott and M. Collins	14c; relative			
Lower	Colorado River	Carpenter, Chavez, Miller, and Lawrence 2006		41BP627 McKinney Roughs Site	C.Crawford and C. Frederick	14c; relative			
Lower	Colorado River	Baker and Penteado-Orellana 1977			V. Baker and M. Penteado-Orellana	Relative			
Lower	Colorado River	Looney 1977		Bastrop and Fayette Counties-Thesis	R. Looney	Relative			
Lower	Pin Oak Creek	Largent 1987		Pin Oak Creek site (41FY53) thesis	F. Largent Jr.				
Lower	Colorado River	Baker and Penteado-Orellana 1978			V. Baker and M. Penteado-Orellana	Relative			
Lower	Colorado River	Blum 1992		Dissertation	M. Blum	14c; relative			
Lower	Colorado River	Blum and Valastro, Jr. 1994		GSA article	M. Blum and S. Valastro	14c			
Lower	Colorado River	Miller et al. 1996		MidTexas Pipeline	M. Waters and L. Nordt	Relative			
Lower	Colorado River	Blum and Aslan 2006		article	M. Blum and A. Aslan	Relative			
	indicates study selected for recalibration								

investigations prior to the 1950s lacked chronometric control, the characterization of the deposits used relative dating techniques. Furthermore, only within the last 30 years have there been concerted geomorphological investigations considering the effects of the drainages on the archaeological sites. Numerous investigations have been performed with some facet of geoarchaeology within the last few decades, but a majority of these was either too narrowly focused on site integrity or restricted to a specific time period (e.g., Brownlow 2004, Carpenter et al. 2006, Lintz et al. 1991, Ricklis and Collins 1994, Quigg and Peck 1995). Fortunately, several projects within the basin, associated with the construction of reservoirs, encompass deposits from the Late Pleistocene to the present and were extensively investigated regarding alluvial history of the basin. Beginning in the upper extent, a review of select projects within the basin will be conducted that will proceed downstream to the gulf.

The most comprehensive investigations within the Colorado River basin were conducted for the O. H. Ivie Reservoir at the confluence of the Concho and Colorado Rivers through the 1980s to early 1990s (Lintz et al. 1993). In part, these investigations extensively examined the alluvial deposits of the Colorado and Concho Rivers as well as several tributaries and generated a series of excellent research (e.g., Blum 1989, 1992; Blum et al. 1989, 1994; Blum and Valastro 1989, 1992). Unfortunately, out of the numerous radiocarbon assays from the O. H. Ivie investigations, only a select few (Tx-5770 and Tx-6293) were corrected for isotopic $\delta^{13}\text{C}$ fractionation (Blum and Valastro 1992:428; Winans 2010). Therefore, a majority of the assays could not be included in the current recalibration study. Despite this impediment, Blum and others (1994) consider these dates as a minimum for each of the characterized deposits. Thus, the chronometric

data will not be calibrated, but the interpretations of these researchers will be reviewed in more detail in the subsequent section.

Situated slightly upstream from the O. H. Ivie Reservoir investigations along the Concho River, archaeological research was conducted near San Angelo at sites 41TG307 and 41TG309 in the early 1990s for a wastewater alignment (Quigg et al. 1996). A component of this research was to characterize the alluvial deposits of the drainage (Frederick 1996:85–110). The deposits along the Concho River at these sites encompass the Holocene and were correlated to previous geoarchaeological work at the O. H. Ivie Reservoir. These investigations utilized relative and absolute dating techniques derived from charcoal, humate, and shell. Although sparse, a series of radiocarbon assays associated with the depositional history of the drainage were collected and corrected for isotopic $\delta^{13}\text{C}$ fractionation (Quigg et al. 1996: Table 12.2).

In the late 1980s, Blum (1987) conducted a series of investigations along the Pedernales River, a tributary of the Colorado River, in order to determine the alluvial history of the drainage. These investigations utilized relative dating techniques with a small suite ($n=12$) of radiocarbon assays (Blum 1987:69). However, no information was provided regarding if the assays were calibrated or corrected for isotopic $\delta^{13}\text{C}$ fractionation. Therefore, as with the O. H. Ivie data, the chronometric data for these investigations were not calibrated for this study, but the interpretations of Blum's research will be reviewed in more detail in the subsequent section.

In the lower extent of the Colorado Basin, downstream from the Balcones Escarpment more research relevant to the depositional history of the basin was conducted. Several previous investigators, in particular Blum (1992:81–102) provide an

in depth review of this research. Most significant to the current study was Blum's (1992) research of the lower extent of the basin, which characterized the depositional history and environments of the Colorado River. These investigations utilized relative and absolute dating techniques and, most importantly, were corrected for isotopic $\delta^{13}\text{C}$ fractionation.

Geomorphic/Alluvial History

The following review of the interpreted geomorphic/alluvial history of the Colorado River basin is based upon select summaries of previous research (Figure 5.2). For the upper extent of the basin, the research conducted along the Colorado and Concho Rivers for several reservoir projects and the Pedernales River on the Edward Plateau. The lower extent of the Colorado River basin is a distillation of the abundant investigations of the region and, particularly, Blum's (1992) Colorado River allostratigraphic research.

Upper Extent

This review of the upper extent of the Colorado River Basin is composed of three closely related projects. Two of these investigations were conducted along a portion of the Concho River and its confluence with the larger Colorado River, while the third is on the Pedernales River. The Pedernales River investigations were conducted within the Edwards Plateau while the other investigations occurred at the margins of the Plateau and the Southern High Plains.

Colorado and Concho Rivers

The largest project within the basin was conducted for the O. H. Ivie Reservoir project in Concho, Coleman, and Runnels Counties. The researchers recognized six




Figure 5.2 Initial Depositional History Colorado River Basin.

allostratigraphic units that extend from the Early-Middle Pleistocene until modern deposits (Blum and Lintz 1993; Blum et al. 1994; Blum and Valastro 1992). Regarding the Late Pleistocene to the Holocene, the Late Pleistocene has two terraces that were associated with this time described as roughly 12–14 and 16–18 m above the modern drainage channel (Blum and Valastro 1992:427). The more recent Late Pleistocene terrace exhibits some truncation from erosion and partially capped by eolian deposits. The only two assays (i.e., Tx-5770 and Tx-6293) from the O. H. Ivie dataset to be corrected for isotopic fractionation were derived from this terrace. These dates indicate the terrace was deposited prior to $11,430 \pm 540$ ^{14}C yr BP.

The next allostratigraphic unit is identified as the Early to Middle Holocene alluvium and described as situated roughly 6 m above the modern channel and ranges from 2–9 m thick (Blum and Valastro 1992:431). This stratigraphic unit was chronometrically dated with 25 assays that suggest it was deposited prior to $9,930$ ^{14}C yr BP and continued until the drainage avulsed sometime after $5,000$ ^{14}C yr BP (Blum and Valastro 1992:431). The researchers note that the terrace was stable for an extended period (possibly 3,000 years) allowing for soil pedogenesis until it was capped by the overlying Late Holocene allostratigraphic unit (Blum and Valastro 1992:431). The Late Holocene deposits are characterized as unconformably overlying the Early to Middle Holocene alluvium and are situated up to 5–6 m above the existing channel. Twenty-two radiocarbon assays suggest that this allostratigraphic unit was deposited between $4,600$ – $1,000$ ^{14}C yr BP (Blum and Valastro 1992:431–434). The final allostratigraphic unit is identified by the researchers as modern aligns the channel and chronometrically suggests deposition sometime prior to 840 ± 70 ^{14}C yr BP (Blum and Valastro 1992:436). These

radiocarbon assays were not corrected for isotopic $\delta^{13}\text{C}$ fractionation and were not calibrated for this study.

Concho River

In contrast, the investigations conducted upstream along the Concho River for the San Angelo wastewater alignment were not as extensive as that employed at the O. H. Ivie investigations. However, this research at sites 41TG307 and 41TG309 succinctly characterized the Concho drainage alluvial history. Frederick (1996) recognized through a series of backhoe trenches and cutbank exposures four alluvial deposits, which he interpreted to correlate with the O. H. Ivie data (Figures 5.3a and 5.3b). The oldest deposits were the Late Pleistocene alluvium only observed in a few locations situated beneath eolian or alluvial deposits (Frederick 1996:91–94). The Late Pleistocene deposit is described as sloping with an undulatory tread implying an erosional event subsequent to deposition. Next, the overlying alluvial deposits designated Early-Middle Holocene is characterized as unconformably situated on the Late Pleistocene deposits and having two distinct alluvial fills (Frederick 1996:91–94). Specifically, the Early-Middle Holocene deposits are composed of fine-grained overbank facies and a more coarse-grained pink colored sandy channel facies. The radiocarbon data for the overbank alluvium indicate deposition between 8,300–5,300 ^{14}C yr BP, which suggests the pink colored channel deposits accumulated between 10,000 and 8,300 ^{14}C yr BP (Frederick 1996; Frederick and Boutton 1996).

Situated above the Early-Middle Holocene unit are the Middle-Late Holocene deposits described as a mix of coarse (e.g., gravel and loamy sands) channel facies and fine-grained overbank facies (Frederick 1996:95–97). Based on three radiocarbon assays (i.e., Beta-69766, Beta-72273, and Beta-69770), the interpreted deposition of the Middle-




Figure 5.3a Profile of 41TG307 on Concho River (adapted from Quigg et al. 1996; Figure 5.2).




Figure 5.3b Profile of 41TG309 on Concho River (adapted from Quigg et al. 1996: Figure 5.3).

Late Holocene unit to roughly coincide with that proposed at the O. H. Ivie data.

However, instead of terminating around 1,000 ^{14}C yr BP, the Middle-Late Holocene ends about 1,300 ^{14}C yr BP.

The most recent unit, the Modern alluvium was characterized as overbank, channel, and bar facies that composes the entire T_0 terrace as well as aprons the adjacent T_{1a} tread (Frederick 1996:97–100). Notably, Frederick (1996:99) indicates that the deposits of the modern alluvium situated over the Middle-Late Holocene unit are almost indistinguishable. Yet, a radiocarbon assay (Beta-70134) indicating a *terminus post quem* of 960 ^{14}C yr BP for the Middle-Late Holocene unit was collected (Frederick 1996: Figure 5.2, Table 12.2). This assay derived from humate was not reviewed in the Middle-Late Holocene unit discussion and was presumably rejected as being stratigraphically inconsistent. However, it may be correct and the allostratigraphic unit may actually be the Modern alluvium instead of the Middle-Late Holocene unit due to the difficulty of discerning the two deposits. Regardless, the Modern alluvium was interpreted to have been deposited between 1,200 ^{14}C yr BP to the present (Frederick 1996: 97–100).

Pedernales River

Located further downstream in the Colorado River basin along the Pedernales River, Blum (1987) investigated a series of drainage cutbanks centered around Fredericksburg, Texas (Figure 5.1). Of note, Blum (1987) did not calibrate the radiocarbon assay results processed at the Radiocarbon Laboratory at the University of Texas (Blum 1987:3). Considering the researcher lists the results with the assay's deviation, the assumption is made that results are uncorrected ^{14}C years that are reported. This research observed seven allostratigraphic units (Units A–G), which encompasses

Early Pleistocene to Modern deposits. Most relevant, Unit D is associated with the Late Pleistocene, Unit E with the Early Holocene, Unit F with the Late Holocene, and Unit G with Modern deposits. Unit D is characterized as a distinct terrace positioned 12–13 m above the modern channel composed of a fining upward of clast materials (gravels to clays). The radiocarbon assays for this unit indicated $17,260 \pm 230$ ^{14}C yr BP and a very dubious $5,200 \pm 340$ ^{14}C yr BP.

Next, the Early Holocene Unit E is described as variable with coarse, weakly cemented gravels-gravelly sands in some exposures and a predominance of finer-grained sediments capped with a weakly developed paleosol (Blum 1987: Figure 34). Based upon select radiocarbon assays and Pleistocene faunal remains (*Equus sp.*), the Unit E alluvium is interpreted to have been deposited between 11,000–7,000 ^{14}C yr BP (Blum 1987:88). Deposited unconformably upon Unit E is the Late Holocene Unit F that is characterized as possibly composed of up to 10 m of gravels and sand capped by a weakly developed soil (Blum 1987:91–93). The sparse chronometric data for this unit suggest deposition occurred at roughly 5,000–800 ^{14}C yr BP. The modern deposits, Unit G, are described as laterally confined with roughly 5 m of sand and gravels. Blum (1987:96) notes that a radiocarbon assay (Tx-5532) was collected from the base of Unit G indicating this allostratigraphic unit was deposited between 900 ^{14}C yr BP and the present. Overall, the depositional history outlined for the Pedernales River correlates with that proposed for the Colorado and Concho Rivers (O. H. Ivie and San Angelo) in the Upper Extent of the basin.

Summary Upper Extent

The researchers interpret the alluvial history of this section of the Colorado River basin to begin with a period of channel aggradation during the Late Pleistocene roughly

coinciding with the Last Glacial Maximum. This period of floodplain construction was followed by a drainage avulsion and an extended phase of erosion around 14,000 ^{14}C yr BP, which deeply incised into bedrock. This erosion continued until the development of the Early-Middle Holocene floodplain between 11,000–5,000 ^{14}C yr BP followed by another erosive event concurrent with a period of stability that formed a soil capping the unit. The Middle-Late Holocene allostratigraphic unit began forming around 4,600 ^{14}C yr BP and continued until approximately 1,000 ^{14}C yr BP. This period ended with a drainage avulsion and an extended phase of erosion concurrent with a period of stability that formed a soil capping the unit. The modern phase of floodplain construction began roughly 800 years ago and continues to the present.

Lower Extent

As previously defined, the lower extent of the basin extends from the Balcones Escarpment downstream to the Gulf Coast. Undoubtedly, the most comprehensive investigations within this region were by Blum (1992). These investigations recognized four Members for the entire lower Colorado River basin, which encompassed the Late Pleistocene to modern times (Blum 1992; Blum and Törnqvist 2000; Blum and Valastro 1994). Blum (1992) examined a series of Colorado River profiles between Austin and Wharton, in part, to characterize the depositional history of the lower basin and correlate it with the upper, determine the chronology of these fluvial events, and examine the influence of climatic and eustatic effects on these fluvial events.

Colorado River

The four identified allostratigraphic members include the Eagle Lake alloformation of the Late Pleistocene and the Columbus Bend alloformation members 1–

3 encompassing the Late Pleistocene to Holocene (Figures 5.3a and 5.3b). The Eagle Lake alloformation is characterized as exhibiting varied facies depending upon position along the drainage (Blum 1992:149–165). In the section identified as transport within the basin (i.e., Interior Coastal Plain), the deposits are primarily gravelly clast materials while the section identified as the depositional (i.e., Coastal Prairies and Marshes) exhibits finer-grained clast materials. The Eagle Lake alloformation varies in thickness about 8–10 m with the base situated on bedrock about 6–8 m above the modern channel (Figures 5.4a and 5.4b). This alloformation was dated with a series of radiocarbon assays, which indicate that accumulation began prior to 20,000 ^{14}C yr BP and ended sometime after 14,000 ^{14}C yr BP (Blum 1992: Table 6.1).

The Columbus Bend allomembers 1–3 are three terrace landforms that compose the Columbus Bend alloformation (Blum 1992: Figure 6.15). The roughly 10–12 m thick Late Pleistocene-Early Holocene deposits of the Columbus Bend member 1 rests unconformably on bedrock and against the Eagle Lake alloformation. The Columbus Bend allomember 1 is described as comprising a variety of channel related deposits ranging from gravels to fine sands. Blum (1992:177–178) interprets the deposition of this allostratigraphic unit to have been predominantly attributed to lateral migration as overbank deposits suggested by thick deposits of finer clast materials were rare. The radiocarbon assays for the Columbus Bend member 1 indicate deposition occurred between 13,000–5,000 ^{14}C yr BP (Blum 1992: Table 6.2).

Inset against and overlapping the Columbus Bend member 1 is the Middle-Late Holocene Columbus Bend member 2. These deposits are typically over 12 m thick extending below modern water levels and characterized as having varied channel facies




Figure 5.4a Idealized Cross-section of Colorado River Basin at the Eagle Lake locality (adapted from Blum 1992: Figure 6.5).




Figure 5.4b Idealized Cross-section of Columbus Bend Members 1–3 at the Austin, West Point, and Columbus locations (adapted from Blum 1992: Figure 6.15).

with a predominance of floodplain facies (Blum 1992:186). The Columbus Bend allomember 2 radiocarbon data suggest deposition occurred prior to 5,000 ^{14}C yr BP and continued until sometime after 1,000 ^{14}C yr BP.

Interestingly, Blum (1992:186–190) observes that the Columbus Bend member 2 seemingly had frequent episodes of high magnitude overbank flooding and ended with a period of stability, which formed a soil that capped the unit. Next, the Columbus Bend member 3 is unconformably inset against the Middle-Late Holocene Columbus Bend member 2 indicating some erosion of this older unit. The dramatic avulsion of the Colorado River that began the modern Columbus Bend member 3 is interpreted to have abandoned its initial course, the existing Caney Creek, and moved eastward to its modern course (Blum 1992:190–193). This avulsion occurred near Wharton, Texas with Caney Creek trending southeast containing the older allostratigraphic units and the Colorado River trending south-southwest containing the modern Columbus Bend member 3 both emptying into the gulf about 32 km (20 miles) apart.

The Columbus Bend member 3 is described as ranging from 1–10 m in thickness and collected radiocarbon assays suggest deposition between 600–100 ^{14}C yr BP. Of note, the assays for the Columbus Bend member 3 were the only samples derived from wood while the remaining assays were humate materials (Blum 1992: Tables 6.1–6.4). Finally, Blum (1992:193) only notes cultural deposits within the Columbus Bend member 3 consist only of historic artifacts. The other allostratigraphic units for the lower extent of the basin have no mention of cultural materials as being present.

Summary Lower Extent

The alluvial history proposed for the lower extent of the basin is similar to that indicated for the upper extent. Beginning at the Last Glacial Maximum, the Colorado

River had a period of extensive deposition between 20,000–14,000 ^{14}C yr BP (Blum 1992; Blum and Valastro 1994). This period of deposition was followed by an extended period of incision eroding underlying bedrock throughout the lower basin to its current levels. Beginning around 12,000 ^{14}C yr BP, the formation of the Columbus Bend alloformation units occurred. Columbus Bend member 1 deposition extended from roughly 12,000–5,000 ^{14}C yr BP followed by a reduction in flood magnitude beginning the deposition of Columbus Bend member 2 (Blum 1992; Blum and Valastro 1994: Figure 10). The reduction in flood magnitude allowed for pedogenesis to occur in the Columbus Bend member 1 unit, which continued until approximately 2,500 ^{14}C yr BP.

The Columbus Bend member 2 floodplain accumulated between 5,000–1,000 ^{14}C yr BP with an increase in flow regime occurring after 2,500 ^{14}C yr BP resulting in the burial of Columbus Bend member 1 (Blum 1992; Blum and Valastro 1994). Subsequent to 1,000 ^{14}C yr BP, the flood magnitude decreased abandoning the Columbus Bend member 2 and at the Caney Creek meanderbelt (Blum 1992; Blum and Valastro 1994).

The modern Columbus Bend member 3 accumulated within the last 600 years. Based on the general thickness of the units and the floodplain facies, Blum (1992:193–197) observes that the flow regime for the basin changed over time. Specifically, from the Late Pleistocene up to the Middle Holocene floodplain construction was predominantly lateral migrations while the latter half of the Holocene had a noticeable increase in overbank flooding (i.e., vertical accretion).

Calibration Results

Select assays of radiocarbon datasets for the previously reviewed Upper and Lower Extents of the Colorado River Basin were recalibrated. As previously mentioned,

only two of the radiocarbon assays (Tx-5770 and Tx-6293) from the O. H. Ivie investigations were corrected for isotopic $\delta^{13}\text{C}$ fractionation (Blum and Valastro 1992:428; Winans 2010). Similarly, the assays for Blum's (1987) investigations along the Pedernales River were not calibrated or corrected for isotopic $\delta^{13}\text{C}$ fractionation. Consequently, neither the O. H. Ivie nor Pedernales River datasets were calibrated for this study. Fortunately, a series of radiocarbon assays from the investigations along the Concho River near San Angelo were corrected for isotopic $\delta^{13}\text{C}$ fractionation (Quigg et al. 1996: Table 12.2). Although sparse, these data are used as a proxy for the upper extent of the Colorado River basin. In contrast, the dataset for the lower extent of the Colorado River basin is robust and adequately represents the depositional history of the basin.

Upper Extent

Five radiocarbon assays were selected from Quigg and others (1996: Table 12.2) chronometric assemblage derived from fluvial sediment (i.e., humate) and wood. These assays from charcoal and humus were selected due to their known stratigraphic context and association with three of the identified stratigraphic units (Figures 5.3a and 5.3b). Other assays were available, but rejected as these samples were derived from mussel shell. One assay represents the Early-Middle Holocene, three assays represent the Middle-Late Holocene, and one assay represents the Modern unit (Table 5.2). Notably, one assay (Beta-70134) was not included due to several troubling factors including its stratigraphic inconsistency as initial radiocarbon analysis indicated (Frederick 1996:97–100). Therefore, due to the questionable validity of this sample it was not included in the recalibration.

Table 5.2 Radiocarbon Data Colorado River Basin

Locality	Lab#	13c Adjusted 14c yr BP	Dev +/-	Depth (m)	$\delta^{13}C$	Material sampled	Initial 68.2%	Initial 95.4%	IntCal09 68.2%	IntCal09 95.4%
Eagle Lake Alloformation (Blum 1992: Table 6.1)										
Eagle Lake	Tx-7011***	18600	700	4	-25	fluvial sediment			22000-20290	22980-19610
Eagle Lake	Tx-7012	18380	3890	7.3	-31	fluvial sediment			21400-19170	22990-18070
West Point	Tx-7225	16180	1170	7	-17.9	fluvial sediment			20780-18990	21910-18130
Eagle Lake	Tx-7013	16090	450	3.9	-23.4	fluvial sediment			19960-18990	20440-18660
Eagle Lake	Tx-7010	15970	1700	1.9	-24.8	fluvial sediment			20960-18930	22270-17890
Eagle Lake	Tx-7011***	15900	810	1.9	-24.9	fluvial sediment			20280-18860	21210-18120
Columbus	Tx-7230	15610	1300	7.5	-25.2	fluvial sediment			20660-18790	21850-17810
Columbus Bend Member 1 (Blum 1992: Table 6.2)										
Columbus	Tx-7326	12970	640	10.5	-23.4	fluvial sediment			15820-14030	16720-13470
West Point	Tx-7224	10910	600	2.8	-26.7	fluvial sediment			13430-11940	14420-11220
Utley	Tx-7328	7970	630	5.5	-24.2	fluvial sediment			9800-8300	10770-7740
Columbus	Tx-6811	7730	150	5.2	-17.8	fluvial sediment			8790-8410	8980-8230
Eagle Lake	Tx-7323	5350	180	4.2	-20.7	fluvial sediment			6310-5930	6530-5730
West Point	Tx-7226	4960	70	1.5	-14.6	2Bkb horizon			5840-5630	5890-5590
Columbus	Tx-7325	4640	120	1.4	-15.5	3Bkb horizon			5560-5260	5640-5060
Columbus Bend Member 2 (Blum 1992: Table 6.3)										
Webberville	Tx-7232	4160	70	10.5m	-19.1	fluvial sediment			4760-4550	4830-4450
Eagle Lake	Tx-7008	4120	110	4.3m	-18.9	fluvial sediment			4750-4460	4840-4310
West Point	Tx-7220	4010	100	10.6m	-21.2	fluvial sediment			4650-4320	4800-4180
Webberville	Tx-7234	3640	60	7.5m	-19.7	fluvial sediment			4060-3880	4140-3780
Webberville	Tx-7233	3390	60	10.4m	-25.8	sediment wood			3720-3560	3820-3480
Webberville	Tx-6809	3440	90	10m	-18.4	fluvial sediment			3830-3580	3950-3480
Columbus	Tx-6810	3440	90	10m	-18.4	fluvial sediment			3830-3580	3950-3480

Table 5.2 Radiocarbon Data Colorado River Basin

West Point	Tx-7222	3380	140	1.2m	-18.9	fluvial sediment		3820-3480	4020-3340
Webberville	Tx-7330	3250	90	5.5	-20.6	fluvial sediment		3590-3390	3700-3270
West Point	Tx-7221	2950	60	6	-20.1	fluvial sediment		3220-3010	3320-2930
Eagle Lake	Tx-7007	1870	60	1.1	-20.8	fluvial sediment		1880-1730	1950-1640
Webberville	Tx-7331	1900	60	1.5	-16.1	fluvial sediment		1910-1750	1980-1700
West Point	Tx-7223	1660	60	2	-20	fluvial sediment		1660-1460	1710-1410
Columbus	Tx-6812	1590	70	4.2	-14.8	fluvial sediment		1560-1400	1680-1340
Columbus	Tx-6813	820	70	0.3	-13.7	A horizon		890-720	920-680
Columbus Bend Member 3 (Blum 1992:Table 6.4)									
Columbus	Tx-7334	490	60	8.5	-28.5	wood		540-350	620-320
Columbus	Tx-7335	350	60	8	-26.4	wood		470-320	510-290
West Point	Tx-7321	190	60	6.2	-28.3	wood		290-100	420-0
Columbus	Tx-7227	70	60	5.5	-27.9	wood		260-60	280-0
Concho River Investigations (Quigg et al. 1996:Table 12.2)									
Early-Middle Holocene Horizon									
41TG307	Beta-70133	7320	120	2.9	-20.4	Humus/soil	5370 B.C.	8270-8010	8360-7890
Middle-Late Holocene Horizon									
41TG309	Beta-69769	4450	60	2.4	-18.5	Charcoal	2500 B.C.	5220-4940	5290-4870
41TG307	Beta-69766	3840	60	3	-25.8	Charred material	1890 B.C.	4370-4150	4420-4050
41TG307	Beta-72273	3200	60	1.9	-17.9	Humus/soil	1250 B.C.	3510-3370	3580-3290
Modern Holocene Horizon									
41TG309	Beta-69770	1230	60	4	-25.4	Charcoal	A.D. 720	123-1060	1280-980
41TG307	Beta-70134	920	60	0.5	-18.2	Humus/soil	A.D. 1030	910-780	960-720
*standard value of -25.0 assumed by previous investigators									
**radiocarbon date not listed on original table									
***duplicate sample number in original text									

The results of this recalibration indicate the *terminus ante quem* for the Early-Middle Holocene terrace is 8,270–8,010 cal yr BP (Table 5.2). The overlying Middle-Late Holocene unit suggests deposition began prior to 5,220 cal yr BP indicating the abandonment of the Early-Middle Holocene unit sometime in the intervening 2,790 years. The *terminus post quem* of the Middle-Late Holocene unit (i.e., assay Beta-72273) is 3,370 cal yr BP while the overlying modern unit began sometime prior to 1,230 cal yr BP. Therefore, the abandonment of the Middle-Late Holocene terrace and formation of the modern terrace occurred in the approximate 2,000-year interval.

Despite the obvious limitations of the dataset, the recalibration of the Concho River chronometric data is informative. Specifically, in regards to the Early-Middle Holocene deposits Frederick (1996:94) notes a horizon of pink colored channel deposits likely deposited between 10,000 and 8,300 yr BP. The recalibration of the data conforms with Frederick's (1996) initial interpretation, but may need to be pushed back in age considering the 8,270–8,010 cal yr BP result. Further, the deposition of the Middle-Late Horizon between 5,220–3,370 cal yr BP conforms with the interval of deposition proposed at O. H. Ivie and the Pedernales River. Similarly, the deposition of the modern deposits also concurs with that proposed by Frederick (1996:97–100).

Lower Extent

Nineteen radiocarbon assays were selected from the Blum's (1992) assemblage derived from fluvial sediment (i.e., humate) and wood. Admittedly, only three of the assays were from wood while the remaining 15 assays were from humate materials (Table 5.2). These 19 assays were selected for calibration because their stratigraphic position at their collection location at the Eagle Lake, Columbus, and West Point

localities was identified (Blum 1992: Figures 6.6a, 6.15). More significantly, all of these samples were corrected for isotopic fractionation (Blum 1992: Tables 6.1, 6.2, and 6.3). Four assays are from the Eagle Lake Member, five assays are from the Columbus Bend Member 1, seven are from the Columbus Bend Member 2, and three are from Columbus Bend Member 3.

The results of the recalibration indicate that the Eagle Lake Member began deposition prior to 22,500 cal yr BP and terminated deposition sometime after 19,000 cal yr BP, which is immediately after the last glacial maximum at ~23,500 cal yr BP (Figure 5.5). The overlying Columbus Bend Member 1 has a *terminus ante quem* of 15,940 cal yr BP suggesting an approximate 3,000-year gap between the two members. This gap (19,000–15,940 cal yr BP) is interpreted to be a dramatic period of deep bedrock erosion-incision, which concurs with the disconformity separating the two units (Blum 1992, Blum and Valastro 1994).

The Columbus Bend Member 1 continued deposition until a soil capping these deposits began to develop between 5,550–5,210 cal yr BP suggesting a period of stability. This period of pedogenesis may indicate when the Colorado River decreased flow and began the deposition of the Columbus Bend Member 2. The *terminus ante quem* of the Columbus Bend Member 2 is 4,570 cal yr BP suggesting the floodplain abandonment of Columbus Bend Member 1 and erosion minimally occurred during the 640-year gap between Columbus Bend Members 1 and 2. During this time, the Colorado River is interpreted to have increased in flood magnitude and overtopped the Columbus Bend member 1 terrace and soil. Initially, this was argued to have occurred around 2,500 ¹⁴C yr BP. Two radiocarbon assays (i.e., Tx-6533 and Tx-6534) indicating an age of




Figure 5.5 Calibrated Depositional History Colorado River Basin.

3340 \pm 90 and 3320 \pm 90 ^{14}C yr BP, respectively were collected from the Columbus Bend Member 2 veneer capping the soil (Figure 5.4b).

Unfortunately, these assays were not corrected for isotopic $\delta^{13}\text{C}$ fractionation. However, a radiocarbon assay (Tx-6810) of equivalent uncorrected radiocarbon age (i.e., 3330 \pm 90 ^{14}C yr BP) was corrected for fractionation and calibrated. This calibrated assay used as a proxy dated to 3830–3580 cal yr BP, which could be inferred to suggest that the increase in flood magnitude on the Colorado River occurred prior to this age (Table 5.2).

The Columbus Bend Member 2 continued deposition until roughly 900 cal yr BP when soil pedogenesis occurred between 890–710 cal yr BP. The overlying Columbus Bend Member 3 has a calibrated *terminus ante quem* of 540 cal yr BP suggesting an approximate 170-year gap between it and Columbus Bend Member 2. Again, the avulsion of the Colorado River, which abandoned the present day Caney Creek meander belt, may have taken place prior to the pedogenesis of the Columbus Bend Member 2 and before the deposition of the Columbus Bend Member 3, which continues up to the present.

The recalibration of the Lower Colorado River basin chronometric data when contrasting the initial investigation exhibits some notable trends. Typically, adjustments are limited to the older assays, but the recalibration of Blum's (1992) dataset exhibits shifts throughout all of the allostratigraphic members. These prevalent adjustments between the initial and recalibrated datasets are in no small part attributed to the fact that the initial ^{14}C assays were not calibrated. The most prominent adjustment occurs between the Eagle Lake and Columbus Bend 1 members (Figure 5.5). The Late Pleistocene Eagle Lake Member terminates sometime after 19,000 cal yr BP much older than previous

interpreted. Similarly, the Columbus Bend Member 1 begins prior to 15,940 cal yr BP much earlier than initially proposed (i.e., ~13,000 ^{14}C yr BP). Also, the termination of Columbus Bend Member 1 occurred about 200–300 years earlier and followed by a gap of roughly 650 years before Columbus Bend Member 2 begins deposition.

The abandonment of the Columbus Bend Member 1 floodplain is interpreted to have been followed by a period of erosion, which seemingly occurred between 5,210–4,570 cal yr BP. The termination of Columbus Bend Member 2 has been shifted to end about 300 years more recently around 710 cal yr BP compared to the initial ~1,000 ^{14}C yr BP (Blum 1992, Blum and Valastro 1994). Finally, the beginning of Columbus Bend Member 3 occurred around 500 cal yr BP as opposed to ~1,000 ^{14}C yr BP separated by an approximate 200 year erosion after the abandonment of the Columbus Bend Member 2 floodplain.

Other observations during the recalibration of the lower extent of the Colorado River basin include several chronological gaps within the radiocarbon assays of Columbus Bend Members 1 and 2 (Figure 5.6). Two chronological gaps were observed in the Columbus Bend Member 1 between 11,940–8,790 cal yr BP and 8,410–5,840 cal yr BP and one recognized hiatus in Columbus Bend Member 2 between 3,010–1,660 cal yr BP. These lulls may be attributed to sampling rather than issues of geomorphic processes or preservation. To investigate this possibility an additional suite of radiocarbon assays from Blum's (1992) chronometric dataset were examined. Additional assays were gathered from Columbus Bend Members 1 and 2 regardless of whether their stratigraphic context could be determined and incorporated into the recalibration study. These data suggest that the 11,940–8,790 cal yr BP is the result of sampling as it




Figure 5.6 Select Calibration Plot of Lower Extent of Colorado River Basin; arrows indicate hiatus periods.

disappears with the introduction of the additional assays. However, the new data demonstrate that gaps at 8,410–6,310 cal yr BP and 3,010–1,880 cal yr BP remain. These chronologic lulls have narrowed down a little, but still suggest a hiatus in Columbus Bend Members 1 and 2. Again, the reason for these phenomena is undetermined if they are attributed to depositional processes, sampling, or a combination of these factors. Interestingly, these chronological gaps do correlate with similar lulls in other drainage basins. These temporal hiatuses, apparent correlations and possible causes are examined further and contrasted with other recalibrated analyses in Chapter 8.

CHAPTER 6

Recalibrated Geoarchaeological Framework within the Brazos River Basin of Texas

The Brazos River is the largest drainage within Texas extending about 1,200 miles (2,000 km) from its headwaters at Blackwater Draw in New Mexico to its terminus at the Gulf of Mexico at Freeport, Texas near Galveston (Figure 6.1). The Brazos River basin encompasses about 44,000 square miles (114, 000 km²) and throughout its course drops in elevation about 4,600 feet (Epps 1973; Hendrickson 2010). The contributory network of drainages within the Brazos River is extensive. From upstream to downstream, several of the most significant contributing drainages of this large basin include Yellowhouse Draw, Blackwater Draw, Running Water Draw, Double Mountain Fork, Salt Fork, Clear Fork, Palo Pinto Creek, Bosque River, Leon River (with Henson Creek, Cowhouse Creek—Table Rock Creek—House Creek), Lampasas River, Little River (San Gabriel River—Brushy Creek and Salado Creek—Buttermilk Creek), Navasota River, and Oyster Creek. As a consequence of this vast network, the Brazos River and its tributaries crosses a diversity of physiographic settings between its genesis and conclusion. Trending south and east from its beginning in the High Plains, the basin crosses the Rolling Plains, the Cross Timbers and Prairies, across the alternating Blackland Prairies and Post Oak Savannah regions, and finally the Gulf Prairies and Marshes.

Previous Investigations

Possibly due to its size, the Brazos River basin is the most extensively investigated basin in Texas. As early as 1901, researchers have been evaluating and




Figure 6.1 Overview of Brazos River Basin: 1) Lubbock Lake site, 2) Fort Hood, 3) and A&M study area.

documenting the basin, but the investigations prior to the 1950s lacked chronometric control (Hill 1901: 345—359). Subsequent to the introduction of ^{14}C analyses, relative temporal characterizations of drainage terrace deposits were then supplemented with absolute dating (Table 6.1). To be sure, there have been some substantial geomorphological investigations previously conducted, but a comparatively few of those truly considered the effects of the drainages on the archaeological sites. Within the Brazos River basin, the incorporation of geomorphic examinations into archaeological investigations (i.e., geoarchaeology) began early. These early concerted efforts employing archaeological geology occurred due to Early Man studies particularly at the Lubbock Lake site (41LU1) on a tributary of the Brazos River. Since that time, numerous significant geoarchaeological investigations have been carried out along the Brazos River and its tributaries. Although more investigations have been performed with some facet of geoarchaeology, but most of these were typically general reviews of the immediate site area focused primarily with site integrity or a similarly narrow focus (e.g., Alexander 2008; Gadus et al. 2006; Gibson 1997; Hilliard 1997; Pearl 1997).

The culmination of these previous geomorphic, geoarchaeological, and archaeological investigations are a collection of Late Quaternary stratigraphic history across the Brazos River basin. Several researchers have compiled a comprehensive review of the previous investigations in the upper extent (Holliday 2009, 2000, 1997; Mandel 1992:53–57), the middle extent (Nordt 1993, 1992), and the lower extent (Abbott 2000). Due to the broad geography of their coverage, the varied focus of those investigations and the span of time, only a select few of those research projects were selected for this study.

Table 6.1 Previous Geoarchaeological Investigations in the Brazos River Basin

Drainage Extent	Drainage	Resource(s)	Project-Site(s)	Geoarchaeologist or Researcher	Chronometric Data
Upper		Hill 1901		Robert Hill	Relative
Upper	Brazos River	Stricklin 1961		F. Stricklin	Relative
Upper	Yellowhouse Draw	Holliday 1985, 1988, 1995, 1997, 2000, 2009; Holliday and Johnson 1983, 1986, 1981; Holliday et al. 1983, 1985, 1999; Johnson and Holliday 1980; Stafford 1981, 1983	Lubbock Lake site (41LU1)	V. Holliday, T. Stafford	14c; relative
Upper	Double Mountain Fork of Brazos River	Blum, Abbott, and Valastro 1992	Justiceburg Reservoir	Blum, Abbott, and Valastro	Radiocarbon
Upper	Clear Fork of the Brazos River	Mandel 1992	Southbend Reservoir	Rolfe Mandel	Radiocarbon
Upper		Ferring 1995b	Southern Plains	C. Reid Ferring	Radiocarbon
Middle	Brazos River-McLennan County	Bronaugh 1950	Thesis	R. Bronaugh	Relative
Middle	Brazos River	Epps 1973	Thesis	L. W. Epps	Radiocarbon
Middle	North Bosque River	Brotherton 1978	Thesis	M. Brotherton	Relative
Middle	Brazos River	Nordt 1983	Thesis	L. Nordt	Relative
Middle	Brazos River	Woolly 1985	Thesis	B. Woolly	Relative
Middle	Brazos, Navasota, and Trinity Rivers	Nordt 1986	Article associated with 1983 thesis	L. Nordt	Relative
Middle	Aquilla Creek	Brown 1987	Aquilla Lake	Peter Patton	Relative
Middle	Leon River	Tharp 1988	Thesis	Tommy L. Tharp	Relative
Middle	Navasota River	Fields 1990	Jewett Mine-Charles Cox, Lamps Creek Knoll, and Buffalo Branch Sites	R. Fields, Bousman, et al	Radiocarbon
Middle	Navasota River	Fields, Klement, Bousman, Tomka, Gadus, and Howard 1991	Jewett Mine-Bottoms, Rena Branch, Moccasin Springs Sites	R. Fields, Bousman, et al	Radiocarbon
Middle	Cowhouse, Table Rock, and Henson Creeks	Nordt 1992	Ft. Hood	L. Nordt	Radiocarbon
Middle	Cowhouse, Table Rock, and Henson Creeks	Nordt 1993	Ft. Hood	L. Nordt	Radiocarbon
Middle	Cowhouse Creek	Nordt, Boutton, Hallmark, and Waters 1994	Ft. Hood	Nordt, Boutton, Hallmark, and Waters	Carbon isotopes; 14c
Middle	Henson Creek	Nordt 1995	Ft. Hood-Henson Creek	L. Nordt	Radiocarbon
Middle	Cowhouse Creek	Nordt 1996	Ft. Hood-Dissertation	L. Nordt	Stable C isotopes

Table 6.1 Previous Geoaerchaeological Investigations in the Brazos River Basin

Middle	Brazos River	Hilliard 1997	Waco Mammoth Site-Thesis	K. Lee Hilliard	Relative
Middle	Buttermilk Creek	Gibson 1997	Thesis	B. D. Gibson	Radiocarbon
Middle	Upper Lampasas	Pearl 1997	Thesis	F. Pearl	Radiocarbon
Middle	Cowhouse Creek	Nordt, Boutton, Hallmark, and Waters 1998	Ft. Hood-Pedogenic carbonate accumulations	Nordt, Hallmark, Wilding, and Boutton	Stable C Isotopes
Middle	Brushy Creek	Collins 1998	Wilson-Leonard (41WM235)	Bousman, Goldberg, Stafford, Collins	Radiocarbon and others
Middle	Owl Creek	Hilliard 2000	Ft. Hood GIS study	using L. Nordt data	Radiocarbon
Middle		Mehalchick, Kleinbach, Boyd, and Kibler 2000	Ft. Hood	L. Nordt	Radiocarbon
Middle	Little River	Mahoney and Tomka 2001; Nordt 2001b	41MM340 and 41MM341	L. Nordt	Radiocarbon
Middle	Brazos River	Prochnow 2001	Horn Shelter No. 2-Thesis	S. Prochnow	Radiocarbon
Middle	Brushy Creek	Abbott 2003	Blackland Prairie 41WM815	J. Abbott	Radiocarbon
Middle	Little River	Mahoney, Tomka, Mauldin, Shafer, Nordt, Greaves, and Galdeano 2003	41MM340	L. Nordt	Radiocarbon
Middle	Cowhouse Creek	Nordt 2004	Ft. Hood-Cowhouse Ck	L. Nordt	Radiocarbon
Middle	Cowhouse Creek	Campbell and Johnston 2004	Ft. Hood	using L. Nordt data	n/a
Middle	Brazos River	Gadus, Fields, and Kibler 2006	J. B. White Site (41MM341)	Karl Kibler ?	Radiocarbon
Middle	Brazos River	Bongino 2007	Waco Mammoth Site-Thesis	J. D. Bongino	Radiocarbon, OSL
Middle	Buttermilk Creek	Alexander 2008	Gault Site (41BL323)-Thesis	using L. Nordt data	Radiocarbon
Lower	Brazos River	Bernard, Major, Parrott, and Leblanc 1970	Bureau of Economic Geology	W. A. Bernard, G. F. Major, B. S. Parrott, and R. J. Leblanc, Jr.	Relative
Lower	Brazos-Coastal	Wilkinson and Basse 1978		Wilkinson and Basse	Radiocarbon
Lower	Brazos-Coastal	Aten 1983		L. Aten	Radiocarbon
Lower	Brazos River	Voelinger 1990	Thesis		Relative
Lower	Brazos River	Waters and Nordt 1995		M. Waters and L. Nordt	Radiocarbon
Lower	Brazos River	Miller 1995	MIDTEXAS pipeline	M. Waters and L. Nordt	Relative
Lower	Brazos River	Husain 1998		Syed Raziuddin Husain	Relative
Lower	Trinity-Coast	Abbott 2001	TXDOT Houston Area PALM	J. Abbott	corrected 14C; relative
Lower	Brazos River	Sylvia and Galloway 2006		D. Sylvia and W. Galloway	Thermoluminescence; Radiocarbon
Lower	Brazos River	Uriata 2009	Vernor Mammoth Site-Thesis	Uriata	Relative

indicates study selected for recalibration

Using a modified basin divisions recognized by Epps (1973) and Nordt (1983), the Brazos River Basin is broken into three parts. Largely attributed to the underlying geology and physiography these sections consist of the Upper Extent, the Middle Extent, and the Lower Extent. Roughly outlined, the Upper Extent begins at Yellowhouse Draw (Lubbock Lake) in the High Plains and trends eastward off of the Llano Estacado across the Osage Plains until about the Parker and Hood County line. From this point, the Middle Extent trends south-southeast across the Cross Timbers and Blackland Prairie stopping just south of the margins of the Balcones Escarpment and the confluence of the Little and Brazos Rivers. At this point, the Lower Extent begins to cross the Gulf Prairies and Marshes of the Coastal Plain and runs southeastward until the Brazos River finally empties into the Gulf of Mexico near Freeport, Texas.

Upper Extent

One project conducted in the Upper Extent of the Brazos River Basin reexamined here is the Lubbock Lake investigations (Holliday 1997). In addition to the ^{14}C dating, this research of this project is relevant due to the extensive investigations of the Lubbock Lake site (41LU1) with the intent of characterizing the depositional history of the channels and surrounding landscape. The extensiveness and implications of the previous research at the Lubbock Lake site (41LU1) is not to be understated. The research at this locality is varied and prolific (e.g., Holliday 1985, 1988, 1995, 1997, 2000, 2009; Holliday and Johnson 1983, 1986, 1981; Holliday et al. 1983, 1985, 1999; Johnson and Holliday 1980; Stafford 1981, 1983) and the following review does not intend to supersede previous research. Rather, this review is a compilation of previously identified stratigraphy and an examination of the chronometric analyses (Figure 6.2).




Figure 6.2 Initial Depositional History Brazos River Basin.

The Lubbock Lake site (41LU1) was first discovered in 1936 during excavations in the Yellowhouse Draw House for construction of a reservoir in Lubbock, Texas (Holliday 1997:76). These construction activities encountered evidence of Paleoindian occupations, which instigated investigations over subsequent decades. Various prominent researchers have comprehensively investigated the deposits at Lubbock Lake and along the Yellowhouse Draw drainage with some minor variations in interpretation. Generally, the stratigraphy at Lubbock Lake is characterized as having five primary strata (Strata 1–5) containing various internal horizons, and paleosols (Holliday 1997, 1985; Stafford 1981).

The oldest deposit recognized at Lubbock Lake is Stratum 1, which is described as alluvial deposits and possibly localized lacustrine deposits that contain Pleistocene fauna and Clovis cultural materials (Holliday 1985:1484–1486, 1997:78–83; Stafford 1981). Radiometric data for this stratum suggests it terminated sometime prior to 11,000 ^{14}C yr BP (Holliday 1985:1484). Above the first horizon is the complex Stratum 2 characterized as containing several internal horizons (Strata 2A, 2B, 2e, 2s, and 2F) composed of lacustrine, marsh, eolian, and possibly spring deposits and capped by the Firstview Soil, a paleosol (Figure 6.3). The horizons of Stratum 2 are interpreted to have been deposited roughly between 11,000–6,300 ^{14}C yr BP (Holliday 1985:1486–1487; Stafford 1981:552). The Firstview Soil is indicated to have developed approximately between 8,500–6,300 ^{14}C yr BP (Holliday 1985:1487). Situated above Stratum 2, is Stratum 3 recognized as having two distinct internal horizons with one composed of eolian deposits (3e) and a lacustrine deposit (3l) and capped by a buried soil identified as the Yellowhouse Soil (Holliday




Figure 6.3 Idealized Profile of Lubbock Lake site (adapted from Haas et al. 1986: Figure 4).

1985:1487). Stratum 3 contains cultural materials from the Archaic period and is interpreted to have been deposited about 6,300 ^{14}C yr BP and capped by the overlying horizon about 5,500 ^{14}C yr BP (Holliday 1985:1488).

Unconformably overlying Stratum 3 is Stratum 4 composed of two internal horizons (i.e., Strata 4A and 4B) and capped by the Lubbock Lake Soil (Holliday 1985:1488–1489). The deposits of Strata 4A and 4B are described as spring and eolian deposits, respectively and contain cultural materials extending from the Middle Archaic to Late Prehistoric. Stratum 4 is interpreted to have been deposited between 5,500–4,500 ^{14}C yr BP followed by an extended period of stability represented by the Lubbock Lake Soil.

Subsequent to the extended period of stasis, the deposition of Stratum 5 began around 750 ^{14}C yr BP (Holliday 1985:1489). Stratum 5 is characterized as containing two internal horizons (Strata 5A and 5B) of lacustrine deposition each capped with the Apache and Singer soils, respectively. The Stratum 5 deposits contain Late Prehistoric to Historic cultural materials and continue up to modern times.

The general depositional sequence interpreted from these deposits argues that a period of incision occurred during the Late Pleistocene followed by the deposition of Stratum 1 when the climate was wetter and cooler (Holliday 1985:1489–1492). A decrease in drainage discharge likely attributed to a reduction in moisture ushered in the lacustrine/marsh environment of Stratum 2. Researchers have varied interpretations for the cause of the impoundment of the Yellowhouse Draw at this time. Regardless of whether the static flow is from eolian deposits damming the channel (Holliday 1985, 1997; Holliday and Johnson 1983) or part of a natural pool (Stafford 1981, 1983) the

Yellowhouse Draw at this time was not flowing. The climate is interpreted to have become gradually warmer and drier in part attributed to eolian deposits at the end of Stratum 2 and throughout Strata 3 and 4 (Holliday 1985: Figure 7). The increasing presence of eolian deposits was interpreted to represent a decrease in vegetative cover and stasis to allow the development of the Yellowhouse Soil. A possible drought is interpreted to have occurred followed by a trend toward modern climatic conditions during Stratum 4. Some localized erosion (i.e., unconformity) is noted at the Strata 3 and 4 boundary that may reflect the return of some moisture and modern conditions. The modern climatic conditions are interpreted to have continued throughout Stratum 4 concurrent with the development of the Lubbock Lake Soil. The presence of colluvial slope wash in Stratum 5 may represent swings toward arid environments beginning about 1,000¹⁴C yr BP.

Middle Extent

For a variety of reasons, the Middle Extent of the Brazos River Basin is the most extensively investigated region through geomorphological and geoarchaeological methods in Texas (Table 6.1). One factor is the prevalence of development within this portion of the basin, but most influential is the presence of the Fort Hood military reserve in Hood County. Archaeological and geoarchaeological research has been conducted for almost two decades within this military reserve. Various research within Fort Hood, which encompasses several significant tributaries of the Brazos River, has spawned numerous reports, articles, masters' theses, and doctoral dissertations (Campbell and Johnson 2004; Hilliard 2000; Mehalchick et al. 2000; Nordt 1992, 1993, 1995, 1996, 2004; Nordt et al. 1994; Nordt et al. 1998). This portion of the Brazos River Basin also

contains two of Texas' more prominent prehistoric sites that have some exceptional stratigraphic records and have similarly influenced extensive research (Alexander 2008; Bousman 1998; Collins 1998; Gibson 1997; Goldberg and Holliday 1998). Specifically, Wilson-Leonard (41WM235) on Brushy Creek in Williamson County and the Gault Site (41BL323) on Buttermilk Creek in Bell County. The current research will consider select investigations conducted within Fort Hood on Cowhouse Creek.

Cowhouse Creek (Fort Hood)

Within Fort Hood, the geoarchaeological investigations have focused upon the Henson Creek, North Nolan Creek, Reese Creek, Cowhouse Creek and its tributaries Table Rock, Owl, and House Creeks and the Leon River, which they all eventually intercept. This research over the last two decades has gradually constructed a comprehensive depositional history for the region as well as systematically evaluated a diversity of settings and drainages ranging in size from upland tributaries to their associated lowland trunk channels.

The culmination of these previous investigations of the Fort Hood drainages has identified four late quaternary landforms (designated T₃ to T₀) that contain six allostratigraphic units (Nordt 1992, 1993, 1995, 2004). From oldest to most recent, Nordt (1992, 2004) recognizes the Pleistocene Reserve alluvium only observed on the Leon River, the middle-late Pleistocene Jackson alluvium, the early Holocene Georgetown alluvium, the middle Holocene Fort Hood alluvium, the late Holocene West Range alluvium, and the recent Ford alluvium (Figure 6.4). Further, the West Range unit is occasionally divided into Upper and Lower West Range units interpreted to be separated by an erosional disconformity (Nordt 1992, 2004). These investigations have also documented several buried soils (i.e., paleosols) within the drainages. Within Cowhouse




Figure 6.4 Idealized Cross-section of Cowhouse Creek (adapted from Nordt 1992: Figure 29).

Creek and its tributaries, the Royalty Paleosol is recorded at the top of the Georgetown unit (Nordt 2004). Similarly, Henson Creek contains the Royalty Paleosol and the Tanktrail Paleosol at the top of the Upper West Range unit (Nordt 1995). The chronology of these stratigraphic units is derived from a series of ^{14}C analyses (Nordt 1992: Appendix J).

The oldest allostratigraphic unit identified in the Fort Hood study area is the Jackson alluvium identified resting on Glen Rose limestone and composes the T₂ terrace along the investigated drainages (Nordt 1992, 1993, 1995, 2004). The chronometric data for the Jackson alluvium is provided by a single bulk sediment humate sample roughly dating 15,000 ^{14}C yr BP indicating deposition occurred during the Late Pleistocene (Nordt 2004: Table 1). Nordt (2004:296–297) indicates that a period of incision-erosion occurred before the construction of the second allostratigraphic unit (Georgetown alluvium).

The Georgetown alluvium, which composes the T₁ terrace in the study area, is identified as early Holocene. Eight radiocarbon samples have been collected from this unit, but only two (Beta-63007 and GX-15762) are uncontaminated charcoal (Table 6.2). The remaining samples are from bulk soil humate and have not been used in this study. The two charcoal samples date to about 8,900 ^{14}C yr BP and 8,300 ^{14}C yr BP, respectively. Capping the Georgetown alluvium is the Royalty paleosol (Nordt 2004). Subsequent and possibly concurrent to the development of the Royalty paleosol, a dramatic decrease in hydrologic flow occurred particularly along Cowhouse Creek that partially eroded this paleosol (Nordt 2004:297).

Table 6.2 Radiocarbon Data for Brazos River Basin

Locality	Lab#	13c Adjusted 14c Yr BP	Dev +/- Depth (m)	$\delta^{13}C$	Material sampled	Initial 68.2%	Initial 95.4%	IntCal09 68.2%	IntCal09 95.4%
Upper Extent Brazos River Basin									
Yellowhouse Draw-Lubbock Lake site (41LU1) Stratum 1 (Haas et al. 1986:Table 1; Holliday et al. 1983, 1985)									
1, Area 2	SMU-548	11100	100		wood	9150 BC		13080-12850	13170-12740
1C	SMU-263	11100	80		wood			13080-12850	13170-12740
Yellowhouse Draw-Lubbock Lake site (41LU1) Stratum 2A (Haas et al. 1986:Table 1; Holliday et al. 1983, 1985)									
2ALB2, Area 6	SI-4976	10195	165		hummin	8245 BC		12190-11740	12400-11560
2ALB4, Area 6	SI-4975	9905	140		hummin	7955 BC		11950-11610	12080-11490
2ALB1, Area 2	SMU-285	10530	90		humic acid	8580 BC		12460-12140	12570-12050
2ALB2, Area 6	SI-3200	10360	80		hummin	8410 BC		12330-12040	12450-11850
2ALB2, Area 3	SMU-251	10060	170		humic acid	8110 BC		12070-11670	12340-11520
2A, upper, Tr90	SI-3203	10015	75		hummin, organic mud	8065 BC		11920-11600	11990-11480
Yellowhouse Draw-Lubbock Lake site (41LU1) Stratum 2B (Haas et al. 1986:Table 1; Holliday et al. 1983, 1985)									
2B, base, Area 6	SI-4974	9605	195		hummin	7655 BC		11170-10660	11380-10390
2B, base, Area 6	SMU-828	9870	140		humic acid	7920 BC		11420-11110	11590-10810
2B, base, Area 5	SMU-829	9170	80		humic acid	7220 BC		10470-10260	10560-10210
2B, base, Area 5	SI-4179	9075	100		hummin	7125 BC		10390-10070	10510-9910
2B, base, Area 6	SMU-728	9990	100		humic acid	8040 BC		11490-11270	11620-11220
2B, base, Area 2	SMU-275	9960	80		humic acid	8010 BC		11470-11270	11600-11230
2B, upper, Area 6	SI-4177	8655	90		hummin, organic mud	6705 BC		9820-9560	10080-9500
2B, upper, Area 6	SMU-830	8210	240		humic acid	6260 BC		9420-8800	9760-8560
2B, upper, Area 3	SMU-302	7890	100		humic acid	5940 BC		8920-8600	9000-8470
2B, upper, Area 3	SMU-262	7970	80		humic acid	6020 BC		8960-8680	9030-8600
2B, upper, Tr90	SI-3204	7255	75		hummin	5305 BC		8160-8000	8270-7950
2B, upper, Area 6	SI-4178	6705	95		hummin	7880-7260 BP		7650-7490	7750-7430

Table 6.2 Radiocarbon Data for Brazos River Basin

2B, upper, Tr108	SMU-544	6400	80			humic acid	7555-7030 BP	7420-7270	7480-7180
Yellowhouse Draw-Lubbock Lake site (41LU1) Stratum 3 (Haas et al. 1986:Table 1; Holliday et al. 1983, 1985)									
3I, Tr49	SMU-1093	5220	50			A Horizon, humic acid	4325-3805 BC	6100-5930	6180-5910
3e, Tr108	SMU-545	5770	80			humic acid	6895-6375 BP	6630-6440	6720-6340
3e, Tr108	SMU-531	4900	60			A Horizon, humic acid	5830-5350 BP	5890-5830	5900-5740
Yellowhouse Draw-Lubbock Lake site (41LU1) Stratum 4A (Haas et al. 1986:Table 1; Holliday et al. 1983, 1985)									
4A, Tr116	SMU-1200	5270	150			humic acid	4420-3780 BC	5810-5720	5850-5660
Yellowhouse Draw-Lubbock Lake site (41LU1) Stratum 4B (Haas et al. 1986:Table 1; Holliday et al. 1983, 1985)									
4B, Tr108E	SMU-1191	2070	130			A Horizon, humic acid	4085-625 BP	2240-1900	2340-1740
4B, Tr108E	SMU-1177	1550	50			A Horizon, humic acid	1465-1365 BP, 1515-1415 BP	1510-1380	1550-1340
4B, Area 7	SMU-1090	1270	40			A Horizon, humic acid	1225-1145 BP, 1240-1160 BP, 1275-1195 BP, 1295-1215 BP	1260-1150	1290-1080
4B, Area 16	SMU-492	4960	50			humic acid	5845-5595 BP	5690-5610	5730-5590
4B, Area 16	SI-4171	4700	65			humic acid	5710-5500 BP	5550-5340	5580-5310
4B, Tr59	SI-4174	1955	75			A horizon, humic acid	2015-1835 BP, 1975-1815 BP	2000-1820	2110-1720
4B, Tr59	SI-3201	1215	65			A horizon, humic acid	1255-1115 BP	1230-1060	1280-980
4B, Tr108	SMU-651	890	70			A horizon, humic acid	870-710 BP	900-750	940-700
4B, Area 8	SI-4169	880	70			A horizon, humic acid	870-710 BP, 835- 696 BP	900-750	930-700
4B, Tr108	SMU-534	870	40			A horizon, humic acid	835-735 BP, 810- 710 BP	890-740	910-700
Yellowhouse Draw-Lubbock Lake site (41LU1) Stratum 5A (Haas et al. 1986:Table 1; Holliday et al. 1983, 1985)									
5A, Area 7	SMU-893	450	50	19		charcoal, GA7-5	550-470 BP	530-450	550-340
5A, Area 7	SMU-970	380	50	21		charcoal, GA7-5	525-425 BP	490-360	520-320
5A, Area 7	SMU-968	440	40	20		charcoal, GA7-5	560-460 BP	520-460	540-340
5A, Area 7	SMU-314	720	40	19		A Horizon, humic acid	720-620 BP	650-560	680-550
5A, Area 8	SI-3208	640	75	19		A horizon, humic acid	730-570 BP, 660- 500 BP, 650-490 BP	610-540	660-510
5A, Area 8	SI-2701	505	55	20		charcoal	590-470 BP	560-500	630-450

Table 6.2 Radiocarbon Data for Brazos River Basin

5A, Area 8	SI-2700	380	40	20	charcoal	525-425 BP	490-360	510-320	
5A, Area 19	SMU-546	320	60	21	charcoal	440-320 BP, 395-255 BP	470-350	500-360	
5A, Area 14	SI-2704	315	50	21	charcoal	475-375 BP, 440-340 BP, 385-265 BP	460-350	500-300	
5A, Area 8	SMU-345	300	60	21	charcoal	475-355 BP, 470-350 BP, 390-250 BP	460-350	500-300	
5A, Area 15	SI-2703	285	60	21	charcoal	390-250 BP	460-350	500-290	
5A, Tr108	SMU-555	220	50	22	A Horizon, humic acid	355-235 BP	430-300	470-270	
5A, Area 7	I-208	160	90	23	bone, bison				
Yellowhouse Draw-Lubbock Lake site (41LU1) Stratum 5B (Haas et al. 1986; Table 1; Holliday et al. 1983, 1985)									
5B, Area 7	SMU-343	160	60		A Horizon, humic acid		230-30	300 - -10	
Middle Extent Brazos River Basin									
Ft Hood- Stratigraphic Unit A - Jackson Alluvium (Nordt 2004: Table 1)									
Cowhouse ck; CB10	Beta-38694	15270	260	2.9	Bulk sediment-humate		18680-18080	18960-17790	
Ft Hood- Stratigraphic Unit B- Georgetown Alluvium (Nordt 2004: Table 1)									
Cowhouse ck; CB2	Beta-63007	8830	70	6.71	Charcoal-hearth		10100-9750	10170-9630	
Ft Hood-Stratigraphic Unit C- Ft Hood Alluvium (Nordt 2004: Table 1)									
Cowhouse ck; CB6	Beta-37618	6850	90	6.25	Charcoal-dispersed	-19.3	7780-7600	7890-7520	
Cowhouse Ck; CB15	Beta-46192	5820	110	6.84	Bulk sediment-humate-hearth		6770-6500	6910-6390	
Cowhouse Ck; CB15	GX-15892	5740	300	3.79	Charcoal-midden	-16.3	6960-6290	7280-5990	
Cowhouse Ck; CB12	Beta-37452	5210	230	1.62	Charcoal-hearth	-16.5	6350-5860	6620-5640	
Ft Hood-Stratigraphic Unit D1 - (Lower) West Range Alluvium (Nordt 2004: Table 1)									
Cowhouse Ck; CB9	TX-6705	4170	100	6	Charcoal-dispersed	-18.5	4790-4520	4890-4420	
Cowhouse Ck; CB13a	TX-6704	3950	290	7.05	Charcoal-dispersed	-18.5	4680-3980	4980-3650	
Cowhouse Ck; TR11	TX-6703	3010	110	2.3	Charcoal-hearth		3330-3030	3440-2900	
Cowhouse Ck; CB13a	Beta-38173	2860	50	1.79	Charcoal-dispersed		3070-2910	3170-2860	

Table 6.2 Radiocarbon Data for Brazos River Basin

Cowhouse Ck; CB17	Beta-37451	2720	110	1.47		Charcoal- hearth		3040-2790	3200-2700
Ft Hood-Stratigraphic Unit D2- (Upper) West Range Alluvium (Nordt 2004: Table 1)									
Cowhouse Ck; CB1	TX-6702	2380	150	8.3		Charcoal- dispersed		2430-2060	2620-1910
Cowhouse Ck; CB18	Beta-37156	1820	80	1.1		Charcoal- hearth		1850-1640	1930-1560
Cowhouse Ck; CB16	Beta-37450	1690	90	2.75	-19.2	Charcoal- hearth		1710-1480	1820-1390
Cowhouse Ck; CB14	Beta-38174	1500	60	5.71		Charcoal- dispersed		1490-1330	1530-1300
Cowhouse Ck; CB4	TX-6701	650	160	4.05		Charcoal- hearth		830-580	940-510
Cowhouse Ck; CB13a	TX-6700	600	140	0.47	-20.3	Charcoal- hearth		750-570	900-500
Ft Hood-Stratigraphic Unit E- Ford Alluvium (Nordt 2004: Table 1)									
Cowhouse Ck; CB3	Beta-38177	390	60	3.14		wood		450-320	500-300
Cowhouse Ck; CB5	TX-6699	370	180	2	-19	Charcoal- dispersed		440-290	500-120
Cowhouse Ck; CB7	TX-6697	300	100	3.1		Charcoal- dispersed		440-290	490-150
Cowhouse Ck; TR18	Beta-37008	190	90	1.8		Charcoal- hearth		420-220	470-70
Lower Extent Brazos River Basin									
Allostratigraphic Unit I (Waters & Nordt 1995: Table 1)									
	SMU-1754	17730	130		-27.5	Wood		21450-20970	21500-20770
Buffalo Ranch- A&M Soil	A-7513	8465	100		-15	Soil	9635- 9272	9550-9400	9660-9260
Hearth in A&M Soil	GX-15417	8390	330		-24.1	Charcoal	9585- 9151	9570-9320	9830-9160
Allostratigraphic Unit II (Waters & Nordt 1995: Table 1)									
County Road Locality	A-7511	8145	75		-29.1	Wood	9265- 8771	9200-8810	9270-8770
Hearth in County Road Locality	A-6401	6480	100		-26.1	Charcoal	7572- 7244	7480-7290	7580-7170
Hearth in Buffalo Soil	A-7510	4185	55		-25.6	Charcoal	4836-4787,4763- 4690,4677-4644	4851- 4570	4860-4570
Allostratigraphic Unit III (Waters & Nordt 1995: Table 1)									
Log in A&M soil	GX-15415	2505	75		-27.1	Wood	2640-2459 (52.3%)	2650-2360	2740-2350
Charcoal at Berger Farm	A-7509	1405	145		-25.4	Charcoal	1520-1429 (19.9%), 1426- 1261 (48.3%)	1520-1170	1620-980

Table 6.2 Radiocarbon Data for Brazos River Basin

Goodland Farm- Asa Soil	GX-15418	1320	170	-16.4	Soil	1301-1054	1415- 916	1390-1010	1610-910
Hearth in Asa soil	A-6400	880	50	-26.1	Charcoal	910-854 (37.7%), 828-813 (7.9%), 799-755 (22.7%)	925-727	910-740	930-720
Allostatiographic Unit IV (Waters & Nordt 1995: Table 1)									
Goodland Farm- log	GX-15416	530	70	-23.7	Wood	556-502	633-466	560-460	630-320
Debris/twigs Goodland Farm	GX-15422	430	70	-26.5	Wood	525-475	550-425	530-340	540-320
Goodland Farm- log	GX-15420	400	70	-25.5	Wood	504-438	520-365	520-330	530-320
Goodland Farm- log	GX-15419	315	70	-28.8	Wood	485-390	497-312	500-390	510-310
Debris/twigs Goodland Farm	GX-15421	270	70	-26.3	Wood	460-358 (58.0%), 319-296 (10.2%)	495-266	500-380	510-290

*standard value of -25.0 assumed by previous investigators

**radiocarbon date not listed on original table

***duplicate sample number in original text

After the brief erosional event, the Fort Hood alluvium began to be deposited upon the Georgetown alluvium (Nordt 2004). This alluvial unit composes the majority of the T₁ terrace and had 12 radiocarbon samples to provide chronometric data. Half of these samples were charcoal and primarily date to 6,900–4,700 ¹⁴C yr BP. One sample (GX-15760) collected from the Leon River investigations dates to 8,600 ¹⁴C yr BP, which temporally overlaps with the older Georgetown alluvium. The construction of the Fort Hood alluvium ended during another change in hydrologic flow coupled with an erosional event.

Above the Fort Hood alluvium on the T₁ terrace on Cowhouse Creek is the West Range alluvium unit. Frequently recognized as two separate units (upper and lower), this allostratigraphic unit has had the most radiocarbon samples (n=29) collected from it within the Fort Hood study area. Sixteen of these samples are derived from charcoal and primarily date from 4,200–600 ¹⁴C yr BP. The division between the upper and lower West Range alluvium is interpreted to be a very brief erosional event and an increase in hydrologic flow occurring around 2,400 ¹⁴C yr BP (Nordt 2004:297). Subsequent to that, the Upper West Range alluvium is indicated to have a coarser bed load from the increased flow. Capping the Upper West Range in some locations (Henson Creek) is a buried soil identified as the Tanktrail paleosol (Nordt 1995). Further, only the Upper West Range division was identified on Henson Creek, which occupied the T₀ landform and not the T₁ as identified on the larger Cowhouse Creek (Nordt 1995, 2004). The absence of the Lower West Range on Henson Creek is likely attributed to a complete removal from the brief erosional event around 2,400 ¹⁴C yr BP (Nordt 1995:214). The West Range alluvium ended during another erosional event around 600 ¹⁴C yr BP, which

incised into underlying bedrock beginning the modern Cowhouse Creek floodplain (Nordt 2004:297).

Finally, above the West Range alluvium forming the modern Cowhouse Creek floodplain and current allostratigraphic unit is the Ford alluvium. This alluvial unit composes the T₀ terrace and has 12 radiocarbon samples to provide chronometric data. These samples, all derived from charcoal or wood, date from 700–200 ¹⁴C yr BP. However, using only those samples from the Cowhouse Creek drainage Nordt (2004) correlates the Ford alluvium to encompass 400 ¹⁴C yr BP to the present.

Lower Extent

Numerous geomorphic examinations have been conducted along the Lower Extent of the Brazos River Basin. One of the most significant is Abbott's (2001) synthesis of regional geoarchaeology, which provides an exceptional review of previous research for the lower extent of the basin as well as the Gulf Coast. This research examined the Late Quaternary stratigraphy and various geomorphic processes of the Houston area. Further, Abbott (2001) cogently characterized the affects of the processes upon the cultural resources within this area and developed a model for evaluating the likelihood for the presence and integrity of archaeological resources. Although these investigations did have chronometric data, it was not a primary component of the research.

Similarly, most of the other geomorphic investigations in the lower basin have not undertaken extensive chronometric analysis (e.g., Husain 1998; Nordt 1983, 1986). One exception is a project conducted in the mid-1990s that did examine a suite of radiocarbon samples with the intent of characterizing the depositional history of the basin. The

research conducted by Waters and Nordt (1995) compared allostratigraphic units they had identified in the Brazos River study area with other drainage basins in the region. The researchers investigated a 75 km segment of the Brazos River between the cities of Hammond and Navasota and west of College Station, Texas. These investigations involved the examination of numerous drainage profiles as well as documentation of six cutbank exposures and the collection of charcoal and bulk sediment samples for chronometric analyses.

Brazos River (A & M Study area)

The culmination of these investigations was the identification of a complex depositional history of the Brazos River that extended into the Late Pleistocene, which exhibited multiple allostratigraphic units (Figure 6.5). The researchers interpreted the stratigraphy in the examined floodplain to have five allostratigraphic units (i.e., Units I–V) bounded by erosional disconformities and buried soils (Waters and Nordt 1995:311–312). The chronometric analyses for this study consisted of 15 radiocarbon samples composed of wood and charcoal and two sediment humate samples (Waters and Nordt 1995:315). Although the researchers calibrated these radiocarbon results to calendar years, they reported the results in radiocarbon years.

The earliest allostratigraphic unit (Unit I) is situated upon Tertiary bedrock and had three radiocarbon samples (two wood and one bulk sediment humate) that ranged from approximately 18,000–8,400 ^{14}C yr BP (Waters and Nordt 1995:Table 1). Two of the radiocarbon samples were collected from a buried soil (A&M soil), which capped Unit I interpreted to be the *terminus post quem* for this allostratigraphic unit at roughly




Figure 6.5 Idealized Cross-section of Brazos River at A&M study area (adapted from Waters and Nordt 1995: Figure 3).

8,400 ¹⁴C yr BP. Notably, one of the samples (GX-15417) came from a cultural feature within the A&M paleosol, which caps Unit I (Waters and Nordt 1995:313).

The second allostratigraphic unit (Unit II) rests unconformably above Unit I suggesting a period of erosion between the end of Unit I and the beginning of Unit II (Waters and Nordt 1995:315–316). The researchers interpreted this break as a decrease in hydrologic discharge along the Brazos River citing a smaller channel and decrease in lateral movement of the drainage. Three radiocarbon samples were collected from Unit II ranging from 8,100–4,200 ¹⁴C yr BP. One of the radiocarbon samples (i.e., AA-12579) was collected from a cultural feature within a buried soil (Buffalo soil) that caps Unit II and marks the *terminus post quem* for this unit at about 4,200 ¹⁴C yr BP.

The third allostratigraphic unit (Unit III) is unconformably situated above Unit II, which in places has eroded the Buffalo soil (Waters and Nordt 1995:314–315). The researchers interpret this erosion as an avulsion event that terminated the stable period of the Buffalo soil and began Unit III. The chronological data for the third allostratigraphic unit consists of four samples that range from 2,500–900 ¹⁴C yr BP. Unit III is capped by a buried soil (Asa soil) from which two radiocarbon samples were collected. One sample at the base of the Asa soil was a bulk soil sample (i.e., GX-15418) dating to roughly 1,300 ¹⁴C yr BP while the second sample (i.e., A-6400) was collected from a cultural feature and marks the *terminus post quem* for Unit II at about 900 ¹⁴C yr BP. Notably, temporally diagnostic artifacts were recognized at both the top and bottom of the Asa paleosol. Near the base, a Middle to Transitional Archaic Gary/Kent projectile point interpreted to range from 4,450–1,450 cal yr BP was observed while Late Prehistoric Scallorn and Perdiz artifacts interpreted to range from 1,250–450 cal yr BP were

observed at the top of the Asa paleosol (Turner and Hester 1999). Simply put, the diagnostic artifacts provide a broad range of 4,450–450 cal yr BP for the Asa paleosol while the radiocarbon analyses indicate a much more refined range of 1,300–900 ^{14}C yr BP for the buried soil.

Overlying Unit III is the fourth allostratigraphic unit (Unit IV), which has five radiocarbon samples all from wood that roughly range from 530–300 ^{14}C yr BP (Waters and Nordt 1995: Table 1). Unit IV is also capped by a buried soil (Katie soil), which the authors describe as ‘weakly developed’ (Waters and Nordt 1995:315).

The final allostratigraphic unit is Unit V and represents the modern floodplain surface (Waters and Nordt 1995:315). This unit is characterized as a thin drape capping Unit IV and is interpreted to have begun deposition approximately 300 years ago (Waters and Nordt 1995:315).

Overall, the authors briefly summarize the history of the Brazos River study area (Waters and Nordt 1995:316). Sometime in the Late Pleistocene around 18,000 ^{14}C yr BP, a large and widely migrating Brazos River deposited Unit I. By the beginning of the Holocene about 8,400 ^{14}C yr BP this unit had a period of stability, which developed the A&M soil. Between 8,400–8,100 ^{14}C yr BP, the Brazos River avulsed and decreased in hydrologic flow and began depositing Unit II. The deposition of this unit continued until roughly 4,200 ^{14}C yr BP when a period of stability occurred developing the Buffalo soil. Possibly lasting until 2,500 ^{14}C yr BP, the stability ended when the Brazos River avulsed again severely eroding the Buffalo soil and began depositing Unit III. The construction of Unit III continued until roughly 1,250 ^{14}C yr BP when the Brazos River entered a period of stability, which developed the Asa paleosol. At approximately 500 ^{14}C yr BP,

the river avulsed again forming Unit IV that lasted until about 300 ¹⁴C yr BP when the Katie paleosol developed. The modern unit (Unit V) began deposition at roughly 300 ¹⁴C yr BP with the latest avulsion of the Brazos River forming the modern drainage channel.

Calibration Results

The radiocarbon datasets for the previously reviewed Upper, Middle, and Lower Extents of the Brazos River Basin were recalibrated. Beginning at the Lubbock Lake site on Yellowhouse Draw, the results are presented from this point in the upper limits of the basin followed by the Fort Hood chronometric results downstream and finally the chronometric data at College Station area. Undeniably, there are an abundance of previous investigations and radiocarbon datasets throughout the Brazos River Basin that could also have been recalibrated. However, these three datasets have been extensively used by other researchers to characterize the depositional history of the Brazos River basin. Further, each study has good stratigraphy that extends to the Late Pleistocene, has cultural deposits in almost all of the recognized stratigraphic units, and has a robust chronometric dataset.

Upper Extent

Forty-eight radiocarbon assays were selected from the Lubbock Lake (41LU1) assemblage derived from humic acid, humin, and charcoal (Haas et al. 1986:Table 1). The selected assays are part of a relatively straightforward profile, which exhibits the stratigraphic context of each of the samples in relation to each other. Consequently, with the vertical relationship and strata information, the recalibration of the samples can be examined and interpreted (Haas et al. 1986: Figure 4). Of note, the overwhelming majority of these radiocarbon assays are from humic acid or humin. Unfortunately,

samples derived from charcoal were the minority in this assemblage. Therefore, due to reasons previously elaborated (e.g., mean residence time), the calibrated results for these samples may trend older than their true temporal context. However, these chronometric data do conform sequentially and associated temporally diagnostic artifacts do correlate with the respective strata.

The five stratigraphic units (Strata 1–5) at the Lubbock Lake site were categorized into eight phases based upon internal horizons within the strata. Specifically, the chronometric assays were grouped from oldest to youngest into Stratum 1, Stratum 2A, Stratum 2B, Stratum 3, Stratum 4A, Stratum 4B, Stratum 5A, and Stratum 5B. The two samples collected from the top of Stratum 1 calibrated to 13,080–12,850 cal yr BP.

The *terminus ante quem* for the overlying Stratum 2A calibrated to 12,460 cal yr BP suggesting a possible 390-year gap between the end of Stratum 1 and the beginning of Stratum 2. Interestingly, Stratum 1 is recorded to contain Clovis cultural materials while Stratum 2A contains a Folsom occupation (Haas et al. 1986: Figure 3). The *terminus post quem* of Stratum 2A is indicated to be about 11,600 cal yr BP while the base of Stratum 2B dates to roughly 11,490 cal yr BP suggesting a very brief gap (100 years) that falls within the margin of deviation. The brief (about 860 years) Stratum 2A contains Folsom deposits while Stratum 2B is recorded to have Plainview cultural materials at the lower portions and Firstview occupations near the top (Haas et al. 1986: Figure 3).

Stratum 2B is capped by a buried soil aptly titled the Firstview Soil (Haas et al. 1986; Holliday 1985). The assays from the Firstview Soil indicating the *terminus post quem* of Stratum 2B calibrate to 8,920–7,270 cal yr BP (Table 6.2). The three assays from the overlying Stratum 3 calibrates to 6,630–5,830 cal yr BP suggesting an

approximate 670 year gap between Strata 2 and 3. Stratum 3 is capped by the Yellowhouse Soil characterized as weakly developed (Holliday 1985:1487). Two of the Stratum 3 assays (i.e., SMU-1093 and SMU-531) come from the Yellowhouse Soil calibrating to 6,100–5,830 cal yr BP almost encompassing the entire span of Stratum 3.

Interestingly, the overlying Stratum 4A calibrates to 5,810–5,720 cal yr BP suggesting a very brief gap between Strats 3 and 4, which is at variance with the observed stratigraphy. Specifically, Stratum 4A is recognized to unconformably rest upon Stratum 3 suggesting an erosive event belying the negligible gap between these strata (Holliday 1985:1488). Consequently, the accuracy of the dates for the Strata 3 and 4 transition should be accepted with prudence.

The calibrated *terminus ante quem* for Stratum 4B is 5,690–5,340 cal yr BP while the Lubbock Lake Soil assays capping Stratum 4B calibrate to 2,240–740 cal yr BP. These results suggest an approximate 3,100-year hiatus between the Lubbock Lake Soil and the base of Stratum 4B (Haas et al. 1986: Figure 4). This hiatus likely attributed more to sampling rather than deposition, but it is interesting that this gap falls within the enigmatic Middle Archaic archaeological period.

The overlying Stratum 5A has a calibrated *terminus ante quem* of 650–560 cal yr BP and is capped by the Apache Soil providing the *terminus post quem*, which dates to 430–300 cal yr BP (Figure 6.4; Table 6.2). Finally, the sole assay for Stratum 5B calibrates to 230–30 cal yr BP suggesting that it is decidedly modern.

Comparing the initial Lubbock Lake radiocarbon calibration results to the recalibration of this study demonstrates some significant differences (Figure 6.6). Notably, these shifts were to be expected considering the initial radiocarbon assays were




Figure 6.6 Calibrated Depositional History Brazos River Basin.

only partially calibrated. Specifically, assays older than 7,240 ^{14}C yr BP (i.e., preceding Stratum 3) were not initially calibrated (Holliday et al. 1983:170, 1985). Rather, the initial dates prior to Stratum 3 were to be considered ‘minimum ages’ for the respective assays (Holliday et al. 1983:171). Therefore, the older assays of the site (particularly in Strata 1 and 2) having been adjusted by more recent calibration data, do exhibit significant adjustment (Figure 6.6). The recalibration results push the ages of Stratum 1 back about 1,500 years and considerably shorten this unit. Similarly, Stratum 2 has also been pushed back to start approximately 12,500 cal yr BP. Further, the beginning of Stratum 4 has been shifted about 500 years older to begin about 5,800 cal yr BP. However, the Strata 3 and 4 transition rests on a disconformity and, as previously mentioned, the accuracy of the dates for the Strata 3 and 4 transition warrants caution. The comparison of the chronometric data for the remaining Lubbock Lake strata aligns exceptionally well. Although there are some refinements in the assays, these differences are negligible.

Another observation from the recalibration of the Lubbock Lake data regards the ‘hiatus’ in Stratum 4B. Specifically, an apparent 3,100-year gap in radiocarbon age reveals itself between the Lubbock Lake Soil and the base of Stratum 4B. None of the Stratum 4B chronometric data overlaps this time period. This temporal gap is probably more a result of sampling rather than depositional or geomorphic processes.

Middle Extent

Twenty-one radiocarbon assays from Cowhouse Creek and one from Tablerock Creek were selected from the investigations in Fort Hood derived from humate and charcoal to be calibrated (Nordt 1992, 2004: Table 1). The selected assays are overwhelmingly

charcoal are from a good stratigraphic context, and compose an assemblage used to characterize the alluvial history of Cowhouse Creek and Fort Hood (Nordt 2004: Figures 5 and 6). As previously indicated, six allostratigraphic units have been identified within the Fort Hood study area, which from oldest to youngest include Jackson alluvium, Georgetown alluvium, Fort Hood alluvium, Lower West Range alluvium, Upper West Range alluvium, and the Ford alluvium.

The one sample collected from the Late Pleistocene Jackson alluvium calibrated to 18,680–18,080 cal yr BP (Table 6.2). This assay (Beta-38694) is the sole humate sample in this assemblage. Above the Jackson alluvium is the Georgetown allostratigraphic unit, which calibrated to 10,100–9,750 cal yr BP. The significant 7,980-year gap between the end of the Jackson alluvium and the beginning of the Georgetown alluvium correlates to a period of incision, which likely eroded significant deposits of the Jackson allostratigraphic unit (Nordt 2004). The Georgetown alluvium is capped by the Royalty paleosol, but none of the selected assays from Cowhouse Creek was from this buried soil. However, a charcoal assay (GX-15762) associated with the Royalty paleosol from Tablerock Creek was used as a proxy and calibrated indicating the buried soil dated to 9,410–9,120 cal yr BP (Nordt 1992). The four assays for the overlying Fort Hood alluvium calibrates to 7,780–5,860 cal yr BP. An approximate 1,300-year gap separates the Georgetown and Fort Hood units, which correlates to a period of incision. The *terminus ante quem* for the Lower West Range alluvium calibrates to 4,790 cal yr BP and terminates sometime after 2,790 cal yr BP.

The overlying Upper West Range calibrates to 2,430–570 cal yr BP indicating an approximate 400-year gap between the Upper and Lower West Range units. Of note, a

hiatus is present in both these West Range units (Figure 6.7). In the Lower West Range, a 650-year hiatus occurs at one sigma deviation between 3,980–3,330 cal yr BP. Similarly, in the Upper West Range a 500-year hiatus occurs between 1,330–830 cal yr BP. At present, it is undetermined if these gaps are reflective of depositional processes, sampling, or a combination of these factors. However, the hiatus of the Lower West Range coincides with the aforementioned gap observed in Stratum 4B at Lubbock Lake (Haas et al. 1986; Holliday 1985). Regardless, the period of incision identified between the Lower and Upper West Range units appears to have occurred between 2,790–2,430 cal yr BP (Nordt 2004: Figure 6). Finally, overlying the Upper West Range unit is the Ford alluvium that calibrates to 450–220 cal yr BP. The identified period of incision between these units appears to have been brief (120 years) and occurred between 570–470 cal yr BP.

Some notable differences are apparent when contrasting the initial Cowhouse Creek at Fort Hood radiocarbon calibration results to the recalibration of this study. Typically, adjustments are limited to the older assays, but the recalibration of the Cowhouse Creek dataset exhibits shifts throughout all of the allostratigraphic units. The largest shift involved the Georgetown unit where the recalibration truncated the beginning and terminus of this unit (Table 6.2). Admittedly, the recalibration did utilize a radiocarbon assay (i.e., GX-15762) from a drainage (i.e., Tablerock Creek) other than Cowhouse Creek as a proxy to date the Royalty paleosol. However, the assay was derived from charcoal and conforms to the Cowhouse Creek chronology and is considered reliable. Another significant shift occurs in the Fort Hood unit, which has been pushed back about 700 years. The shift suggests that the Fort Hood unit is almost




Figure 6.7 Select Calibration Plot of Cowhouse Creek assays; arrows illustrate hiatus periods.

entirely within the Early Archaic instead of spanning the Early and Middle Archaic periods. The beginning of the Lower West Range unit is shifted about 500 years to begin at roughly 4,800 cal yr BP. Interestingly, a 650-year gap in the radiocarbon assays (i.e., 3,980–3,330 cal yr BP) is revealed in the Lower West Range unit and a similar 500-year gap occurs in the Upper West Range.

Lower Extent

Fourteen radiocarbon assays from the investigations along the Brazos River west of College Station was selected for calibration (Table 6.2). Only two of the fourteen assays were derived from humate, the majority of the samples came from charcoal and wood. Also of note, Waters and Nordt (1995) encountered cultural features or artifacts within each of the buried soils at the College Station study area. Specifically, evidence was observed in the A&M soil of Unit I, the Unit II Buffalo soil, the Unit III Asa soil, and in the Katie soil of Unit IV. The presence of these cultural deposits provided data (i.e., hearth charcoal) from which to securely date each of the paleosols and by extension, periods of environmental stability.

Unfortunately, one of the three radiocarbon assays available for Unit I is derived from humate. However, the result of the calibration of the humate assay (i.e., A-7513) conforms with the charcoal radiocarbon result from Unit I and is considered reliable. Notably, Waters and Nordt (1995: Table I) provide an assay (SMU-1754) derived from wood reportedly in the Unit I channel facies, which is the *terminus ante quem* for this unit. The calibration of this assay dates to 21,450–20,970 cal yr BP with the next recalibration date is 9,570 cal yr BP. Considering Waters and Nordt (1995) do not indicate a disconformity between these two assays, the implications are that the erosive

pre-Holocene event is not represented at this study area. Further, if the erosive event is not present then the enigmatic Pre-Clovis timeframe should be intact. However, there are multiple reasons that this assay (SMU-1754) should be regarded with caution. The assay was collected from a gravel pit on the edge of the floodplain and its vertical position to the other Unit I samples is not indicated. Therefore, this assay is interpreted with some prudence.

The calibration of the A&M soil that caps Unit I and provides the *terminus post quem* suggests that this unit ceased deposition sometime after 9,320 cal yr BP. The overlying Unit II began deposition prior to 9,200 cal yr BP indicating an extremely short (80 year) transition between the two units. The deposition of Unit II continued until the Middle Holocene ending sometime after 4,640 cal yr BP. The calibration of the Buffalo Soil that caps Unit II suggests a period of stability occurred prior to 4,840 cal yr BP. Notably, two gaps in the chronometric data of Unit II are evident within this unit. One gap exhibits a 1,300-year hiatus between 8,810–7,480 cal yr BP while the second 2,500-year gap occurs between 7,290–4,840 cal yr BP (Table 6.2 and Figure 6.6). These gaps are partially attributed to sampling since only three radiocarbon assays represent Unit II. Another possibility may be attributed to a base level rise in sea level (i.e., transgression). As previously mentioned in the Nueces River basin study, a rapid rise in sea level occurred along the Gulf at 6,800–5,900 cal yr BP and 4,200–3,000 cal yr BP (Ricklis and Blum 1997; Ricklis and Cox 1998).

Unconformably resting on Unit II is the third allostratigraphic unit (i.e., Unit III) (Figure 6.5). The recalibrated *terminus ante quem* for Unit III suggests deposition began prior to 2,650 cal yr BP when the Brazos River is interpreted to have avulsed and

severely eroded the Buffalo soil of Unit II. The erosive event occurred sometime between 4,640–2,650 cal yr BP, which not coincidentally squarely falls within the second sea level transgression (4,200–3,000 cal yr BP) identified by Ricklis and others (1997, 1998). The Asa Soil that caps Unit III provides the *terminus post quem* that suggests the Brazos River entered a period of stability prior to 1,390 cal BP and ends sometime after 740 cal yr BP. Finally, sometime prior to 560 cal yr BP an avulsion occurred that began the construction of Unit IV. The Brazos River entered a brief period of stability forming the Katie Soil around 300 cal yr BP followed by the most recent avulsion, which formed the modern channel (Unit V).

In aggregate, this depositional history of the Brazos River basin identified from the recalibration of previous research will be compared with those in other drainage basins and correlated with extrinsic factors in Chapter 8.

CHAPTER 7

Recalibrated Geoarchaeological Framework with the Trinity River Basin

The Trinity River is solely contained within Texas and is generally recognized to have an upper and lower extent (Gard 2010). The Upper Trinity River basin is situated in the North Central Plains region and encompasses the headwaters region of the basin bounded by the Brazos River basin to the south and the Red River basin to the north. The Trinity River basin is the only basin within this study that does not cross the Edwards Plateau. Instead, the Upper Trinity River is recognized to cross forested rolling topography with narrow stream channels with three main headwater branches, the Elm Fork, the West Fork, and the Clear Fork Rivers (Ferring 1991; Gard 2010). In contrast, the Lower Trinity River basin crosses the grass prairies of the Gulf Coastal Plain beginning just between Dallas, Texas and the Trinity's confluence with the Elm Fork River and trends southeastward to terminate at Trinity Bay on the coast (Figure 7.1). Overall, the basin encompasses a total 17, 969 mile² (46,500 km²) area with the three Upper Trinity River branches averaging 114 miles (183 km) in length while the Lower Trinity River basin is about 260 miles (420 km) long (Ferring 1991; Gard 2010; Garvin 2008; Prikryl 1990). Some of the prominent contributory drainages in the Trinity River basin include Elm Fork, East Fork, West Fork, and Clear Fork Rivers. Less prominent tributaries include Ten Mile, Five Mile, White Rock, Keechi, Clear, Hickory, and Cedar creeks as well as Cedar Bayou. The easternmost tributary is the East Fork that is about 78 miles (125 km) long and extends through Grayson, Dallas, and Kaufman Counties. The central tributary drainage is the roughly 85 mile (137 km) long Elm Fork River,




Figure 7.1 Overview of Trinity River Basin: 1) Ray Roberts-Upper Trinity River study area.

which runs through Montague, Cooke, and Denton Counties. The westernmost tributary drainage is the 180 mile (290 km) long West Fork River that runs through Archer, Jack, Tarrant, and Dallas counties. The Trinity River within the Lower extent of the basin runs generally southeast through Kaufman, Ellis, Henderson, Navarro, Freestone, Anderson, Leon, Houston, Madison, Walker, Trinity, San Jacinto, Polk, Liberty and Chambers Counties where it empties into Trinity Bay near Anahuac, Texas.

Previous Investigations

The Trinity River and its deposits have been of interest to geologists and archaeologists for over a hundred years (Ferring 2000). Arguably, the first geoarchaeological investigation to have been conducted in Texas occurred in 1920 in the Trinity River basin (Table 7.1). Specifically, Robert Hill and Ellis Shuler examined a human skeleton discovered at the Lagow Sand Pit along the Trinity River in Dallas County to determine its association with Pleistocene fauna (e.g., mammoth, camel, and horse) also discovered there (Ferring 2000:47). Hill and Shuler interpreted the human remains to be contemporaneous with the Pleistocene fauna, although subsequent analyses in the late 1960s determined that the remains were actually much younger.

Despite the fact that these early researchers (e.g., Robert Hill, Cyrus Ray, or Ellis Shuler) did not benefit from radiocarbon dating, their research attempted to determine the age of the Trinity River terraces and its deposits having some success. The first researchers to characterize the terraces of the Trinity River in combination with reconstructing the paleoenvironment were Stovall and McAnulty (1950) in Henderson

Table 7.1 Previous Geoaarchaeological Investigations in the Trinity River Basin

Drainage Extent	Drainage	Resource	Project-Site(s)	Geoarchaeologist or Chronometric Data	
				Researcher	Data
Upper	Trinity River	Shuler 1935; article		E. Shuler	Relative
Upper	Trinity River	Stovall and McAnulty 1950; article	Henderson, Navarro, and Freestone Counties	J. Stovall and W. McAnulty	Relative
Lower	Trinity River-Coast	Aten 1983	Synthesis Gulf Coast	L. Aten	14c; relative
Upper	Trinity River	Prikryl and Yates (editors) 1987	41CO141 Ray Roberts Reservoir	C.R. Ferring	14c; relative
Upper	Elm Fork	Ferring 1994	Upper Trinity River basin; Dissertation	C.R. Ferring	14c; relative
Upper	Elm Fork	Ferring 1995a; 2001	Aubrey Clovis Site	C.R. Ferring	14c; relative
Upper	Elm Fork	Ferring and Yates 1997	Ray Roberts Reservoir	C.R. Ferring	14c; relative
Upper	Mill Creek	Byers 2007	Dickie Carr Site (41PR26) Dissertation	J. Byers	Relative
Lower	Trinity River	Garvin 2008	Trinity River incised valley; thesis	M. Garvin	OSL; relative
indicates study selected for recalibration					

County. Similarly, the lower extent of the basin and gulf coast has been extensively considered by numerous researchers (e.g., Bernard et al. 1962, 1970). Concerted geoarchaeological investigations appear to have begun in the late 1960s in the lower extent of the Trinity River basin and in the 1980s for the upper extent. In the Lower Trinity River basin, Aten (1983) conducted a series of archaeological investigations along the coast with a research focus on geomorphic processes.

For the Upper Trinity River basin, the research that occurred for Lake Ray Roberts involved an extensive geoarchaeological component (Prikryl and Yates 1987). Since then several significant geoarchaeological investigations have been carried out in the Trinity River Basin. Interestingly, the research in the Trinity River basin has largely been conducted either in the upper extent or in the extreme lower extent along the Gulf Coast. Unfortunately, geoarchaeological research along the middle region of the Trinity River has been sparse. Despite the limited amount of geoarchaeological investigations within the basin, the research that has been conducted is thorough and far-reaching.

Some of the first archaeological investigations within the Trinity River basin with a focus on geomorphic processes was conducted by Aten (1983). The initial research was associated with the Wallisville Reservoir project in Chambers County, which later developed into a much larger synthesis. Aten (1983:104–162) supplemented previous archaeological investigations with geologic investigations along the Trinity River and produced a synthesis of Late Quaternary stratigraphy for the drainage and the coast. Within the basin, Aten (1983:105) characterized the Trinity River as having a series of fluvial terraces encompassing the Pleistocene to modern times. From oldest to youngest

these include terraces T₄–T₀ with terraces T₄ and T₃ associated with the Pleistocene while T₂–T₀ encompassing the Late Pleistocene to modern (Aten 1983:Table 8.3).

The most extensive geoarchaeological investigations within the Trinity River basin are associated with the Ray Roberts-Lewisville Reservoir project in Cooke, Denton, and Grayson Counties (Prikryl and Yates 1987; Ferring and Yates 1997). The creation of these reservoirs from the impoundment of segments of the Elm Fork River generated archaeological investigations extending over two decades by a variety of interdisciplinary researchers (e.g., SMU, Environmental Consultants Inc., USACE-FW, and University of North Texas). Among the many notable accomplishments from this research was the discovery and investigation of the Aubrey Clovis site (41DN479). The Aubrey Clovis site situated on the Elm Fork River was identified to have an intact stratigraphy extending back into the Pleistocene (Ferring 1990a, 1990b, 1991, 1992, 1994, 1995a, 1995b, 2000, 2001; Humphrey and Ferring 1994). Most significant, the site has a Paleoindian occupation with a well-dated stratigraphy that suggests it is the oldest Clovis site currently known (Ferring 2001). In addition, the extensive investigations at Aubrey Clovis have provided a robust radiocarbon dataset and information from past environments.

Finally, a geoarchaeological examination was conducted at the Dickie Carr site (41PR26) also in the upper extent of the Trinity River basin (Byers 2007). The Dickie Carr site is located in eastern Parker County situated on a terrace of Mill Creek, a tributary of West Fork River. Byers (2007:57–72) identified three stratigraphic units (Units I–III) that contained a complex stratigraphy extending to the Late Pleistocene with the remains of a *Mammuthus columbi* (Unit Ib), a Late Paleoindian component (Unit IIa),

and a Late Archaic component (Unit IIb). The researcher compared the site deposits and setting with other archaeological sites in the region. Unfortunately, no chronometric analyses were conducted for this research.

Geomorphic/Alluvial History

The following review of the depositional history of the Trinity River basin is composed of an intensive review of the interpretations associated with the Ray Roberts-Lewisville Reservoir project and particularly the Aubrey Clovis site (41DN479). As mentioned previously, there is a paucity of geoarchaeological research in the interior or middle portions of the Trinity River basin followed by the coastal region. Although this depositional history is in the upper limits of the basin, the data is applicable to the rest of the basin.

Upper Extent Trinity River Basin

In summarizing the alluvial history of the Upper Extent of the Trinity River basin, Ferring (1994) recognizes four morphostratigraphic units (i.e., landforms) composed of deposits from six alloformations (Figure 7.2). The morphostratigraphic units identified from oldest to youngest include the Stewart Creek Terrace, Hickory Street Terrace, Denton Creek Terrace, and the Floodplain that are interpreted to encompass the Middle Pleistocene up to the present.

The Stewart Creek Terrace composed of Irving alloformation deposits and the Hickory Street Terrace, which is composed of the Coppell alloformation deposits are indicated to date to the Pleistocene sometime around 30,000 years ago (Ferring 1994; Ferring and Yates 1997). The more recent Denton Creek Terrace is composed of deposits from the Carrollton alloformation that contain Pleistocene faunal remains (e.g.,




Figure 7.2 Stratigraphic Columns from Aubrey Clovis site (adapted from Ferring 1994: Figure 3.9).

Bison antiquus) and is interpreted to date to roughly 30,000–20,000 years ago (Ferring 1994:47–48). Most relevant to the archaeology of the region is the landform identified as the Floodplain. The Floodplain contains deposits from three alloformations that from oldest to youngest include the Aubrey, Sanger, and Pilot Point alloformations.

Ferring (1994) indicates that the Aubrey alloformation dates to the Late Pleistocene-Holocene transition. The *terminus ante quem* for this alloformation is from the Aubrey Clovis site (41DN479), which initially calibrated to 17,030 cal yr BP (SMU-2236) and terminates sometime after 14,410 cal yr BP (Ferring 1994; Ferring and Yates 1997). The initial calibration of the overlying Sanger alloformation begins in the Late Pleistocene sometime prior to 13,460 cal yr BP (AA-5274) and extends to sometime after 7,550 cal yr BP (SMU-2339) (Ferring 2001: Table 3.2). Ferring (1994:58–59) notes that the Sanger alloformation is capped by a moderately developed buried soil that in several locations has been eroded by the Pilot Point alloformation. The Pilot Point deposits began prior to 4,470 cal yr BP (SMU-2401) and terminated sometime after 1,676 cal yr BP (Beta-14963). A well-developed cumulic soil identified as the West Fork soil frequently caps the Pilot Point alloformation interpreted to encompass the last 4,000 years.

Beginning in the Late Pleistocene, the alluvial history of the Upper Trinity River basin as defined by Ferring (1994:147–149) has a period of stasis with no discernable geomorphic activity occurring between 14,000–11,000 years ago (17,000–13,000 calibrated). Specifically, Ferring (1994:147) notes that no alluvial or colluvial deposition or evidence of a disconformity is associated with this period. Subsequent to 11,000 years (13,000 calibrated) ago, a period of rapid alluviation occurs in the basin initiating a phase

of valley filling (Figure 7.3). Interestingly, this period of rapid alluviation occurred shortly after the Clovis occupation at the Aubrey Clovis site and continued until 7,500 years ago (7,550 calibrated) (Ferring 1994:148). This event is inferred to be associated with a shift to a moister (i.e., mesic) climate. By the middle of the Holocene, a shift to a drier (xeric) climate occurred as evidenced by soil development in the Sanger alloformation and eolian deposits in some of the uplands of the upper basin of the Trinity River (Ferring 1994:148). These xeric conditions seemingly continued until roughly 4,500 years ago when a period of rapid alluviation occurred and the deposition of the Pilot Point alloformation began (Ferring 1994:148–149). The rapid alluviation is particularly apparent between 3,000–2,000 years ago (Ferring 1994:149).

In contrast to other researchers, Ferring (1994) notes that alluviation in the Upper Trinity River basin does not correlate with arid conditions. Instead, Ferring (1994:150–153) noted in this basin that the fluvial response to moist conditions was rapid alluviation and/or erosion while drier conditions led to stability and soil development and that the primary internal influence on a landscape's evolution is the underlying bedrock, which affects the vegetation. Specifically, the calcareous loams and clay loams derived from weathered limestone support a prairie environment that is more resistant to erosion. Conversely, the non-calcareous soils derived from sandstone and shale supports a mixed forest environment that has a lower threshold for erosion. Two primary external influences identified are the climate and changes in sea level. Although no examples are provided, eustatic influence is proposed as a possible influence to the upper extent of the basin.




Figure 7.3 Initial Depositional History Trinity River Basin.

Regarding overall archaeological potential within the upper basin, Ferring and Yates (1997) indicate that archaeological sites occur in two principal contexts (i.e., terraces and upon or below floodplains). For the terraces, Ferring and Yates (1997) propose that the Trinity River terraces between Dallas and Valley View are Late Pleistocene in age. As such, any archaeological sites less than 11,000 years old (~13,000 calibrated) could occur on the terrace surfaces, which are supported by numerous surveys in the area particularly in Denton, Dallas, and Tarrant Counties (Ferring and Yates 1997:279).

For the floodplains in the Upper Trinity River basin, the age of the sites on the floodplains can be predicted by their stratigraphic location (Ferring and Yates 1997: Table 18.2). Unfortunately, Ferring and Yates (1997) note that cutbank exposures along Elm Fork River are “poor” and alluvial units thicken as they progress downstream. Consequently, older sites will become increasingly difficult to discern downstream. This will be particularly evident for Paleoindian–Middle Archaic sites, which will likely require mechanical excavation for their discovery. In contrast, the Pilot Point alloformation is exposed within the cutbanks of the entire upper extent of the basin and is characterized as thick, dark and clayey with a buried soil (West Fork soil) located beneath recently deposited sand (Ferring and Yates 1997:280).

Calibration Results

The radiocarbon datasets for the previously reviewed investigations associated with the Ray Roberts-Lewisville Reservoir project in the upper Trinity River basin were recalibrated (Table 7.2). The chronometric data consists of 57 samples derived from a

Table 7.2 Radiocarbon Data Trinity River Basin

Locality	Lab#	¹³ C Adjusted ¹⁴ C yr BP	Dev +/- (m)	Depth (m)	δ ¹³ C	Material sampled	Initial 68.2%	Initial 95.4%	IntCal09 68.2%	IntCal09 95.4%
Aubrey alloformation (n=9)										
Aubrey Clovis site										
Strat B1b Trench 2A and 13	SMU-2236	14200	220	85.5	-28	Peat	17030		17340- 16910	17630- 16790
Strat B2d Trench 1	SMU-2199	13810	880	87.95	-27.6	Humates	16560		17030- 15390	17590- 14660
Strat B2d Trench 1	SMU-2195	13710	80	87.95	-27.9	Humates	16440		16930- 16740	17030- 16620
Strat B2a Trench 1	SMU-2302	13665	170	87.2	-28.3	Humates	16380		16930- 16560	17110- 16060
Strat B1 Trench 1	SMU-2304	13570	400	87.35	-28	Peat	16260		16900- 15680	17310- 15180
Strat B2b Trench 1	SMU-2303	13575	100	87.95	-28.3	Peat	16260		16840- 16570	16940- 16320
Strat B2d Trench 1	SMU-2202	13340	410	87.95	-23.4	Peat	15940		16710- 15440	17090- 14950
Strat 2A and 13	SMU-2305	13260	105	86.75	-29	Peat	15830		16540- 15750	16730- 15350
Strat C2 Trench 2A and 13	SMU-2478	12330	170	87.2	-24.2	Humates	14410		15100- 14490	16010- 14140
Sanger alloformation (n=16)										
Strat G/A Trench 2A and 13	AA-5271	11540	110			Charcoal	13460		13510- 13280	13650- 13180
Strat G/A Trench 2A and 13	AA-5274	11590	90			Charcoal	13520		13550- 13330	13670- 13260
Aubrey Clovis; Strat E1 Trench 2A and 13	SMU-2194	10940	80	87.8	-18.3	Humates	12860		12940- 12710	13070- 12640
Aubrey Clovis; Strat G Trench 25	SMU-2338	10720	90	89.05	-17	Humates	12650		12720- 12570	12860- 12450
Aubrey Clovis; Strat E3 Trench 2A and 13	SMU-2406	10390	80	88.45	-16.9	Humates			12410- 12110	12540- 11990
Aubrey Clovis	Beta-32002	10360	150			Sediment Humate			12420- 11890	12570- 11540

Table 7.2 Radiocarbon Data Trinity River Basin

Aubrey Clovis; Strat G Trench 25	SMU-2398	10080	80	89.65	-18.6	Humates	11475	11850- 11440	11990- 11310
Aubrey Clovis; Strat G Trench 25	SMU-2399	9570	130	90.3	-18.9	Humates	10750	11100- 10710	11210- 10530
Fullingham Locality 1	GX-18281	9455	255			Charcoal		11120- 10410	11580- 10110
Gateway Park, Fort Worth	Beta-14905	8940	185			Charcoal		10260-9750	10530-9570
Denton Creek Profile 1	Beta-46150	8510	120			Sediment Humate		9640-9360	9870-9180
Aubrey Clovis; Strat G Trench 25	SMU-2400	7460 (8275 in Ferring 1994)	70	91.55	-18.5	Humates	8210	8350-8200	8390-8090
Aubrey Clovis; Strat G Trench 25	SMU-2339	6740 (7580 in Ferring 1994)	320	92.3	-18.1	Humates	7550	7930-7320	8270-6960
Village Creek	Beta-34049	6445	110			Sediment Humate		7460-7260	7560-7100
Denton Creek Profile 1	Beta-46151	6410	100			Sediment Humate		7420-7230	7510-7070
Village Creek	Beta-34051	5680	80			Soil Humate		6610-6400	6680-6320
Pilot Point alloformation (n=32)									
Village Creek	Beta-26738	4150	100			Soil Humate		4710-4460	4820-4340
Aubrey Clovis; Strat H Trench 25	SMU-2401	4030	60	93.05	-16.3	Humates	4470	4580-4430	4770-4310
Aubrey Clovis; Strat H Trench 25	SMU-2402	3260	70	93.55	-16.4	Humates	3470	3580-3410	3670-3360
41CO150	Beta-16526	2910	250			Charcoal		3380-2800	3700-2450
41CO141-1	Beta-16418	2800	90			Charcoal		3060-2820	3200-2760
41CO150	Beta-32514	2750	60			Charcoal		2930-2790	3020-2760
41CO150	Beta-32515	2600	50			Charcoal		2760-2580	2830-2500
41CO150	Beta-32516	2320	70			Charcoal		2470-2210	2680-2150
Aubrey Clovis; Strat H Trench 25	SMU-2403	2080	70	94.35	-16.8	Humates	2008	2150-1960	2300-1890
41CO150	Beta-32513	1985	70			Charcoal		2040-1860	2130-1760

Table 7.2 Radiocarbon Data Trinity River Basin

Village Creek	Beta-26737	1825	70					1840-1650	1920-1570
41CO141-1	Beta-16417	1740	90					1780-1550	1870-1430
Aubrey Clovis; Strat H Trench 25	SMU-2404	1730	50	95.05	-16.6		1676	1710-1570	1800-1530
41CO150	Beta-32982	1680	70					1680-1490	1780-1410
41DN99	Beta-32524	1410	80					1400-1260	1510-1160
Gateway Park, Fort Worth	Beta-14904	1410	100					1430-1220	1530-1090
Denton Creek Profile 1	GX-18282	1350	120					1380-1120	1510-1000
Locality RRG-1	Beta-14908	1300	60					1280-1150	1320-1070
41CO141-2	Beta-32530	1280	90					1280-1090	1350-990
41CO144	Beta-32986	1280	90					1280-1090	1350-990
41DN99	Beta-32523	1270	70					1270-1100	1300-1010
41DL149	Beta-13962	1100	110					1160-910	1260-780
41TR68	Beta-22028	980	70					960-800	1050-740
41CO141-1	Beta-16416	965	60					930-800	1000-740
41CO141-2	Beta-32529	760	70					770-660	890-560
41CO144	Beta-32521	760	70					770-660	890-560
41DL12	Beta-14907	730	80					750-590	880-550
41TR68	Beta-22487	680	50					670-570	710-550
41DL149	TX-4001	680	200					870-550	1060-430
41TR68	Beta-22488	590	100					660-530	730-480
41DN99	Beta-32522	530	70					630-520	660-480
41DL149	Beta-14963	510	80					630-500	660-420

*standard value of -25.0 assumed by previous investigators

**radiocarbon date not listed on original table

***duplicate sample number in original text

dozen locations (Ferring 1994: Table 3.1; Ferring 2001: Table 3.2). Of note, the radiocarbon assays associated with the Coppel and Carrollton alloformations were not recalibrated for this study due to their early temporal setting (i.e., Pleistocene). Rather, only the assays associated with the Aubrey, Sanger, and Pilot Point alloformations were recalibrated (Table 7.2). These alloformations were selected due to their associations with cultural materials and the implications regarding paleoenvironmental interpretation across the upper extent of the Trinity River basin. The materials composing the radiocarbon assays include charcoal, soil humate, peat organic residue, and sediment humate (Ferring 1994: Table 3.1). Unfortunately, none of the assays for the Aubrey alloformation were derived from charcoal, but the *terminus ante quem* of the overlying Sanger alloformation are from charcoal. The Pilot Point alloformation contained the most radiocarbon assays (n=32) of the three and also had the most samples derived from charcoal (n=22). Thus, the Pilot Point alloformation is the most securely dated of the three.

The results of the recalibration of all the assays revealed an adjustment in the Aubrey alloformation, but admittedly not that dramatic considering the initial calibration (Ferring 2001: Table 3.2). Specifically, the Aubrey alloformation is indicated to have begun sometime prior to 17,340 cal yr BP and terminated sometime after 14,490 cal yr BP (Figure 7.4). The stratigraphic spread of the Aubrey alloformation assays suggests a good, continuous coverage (Figure 7.5). The overlying Sanger alloformation seems to have begun deposition prior to 13,510 cal yr BP and terminated sometime after 6,400 cal yr BP. The *terminus ante quem* for this alloformation are from the two charcoal samples (AA-5271 and AA-5274) recovered from the Aubrey Clovis site.




Figure 7.4 Calibrated Depositional History Trinity River Basin.




Figure 7.5 Select Calibration Plot of Upper Trinity River Basin assays; arrow indicates hiatus period.

These dates (13,550–13,280 cal yr BP) are significant in that they are at the boundary between the Aubrey and Sanger alloformations and date the Clovis occupation at the site (Ferring 2001:50). Further, the assay (SMU-2194) that caps the Clovis occupation for the site recalibrated to an age of 12,940–12,710 cal yr BP (Table 7.2). The Clovis occupation at Aubrey Clovis seems to have securely occurred between 13,280–12,940 cal yr BP, which makes them the oldest North American Clovis site (Ferring 2001:50–51). Remarkably, these dates do not change from the initial calibration conducted by Ferring (2001: Table 3.2). Consequently, Ferring's (1994, 2001) initial interpretation appears to be unchanged in light of the most recent calibration curve. However, for the remainder of the Sanger alloformation, there appears to be two significant gaps that occurred between 9,360–8,350 cal yr BP and 7,230–6,610 cal yr BP. The overlying Pilot Point alloformation began sometime prior to 4,710 cal yr BP and continued until sometime after 500 cal yr BP. In general, the Pilot Point alloformation has an excellent stratigraphic spread suggesting continuous accumulation, but there is one temporal gap between 4,400–3,580 cal yr BP (Figure 7.6).

Comparing these recalibrated results to the initial investigations, the most prominent changes are in the older dates. As mentioned previously, the adjustments from the current Incal09 curve is negligible. The statistical analyses using MCMC did refine the temporal spread of some of the assays, particularly in the Aubrey alloformation. Possibly as an indicator of the good, continuous coverage of the assays, the adjustments from the MCMC iterations were not drastic. Of note, another reason is likely to be that due to the incorporation of assays from a dozen locations, the stratigraphical arrangement of the assays within the alloformations was commonly unknown and sorted solely by age.

Regardless, the overall temporal extent encompassed within the alloformations is informative.

The results of the recalibrated chronometric data within the Trinity River basin are examined further and compared with other recalibrated data in the following Chapter 8.

CHAPTER 8

Patterns and Correlations across Texas River Basins and Region

This chapter consolidates the results from the previous basin recalibration studies. One of the primary objectives is to determine any depositional patterns within and between drainage basins and, by extension, attempt to correlate them with extrinsic factors (e.g., climate and eustasy). As with any search for patterns, the researcher will inevitably find them in abundance. At issue is the relevance and validity of identified relationships. Simply put, when is a pattern an *a priori* construct made in the researchers' mind and when does it truly reflect the effect of an external agent? This is a particularly apt question in regards to comparing multiple drainage basins over an expansive region using data from disparate researchers each with distinct research foci. The radiocarbon recalibrations for this study have provided a chronological baseline for all of the selected study areas. This recalibrated chronological framework is a factor that previous investigations did not have.

This chapter begins by reviewing *intra*-basinal relationships of each of the drainage basins followed by *inter*-basinal connections, and finally regional patterns. The chapter ends with a review of extrinsic factors that may be attributed to these proposed patterns. For the basin comparisons, there are three basic categories that are used indicate general activity within the drainages: periods of aggradation, periods of stability, and periods of instability. Aggradation is interpreted to be periods when deposition was occurring within the basin represented by recorded allostratigraphic units (e.g., Ford alluvium and Columbus Bend 2). Stability is interpreted to be a period when neither significant erosion nor aggradation is recorded and is associated with pedogenesis (e.g.,

Royalty and Asa paleosols) (Holliday 1990). Most importantly, instability in a basin is interpreted to be when periods of erosion/incision, a hiatus, or data gap occurs. Examples of erosion/incision are the erosional events noted by Blum (1987, 1992) on the Pedernales and Colorado Rivers. The hiatus periods refers to chronological gaps observed in the recalibration of the radiocarbon assays in allostratigraphic units (Figure 8.1). Some examples of hiatus events are in the Columbus River allostratigraphic units Columbus Bend 1 and Columbus Bend 2 (Figures 5.5 and 5.6). Finally, the data gaps refer to separations between allostratigraphic units where no erosion or aggradation is recorded. An example of this is in the Brazos River between Units II and III (Figures 6.2 and 6.6). The data gaps are likely attributable to erosion.

Intra-Basinal Patterns

Some of the drainage basins in this study are more appropriate for internal comparisons than others. The basins most useful for internal comparison are the Brazos River Basin and to some degree the Nueces and the Guadalupe River Basins. Despite the extensive investigations along the Trinity and Colorado River Basins, only one dataset in each of the drainage basins could be recalibrated.

Nueces River Basin

The alluvial history of the Upper Dry Frio River is seemingly more comparable to the Frio River valley downstream than to the adjacent Sabinal River valley (Figure 8.2). Two of the gaps (i.e., 6,450–5,750 yr BP and 4,050–3,650 yr BP) in the radiocarbon analysis during Phase II investigations at Choke Canyon Reservoir roughly





Figure 8.1 Examples of a Hiatus and Chronometric Gap.



*Calibration Version 4.1.6 Oxcal

Figure 8.4 "Rwgtpu'cmqi "Vtkdwct { 'F tckpci gu'lp'Uwf { .

correlate with some of the gaps in the Units IIa–IIc (Hall et al. 1986:586–588). While the Choke Canyon Reservoir chronology is crude the comparison may suggest synchronous mechanisms affecting the Dry Frio and Frio Rivers. Unfortunately, the Choke Canyon Reservoir radiocarbon analyses were not corrected for isotopic fractionation and were not recalibrated here.

Similarly, the occupation hiatuses (i.e., 6,800–5,900 and 4,200–3,000 yr BP) identified by Ricklis and Blum (1997) and recalibrated here also seem to correlate with the erosional disconformities-lateral migration of the Dry Frio River at Woodrow Heard (Figure 8.2). If the previously mentioned humate sample (Beta-112981) is omitted from the Woodrow Heard assays, then the disconformity between Unit IIa and Unit IIb dates to 6,880–5,580 cal yr BP (Figure 8.2). This disconformity overlaps the first occupation hiatus identified as a period of rapid sea level rise by Ricklis and Blum (1997). Similarly, the disconformity between Unit IIb and Unit IIc occurred between 5,080–3,570 cal yr BP, which roughly coincides with the second occupation hiatus. Again, the only chronometric data available for Unit IIc is a humate sample (Beta-112980) and may likely date more recent than indicated.

Guadalupe-San Antonio River Basin

For the Guadalupe-San Antonio River Basin, the dissimilarities may be more informative than the similarities. Although the internal comparisons suggest more similarities between the upper and lower extent, than the middle extent these are generally rare. The Jonas Terrace site exhibited similar periods of erosion-incision as those reported at Copano Bay during 4,250–3,000 and 2,500–2,250 cal yr BP (Figure 8.2). Surprisingly, none of the periods of instability or stability seemingly overlap

between the Richard Beene site and either the Jonas Terrace site or the Copano Bay study area. The depositional history at the Richard Beene site appears to be unique.

Brazos River Basin

The recalibration results in the Brazos River basin seemingly demonstrated the most intra-basinal patterns of this study. This may be attributed to the abundance of data over a broad geographical range within the basin. Regardless, several phenomena became apparent when the upper, middle, and lower extents were compared.

Beginning with the Late Pleistocene, the proposed period of erosion-incision is present in two of the Brazos River basin study areas. The exception is the A&M study area in the lower extent. This dataset had a suspicious assay dating to roughly 21,000 cal yr BP with no observed disconformities until after 9,300 cal yr BP. Regarding the upper and middle extents, the ending of this Early Holocene erosive event has a different time in each of the study areas, these differences may be due to sampling. However, it is interesting to note that this erosive cycle ended at 13,080 cal yr BP at Lubbock Lake, while Fort Hood has a more recent terminus of 10,100 cal yr BP.

Comparing the three study areas, the most prominent phenomena are periods of incision, hiatus, or data gaps in the radiocarbon record. The middle and lower extents exhibit the most similarity with two periods of overlap at 8,750–7,750 cal yr BP and at 5,750–5,250 cal yr BP. Notable, are overlaps in all three basin areas at 5,250–5,000, 4,000–3,250, and 2,750–2,500 cal yr BP. While these phenomena may in part be attributable to sampling the pervasiveness of the similarities suggests there may be a common synchronous mechanism.

Regarding periods of pedogenesis across the basin, the similarities are surprisingly few (Figure 8.2). At Fort Hood, the Royalty Paleosol roughly correlates

with the A&M paleosol at the A&M study area. At Lubbock Lake, the Lubbock Lake soil of Stratum 4B and the Apache soil of Stratum 5A roughly overlap with the Asa and Katie soils, respectively at the A&M study area. However, these are the only similarities suggesting a regional period of stability.

Inter-Basinal and Regional Patterns

Widening the examination, the comparison of the different drainage basins within Texas and region demonstrates some interesting patterns. The inter-basinal comparisons were examined by three different categories (i.e., tributaries, trunk systems, and regional). First, the tributary systems of each of the basins were contrasted for any apparent patterns (Figure 8.2). All of these tributaries were on the Edwards Plateau, affording a comparison with the next comparison category, trunk systems. Trunk systems are the main channels of each of the basins (Figure 8.3). The study areas of each trunk system were on the Coastal plain and off the Edwards Plateau. The third category is a cumulative overview of all the depositional histories of all the Texas drainage basins in this study.

An examination of the tributary drainages in this study suggests some patterns in depositional history in the Edwards Plateau (Figure 8.2). Only three study areas were compared for this tributary comparison, the Cowhouse Creek (Brazos River basin), the Concho River (Colorado River basin), and the South Fork San Geronimo Creek (Guadalupe-San Antonio River basin). Unfortunately, the chronology at the Jonas Terrace site on the South Fork San Geronimo Creek begins at 4,250 cal yr BP and does not extend as far back as the other tributary study areas. Two earlier periods of instability are apparent at the other tributaries. At the Concho River and the Cowhouse Creek study




Figure 8.5 Patterns along Trunk Portions of Texas Drainages in Study.

areas, periods of instability occurs at 8,000–7,750 and 5,750–5,250 cal yr BP (Figure 8.2). At all three drainages two periods of instability are inferred at 3,750–3,250 and 2,750–2,250 cal yr BP. Notably, no periods of synchronous stability seem to have occurred at anytime among these tributaries on the Edwards Plateau.

Four study areas were selected for the comparison of Texas drainage basin trunk systems, the Brazos River, Colorado River, Medina River (Guadalupe-San Antonio basin), and the Trinity River. Beginning with the Late Pleistocene, the Colorado and Trinity Rivers exhibit a period of erosion-incision between 19,000–15,940 and 14,500–13,500 cal yr BP, respectively. Interestingly, neither the Brazos nor Medina Rivers exhibit this period of incision prior to the Holocene. The Medina River system may have an incision event prior to 15,900 cal yr BP, which is the *terminus ante quem* for Unit A3 (Perez Horizon). However, no definitive indication is apparent. The Brazos River exhibits continuous deposits up to 21,000 cal yr BP. However, this is based on a single radiocarbon assay from the margins of the floodplain (Waters and Nordt 1995). The applicability of the assay as representative of the Late Pleistocene is dubious. Regardless, deposition is occurring in all four basins by 13,500 cal yr BP and continues until 9,750 cal yr BP.

The first period of synchronous instability (Synchronous Event I) represented in all four basins occurred around 8,750–8,250 cal yr BP. This instability likely encompassed a more extensive period, but the Medina and Trinity River basins suggest normal deposition after 8,250 cal yr BP. The next period of synchronous instability (Synchronous Event II) in all four basins begins at roughly 7,000 cal yr BP and extends until 6,250 cal yr BP. As previously mentioned, several periods of transgression are

noted to have occurred in the Early-Middle Holocene. These occurred at roughly 6,800–5,900 cal yr BP (Ricklis and Blum 1997; Paine 1991). Although the period of instability overlaps the period of sea level rise, the concurrence of the two phenomena may imply a correlation.

The next period of synchronous instability (Synchronous Event III) is not as evenly distributed as the other trends and the timing of it is approximate. Specifically, between roughly 5,250–5,000 cal yr BP, there is a period of instability occurring in these four basins. The one tenuous exception may be the Medina River basin, which has a period of stability ending about that time followed by a period of instability. While there does appear to be instability centered around 5,100 cal yr BP, but the duration of it in each basin varies considerably. At this time, the Trinity and the Brazos River basins seem to exhibit the most instability with the Medina and Colorado River basins the least. Previous researchers have noted a paucity of alluvial deposits at roughly this time in Texas and the Southern Plains (Baker et al. 2000; Hall 1990a:343). The scarcity of deposits dating to this time may imply a similar period of instability in those areas. Also, northward in the Great Plains, Schmieder (2008) examining lake sediments in the Sand Hills of Nebraska and observed an extensive period of drought beginning at this time. More locally, Nordt (2004) and other researchers in their examination of C₃ and C₄ isotopes at the Richard Beene site interpreted a decrease in C₄ and a brief cool period at this time. While Cooke (2003) and other researchers propose that the mantle in the uplands of the Edwards Plateau had an intense period of erosion. Not coincidentally, this phenomenon also is squarely in the midst of the enigmatic Middle Archaic, which has long been recognized to have a comparatively lower frequency of sites than the other

Years BP*	Northern Plains Bettis 2003	Republican River SW Nebraska (Daniels and Knox 2005)	C ₄ and Climate South-Central Texas-Richard Beene (Nordt et al. 2002)	Midwest-Climate Change Alluvial Settings (Baker et al. 2001)	OK Panhandle Bement et al. 2007	Midwest-Climate Multi-proxy (Baker et al. 1998)	Texas Bog Pollen (Bousman 1998)	North-Central Texas Climate from Shell Isotopes (Brown 1998)	Paleoclimate Review Mayewski et al. 2004	Bison in Texas (Dillehay 1974)	
0 k		Incision					open woodlands		rapid climate change	bison absence; Absence Period III	
		Rapid alluviation								bison presence; Presence Period III	Synchronous Event IV
		Incision					open woodlands				
1 k		Primarily 1100 cal yr BP							rapid climate change	bison absence; Absence Period II	
			Increase in C ₄ plants; cool interval correlates with regional tree pollen				non-arboreal plant pollen predominate				
2 k	Major aggradation					increase in C ₄ ; forest to prairie		cooler and wetter conditions prevail			
									rapid climate change	bison presence; Presence Period II	
3 k						decrease in C ₄ ; prairie to forest		brief cool-wet period			
									rapid climate change		
4 k	Major aggradation										
			brief cool period; decrease in C ₄				gradual increase in non-arboreal pollen (i.e., grasslands); approx. peak 5,500 14c yr BP				
5 k									rapid climate change		Synchronous Event III
			Increase in C ₄ ; warm & dry Period			increase in C ₄ ; forest to prairie				bison absence; Absence Period I	
6 k							brief arboreal pollen increase				Synchronous Event II
							non-arboreal pollen increase				
7 k	Major aggradation				Stability						
8 k			Drastic decrease isotopic values suggesting cooler conditions		Stability		pollen suggests open woodlands		rapid climate change		Synchronous Event I
9 k					Stability		non-arboreal plant pollen predominate				
							dramatic change				
10 k				stability			pollen suggests open woodlands			bison presence; Presence Period I	
11 k			Higher temps; decrease in moisture	discontinuity	Stability						

Figure 8.4 Climatic Patterns

cultural periods (Collins 2004). Cumulatively, this all suggests a pervasive synchronous event of instability across the region.

The last synchronous event (Synchronous Event IV) exhibited in all four basins is a period of stability (i.e., pedogenesis). Although periods of soil development are recognized to have occurred sporadically in all of the basins, none of these align except at 1,000–750 cal yr BP (Figure 8.3). Longer periods of stability seemingly occurred in the Brazos and Trinity River basins beginning as early as 1,500 cal yr BP while the Colorado River basin exhibits the weakest correlation. A period of stability at roughly 1,000 cal yr BP is very widespread and has been noted by previous researchers in alluvial settings in Iowa and Missouri (Bettis 2003; Bettis and Mandel 2002:145), in the Kansas River system (Arbogast and Johnson 1994), and possibly in alluvial settings across Oklahoma and Texas (Hall 1990a). Unfortunately, the dataset for Hall's (1990a: Table A) investigations were not corrected for isotopic fractionation and was not calibrated for this study. However, all other descriptions of these depositional histories are in accord with the period of stability observed in the four basins. Furthermore, Collin's compared archaeological sites with stratified cultural and geological horizons (i.e., gisements) compared in nine locations (Collins 1995:374, 2004:111). This comparison revealed a pervasive episode of stability around this time (Collins 1995: Table 2, 2004: Figures 3.9a and 3.9b). While Collin's (1995, 2004) data are reported in radiocarbon years BP, most of these nine study areas are part of the current recalibration study (e.g., Fort Hood and Richard Beene) suggesting an equivalency can be made for the areas not recalibrated (e.g., South Bend Reservoir on Clear Fork of the Brazos River). Thus, there appears to be a widespread synchronous episode of stability at 1,000–750 cal yr BP across these

four basins and into alluvial settings on the Great Plains.

Causal Factors

For over a century researchers have examined the relationship between reactions in fluvial systems in response to external effects (e.g., Bull 1991, 2000; Knighton 1998; Knox 2000; Schumm 1993, 2003). See Blum and Törnqvist (2000) for a detailed examination of climate and sea level effects on drainages. However, a brief review follows to provide a general framework for the processes of these interrelated causal relationships.

Four factors are generally recognized to be the primary external influences on fluvial systems: tectonic activity, glaciation, climate change, and eustasy (Blum 1993: Table 1; Blum and Straffin 2001:195; Bull 1991, 2000; Knighton 1998; Schumm 2003). These factors can operate individually or in convergence and the sensitivity of the basins to these factors are filtered through a variety of controls including localized geology. At its simplest level, these factors influence the fluvial response stratigraphically (i.e., the storage or removal of sediment), morphologically (e.g., channel width, sinuosity), and deposition (i.e., bedload) (Blum 2007; Blum and Straffin 2001). An often-overlooked factor influencing drainages is anthropic mechanisms such as cultivation or vegetation removal (Frederick 1995). However, in Texas only the historic era to the present would appear to be affected by this factor.

Considering tectonic activity and glaciation have had little effect to the Texas drainage networks of the Late Quaternary, this leaves climate and eustatic influences as likely causes of the observed patterns. To be sure, tectonism is a factor of influence in Texas basins, but it almost exclusively takes form as a slow subsidence (i.e., 0.05 mm a

year) of the continental margin (Paine 1993). Similarly, glaciation has not been proposed to affect the Texas waterways other than as melt water pulses, which ultimately defaults to sea levels and/or the climate. Consequently, researchers almost exclusively focus on climate and eustasy and the responses of drainages in Texas and the region. The influence of these two factors will be considered by reviewing the interpreted paleoenvironmental record and sea levels proposed by previous researchers.

Climate

The review of the climate was accomplished by compiling proxy data (e.g., bog pollen, speleothems, and various isotopic analyses) from several regional and global studies used to reconstruct the paleoenvironment for the late Pleistocene and Holocene in Texas (Bousman 1992, 1994, 1998a; Cooke et al. 2003; Nordt et al. 2002; Toomey et al. 1993). Admittedly, not all data are unanimously accepted and there are gaps in the record, but a preponderance of researchers generally accept the review provided here.

Subsequent to the last glacial maximum approximately 23,500 cal yr BP, the climate has been interpreted to have had cooler temperatures and more mesic conditions for the Central Texas region, South Texas Plains, and Texas coastal plain (Bousman 1998; Bryant and Holloway 1985; Bryant and Shafer 1977; Hudler 2000; Musgrove et al. 2001; Nordt et al. 2002, 2007; Sylvia and Galloway 2006; Toomey et al. 1993). These cool and mesic conditions prevailed until 15,000 cal yr BP and again around 12,000 cal yr BP, when pollen and isotopic analyses data suggest that glacial melt waters entered the Gulf of Mexico and triggered arid and presumably cooler conditions in southern and Central Texas (Bousman 1998:214; Nordt et al. 2002:182). This assessment is further supported by low growth rates on speleothems from dated stalagmites in several central

Texas caves, implying more xeric conditions (Musgrove 2000; Musgrove et al. 2001). This also correlates with investigations in the Southern Plains where Holliday (2000) argues that climatic oscillations occurred around the Late Pleistocene-Holocene transition with cooler and moister conditions for the Clovis period and particularly arid and warmer conditions during the Folsom period. Based upon the recalibration of the Lubbock Lake data for this study (see Chapter 6), this suggests that the climate was cooler and mesic at roughly 13,250–12,750 cal yr BP and conditions were most arid between 12,500–11,500 cal yr BP. The more recent xeric period was followed by a shift back to cooler temperatures and moist conditions in central and southern Texas, which continued into the Early Holocene (Bousman 1998:214).

From the Early to Middle Holocene (~ 11,500–5,000 cal yr BP), the proxy data suggest that the climate became gradually warmer and more xeric (Bryant and Shafer 1977; Toomey et al. 1993). These data include pollen evidence suggesting a decrease in arboreal canopy and open grassland for central and south central Texas (Bousman 1998), various fauna indicator species from cave deposits (Hudler 2000; Toomey et al. 1993), the presence or extinction of various Molluscan fauna (Neck 1983, 1987), and shifts in C₃–C₄ plant production (Nordt et al. 1994, 2002). At this time, Greenland ice core data suggests abrupt climatic changes in the climate at 8,900–8,300 and 8,200 cal yr BP (Hu et al. 1999; Yu and Wright 2001). In conjunction, Barber and others (1999) argue that the Laurentide ice sheet had an abrupt reduction in size and flushed massive amounts of freshwater into the Labrador Sea around 8,400 cal yr BP, which they argue triggered the climatic changes between 8,400–8,000 cal yr BP.

Further, Mayewski and others (2004) examined over 50 paleoclimate records in

the Northern and Southern hemispheres from around the world. These researchers observe a period of rapid climate change between 9,000–8,000 cal yr BP, which in North America is exhibited by rapid glacial advances in the northwest and the previously mentioned surge of melt water (Mayewski et al. 2004:248–249). In an alluvial setting (South Fork of the Big Nemaha River) in southeastern Nebraska on the Great Plains, Baker and others (2000) propose that a disappearance of upland forests and an extended period of dry conditions occurred between 9,200–6,500 cal yr BP. More locally, Dillehay (1974) in researching the presence or absence of bison in the Southern Plains inferred an extended period of absence beginning around ~7,900 yr BP (6000 BC) that coincides with this warming period. These warming and xeric conditions existed throughout this time with some minor deviations and probably localized variations (Hudler 2000:88–89).

One anomaly of note is a very brief episode of moister conditions in southern and central Texas occurring ~6,000 yr BP as evidenced by an increase in arboreal pollen and data from isotopic composition of organic and inorganic carbon (Bousman 1998; Nordt et al. 2002:186). This brief cool and moist episode was immediately followed by an extremely arid and warm climate (Bousman 1998; Nordt et al. 2002). This xeric period lasting roughly 1,000 years, was exhibited by a drastic reduction in arboreal pollen and an increase in grassland pollen (Bousman 1998). Further evidence of these xeric conditions, is the reappearance of bison on the Southern Plains beginning around ~4,500 cal yr BP or 2500 BC (Dillehay 1974). Also in the Great Plains, Baker and others (1998) in their examination of pollen and plant macrofossils in northeastern Iowa, note a rapid change in vegetation from a forest to prairie setting at 6,000 cal yr BP and they observed the percentage of C₄ values reached a peak around 5,000 cal yr BP. More broadly,

Mayewski and others (2004:250) observe rapid climate change globally between 6,000–5,000 cal yr BP where central North America experiences a strengthening of westerly winds among other widespread climatic events.

After this arid and warm period extreme, the Late Holocene climate is described as gradually increasing in moisture and cooling in temperature (Bousman 1998; Nordt et al. 2002, 2007; Butzer et al. 2008). Johnson and Goode (1994) in their examination of the Jonas Terrace site also propose that conditions were becoming more mesic and cooler, but they have it occurring around 3,850 cal yr BP (1900 BC) and roughly extending until 1,950 cal yr BP (0 BC). The mesic indicators of this time were exhibited through a gradual increase in woodland canopy and data from stable isotope analyses in buried soils (Bousman 1998; Nordt et al. 2002, 2007). Nordt and others (2007:159) characterize this period as a ‘cool interlude’ before conditions again transition into a more xeric and warm climate. This arid interval extends from roughly 2,600–1,000 cal yr BP before again becoming slightly more mesic and continuing as such up to the present (Nordt et al. 2007). Coinciding with these swift transitions from mesic to xeric conditions includes the absence of bison in the Southern Plains between roughly (~1,500–950 cal yr BP (AD 500–1200) and subsequent reappearance between ~950–400 cal yr BP or AD 1200–1550 (Dillehay 1974).

More broadly, at 4,200–3,800 cal yr BP glaciers advanced in western North America and central North America had intense westerly winds, which weakened at 3,500–2,500 cal yr BP (Mayewski et al. 2004:250). Rapid climate changes are also indicated globally at 1,200–1,000 cal yr BP manifesting as cooler temperatures in the Sierra Nevada mountains based on tree ring data while between 600–150 cal yr BP a

period of polar cooling and increased moisture in the tropics occurred (Mayewski et al. 2004). In an alluvial setting in southeastern Nebraska, an abrupt, but brief disappearance of riparian trees indicates arid conditions at 3,360–2,910 cal yr BP (Baker et al. 2000). Further, Huckleberry and Duff (2008) note in western New Mexico that increased moisture caused widespread valley entrenchment around ~1050–900 yr BP (AD900–1050) and 650–550 cal yr BP (AD 1300–1400) with the latter followed by an extended period of arid conditions. Locally, the last 1,000 years are indicated to have some brief fluctuations of arid conditions occurring around before trending toward modern climates of the present (Bousman 1998:216).

Eustasy

The effects of changes in worldwide sea levels (eustasy) are more limited on drainage systems than that of climatic changes. Researchers have investigated various aspects of rising and falling sea levels at global and local levels and the distance upstream of those influences have on drainage basins. There is considerable debate regarding the influence eustasy has on a drainage system and to what degree (Schumm 1993). At its simplest form, the influence is generally interpreted to result in as drainage incision (down cutting) for lowering sea levels (regression) and avulsion and aggradation for a rise in sea levels (transgression) (Anderson et al. 2004; Banfield and Anderson 2004; Blum 1993; Blum and Aslan 2006; Blum et al. 2001; Blum and Price 1998; Blum and Törnqvist 2000; Durbin 1999; González 2008; González and Törnqvist 2009; Leeder and Stewart 1996; Schumm 2003; Thomas and Anderson 1994; Törnqvist et al. 2004; Van de Plassche et al. 1998; Zaitlin et al. 1994). Regarding the distance upstream the influence of sea level has on a drainage basin, one of the primary factors is the slope of the coastal

plain. In periods of regression when the coastal plain is roughly equal to that of the coastal shelf, there should be a slight extension of the channel onto the coastal shelf. If the coastal plain is noticeably steeper than the coastal shelf the channel will extend and aggradation of deposits will occur, and if the coastal plain is shallower than the coastal shelf, then the channel will extend in conjunction with dramatic incision (Blum and Törnqvist 2000; Schumm 1993:281–282). For Texas, the coastal plain is generally steeper than the coastal shelf.

Furthermore, the effects of sea level changes may be severely limited in coverage to the drainage basins. Specifically, previous researchers indicate that eustatic effects on drainages are generally limited to within 100 km (62 miles) of the coast, which falls within the lower extent of all the examined basins of this study (Blum 1993; Durbin 1999; Etheridge et al. 1998). Most confounding is discerning the difference between influences of eustatic effects versus climate within a basin, particularly within the lower extent. Researchers Blum and Aslan (2006) have proposed criteria for determining the influences of climate versus sea level change on drainage valleys. They indicate that climatic influences should be exhibited by stratigraphic units that extend from mixed-bedrock valleys (upstream) across the coastal plain to the distal reaches (downstream) of a basin. There is a recognized continuity of facies architecture throughout the drainage basin particularly if the climatic influence is major. In contrast, the signature of sea level influences on a drainage basin is more complex. During a drop in sea levels, there is incision within the drainage, which may result in a valley separation on the coastal plain concurrent with lateral migration and meander belt construction. The mixed-bedrock valleys will incise with periodic lateral migration and creation of terraces while a

rise in sea levels trigger a shortening of the channel and expansion of the deltas.

Notably, Blum and Aslan do not identify characteristics for the mixed-bedrock valleys upstream during the periods of sea level rise.

Previous investigations within the Gulf of Mexico have characterized the Late Quaternary stratigraphy, but unfortunately, most of these focus on periods much older than the latest Pleistocene. As with the paleoclimate interpretations, not all data for sea levels are widely accepted and are frequently contradictory. The following review includes some of the more recent investigations in the region, which largely concur in their interpretations.

Since the Last Glacial Maximum (~23,500 cal yr BP) in the Gulf of Mexico, researchers have identified at least two pulses of glacial melt water that entered the gulf prior to the Holocene (Figure 8.5). Fairbanks and others (1989) examined coral reefs in the Caribbean and argue that these pulses occurred at 13,500–13,000 cal yr BP and 11,000–10,500 cal yr BP. Off the Texas coast, Snow (1998) examined near shore core samples of the Colorado River delta and radiocarbon data from previous investigations and observed the first melt water pulse (MWP 1A) occurred at roughly 14,500–13,750 yr BP. This first pulse produced a rise in sea level of roughly 36 mm a year. The second melt water pulse (MWP 1B) occurred at roughly 12,000–11,500 yr BP and produced a sea level rise of approximately 16 mm a year. Snow (1998:129–131) characterized these pulses as producing high sediment yields that were primarily controlled by climate. Following the second melt water pulse, the sea level continued to gradually rise at a roughly constant rate of 10 mm a year between 11,500–5,000 cal yr BP. Snow (1998) observed a transition from fluvial dominated deposits of the Colorado River to wave

Years BP*	Nueces Bay data (Ricklis and Blum 1997)	IntCal 09 Curve Copano Bay Area (Paine 1991)	Colorado Delta data (Snow 1998)**	Mississippi Delta (Gonzalez 2008; Gonzalez & Tornqvist 2009)	Mississippi Delta (Tornqvist et al. 2004)	Colorado River Delta-Copano Bay (Blum et al. 2001)	Ice Core Data (Alley et al. 1997)	Paleoclimate Review Mayewski et al. 2004	Carribbean Coral reefs (Fairbanks et al. 1989)	
0 k				sea level drop				rapid climate change		
				sea rise peak						Synchronous Event IV
1 k				Slow sea level rise begins 1350 yr BP				rapid climate change		
2 k		Incision								
		Incision								
3 k	rapid sea level rise; coincides with coastal occupation hiatus	Incision			Gradual sea level rise about 1.5 mm/yr			rapid climate change		
4 k		Drop in Sea Level; Regression						rapid climate change		
5 k	rapid sea level rise; coincides with coastal occupation hiatus	Third transgressive pulse of sea level rise	Eustatically influenced Phase 2: Wave deposits dominate; sea level rise 10mm/yr; climate continues to be warm-dry			Younger 14c data suggests sea level rise about 3-4 mm yr; rose above modern sea level by 2 m		rapid climate change		Synchronous Event III
6 k										Synchronous Event II
7 k						Generally gradual rise about 3.5 mm/yr; interpreted to be end of glacial melt water pulse	Older 14c data suggest sea level rise interpreted to be 9-11 mm/yr			
8 k							extensive dramatic cold-dry cycle			Synchronous Event I
9 k										
10 k			Climatic influenced Phase 1: Fluvial deposits dominate; sea level rise 10mm/yr; transitions from cool-moist to warm-dry climate							
									large glacial melt water pulse	
11 k			Meltwater Pulse 1B; sea rise 16mm/yr							

Figure 8.5 Eustatic Patterns

dominated deposits (i.e., eustatic) around 9,500 cal yr BP. Considering the rates of sea level between 11,500–5,000 cal yr BP were rising at an equivalent level (i.e., 10mm/yr) this transition is interpreted as evidence of a shift from climate influence to eustatic influence.

More controversially, Blum and others (2001) in their examination of the Colorado River delta and nearby Copano Bay interpreted a dramatic rise in sea level exceeding modern mean sea levels (msl) by over 2 m. The researchers interpreted this transgression to have occurred between either 7,800–6,800 cal yr BP or 7,800–4,800 cal yr BP depending on the dataset used (Blum et al. 2001). Specifically, a few samples are indeterminate if they are affected by younger calcite thus providing two datasets (Blum et al. 2001:586). In contrast, Törnqvist and others (2004) analyzed deposits of basal peat in the Mississippi delta, which are typically deposited in coastal settings between the msl and high water mark. These researchers encountered deposits between 8,000–3,000 cal yr BP that exhibited a gradual rise of sea level with no drastic jumps in sea level. A slight bump in sea level rise (3.5 mm/yr) did occur between 8,000–7,000 cal yr BP followed by a decrease with a sea level rise of 1.5 mm a yr up to 3,000 cal yr BP. Further, no indications of sea levels exceeding modern msl were observed, but they did not exclude the possibility of it occurring between 3,000 cal yr BP and the present.

In the Middle Holocene, the previously mentioned research by Ricklis and Blum (1997) interpret a rise in sea levels that coincides with a hiatus in coastal occupations by native groups. The recalibration of these data (see Chapter 3) did not adjust the initial conclusions of sea level rise occurring at 6,800–5,900 and 4,200–3,000 cal yr BP (Ricklis and Blum 1997). In the Copano Bay area of the Guadalupe-San Antonio River basin, the

previously mentioned research of Paine (1991) examined a variety of datasets (e.g., sea cores, trench profiles, archaeological investigations) to characterize the Late Quaternary deposits of the bay. Unfortunately, only a portion of these data could be recalibrated for this study. Therefore, Paine's (1991) interpretations prior to 6000 cal yr BP are not calibrated. Regardless, three transgressive pulses are recognized to have occurred during the Holocene. With only the last transgressive pulse could be recalibrated (see Chapter 4). Briefly, the first pulse is described as a transition from fluvial (i.e., stream) to marine (i.e., sea) influenced deposition implying a rapid rise in sea level followed by a return to fluvial deposition interpreted to be a period of stillstand or possible drop in sea level (Paine 1991:61–64). The second pulse is again suggested by a transition from stream deposition to marine deposition inferring another sea level rise shortly followed by a transition back to a dominant stream deposition suggesting another stillstand.

The third pulse was recalibrated for this study and dates to 5,750–4,750 cal yr BP. Interestingly, this roughly correlates with the sea level rise interpreted by Ricklis and Blum (1997). Paine (1991:64) characterized this last transgressive pulse as a transition from stream to marine deposition suggesting a slow sea level rise. Of note, this rise in sea levels rose approximately 0.9 m above modern sea levels (Paine 1991:170–171). However, this interpretation is in the minority, as most researchers do not interpret a sea level rise of that magnitude at this time. Subsequent to the last transgressive pulse a sea level stillstand begins that continues to the present (Paine 1991:57). Simply put, these three pulses are argued to be evidence for eustatic effects. If correct, only the lower portions of the basins would be affected.

For the Late Holocene, extensive investigations have recently been conducted in the Mississippi River delta (e.g., González 2008; González and Törnqvist 2009). These investigations analyzed basal deposits of peat, which, as previously mentioned, are interpreted to be deposited between the mean sea level and high water mark (González and Törnqvist 2009:1739). Additionally, the peat provided the source for the radiocarbon assays that enabled high-resolution chronological control. These researchers observed a gradual rise in sea levels beginning around 1,350 cal yr BP (AD 600) that peaked around 850 yr BP (AD 1100) and declined until roughly 450 cal yr BP or AD 1500 (González 2008; González and Törnqvist 2009: Figure 11).

Correlations

A comparison of the recalibrated data from this study will now be conducted with the available information of climate and eustatic factors. Understandably, the recognition of contemporaneous events does not confirm a correlation. However, the intent of this component of the study is to identify areas and temporal periods where more research may be warranted.

The first correlation study is of the four synchronous events observed within the four basins identified during the Inter-basinal comparisons. These synchronous events consist of three apparent periods of instability at 8,750–8,250, 7,000–6,250, and 5,250–5,000 cal yr BP and one period of stability between 1,000–750 cal yr BP.

Synchronous Event I

For the first period of instability (8,750–8,250 cal yr BP), several studies indicate pervasive events occurring immediately preceding or contemporaneous with this phenomenon. In particular, an abrupt change in the global climate occurred between

9,000–8,000 cal yr BP likely triggered by a large pulse of glacial melt water (Alley et al. 1997; Barber et al. 1999; Hu et al. 1999; Mayewski et al. 2004). In the paleoenvironmental research of the Oklahoma Panhandle along an alluvial setting (Bull Creek), Bement and others (2007) note a period of pedogenesis suggesting stability at roughly 8,460 cal yr BP. Similarly, in an alluvial setting (South Fork of the Big Nemaha River) in southeastern Nebraska on the Great Plains, Baker and others (2000) propose that a disappearance of upland forests occurred between 9,200–6,500 cal yr BP. Also, bog pollen in Texas exhibits a transition from more diverse forests to open woodlands while C₄ isotopic values at the Richard Beene site decreased, all inferring a cool, moist climate at this time (Bousman 1998; Bousman and Oksanen in press; Nordt et al. 2002). Interestingly, this is also roughly the end of Dillehay's (1974) Presence Period I for bison in the Southern Plains. Regarding eustatic effects, an abrupt increase of sea level occurs at roughly this time, but this interpretation is not widely accepted (Blum et al. 2001; Törnqvist et al. 2004).

Considering the archaeological record in Texas at this time, this synchronous event falls at the transition from Late Paleoindian to Early Archaic archaeological period or at the beginning of the Early Archaic archaeological period (Figure 8.6). A pervasive geomorphic period of instability seemingly occurs roughly simultaneous to a period of transition within the archaeological record. McKinney (1981:114) does interpret changes in Central Texas Archaic lithic technology as adaptation to environmental changes. Contemporaneity of events does not indicate causality and reaction, but these concurrent phenomena do warrant some attention.




Figure 8.6 Calibrated cultural chronology contrasted with Synchronous Events I-IV (adapted from Mehalchick et al. 2000).

Synchronous Event II

The second period of synchronous instability (7,000–6,250 cal yr BP) is the longest of those identified. This period immediately precedes an abrupt climate change in the global record (Mayewski et al. 2004). In Nebraska of the Central Plains, an extended period of drought occurred at this time while in the Midwest, various data (e.g., speleothems and alluvium) suggest a major climatic transition near the end of this second period of instability (Baker et al. 1998; Baker et al. 2001). Similarly, in alluvial settings in Iowa and Illinois an extended period of drainage aggradation is indicated for this time (Bettis 2003). Within Texas, the bog pollen data suggests a transition to non-arboreal plants inferring grassland prairies and by extension xeric conditions (Bousman 1998).

Notably, the previously mentioned ‘cool interlude’ associated with a short-lived increase in arboreal pollen and data from isotopic composition of organic and inorganic carbon occurred at ~6,000 cal yr BP (Bousman 1998:210; Nordt et al. 2002:186). However, this interlude was preceded by a period increased of $\delta^{13}\text{C}$ values denoting xeric conditions, which concurs with the second period of instability (Nordt et al. 2002). Interestingly, this period is noted as a time of alleged bison absence (Absence Period I) (Dillehay 1974). However, occurrences of bison have been encountered on archaeological sites (41HY160 and 41HY165) in Central Texas clustering around 5,900–5,700 cal yr BP (Lohse 2010). The presence of bison at that time implies xeric conditions, which concurs with the pollen and isotopic data. Concerning eustatic effects, no prominent rise or drop in sea levels are recorded for this time.

Synchronous Event II occurs roughly in the middle of the Early Archaic (Figure 8.6). This period of instability does not appear to coincide with any obvious widespread cultural change.

Synchronous Event III

The third identified period of synchronous instability (5,250–5,000 cal yr BP) has long been recognized as a time of geomorphic change likely accounting for the paucity of Middle Archaic data (Collins 2004:115; McKinney 1981). Mayewski and others (2004) have identified widespread evidence of abrupt climate change occurring at this time. Some indications of these include large-scale glacial advances in the northern and southern hemispheres and an increase in aridity in the Maya lowlands of Central America among others (Hodell et al. 2001; Mayewski et al. 2004). Further, considering that no glacial melt water pulses are interpreted to occur at this time, researchers propose that these climatic fluctuations are possibly attributed to solar variability (Hodell et al. 2001; Mayewski et al. 2004:251). In northeastern US, dramatic shifts in pollen are noted, which are attributed to repeated droughts (Shuman et al. 2009). More locally in the Great Plains, lake sediments suggest a dramatic shift from a wet cycle to drought conditions while assessments of the stable isotopic compositions of buried soils indicate warmer temperatures at roughly this period (Nordt et al. 2007; Schneider 2009).

Within Texas, the bog pollen data indicates the lowest percentage of arboreal pollen occurred at roughly 5,500 yr BP suggesting grassland prairies and xeric conditions (Bousman 1998:210). Further, the soil mantle in the uplands of the Edwards Plateau is recorded to be ending an extended period of erosion at roughly this time (Cooke 2006; Cooke et al. 2003). Using data from Hall's Cave in Kerr County, the researchers propose

that the Central Texas uplands were generally emptied of its soil mantle beginning in the Late Pleistocene and ending roughly concurrent with this third period of synchronous instability (Cooke 2006; Cooke et al. 2003). One corroboration of this may be present as Nordt (1996:16–17) recorded a depositional history at Leon Creek (Guadalupe-San Antonio River basin) situated at the base of the Edwards Plateau, which prior to this time was composed of both alluvial and colluvial deposits and switched to become more gravelly with no evidence of colluvium afterwards.

Large fluctuations in sea level are also interpreted to have occurred at this time. Ricklis and Blum (1997) recognize a short-lived rise in sea level that overlaps with the third period of synchronous instability (Figure 8.5). Further, this sea level rise coincides with a hiatus in cultural occupations along the coast (Ricklis and Blum 1997). Similarly, the recalibrated data of Paine's (1991) Copano Bay research also indicates a period of sea level rise at this time. As previously mentioned in the Nueces River basin (Chapter 3), similar gaps in chronometric data were recognized during the Choke Canyon Reservoir investigations. These gaps are roughly contemporaneous with this period of sea level rise. However, as previously indicated eustatic effects on drainages are generally limited to within 100 km (62 miles) of the coast, which puts the Choke Canyon Reservoir right at the limits of eustatic effects (Blum 1993; Durbin 1999; Etheridge et al. 1998).

Concerning the Texas archaeological record, this synchronous event falls roughly at the beginning of the Middle Archaic (Figure 8.6). Thus, another pervasive geomorphic period of instability coincides with a period of transition within the archaeological record. Further, examining Black and Creel's (1997) radiocarbon chronology of burned rock middens, this event is situated near the beginning of an extended period of increased

burned rock midden use (Figure 8.7). However, the assays of the burned rock midden study have not been calibrated for this study and this comparison is approximate. It is probable, that Synchronous Event III will fall within the radiocarbon gap preceding the increase in burned rock midden exploitation after the assays have been calibrated with IntCal09.

Synchronous Event IV

The final period of synchronous geomorphic activity is a period of pervasive stability between 1,000–750 cal yr BP. Oddly, this period has the most contradictory climatic data possibly because of the plethora of investigations (Figure 8.4). In the western US, a period of cooler temperatures and drought are indicated by tree ring data in the Sierra Nevada Mountains occurring at AD 892–1112 (1,058–838 cal yr BP) and AD 1209–1350 (741–600 cal yr BP), which are argued to coincide with the Mediaeval Warm Period (Mayewski et al. 2004; Stine 1994). In western New Mexico widespread valley entrenchment from an increase in moisture is indicated between roughly 1,050–900 cal yr BP (Hall 1990b; Huckleberry and Duff 2008). Similarly, on the Republican River in Nebraska, after a period of pedogenesis ending at roughly 1,100 cal yr BP, a period of drainage incision occurred between 1,100–750 cal yr BP (Daniels and Knox 2005). However, a period of prolonged pedogenesis is recognized at this time in alluvial settings in the east-central Plains (Johnson and Martin 1987), in the Kansas River basin in the central Plains (Arbogast and Johnson 1994; Johnson and Logan 1990: Figure 9), and in the previously mentioned 15 alluvial settings studied by Hall (1990a).

Locally in Texas, the bog pollen data suggests a period of open woodlands while bison are indicated to have been present during Presence Period III (Bousman 1998;




Figure 8.7 Radiocarbon dates from burned rock middens on Edwards Plateau suggesting frequency of use (adapted from Black and Creel 1997).

Dillehay 1974). Regarding eustatic effects, only recent investigations in the Mississippi River delta provide any information from this time (González 2008; González and Törnqvist 2009). These data from radiocarbon dating basal peat deposits enabling high-resolution chronological control suggested a gradual sea level rise beginning around 1,350 cal yr BP and peaked between 1,000–750 cal yr BP (González 2008; González and Törnqvist 2009).

Similar to the diverse paleoenvironmental results, the archaeological record in Texas at this time is extremely varied (Pertulla 2004: Table 1.1). Regardless of archaeological region, the synchronous period of stability seemingly occurs at a time of transition across Texas. Furthermore, comparing Synchronous Event IV again with Black and Creel's (1997) burned rock midden frequency, the peak of midden use appears to coincide with this period of stability (Figure 8.7). This high frequency of burned rock midden use may be reflective of improved integrity of this period of geomorphic stability. Again, these phenomena warrant further investigation.

Summary

The results of the recalibrated datasets from the drainage basins were consolidated and examined for patterns both within and between the Texas basins and, to a lesser degree, the region. A review of extrinsic factors that possibly influenced the depositional history of these basins was conducted. Finally, the recognized patterns (i.e., Synchronous Events I–IV) of the drainage basins were compared with the extrinsic factors (i.e., climate and eustatic effects) identified within Texas and the region. The contemporaneous occurrence of events does not verify a correlation or causation between the incidents, but the synchronous occurrences may imply valid relationships or minimally a shared causal

mechanism. Therefore, these interpretations are malleable and may be adjusted when new data is encountered.

In sum, based on the recognized patterns within the drainage basins and the reviewed paleoenvironmental and sea levels through the Late Pleistocene-Holocene, it appears that climate was the primary forcing mechanism on the Texas drainage systems. This is particularly evident for Synchronous Events I–III that have the most robust evidence for climatic data within Texas and the region. However, all four of the recognized patterns appear to have been triggered by climatic influences. The clearest evidence of this exists in the depositional histories of alluvial settings in the Great Plains and Midwest far removed from the influence of eustatic effects, but are similar to those exhibited in Texas during these periods.

CHAPTER 9

Conclusions and Future Research

The primary goal of this study was to provide a chronological baseline for the comparison of archaeological sites in Texas drainage basins. To accomplish this objective, an extensive archival review of predominantly geoarchaeological research was undertaken to gather as much data as possible that met several criteria. There were three main criteria used for selecting the radiocarbon assays for this recalibration study. First, assays of charcoal were given priority over other dated materials; second, samples that have been previously ‘corrected’ for isotopic fractionation; and finally, datasets composed of samples in good stratigraphic context. With these criteria, the archival research was scrutinized for investigations that had been conducted in alluvial settings and, most importantly, had at least minimally considered chronometrically dating depositional stratigraphy, which was actually more difficult than initially envisioned. Furthermore, samples from humate materials were used more than initially intended, but these were reluctantly accepted. As discussed in Chapter 2, all calibrations of samples derived from humate materials tend to date inconsistently, sometimes drastically older (approximately 1,000–1,500 years) than comparable charcoal samples. Thus, it was only out of necessity that these samples were utilized and the results of these data should be used with caution.

Subsequent to the recalibration of the various datasets within the select Texas drainage basins, the data were consolidated and examined for any intra-basinal or inter-basinal patterns in depositional history. These comparisons recognized four contemporaneous events that occurred within most, if not all, of the Texas drainage

basins. These events include three periods of apparent instability (i.e., Synchronous Events I–III) and one of stability (i.e., Synchronous Event IV). These events were then compared to possible external mechanisms (e.g., climate and eustatic effects) that may have contributed to these occurrences. Based on similar depositional histories in the Great Plains and Midwest, the most probable trigger for the four synchronous events is climate. Furthermore, when the four events were compared to the archaeological record in Texas, three of them (i.e., Synchronous Events I, II, and IV) aligned during periods of cultural transition while the remaining one (i.e., Synchronous Event III) not surprisingly occurred during the enigmatic Middle Archaic, long recognized for poor integrity. This begs the question, were these adjustments in cultural lifeways a result of the changing environment or something else (e.g., social)? Although not definitive, these data do appear to corroborate the argument that the changing environment is the significant contributing factor to these transitions.

Interpretations

Several general interpretations developed from the review of the archival data and the results of the analyses. One general interpretation is that depositional landforms will exhibit more integrity as the distance downstream from the Edwards Plateau increases. This is almost assuredly due to the more confining channels that have incised into limestone bedrock in the uplands. The fluvial history of these areas demonstrates that they are periodically flushed out removing much of the deposition while in the prairies where the channels become more sinuous, the fluvial deposits are more complete and intact providing a better depositional history. Some examples of this include the Jonas Terrace site (upper Guadalupe-San Antonio River Basin) and the Woodrow Heard site

(upper Nueces River Basin), which have truncated depositional histories.

Comparatively, the Richard Beene site (Guadalupe-San Antonio River Basin) has exceptional stratigraphy and integrity. To be sure, outside bends in higher sinuosity channels are eroded, but the interior bends preserve their deposits. Also, smaller tributary channels are more noticeably affected by changes in deposition than larger waterways. Thus, if the research focus is attempting to characterize changes in past climates then smaller channels should be analyzed (e.g., Dry Frio River, South Fork San Geronimo, Cowhouse, Salado, and Medio Creeks), but if the research is investigating continuous alluvial stratigraphy with possibly robust archaeological deposits then more substantial drainages should be explored (e.g., Brazos, Colorado, Medina and San Antonio Rivers).

Another interpretation from this study includes the issue of integrity of particular time periods and drainages. First, the pervasive period of instability that occurred during the Middle Archaic in Central Texas (i.e., Synchronous Event III) where archaeological sites and cultural activities associated with this temporal setting are expected to be sparse. Thus, any modeling attempting to attribute the paucity of Middle Archaic groups to cultural influences (e.g., low population density or settlement) should first consider the site's location and geomorphic causes. Conversely, any Middle Archaic deposits encountered should be regarded as significant and warranting extensive investigation. Similarly, the low frequency or paucity of radiocarbon data should rarely be used to infer the absence or limited occurrence of cultural activities—to do so is almost assuredly an incorrect interpretation. An example of this (and there are many) include research at Fort Hood where the low frequency of radiocarbon data had been used to imply limited cultural activities during the beginning and end of the Early Archaic, and the Middle

Archaic, while the Late Prehistoric period is indicated to be an era of high cultural activity (Thoms and Olive 1993: Figure 12). Not coincidentally, these periods all align with the four synchronous events identified in the study. Specifically, three periods identified as having low cultural activity coincide with the three periods of instability (i.e., Synchronous Events I–III) identified in this study. Conversely, the era identified as having high cultural activity occurred during the time of pervasive stability (i.e., Synchronous Event I) (Figure 9.1). The initial interpretation of the researchers may be correct in attributing the low frequency of radiocarbon dates at Fort Hood to cultural reasons and not geomorphic integrity (i.e., instability or stability). However, the geomorphic factor of integrity should be seriously considered first.

Another observation derived from the study regards the magnitude of adjustment of radiocarbon assays after calibration. A predominance of recalibrated assays younger than 7,000 cal yr BP exhibited no significant adjustments from the initial calibration. Specifically, the changes usually fell within the standard deviation. This was particularly evident the more recent the initial calibration occurred. Those assays older than 7,000 cal yr BP always demonstrated some significant change, which was more pronounced the older the assay. This phenomenon is attributed to the continued refinement of the calibration curve where more data are added. Presently, Intcal09 has abundant data points that extend to 18,000 cal yr BP. Thus, subsequent calibrations of radiocarbon assays calibrated with the Intcal09 curve should likely not demonstrate any significant adjustments in the future. There will certainly be refinements, but just as the pre-7,000 cal yr BP recalibrations do not noticeably adjust, the assays encompassing the last 12,000 years should not alter significantly.




Figure 9.1 Radiocarbon Frequency at Fort Hood Equated with Cultural Activity (adapted from Thoms and Olive 1993: Figure 12).

Conversely, the use of the statistical calibration models within the OxCal programs (i.e., Bayesian statistics) proved very informative and useful. As previously noted in Chapter 2, each set of assays would be analyzed by a minimum of 30,000 iterations, but frequently went over 3 million iterations to get the most probable outcome with the available data (i.e., radiocarbon age and stratigraphic position). The most informative implementation of these statistical procedures occurred when samples were within a well-defined stratigraphy and were temporally close. The Fort Hood dataset are a good example of this situation. In contrast, the statistical models typically did not assist assays collected from multiple profiles with a vague stratigraphy and/or broad temporal range. Simply put, those investigations that had a research focus aimed at characterizing the chronology of geomorphic stratigraphy are perfectly suited for this application. Whereas, those investigations that have little or no focus on geomorphic stratigraphy are minimally enhanced if at all.

One constructive aspect garnered from this study may be the use of a chronostratigraphic marker in future research. At several locations in the review of the Guadalupe-San Antonio River basin (e.g., Unit II on Leon Creek and Unit 2 San Antonio River) periods of stability that preceded a dynamic erosion-incision event were identified (Nordt 2001a). The combination of a marked phase of stability followed by a period of very dynamic flow may be indicative of the 5,000–4,000 cal yr BP (calibrated IntCal 09 curve) time period. Coupled with this, the paucity of colluvial deposits (e.g., Unit IV at Culebra Creek) previously attributed to an exhausted supply of upland plateau sediments (Nordt 2001a:42). Therefore, geoarchaeological investigations could possibly use this suite of attributes for chronostratigraphic purposes while investigating the drainages (e.g.,

Leon Creek, Salado Creek, Culebra Creek, Medina River, and San Antonio River) along the margins of the Edwards Plateau in the Guadalupe-San Antonio River basin. Numerous researchers have previously proposed using chronostratigraphic markers within and around Texas (e.g., Abbott 2001, Blum 1992, and Bousman and Skinner 2007), but the implementations of these are used by a select few. This is a resource that should be fostered and utilized more in the future by archaeologists aware of these possibilities.

Future Research

Similar to a variety of interpretations that developed over the course of these investigations are several avenues of future research that have also been identified. The first and obvious area of future investigations is testing the validity of the four recognized synchronous events. The most tenuous component of the previous interpretations concerns the data gaps and ‘hiatuses’ within each of the investigated basins. Have these phenomena been affected or made more substantial through sampling or interpretive bias (e.g., poor chronological controls)? This research avenue can be achieved by attempting a more robust chronometric study in each of these basins. Also, the comparison of the drainage basin depositional histories was conducted at 250-year intervals. Future research should focus on tightening the intervals to determine if the synchronous events become more pronounced or vanish.

Furthermore, a portion of the assays from the current study were not corrected for isotopic fractionation. As with this study’s use of select radiocarbon samples derived from humate, the assays that were not ‘normalized’ were used out of necessity. Consequently, future research should adjust for these discrepancies and combine all of

the radiocarbon samples thus providing for a more robust dataset. The result of this effort can also be used for testing the interpretation of the four recognized synchronous events.

A corollary of this concerns filling the gaps of investigation across Texas. Specifically, there are several large regions in Texas that have little or no geomorphically slanted investigations. One of these regions is East Texas, where targeted research evaluating the drainages has not been done to any significant degree. This includes the middle extent of the Trinity River, the Neches River, Angelina River, the Sabine River, and their tributaries. Granted, geoarchaeological investigations have been conducted in this region (e.g., Phillips and Marion 2001), but these are in upland settings, are very focused, and/or do not truly regard depositional histories. The limited geoarchaeological research that has been conducted in the region suggests some significant deposits. Specifically, at the northern end of this region in the Sulphur River basin several researchers have devoted considerable attention to the area (Bousman et al. 1988; Bousman and Skinner 2007; Darwin et al. 2007; Rainey 1974). These researchers have encountered some promising evidence of Pre-Clovis deposits, which warrant further investigations (Bousman and Skinner 2007).

Similarly, select sections of drainage basins included within this study should be investigated. In particular, the upper extent of the Colorado River basin and the lower extents of the San Antonio, Guadalupe River, and Nueces River basins. The closing of these gaps in select areas could add significant data and immeasurably refine the interpretations (e.g., geochronological) held today. The enhancement of these drainage depositional histories could more effectively examine the response of a drainage to

external factors. A limitation of this study was utilizing the stratigraphy of archaeological site investigations and equating that with drainage depositional history. Again, this was done largely out of necessity, but future research can fill the gaps in coverage and refine or replace the interpretations provided here.

Additionally, some consideration should be put into the construction of a radiocarbon database for the use of Texas researchers. A system such as this has been in operation for years in Europe with some very intriguing developments (e.g., Chiverrell et al. 2009; Howard et al. 2009; Johnstone et al. 2006; Lewin et al. 2005; Macklin et al. 2002, 2005, 2006, 2010; Macklin and Lewin 2008). Admittedly, there are some significant obstacles that would likely arise, but the compilation and ready access of chronometric data from archaeological sites and depositional settings for the use of archaeological research is an attainable goal.

Finally, on a related note, some efforts should be put into developing an accepted form of reporting ^{14}C results. The archival research for this study has encountered a multitude of disparate ways of reporting what should be a straightforward dataset. Granted, there is prevalent confusion on the use and interpretation of chronometric data, but this would be made simpler if the presentation of data had some standardization.

Glossary

Aggradation: refers to the addition of sediment to drainage floodplain elevating the height of terraces.

Allostratigraphic unit: a mappable body of sedimentary rock bounded by a discontinuity (NACSN 2005:1578). The use of allostratigraphic unit in this study is considered a recognizable system for characterizing fluvial deposits of previous investigations.

Avulsion: refers to the rapid abandonment of a river channel and the formation of a new river channel.

Calibration: when a radiocarbon date is converted to a calendrical format (McCormac and Baillie 1993; Mook and Waterbolk 1985:20; Lowe and Walker 1997:243; Ramsey 2009:337; Stuiver and Suess 1966; Taylor 1997:68).

cal yr BP: in this study indicates the dates have been calibrated with IntCal 2009 calibration curve using A. D. 1950 as date before present.

Corrected: assays that have been adjusted for isotopic fractionation (i.e., $\delta^{13}\text{C}$ value of -25.0‰) (Hua 2009). Sometimes identified as conventional or normalized.

Data Gap: in the depositional history tables, this refers to a separation between allostratigraphic units where no erosion or aggradation is recorded. The data gaps are likely attributable to erosion, but more data is needed.

Hiatus: in the depositional history tables, this refers to chronological gaps within allostratigraphic units between radiocarbon assays.

Likelihood: in Bayesian analysis, the likelihood is the measured data (absolute dates) that is compared with the prior probability (Ramsey 2009).

MCMC: Markov chain Monte Carlo method that randomly examines each event across a defined distribution gradually increasing the confidence of the result. Also, allows for the inclusion of the uncertainty of multiple factors that can allow for the comparison of points as well as their deviations on a curve (Breyer 2009; Buck and Blackwell 2004:1101; Everitt 2002; Heaton et al. 2009; Ramsey 2009; Upton and Cook 2006).

Prior: in Bayesian analysis, the prior probability is inferred from relative dates, which for this study is stratigraphy and compared with the likelihood probability culminating in the determination of the posterior probability (Ramsey 2009).


^{14}C yr BP: refers to the uncalibrated radiocarbon age and is in radiocarbon years.

APPENDICES

APPENDIX I-Nueces River Basin OxCal Results
Woodrow Heard assays (Decker et al. 2000)

Name Show all Show structure	Unmodelled (BP)						Modelled (BP)						Indices Amodel=98.7 Aoverall=98			Select	Page break		
	from	to	%	from	to	%	from	to	%	from	to	%	Acomb	A	L P C	All Visible			
End 4							3570	3140	68.2	3650	2260	95.4				95.8	<input type="checkbox"/>	<input type="checkbox"/>	
Beta-112980	3570	3440	100.0	3640	3380	100.0	3570	3440	68.2	3630	3390	95.4		99		99.7	<input checked="" type="checkbox"/>	19	<input type="checkbox"/>
▲ 4																	<input checked="" type="checkbox"/>	18	<input type="checkbox"/>
Start 4							4100	3450	68.2	4800	3410	95.4				98.5	<input type="checkbox"/>		<input type="checkbox"/>
End 3							5260	4700	68.2	5310	4050	95.4				98.7	<input type="checkbox"/>		<input type="checkbox"/>
CAMS-9060	5310	5040	100.0	5320	4970	100.0	5310	5080	68.1	5440	4990	95.4		100		99.4	<input checked="" type="checkbox"/>	17	<input type="checkbox"/>
CAMS-9063	5580	5310	100.0	5590	5290	100.0	5470	5310	68.2	5590	5300	95.4		100.3		99.5	<input checked="" type="checkbox"/>	16	<input type="checkbox"/>
CAMS-9054	5580	5320	100.0	5590	5310	100.0	5580	5320	68.3	5590	5310	95.4		99.8		99.3	<input checked="" type="checkbox"/>	15	<input type="checkbox"/>
Beta-112981	6490	6390	100.0	6550	6300	100.0	6480	6320	68.1	6530	6300	95.4		99.2		99.7	<input checked="" type="checkbox"/>	14	<input type="checkbox"/>
▲ 3																	<input checked="" type="checkbox"/>	13	<input type="checkbox"/>
Start 3							6690	6410	68.2	6850	6340	95.4				99.6	<input type="checkbox"/>		<input type="checkbox"/>
End 2							6950	6730	68.2	7030	6530	95.4				99.4	<input type="checkbox"/>		<input type="checkbox"/>
Beta-112974	7010	6880	100.0	7160	6790	100.0	7150	6880	68.2	7160	6800	95.4		99.5		99.5	<input checked="" type="checkbox"/>	12	<input type="checkbox"/>
Beta-112977	7010	6890	100.0	7160	6790	100.0	7150	6890	68.2	7160	6850	95.4		100.4		99.6	<input checked="" type="checkbox"/>	11	<input type="checkbox"/>
Beta-112978	7160	6890	100.0	7170	6790	100.0	7160	6910	68.2	7170	6850	95.4		102.8		99.6	<input checked="" type="checkbox"/>	10	<input type="checkbox"/>
Beta-112976	7430	7310	100.0	7430	7260	100.0	7420	7310	68.2	7430	7270	95.4		95.1		99.6	<input checked="" type="checkbox"/>	9	<input type="checkbox"/>
▲ 2																	<input checked="" type="checkbox"/>	8	<input type="checkbox"/>
Start 2							7610	7300	68.2	8120	7270	95.4				97.9	<input type="checkbox"/>		<input type="checkbox"/>
End 1							8950	8590	68.2	9010	8120	95.4				97.9	<input type="checkbox"/>		<input type="checkbox"/>
CAMS-14496	9010	8770	100.0	9030	8640	100.0	9010	8810	68.2	9030	8720	95.4		104.1		99.1	<input checked="" type="checkbox"/>	7	<input type="checkbox"/>
CAMS-9057	9020	8770	100.0	9130	8640	100.0	9030	8790	68.2	9130	8660	95.4		103.3		99.3	<input checked="" type="checkbox"/>	6	<input type="checkbox"/>
CAMS-14501	9400	9120	100.0	9440	9020	100.0	9380	9120	68.2	9410	9030	95.4		100.9		99.5	<input checked="" type="checkbox"/>	5	<input type="checkbox"/>
Beta-112979	9440	9280	100.0	9480	9140	100.1	9430	9280	68.2	9470	9140	95.4		99.7		99.6	<input checked="" type="checkbox"/>	4	<input type="checkbox"/>
CAMS-14500	9490	9400	100.0	9530	9290	100.0	9480	9310	68.2	9510	9280	95.4		89.3		99.5	<input checked="" type="checkbox"/>	3	<input type="checkbox"/>
▲ 1																	<input checked="" type="checkbox"/>	2	<input type="checkbox"/>
Start 1							9610	9360	68.2	9930	9300	95.4				96.1	<input type="checkbox"/>		<input type="checkbox"/>
▲																	<input type="checkbox"/>		<input type="checkbox"/>


OxCal v4.1.8 Bronk Ramsey (2010); r.5 intr Atmospheric data from Reimer et al. (2009);



Choke Canyon assays

Name	Unmodelled (BP)						Modelled (BP)						Indices Amodel=90 Aoverall=90.5			Select All Visible	Page break				
	from	to	%	from	to	%	from	to	%	from	to	%	Acomb	A	L			P	C		
End 4							430	290	68.2	470	250	95.4						99.4	<input type="checkbox"/>	<input type="checkbox"/>	
TX4668	460	300	68.2	510	150	95.4	460	340	68.2	490	310	95.4		107.8				99.9	<input checked="" type="checkbox"/>	24	<input type="checkbox"/>
TX-4667	490	320	68.2	510	310	95.4	470	330	68.2	500	310	95.4		103.4				99.9	<input checked="" type="checkbox"/>	23	<input type="checkbox"/>
TX-4678	460	310	68.2	510	280	95.4	470	330	68.2	490	310	95.4		107.4				99.9	<input checked="" type="checkbox"/>	22	<input type="checkbox"/>
TX-4677	540	320	68.3	630	300	95.4	480	330	68.2	510	310	95.4		96.1				99.9	<input checked="" type="checkbox"/>	21	<input type="checkbox"/>
TX-4682	550	330	68.2	630	310	95.4	490	330	68.2	520	310	95.4		62.5				99.9	<input checked="" type="checkbox"/>	20	<input type="checkbox"/>
TX-4683	440	290	68.2	500	-10	95.3	460	350	68.2	480	300	95.4		101.2				99.9	<input checked="" type="checkbox"/>	19	<input type="checkbox"/>
TX-4684	460	300	68.2	500	290	95.4	460	340	68.2	490	310	95.4		104.3				99.9	<input checked="" type="checkbox"/>	18	<input type="checkbox"/>
▲ 4																		<input checked="" type="checkbox"/>	17	<input type="checkbox"/>	
Start 4							530	370	68.2	570	330	95.4						99.8	<input type="checkbox"/>		<input type="checkbox"/>
End 3							700	540	68.2	760	430	95.4						99.9	<input type="checkbox"/>		<input type="checkbox"/>
TX-4686	750	570	68.2	900	550	95.4	770	650	68.2	910	560	95.4		99.9				99.9	<input checked="" type="checkbox"/>	16	<input type="checkbox"/>
TX-4685	760	660	68.2	900	560	95.4	770	670	68.2	910	650	95.5		97.8				99.9	<input checked="" type="checkbox"/>	15	<input type="checkbox"/>
TX-4687	1070	930	68.2	1180	920	95.4	1040	930	68.2	1150	790	95.4		102				99.8	<input checked="" type="checkbox"/>	14	<input type="checkbox"/>
▲ 3																		<input checked="" type="checkbox"/>	13	<input type="checkbox"/>	
Start 3							1170	970	68.2	1350	840	95.4						99.9	<input type="checkbox"/>		<input type="checkbox"/>
End 2							1410	1160	68.2	1510	1050	95.4						99.9	<input type="checkbox"/>		<input type="checkbox"/>
TX-4680	1410	1280	68.2	1530	1180	95.4	1520	1300	68.2	1550	1260	95.4		87.1				99.7	<input checked="" type="checkbox"/>	12	<input type="checkbox"/>
TX-4679	2060	1880	68.2	2290	1820	95.4	2060	1880	68.2	2160	1820	95.4		99.9				99.8	<input checked="" type="checkbox"/>	11	<input type="checkbox"/>
TX-4681	1870	1720	68.2	1930	1620	95.4	1870	1720	68.2	1930	1620	95.4		99.9				99.8	<input checked="" type="checkbox"/>	10	<input type="checkbox"/>
TX-4669	2150	1950	68.2	2310	1890	95.4	2150	1950	68.2	2310	1890	95.5		100				99.9	<input checked="" type="checkbox"/>	9	<input type="checkbox"/>
TX-4665	2700	2360	68.2	2720	2350	95.4	2450	2350	68.2	2560	2330	95.4		99.9				99.9	<input checked="" type="checkbox"/>	8	<input type="checkbox"/>
▲ 2																		<input checked="" type="checkbox"/>	7	<input type="checkbox"/>	
Start 2							2550	2380	68.2	2650	2360	95.4						99.9	<input type="checkbox"/>		<input type="checkbox"/>
End 1							2690	2500	68.2	2740	2420	95.4						99.9	<input type="checkbox"/>		<input type="checkbox"/>
TX-4672	2860	2750	68.2	2950	2740	95.4	2860	2750	68.2	2950	2740	95.4		100				99.8	<input checked="" type="checkbox"/>	6	<input type="checkbox"/>
TX-4673	2750	2490	68.2	2750	2360	95.4	2760	2610	68.2	2780	2500	95.4		104.2				99.9	<input checked="" type="checkbox"/>	5	<input type="checkbox"/>
TX-4691	6780	6560	68.1	6890	6480	95.4	6780	6560	68.2	6890	6480	95.4		100				99.8	<input checked="" type="checkbox"/>	4	<input type="checkbox"/>
TX-4690	7420	7170	68.3	7440	7020	95.4	7420	7160	68.2	7440	7000	95.4		94.2				99.6	<input checked="" type="checkbox"/>	3	<input type="checkbox"/>
▲ 1																		<input checked="" type="checkbox"/>	2	<input type="checkbox"/>	
Start 1							8120	7170	68.2	9560	6980	95.4						96.6	<input type="checkbox"/>		<input type="checkbox"/>
▲																		<input type="checkbox"/>		<input type="checkbox"/>	

OxCal v4.1.6 Bronk Ramsey (2010); r:5 Atmospheric data from Reimer et al. (2009);




Ricklis and Blum 1997 data


Name	Unmodelled (BP)						Select	Page break
	from	to	%	from	to	%		
Warning! Duplicate names - Beta-80009								
TX-6125	780	660	68.2	910	560	95.4	<input checked="" type="checkbox"/>	69
TX-6127	900	560	68.2	930	530	95.4	<input checked="" type="checkbox"/>	68
TX-7306	920	760	68.2	940	690	95.4	<input checked="" type="checkbox"/>	67
TX-7305	920	770	68.2	940	690	95.4	<input checked="" type="checkbox"/>	66
TX-6925	960	790	68.2	1060	730	95.4	<input checked="" type="checkbox"/>	65
TX-6919	1060	790	68.2	1080	740	95.4	<input checked="" type="checkbox"/>	64
TX-6926	1060	800	68.2	1170	770	95.4	<input checked="" type="checkbox"/>	63
TX-6924	1170	980	68.1	1240	960	95.4	<input checked="" type="checkbox"/>	62
TX-522	1060	970	68.2	1170	930	95.5	<input checked="" type="checkbox"/>	61
TX-521	1240	1070	68.2	1280	1010	95.4	<input checked="" type="checkbox"/>	60
TX-523	1240	1070	68.2	1280	1010	95.4	<input checked="" type="checkbox"/>	59
TX-7312	1180	980	68.2	1270	960	95.4	<input checked="" type="checkbox"/>	58
TX-7313	1350	1180	68.2	1390	1170	95.4	<input checked="" type="checkbox"/>	57
TX-7304	1400	1290	68.2	1520	1190	95.4	<input checked="" type="checkbox"/>	56
Beta-77683	1610	1410	68.3	1700	1400	95.4	<input checked="" type="checkbox"/>	55
TX-5892	1540	1390	68.2	1690	1320	95.4	<input checked="" type="checkbox"/>	54
Beta-80016	1690	1520	68.2	1710	1410	95.4	<input checked="" type="checkbox"/>	53
TX-6062	1690	1420	68.2	1710	1410	95.4	<input checked="" type="checkbox"/>	52
TX-5893	1700	1420	68.2	1730	1390	95.4	<input checked="" type="checkbox"/>	51
TX-5891	1880	1710	68.2	1950	1600	95.4	<input checked="" type="checkbox"/>	50
Beta-57911	1700	1550	68.2	1820	1510	95.4	<input checked="" type="checkbox"/>	49
TX-7084	1780	1560	68.2	1870	1520	95.4	<input checked="" type="checkbox"/>	48
Beta-77684	1830	1630	68.2	1880	1570	95.4	<input checked="" type="checkbox"/>	47
UGA-6151***	1810	1560	68.2	1890	1410	95.4	<input checked="" type="checkbox"/>	46
Beta-80008	1930	1740	68.2	1990	1710	95.4	<input checked="" type="checkbox"/>	45
UGA-6151	2110	1870	68.2	2300	1730	95.4	<input checked="" type="checkbox"/>	44
Beta-77685	2310	2060	68.3	2330	2000	95.4	<input checked="" type="checkbox"/>	43
TX-7303	2330	2150	68.2	2350	2060	95.4	<input checked="" type="checkbox"/>	42
Beta-77686	2700	2360	68.2	2720	2350	95.4	<input checked="" type="checkbox"/>	41
Beta-57915	2720	2460	68.2	2730	2360	95.4	<input checked="" type="checkbox"/>	40
Beta-80012	2770	2540	68.1	2790	2480	95.4	<input checked="" type="checkbox"/>	39
Beta-77687	2840	2540	68.2	2860	2490	95.4	<input checked="" type="checkbox"/>	38
TX-5664	2840	2540	68.2	2860	2490	95.4	<input checked="" type="checkbox"/>	37
UGA-6152	2850	2510	68.2	2920	2360	95.4	<input checked="" type="checkbox"/>	36
Beta-80005	3320	3070	68.2	3350	2990	95.4	<input checked="" type="checkbox"/>	35
Beta-47105	3070	2860	68.1	3160	2790	95.4	<input checked="" type="checkbox"/>	34
TX-7311	4530	4290	68.2	4810	4150	95.4	<input checked="" type="checkbox"/>	33
TX-7310	4850	4580	68.3	4960	4440	95.4	<input checked="" type="checkbox"/>	32
TX-6881	4850	4620	68.2	4950	4520	95.4	<input checked="" type="checkbox"/>	31
Beta-80009	4960	4640	68.1	5040	4570	95.4	<input checked="" type="checkbox"/>	30
TX-7309	4960	4150	68.2	5440	3860	95.4	<input checked="" type="checkbox"/>	29
Beta-80013	5030	4860	68.2	5270	4830	95.5	<input checked="" type="checkbox"/>	28
Beta-80006	5040	4860	68.2	5280	4840	95.4	<input checked="" type="checkbox"/>	27
TX-7307	4960	4150	68.2	5440	3860	95.4	<input checked="" type="checkbox"/>	26

Beta-53198	5280	4870	68.2	5290	4860	95.4	<input checked="" type="checkbox"/>	25
TX-6963	5280	4870	68.2	5290	4860	95.4	<input checked="" type="checkbox"/>	24
TX-7083	5300	5040	68.2	5440	4880	95.4	<input checked="" type="checkbox"/>	23
Beta-80019	5590	5330	68.2	5590	5320	95.4	<input checked="" type="checkbox"/>	22
TX-7081	5590	5320	68.2	5660	5300	95.4	<input checked="" type="checkbox"/>	21
TX-5303	5600	5330	68.2	5650	5320	95.4	<input checked="" type="checkbox"/>	20
TX-5265	5650	5330	68.2	5730	5320	95.4	<input checked="" type="checkbox"/>	19
TX-5263	5710	5470	68.2	5860	5320	95.4	<input checked="" type="checkbox"/>	18
TX-5264	5900	5660	68.2	5940	5590	95.4	<input checked="" type="checkbox"/>	17
Beta-53072	5720	5580	68.2	5890	5470	95.4	<input checked="" type="checkbox"/>	16
Beta-57912	5740	5600	68.2	5900	5580	95.4	<input checked="" type="checkbox"/>	15
TX-7308	5890	5640	68.2	5900	5600	95.4	<input checked="" type="checkbox"/>	14
Beta-80017	5900	5740	68.2	5910	5660	95.4	<input checked="" type="checkbox"/>	13
Beta-53073	6910	6730	68.2	7000	6660	95.4	<input checked="" type="checkbox"/>	12
TX-7302	7160	6780	68.2	7250	6670	95.4	<input checked="" type="checkbox"/>	11
Beta-57043	7160	6890	68.2	7250	6750	95.4	<input checked="" type="checkbox"/>	10
TX-7082	7250	7000	68.2	7320	6880	95.4	<input checked="" type="checkbox"/>	9
Beta-80007	7280	7020	68.1	7420	6980	95.4	<input checked="" type="checkbox"/>	8
Beta-80014	7420	7260	68.2	7430	7170	95.4	<input checked="" type="checkbox"/>	7
Beta-80009	7420	7260	68.2	7430	7170	95.4	<input checked="" type="checkbox"/>	6
Beta-53647	7420	7320	68.2	7430	7260	95.4	<input checked="" type="checkbox"/>	5
Beta-80015	7470	7320	68.2	7560	7280	95.4	<input checked="" type="checkbox"/>	4
TX-7024	7570	7320	68.2	7660	7250	95.4	<input checked="" type="checkbox"/>	3
Beta-80018	7560	7430	68.2	7580	7420	95.4	<input checked="" type="checkbox"/>	2


Cal v4.17.0 (beta) (Revised 2009) - 5. Absolute date from Reimer et al. (2009)




OxCal v4.1.7 Bronk Ramsey (2010); r.5 Atmospheric data from Reimer et al (2009)



OxCal v4.1.7 Bronk Ramsey (2010); r:5 Atmospheric data from Reimer et al. (2009)



OxCal v4.1.7 Bronk Ramsey (2010); r:5 Atmospheric data from Reimer et al (2009);




APPENDIX II-San Antonio River Basin OxCal Results

Jonas Terrace assays (Johnson 1995: Table 1)


Name	Unmodelled (BP)			Modelled (BP)			Indices			Select	Page break									
	from	to	%	from	to	%	from	to	%			A _{model} =5	A _{overall} =10	A _{comb} A	L	P	C	All Visible		
Warning! Poor agreement - A= 10.0%(A'c= 60.0%)																				
Warning! Poor agreement - A= 5.0%(A'c= 60.0%)																				
End 5				1010	640	68.2	1120	130	95.4							98.3	<input type="checkbox"/>	<input type="checkbox"/>		
Beta-75905	1060	920	100.0	1170	790	100.0	1060	920	68.2	1170	810	95.4				100.6	99.8	<input checked="" type="checkbox"/>	²³ <input type="checkbox"/>	
▲ 5																		<input checked="" type="checkbox"/>	²² <input type="checkbox"/>	
Start 5				1230	980	68.2	1310	870	95.4									99.6	<input checked="" type="checkbox"/>	²¹ <input type="checkbox"/>
End 4				1340	1250	68.2	1390	1120	95.4									99.5	<input checked="" type="checkbox"/>	²⁰ <input type="checkbox"/>
Beta-26345	1380	1290	100.0	1420	1270	100.0	1360	1300	68.2	1410	1270	95.4				118.3	99.7	<input checked="" type="checkbox"/>	¹⁹ <input type="checkbox"/>	
Beta-11250	1890	1610	100.0	2000	1520	100.0	1380	1300	68.2	1450	1290	95.4				1.3	99.6	<input checked="" type="checkbox"/>	¹⁸ <input type="checkbox"/>	
Warning! Poor agreement - A= 1.3%(A'c= 60.0%)																				
▲ 4																		<input checked="" type="checkbox"/>	¹⁷ <input type="checkbox"/>	
Start 4				1400	1310	68.2	1470	1290	95.4									99.5	<input type="checkbox"/>	<input type="checkbox"/>
End 3				1450	1320	68.2	1500	1300	95.4									99.4	<input type="checkbox"/>	<input type="checkbox"/>
Beta-62339	1290	1170	100.0	1310	1070	100.0	1480	1330	68.2	1520	1320	95.4				1.8	99	<input checked="" type="checkbox"/>	¹⁶ <input type="checkbox"/>	
Warning! Poor agreement - A= 1.8%(A'c= 60.0%)																				
Beta-62346	2700	2350	100.0	2730	2330	100.0	2660	2380	68.2	2710	2350	95.4				99.8	99.6	<input checked="" type="checkbox"/>	¹⁵ <input type="checkbox"/>	
Beta-62342	2690	2340	100.0	2730	2320	100.0	2650	2370	68.2	2710	2310	95.4				99.8	99.5	<input checked="" type="checkbox"/>	¹⁴ <input type="checkbox"/>	
Beta-62338	2760	2610	100.0	2780	2470	100.0	2740	2520	68.2	2780	2410	95.4				99.4	99.6	<input checked="" type="checkbox"/>	¹³ <input type="checkbox"/>	
Beta-62349	2770	2710	100.0	2850	2490	100.0	2780	2540	68.2	2850	2430	95.4				99.7	99.6	<input checked="" type="checkbox"/>	¹² <input type="checkbox"/>	
Beta-62348	3450	3260	100.0	3560	3160	100.0	3420	3210	68.2	3520	3080	95.4				93.8	99.4	<input checked="" type="checkbox"/>	¹¹ <input type="checkbox"/>	
▲ 3																		<input checked="" type="checkbox"/>	¹⁰ <input type="checkbox"/>	
Start 3				3800	3340	68.2	4100	3180	95.4									99.4	<input type="checkbox"/>	<input type="checkbox"/>
End 2				4280	3790	68.2	4380	3480	95.4									99.4	<input type="checkbox"/>	<input type="checkbox"/>
Beta-62347	4420	4150	100.0	4500	4090	100.0	4380	4180	68.2	4430	4100	95.4				99.6	99.8	<input checked="" type="checkbox"/>	⁹ <input type="checkbox"/>	
▲ 2																		<input checked="" type="checkbox"/>	⁸ <input type="checkbox"/>	
Start 2				4580	4280	68.2	4730	4170	95.4									99.7	<input checked="" type="checkbox"/>	⁷ <input type="checkbox"/>
End 1				4790	4520	68.2	4850	4360	95.4									99.6	<input checked="" type="checkbox"/>	⁶ <input type="checkbox"/>
Beta-62341	4840	4580	100.0	4860	4520	100.0	4840	4680	68.2	4870	4590	95.4				100.2	99.7	<input checked="" type="checkbox"/>	⁵ <input type="checkbox"/>	
Beta-62343	5220	4840	100.0	5300	4820	100.0	5040	4860	68.2	5240	4740	95.4				113.5	99.7	<input checked="" type="checkbox"/>	⁴ <input type="checkbox"/>	
Beta-62340	5310	4970	100.0	5440	4850	100.0	5140	4890	68.2	5280	4850	95.4				82.7	99.6	<input checked="" type="checkbox"/>	³ <input type="checkbox"/>	
▲ 1																		<input checked="" type="checkbox"/>	² <input type="checkbox"/>	
Start 1				5350	4960	68.2	5770	4890	95.4									95.9	<input type="checkbox"/>	<input type="checkbox"/>
▲																		<input type="checkbox"/>	<input type="checkbox"/>	


OxCal v4.1.6 Bronk Ramsey (2010); r.5 intr whole Atmospheric data from Reimer et al (2009);



Richard Beene assays (Mandel and Thoms 2007)

Name	Unmodelled (BP)						Modelled (BP)						Indices				Select	Page break		
	from	to	%	from	to	%	from	to	%	from	to	%	A _{comb}	A	L	P			C	All Visible
End 4							3370	2590	68.2	3450	1170	95.4					99.2	<input type="checkbox"/>	<input type="checkbox"/>	
Beta-36702	3390	3210	100.0	3450	3080	99.9	3400	3230	68.2	3470	3080	95.4		101.1			99.9	<input checked="" type="checkbox"/>	24	<input type="checkbox"/>
Beta-43330	4830	4520	100.0	4850	4430	100.0	4700	4440	68.2	4810	4430	95.4		92.1			99.9	<input checked="" type="checkbox"/>	23	<input type="checkbox"/>
▲ 4																		<input checked="" type="checkbox"/>	22	<input type="checkbox"/>
Start 4							4870	4600	68.2	5000	4480	95.4					100	<input type="checkbox"/>	<input type="checkbox"/>	
End 3							5030	4780	68.2	5160	4650	95.4					99.9	<input type="checkbox"/>	<input type="checkbox"/>	
AA-20401	5260	4840	99.9	5310	4720	100.0	5290	4930	68.1	5310	4850	95.4		95.3			99.8	<input checked="" type="checkbox"/>	21	<input type="checkbox"/>
GX-21746	5270	4880	99.9	5300	4850	100.0	5280	4960	68.2	5290	4880	95.4		98			99.9	<input checked="" type="checkbox"/>	20	<input type="checkbox"/>
AA-20402	5320	4970	100.0	5580	4850	100.0	5320	5030	68.2	5470	4880	95.4		102.9			99.9	<input checked="" type="checkbox"/>	19	<input type="checkbox"/>
Beta-38700	5440	5070	100.0	5470	4970	100.0	5450	5060	68.2	5470	4980	95.4		100.5			99.9	<input checked="" type="checkbox"/>	18	<input type="checkbox"/>
AA-20400	7670	7470	100.0	7750	7420	100.0	7660	7480	68.2	7790	7420	95.4		100.1			99.9	<input checked="" type="checkbox"/>	17	<input type="checkbox"/>
Beta-47524	7820	7660	100.0	7930	7600	100.0	7830	7660	68.2	7930	7600	95.4		101			99.9	<input checked="" type="checkbox"/>	16	<input type="checkbox"/>
Beta-47525	7840	7680	100.0	7940	7650	100.0	7830	7680	68.2	7930	7620	95.4		101.2			99.9	<input checked="" type="checkbox"/>	15	<input type="checkbox"/>
Beta-47523	7930	7730	100.0	7960	7670	100.0	7920	7730	68.2	7940	7680	95.4		100.5			99.9	<input checked="" type="checkbox"/>	14	<input type="checkbox"/>
Beta-47530	7940	7740	100.0	7970	7670	100.0	7930	7740	68.2	7950	7680	95.4		100.1			99.9	<input checked="" type="checkbox"/>	13	<input type="checkbox"/>
▲ 3																		<input checked="" type="checkbox"/>	12	<input type="checkbox"/>
Start 3							8160	7850	68.2	8370	7800	95.4					99.9	<input type="checkbox"/>	<input type="checkbox"/>	
End 2							8520	8290	68.2	8570	8090	95.4					99.9	<input type="checkbox"/>	<input type="checkbox"/>	
Beta-47529	8520	8380	100.0	8590	8340	100.0	8550	8410	68.2	8600	8370	95.4		93			99.9	<input checked="" type="checkbox"/>	11	<input type="checkbox"/>
Beta-78657	8590	8440	100.0	8610	8410	100.0	8590	8470	68.2	8600	8420	95.4		100.5			100	<input checked="" type="checkbox"/>	10	<input type="checkbox"/>
Beta-78656	8980	8600	100.0	9000	8580	100.0	8770	8590	68.2	8960	8550	95.4		103.9			99.9	<input checked="" type="checkbox"/>	9	<input type="checkbox"/>
Beta-44386	9140	8770	100.0	9410	8590	100.0	8880	8590	68.2	9100	8530	95.4		82.5			99.9	<input checked="" type="checkbox"/>	8	<input type="checkbox"/>
▲ 2																		<input checked="" type="checkbox"/>	7	<input type="checkbox"/>
Start 2							9030	8660	68.2	9310	8600	95.4					99.8	<input type="checkbox"/>	<input type="checkbox"/>	
End 1							9600	9130	68.2	9680	8820	95.4					99.9	<input type="checkbox"/>	<input type="checkbox"/>	
Beta-80687	9660	9530	100.0	9740	9520	100.0	9670	9540	68.2	9890	9510	95.4		96.3			99.9	<input checked="" type="checkbox"/>	6	<input type="checkbox"/>
Beta-47527	10120	9680	100.0	10180	9550	100.0	10120	9680	68.2	10160	9600	95.4		100.3			99.9	<input checked="" type="checkbox"/>	5	<input type="checkbox"/>
Beta-80974	10120	9690	100.0	10170	9600	100.0	10120	9700	68.2	10160	9630	95.4		100.2			99.9	<input checked="" type="checkbox"/>	4	<input type="checkbox"/>
Beta-47526	15460	14900	100.0	16190	14180	100.0	15190	14190	68.2	15780	14030	95.4		91.5			99.5	<input checked="" type="checkbox"/>	3	<input type="checkbox"/>
▲ 1																		<input checked="" type="checkbox"/>	2	<input type="checkbox"/>
Start 1							15950	14340	68.2	18030	14110	95.4					95.7	<input type="checkbox"/>	<input type="checkbox"/>	
▲																		<input type="checkbox"/>	<input type="checkbox"/>	






Copano Bay-Egery Island assays (Paine 1991:Table 5)

Name	Unmodelled (BP)			Modelled (BP)			Indices			Select	Page break						
	from	to	%	from	to	%	A	L	P			C	Visible				
End 4				2260	1920	68.2	2340	1410	95.4			97.8	<input type="checkbox"/>	<input type="checkbox"/>			
TX-6058	2350	2150	100.0	2360	2060	100.0	2320	2150	68.2	2360	2060	95.4	101.1	99.8	<input checked="" type="checkbox"/> 17	<input type="checkbox"/>	
▲ 4															<input checked="" type="checkbox"/> 18	<input type="checkbox"/>	
Start 4				2410	2200	68.2	2550	2120	95.4						99.7	<input type="checkbox"/>	<input type="checkbox"/>
End 3				2590	2320	68.2	2670	2220	95.4						99.7	<input type="checkbox"/>	<input type="checkbox"/>
TX-6059	2660	2340	100.0	2720	2210	100.0	2670	2400	68.2	2710	2340	95.4	94.3	99.8	<input checked="" type="checkbox"/> 13	<input type="checkbox"/>	
▲ 3															<input checked="" type="checkbox"/> 12	<input type="checkbox"/>	
Start 3				2830	2470	68.2	3040	2370	95.4						99.5	<input type="checkbox"/>	<input type="checkbox"/>
End 2				3130	2770	68.2	3250	2550	95.4						99.6	<input type="checkbox"/>	<input type="checkbox"/>
TX-6060	3220	3000	100.0	3340	2940	100.0	3230	3020	68.2	3320	2950	95.4	100.4	99.8	<input checked="" type="checkbox"/> 9	<input type="checkbox"/>	
▲ 2															<input checked="" type="checkbox"/> 8	<input type="checkbox"/>	
Start 2				3750	3150	68.2	4170	3020	95.4						99.3	<input type="checkbox"/>	<input type="checkbox"/>
End 1				4350	3720	68.2	4550	3320	95.4						99.1	<input type="checkbox"/>	<input type="checkbox"/>
TX-6061	4530	4240	100.0	4810	4090	100.0	4500	4180	68.2	4700	4020	95.4	97.2	99.6	<input checked="" type="checkbox"/> 5	<input type="checkbox"/>	
▲ 1															<input checked="" type="checkbox"/> 4	<input type="checkbox"/>	
Start 1				4800	4280	68.2	5540	4110	95.4						95.5	<input type="checkbox"/>	<input type="checkbox"/>
▲															<input type="checkbox"/>	<input type="checkbox"/>	


OxCal v4.1.6 Bronk Ramsey (2010); r:5 intr whole Atmospheric data from Reimer et al (2009):



APPENDIX III-Colorado River Basin OxCal Results
Concho River Results (Quigg et al. 1996: Table 12.2)

Name	Unmodelled (BP)						Modelled (BP)						Indices			Select	Page break				
	from	to	%	from	to	%	from	to	%	from	to	%	A _{comb}	A	L			P	C		
End 3							1130	150	68.2	1230	-1450	95.4						98.2	<input type="checkbox"/>	<input type="checkbox"/>	
B-69770	1270	1060	100.0	1290	980	100.0	1250	1080	68.2	1280	1000	95.4		100.4				99.9	<input checked="" type="checkbox"/>	15	<input type="checkbox"/>
▲ 3																			<input checked="" type="checkbox"/>	14	<input type="checkbox"/>
Start 3							2300	1230	68.2	3020	1100	95.4						99.4	<input type="checkbox"/>	<input type="checkbox"/>	
End 2							3320	2260	68.2	3470	1560	95.4						99.5	<input type="checkbox"/>	<input type="checkbox"/>	
B-72273	3470	3360	100.0	3570	3270	100.0	3510	3370	68.2	3580	3290	95.4		98.8				99.8	<input checked="" type="checkbox"/>	11	<input type="checkbox"/>
B-69766	4410	4150	100.0	4430	4080	100.0	4370	4160	68.2	4420	4050	95.4		99.9				99.8	<input checked="" type="checkbox"/>	10	<input type="checkbox"/>
B-69769	5290	4960	100.0	5310	4860	100.0	5230	4940	68.2	5290	4870	95.4		98.7				99.8	<input checked="" type="checkbox"/>	9	<input type="checkbox"/>
▲ 2																			<input checked="" type="checkbox"/>	8	<input type="checkbox"/>
Start 2							6510	5200	68.2	7520	4990	95.4						99.2	<input type="checkbox"/>	<input type="checkbox"/>	
End 1							8040	6580	68.2	8250	5610	95.4						99.1	<input type="checkbox"/>	<input type="checkbox"/>	
B-70133	8310	8000	100.0	8390	7930	100.0	8270	8010	68.2	8360	7890	95.4		99.8				99.8	<input checked="" type="checkbox"/>	5	<input type="checkbox"/>
▲ 1																			<input checked="" type="checkbox"/>	4	<input type="checkbox"/>
Start 1							9280	8150	68.2	11770	8000	95.4						96.9	<input type="checkbox"/>	<input type="checkbox"/>	
▲																			<input type="checkbox"/>	<input type="checkbox"/>	


OxCal v4.1.7 Bronk Ramsey (2010); r:5 intr whole Atmospheric data from Reimer et al (2009):




Lower Extent Colorado River (Blum 1992: Tables 6.1-6.4)

Name	Unmodelled (BP)						Modelled (BP)						Indices			Select	Page break			
	from	to	%	from	to	%	from	to	%	from	to	%	Acomb	A	L			P	C	All Visible
End 4							130	-360	68.2	230	-930	95.4						99.9	<input type="checkbox"/>	<input type="checkbox"/>
Tx-7227	260	20	100.0	290	0	100.0	250	50	68.2	280	-10	95.4		95				100	<input checked="" type="checkbox"/>	32 <input type="checkbox"/>
Warning! Date may extend out of range - 70+/-60BP																				
Tx-7321	300	-10	100.0	430	-10	100.0	290	90	68.2	410	0	95.4		99.5				100	<input checked="" type="checkbox"/>	31 <input type="checkbox"/>
Tx-7334	550	500	100.0	640	460	100.0	540	340	68.2	620	320	95.4		82.7				100	<input checked="" type="checkbox"/>	30 <input type="checkbox"/>
▲ 4																			<input checked="" type="checkbox"/>	29 <input type="checkbox"/>
Start 4							660	440	68.2	770	350	95.4						100	<input type="checkbox"/>	<input type="checkbox"/>
End 3							800	580	68.2	880	460	95.4						100	<input type="checkbox"/>	<input type="checkbox"/>
Tx-6813	790	670	100.0	920	660	100.0	890	710	68.2	920	670	95.4		86.4				100	<input checked="" type="checkbox"/>	28 <input type="checkbox"/>
Tx-6812	1550	1390	100.0	1690	1330	100.0	1560	1400	68.2	1680	1340	95.4		99.9				100	<input checked="" type="checkbox"/>	25 <input type="checkbox"/>
Tx-7223	1690	1520	100.0	1710	1400	100.0	1660	1460	68.2	1710	1410	95.4		99.8				100	<input checked="" type="checkbox"/>	24 <input type="checkbox"/>
Tx-7221	3220	3000	100.0	3340	2940	100.0	3220	3010	68.2	3320	2930	95.4		100				100	<input checked="" type="checkbox"/>	23 <input type="checkbox"/>
Tx-7222	3840	3450	100.0	3990	3350	100.0	3820	3480	68.2	4020	3340	95.4		100				99.9	<input checked="" type="checkbox"/>	22 <input type="checkbox"/>
Tx-6809	3840	3570	100.0	3960	3460	100.0	3830	3580	68.2	3950	3480	95.4		99.9				100	<input checked="" type="checkbox"/>	21 <input type="checkbox"/>
Tx-7220	4790	4410	100.0	4830	4150	100.0	4570	4260	68.2	4770	4120	95.4		94.5				99.9	<input checked="" type="checkbox"/>	20 <input type="checkbox"/>
▲ 3																			<input checked="" type="checkbox"/>	19 <input type="checkbox"/>
Start 3							4920	4420	68.2	5210	4240	95.4						100	<input type="checkbox"/>	<input type="checkbox"/>
End 2							5330	4760	68.2	5510	4490	95.4						100	<input type="checkbox"/>	<input type="checkbox"/>
Tx-7325	5580	5080	100.0	5610	4970	100.0	5550	5210	68.2	5630	5040	95.4		105.1				100	<input checked="" type="checkbox"/>	18 <input type="checkbox"/>
Tx-7226	5840	5600	100.0	5910	5580	100.0	5840	5630	68.2	5890	5590	95.4		99.9				100	<input checked="" type="checkbox"/>	15 <input type="checkbox"/>
Tx-6811	8640	8380	100.0	9000	8210	100.0	8790	8410	68.2	8980	8230	95.4		100				99.9	<input checked="" type="checkbox"/>	14 <input type="checkbox"/>
Tx-7224	13380	12100	100.0	14050	11210	100.0	13430	11940	68.2	14450	11220	95.4		100.6				99.8	<input checked="" type="checkbox"/>	13 <input type="checkbox"/>
Tx-7326	16820	14190	100.0	17540	13640	100.0	15940	14090	68.2	16790	13510	95.4		93.9				99.8	<input checked="" type="checkbox"/>	12 <input type="checkbox"/>
▲ 2																			<input checked="" type="checkbox"/>	11 <input type="checkbox"/>
Start 2							17570	14930	68.2	18870	14080	95.4						99.8	<input type="checkbox"/>	<input type="checkbox"/>
End 1							19460	17330	68.2	20270	15840	95.4						99.8	<input type="checkbox"/>	<input type="checkbox"/>
Tx-7010	21400	17580	100.0	24380	15920	100.0	21470	19000	68.2	23190	17790	95.4		117.7				99.8	<input checked="" type="checkbox"/>	8 <input type="checkbox"/>
Tx-7013	19760	18800	100.0	20290	18540	100.0	20060	19000	68.2	20840	18660	95.4		90.2				99.8	<input checked="" type="checkbox"/>	7 <input type="checkbox"/>
Tx-7012	28480	18540	100.0	49730	15230	100.0	22300	19380	68.2	24770	18050	95.4		143.3				99.8	<input checked="" type="checkbox"/>	6 <input type="checkbox"/>
Warning! Date may extend out of range - 18380+/-3890BP																				
Tx-7011	23270	21410	100.0	24010	20440	100.0	22500	20630	68.2	23470	19940	95.4		86.2				99.8	<input checked="" type="checkbox"/>	5 <input type="checkbox"/>
▲ 1																			<input checked="" type="checkbox"/>	4 <input type="checkbox"/>
Start 1							24500	21270	68.2	28130	20270	95.4						95.6	<input type="checkbox"/>	<input type="checkbox"/>
▲																			<input type="checkbox"/>	<input type="checkbox"/>

OxCal v4.1.7 Bronk Ramsey (2010); r.5 intr whole Atmospheric data from Reimer et al (2009);



OxCal v4.1.7 Bronk Ramsey (2010); r.5 intr whole Atmospheric data from Reimer et al (2009);




APPENDIX IV-Brazos River Basin OxCal Results


Lubbock Lake site (Haas et al. 1986:Table 1;Holliday et al. 1983, 1985)

Name	Unmodelled (BP)						Modelled (BP)						Indices			Select	Page break				
	from	to	%	from	to	%	from	to	%	from	to	%	A _{comb}	A	L			P	C	All Visible	
Warning! Poor agreement - A= 48.3%(A'c= 60.0%)																					
Warning! Poor agreement - A= 50.1%(A'c= 60.0%)																					
End 8							170	-200	68.2	260	-490	95.4						98.3	<input checked="" type="checkbox"/>	74	<input type="checkbox"/>
SMU-343	290	-10	100.0	310	-10	100.0	230	30	68.2	290	0	95.4		101				99.6	<input checked="" type="checkbox"/>	73	<input type="checkbox"/>
Warning! Date may extend out of range - 160+-60BP																					
▲ 8																			<input checked="" type="checkbox"/>	72	<input type="checkbox"/>
Start 8							300	110	68.2	360	30	95.4						99.8	<input checked="" type="checkbox"/>	71	<input type="checkbox"/>
End 7							390	270	68.2	420	200	95.4						99.5	<input checked="" type="checkbox"/>	70	<input type="checkbox"/>
SMU-555	310	0	100.0	430	-10	100.0	430	300	68.2	470	270	95.4		48.2				99.7	<input checked="" type="checkbox"/>	69	<input type="checkbox"/>
Warning! Poor agreement - A= 48.2%(A'c= 60.0%)																					
SI-2703	440	280	100.0	500	-10	100.0	460	350	68.2	500	290	95.4		107.4				99.8	<input checked="" type="checkbox"/>	68	<input type="checkbox"/>
SMU-345	470	290	100.0	510	-10	100.0	460	350	68.2	500	300	95.4		106.2				99.8	<input checked="" type="checkbox"/>	67	<input type="checkbox"/>
SI-2704	470	300	100.0	510	150	100.0	460	350	68.2	500	300	95.4		101.7				99.8	<input checked="" type="checkbox"/>	66	<input type="checkbox"/>
SMU-546	490	300	100.0	510	150	100.0	470	350	68.2	500	300	95.4		103.8				99.8	<input checked="" type="checkbox"/>	65	<input type="checkbox"/>
SMU-970	510	320	100.0	520	300	100.0	490	360	68.2	520	320	95.4		101.5				99.8	<input checked="" type="checkbox"/>	64	<input type="checkbox"/>
SI-2700	510	330	100.0	520	310	100.0	490	360	68.2	510	320	95.4		102				99.9	<input checked="" type="checkbox"/>	63	<input type="checkbox"/>
SMU-968	520	490	100.0	540	340	100.0	520	460	68.2	540	340	95.4		102.9				99.9	<input checked="" type="checkbox"/>	62	<input type="checkbox"/>
SMU-893	530	490	100.0	550	330	100.0	530	450	68.2	550	340	95.4		104.9				99.8	<input checked="" type="checkbox"/>	61	<input type="checkbox"/>
SI-2701	550	500	100.0	640	490	100.0	560	500	68.2	630	450	95.4		109.2				99.8	<input checked="" type="checkbox"/>	60	<input type="checkbox"/>
SI-3208	680	540	100.0	700	510	100.0	610	540	68.2	660	510	95.4		101.1				99.8	<input checked="" type="checkbox"/>	59	<input type="checkbox"/>
SMU-314	690	660	100.0	730	560	100.0	650	560	68.2	680	550	95.4		47.1				99.9	<input checked="" type="checkbox"/>	58	<input type="checkbox"/>
Warning! Poor agreement - A= 47.1%(A'c= 60.0%)																					
▲ 7																			<input checked="" type="checkbox"/>	57	<input type="checkbox"/>
Start 7							680	580	68.2	720	580	95.4						99.8	<input checked="" type="checkbox"/>	56	<input type="checkbox"/>
End 6							760	630	68.2	820	590	95.4						99.9	<input checked="" type="checkbox"/>	55	<input type="checkbox"/>
SMU-534	900	730	100.0	920	690	100.0	890	740	68.2	910	700	95.4		98.3				99.8	<input checked="" type="checkbox"/>	54	<input type="checkbox"/>
SI-4169	920	720	100.0	940	670	100.0	900	750	68.2	930	700	95.4		101.7				99.8	<input checked="" type="checkbox"/>	53	<input type="checkbox"/>
SMU-651	920	730	100.0	940	670	100.0	900	750	68.2	940	700	95.4		102.3				99.6	<input checked="" type="checkbox"/>	52	<input type="checkbox"/>
SI-3201	1260	1050	100.0	1290	970	100.0	1230	1060	68.2	1280	980	95.4		99.9				99.6	<input checked="" type="checkbox"/>	51	<input type="checkbox"/>
SMU-1090	1280	1170	100.0	1290	1070	100.0	1260	1150	68.2	1290	1080	95.4		99.9				99.7	<input checked="" type="checkbox"/>	50	<input type="checkbox"/>
SMU-1177	1530	1380	100.0	1550	1330	100.0	1510	1380	68.2	1550	1340	95.4		100				99.2	<input checked="" type="checkbox"/>	49	<input type="checkbox"/>
SI-4174	2000	1820	100.0	2120	1710	100.0	2000	1820	68.2	2110	1720	95.4		100				99.9	<input checked="" type="checkbox"/>	48	<input type="checkbox"/>
SMU-1191	2300	1880	100.0	2350	1720	100.0	2240	1900	68.2	2340	1740	95.4		99.9				99.9	<input checked="" type="checkbox"/>	47	<input type="checkbox"/>
SI-4171	5580	5320	100.0	5590	5300	100.0	5550	5340	68.2	5580	5310	95.4		99.9				99.6	<input checked="" type="checkbox"/>	46	<input type="checkbox"/>
SMU-492	5740	5610	100.0	5890	5590	100.0	5690	5610	68.2	5730	5590	95.4		102.5				99.6	<input checked="" type="checkbox"/>	45	<input type="checkbox"/>
▲ 6																			<input checked="" type="checkbox"/>	44	<input type="checkbox"/>
Start 6							5750	5640	68.2	5790	5610	95.4						99.9	<input checked="" type="checkbox"/>	43	<input type="checkbox"/>
End 5							5790	5690	68.2	5830	5640	95.4						99.8	<input checked="" type="checkbox"/>	42	<input type="checkbox"/>
SMU-1200	6280	5900	100.0	6400	5660	100.0	5810	5720	68.2	5850	5660	95.4		46.2				99.9	<input checked="" type="checkbox"/>	41	<input type="checkbox"/>
Warning! Poor agreement - A= 46.2%(A'c= 60.0%)																					
▲ 5																			<input checked="" type="checkbox"/>	40	<input type="checkbox"/>
Start 5							5830	5740	68.2	5870	5670	95.4						99.8	<input checked="" type="checkbox"/>	39	<input type="checkbox"/>
End 4							5860	5780	68.2	5890	5710	95.4						99.9	<input checked="" type="checkbox"/>	38	<input type="checkbox"/>
SMU-531	5660	5580	100.0	5750	5480	100.0	5890	5830	68.2	5900	5730	95.4		9.6				99.8	<input checked="" type="checkbox"/>	37	<input type="checkbox"/>
Warning! Poor agreement - A= 9.6%(A'c= 60.0%)																					
SMU-1093	6000	5920	100.0	6180	5900	100.0	6100	5930	68.2	6180	5910	95.4		100.3				99.7	<input checked="" type="checkbox"/>	36	<input type="checkbox"/>
SMU-545	6670	6470	100.0	6750	6400	100.0	6630	6440	68.2	6720	6340	95.4		96.8				99.5	<input checked="" type="checkbox"/>	35	<input type="checkbox"/>


OxCal v4.1.7 Bronk Ramsey (2010); r.5 intr whole Atmospheric data from Reimer et al (2009);




OxCal v4.1.7 Bronk Ramsey (2010); r.5 intr whole Atmospheric data from Reimer et al. (2009);



OxCal v4.1.7 Bronk Ramsey (2010); r.5 intr whole Atmospheric data from Reimer et al (2009);




OxCal v4.1.7 Bronk Ramsey (2010); r.5 intr whole Atmospheric data from Reimer et al (2009);




Fort Hood- Cowhouse Creek (Nordt 2004: Table 1)

Name	Unmodelled (BP)						Modelled (BP)						Indices			Select	Page break				
	from	to	%	from	to	%	from	to	%	from	to	%	A _{model} =91.9	A _{overall} =92.7	A _{comb}			L	P	C	All Visible
End 6							370	90	68.2	440	-210	95.4						99	<input checked="" type="checkbox"/>	41	<input type="checkbox"/>
Beta-37008	310	-10	100.0	480	-10	100.0	420	220	68.2	470	70	95.4	71.8					99.8	<input checked="" type="checkbox"/>	40	<input type="checkbox"/>
Warning! Date may extend out of range - 190+/-90BP																					
TX-6697	500	150	100.0	530	-10	100.0	440	290	68.2	490	150	95.4	117.8					99.9	<input checked="" type="checkbox"/>	39	<input type="checkbox"/>
Warning! Date may extend out of range - 300+/-100BP																					
TX-6699	550	0	100.0	680	-10	100.0	440	290	68.2	500	120	95.4	123.3					99.9	<input checked="" type="checkbox"/>	38	<input type="checkbox"/>
Warning! Date may extend out of range - 370+/-180BP																					
Beta-38177	510	320	100.0	530	300	100.0	450	320	68.2	500	300	95.4	92.9					99.9	<input checked="" type="checkbox"/>	37	<input type="checkbox"/>
▲ 6																			<input checked="" type="checkbox"/>	36	<input type="checkbox"/>
Start 6							510	360	68.2	590	320	95.4						99.9	<input checked="" type="checkbox"/>	35	<input type="checkbox"/>
End 5							630	460	68.2	740	380	95.4						100	<input checked="" type="checkbox"/>	34	<input type="checkbox"/>
TX-6700	680	510	100.0	790	310	100.0	750	570	68.2	900	500	95.4	99.3					99.9	<input checked="" type="checkbox"/>	33	<input type="checkbox"/>
TX-6701	730	520	100.0	930	330	100.0	830	580	68.2	940	510	95.4	100.9					99.8	<input checked="" type="checkbox"/>	32	<input type="checkbox"/>
Beta-38174	1420	1320	100.0	1530	1290	100.0	1490	1330	68.2	1530	1300	95.4	99.9					99.9	<input checked="" type="checkbox"/>	31	<input type="checkbox"/>
Beta-37450	1710	1520	100.0	1830	1380	100.0	1710	1480	68.2	1820	1390	95.4	99.9					99.9	<input checked="" type="checkbox"/>	30	<input type="checkbox"/>
Beta-37156	1870	1620	100.0	1930	1540	100.0	1850	1640	68.2	1930	1560	95.4	100					99.9	<input checked="" type="checkbox"/>	29	<input type="checkbox"/>
TX-6702	2720	2200	100.0	2780	2040	100.0	2430	2060	68.2	2620	1910	95.4	75.7					99.8	<input checked="" type="checkbox"/>	28	<input type="checkbox"/>
▲ 5																			<input checked="" type="checkbox"/>	27	<input type="checkbox"/>
Start 5							2620	2190	68.2	2800	2010	95.4						99.9	<input checked="" type="checkbox"/>	26	<input type="checkbox"/>
End 4							2870	2510	68.2	2990	2270	95.4						99.9	<input checked="" type="checkbox"/>	25	<input type="checkbox"/>
Beta-37451	2950	2740	100.0	3140	2540	100.0	3040	2790	68.2	3200	2700	95.4	96.3					99.9	<input checked="" type="checkbox"/>	24	<input type="checkbox"/>
Beta-38173	3070	2880	100.0	3160	2850	100.0	3070	2910	68.2	3170	2860	95.4	100					99.9	<input checked="" type="checkbox"/>	23	<input type="checkbox"/>
TX-6703	3370	3000	100.0	3450	2870	100.0	3330	3030	68.2	3440	2900	95.4	100.6					99.8	<input checked="" type="checkbox"/>	22	<input type="checkbox"/>
TX-6704	4840	3980	100.0	5300	3630	100.0	4680	3980	68.2	4980	3650	95.4	106.2					99.6	<input checked="" type="checkbox"/>	21	<input type="checkbox"/>
TX-6705	4850	4520	100.0	4960	4420	100.0	4790	4520	68.2	4890	4420	95.4	97.5					99.8	<input checked="" type="checkbox"/>	20	<input type="checkbox"/>
▲ 4																			<input checked="" type="checkbox"/>	19	<input type="checkbox"/>
Start 4							5340	4710	68.2	5800	4530	95.4						99.8	<input checked="" type="checkbox"/>	18	<input type="checkbox"/>
End 3							6100	5330	68.2	6370	4910	95.4						99.7	<input checked="" type="checkbox"/>	17	<input type="checkbox"/>
Beta-37452	6280	5710	100.0	6470	5480	100.0	6350	5860	68.2	6620	5640	95.4	95.1					99.6	<input checked="" type="checkbox"/>	16	<input type="checkbox"/>
GX-15892	6900	6270	100.0	7280	5910	100.0	6960	6290	68.2	7280	5990	95.4	102					99.7	<input checked="" type="checkbox"/>	15	<input type="checkbox"/>
Beta-46192	6750	6490	100.0	6900	6390	100.0	6770	6500	68.2	6910	6390	95.4	100.1					99.8	<input checked="" type="checkbox"/>	14	<input type="checkbox"/>
Beta-37618	7790	7600	100.0	7930	7560	100.0	7780	7600	68.2	7890	7520	95.4	101.6					99.9	<input checked="" type="checkbox"/>	13	<input type="checkbox"/>
▲ 3																			<input checked="" type="checkbox"/>	12	<input type="checkbox"/>
Start 3							8580	7760	68.2	9310	7630	95.4						99.7	<input checked="" type="checkbox"/>	11	<input type="checkbox"/>
End 2							9790	8550	68.2	10050	7980	95.4						99.7	<input checked="" type="checkbox"/>	10	<input type="checkbox"/>
Beta-63007	10160	9700	100.0	10190	9600	100.0	10100	9750	68.2	10170	9630	95.4	100.2					99.9	<input checked="" type="checkbox"/>	9	<input type="checkbox"/>
▲ 2																			<input checked="" type="checkbox"/>	8	<input type="checkbox"/>
Start 2							13970	10130	68.2	16920	9810	95.4						98.8	<input checked="" type="checkbox"/>	7	<input type="checkbox"/>
End 1							18160	14170	68.2	18640	11310	95.4						98.5	<input checked="" type="checkbox"/>	6	<input type="checkbox"/>
Beta-38694	18720	18090	100.0	18920	17930	100.0	18680	18080	68.2	18960	17790	95.4	97.6					99.6	<input checked="" type="checkbox"/>	5	<input type="checkbox"/>
▲ 1																			<input checked="" type="checkbox"/>	4	<input type="checkbox"/>
Start 1							20610	18370	68.2	24020	18010	95.4						95.9	<input checked="" type="checkbox"/>	3	<input type="checkbox"/>
▲																			<input checked="" type="checkbox"/>	2	<input type="checkbox"/>

OxCal v4.1.7 Bronk Ramsey (2010); r.5 intr whole Atmospheric data from Reimer et al. (2009).




OxCal v4.1.7 Bronk/Ramsey (2010); r.5 intr. whole Atmospheric data from Reimer et al. (2009);



A & M Study Area (Waters & Nordt 1995: Table 1)


Name	Unmodelled (BP)						Modelled (BP)						Indices			Select	Page break				
	from	to	%	from	to	%	from	to	%	from	to	%	Acomb	A	L			P	C	All Visible	
End 4							460	290	68.2	500	140	95.4						98.1	<input checked="" type="checkbox"/>	28	<input type="checkbox"/>
GX-15421	440	150	100.1	500	-10	100.0	500	380	68.2	510	290	95.4		96.4				99.8	<input checked="" type="checkbox"/>	27	<input type="checkbox"/>
GX-15419	490	290	100.0	510	-10	99.9	500	390	68.2	510	310	95.4		102.3				99.8	<input checked="" type="checkbox"/>	26	<input type="checkbox"/>
GX-15420	520	320	100.0	540	300	100.0	520	330	68.2	530	320	95.4		108.1				99.9	<input checked="" type="checkbox"/>	25	<input type="checkbox"/>
GX-15422	530	340	100.0	560	300	100.0	530	340	68.2	540	320	95.4		109.3				99.9	<input checked="" type="checkbox"/>	24	<input type="checkbox"/>
GX-15416	630	510	100.0	660	480	100.0	560	460	68.2	630	320	95.4		75.1				99.8	<input checked="" type="checkbox"/>	23	<input type="checkbox"/>
▲ 4																			<input checked="" type="checkbox"/>	22	<input type="checkbox"/>
Start 4							610	470	68.2	700	340	95.4						99.7	<input checked="" type="checkbox"/>	21	<input type="checkbox"/>
End 3							810	590	68.2	890	510	95.4						99.9	<input checked="" type="checkbox"/>	20	<input type="checkbox"/>
A-6400	910	730	100.0	930	680	100.0	910	740	68.2	930	720	95.4		98.4				99.8	<input checked="" type="checkbox"/>	19	<input type="checkbox"/>
GX-15418	1380	1060	100.0	1560	920	100.0	1390	1010	68.2	1610	910	95.4		100.2				99.5	<input checked="" type="checkbox"/>	18	<input type="checkbox"/>
A-7509	1420	1180	100.0	1610	1000	100.0	1520	1170	68.1	1620	980	95.4		99.9				99.7	<input checked="" type="checkbox"/>	17	<input type="checkbox"/>
GX-15415	2750	2460	100.0	2760	2350	100.0	2650	2360	68.2	2740	2350	95.4		97.3				99.8	<input checked="" type="checkbox"/>	16	<input type="checkbox"/>
▲ 3																			<input checked="" type="checkbox"/>	15	<input type="checkbox"/>
Start 3							3150	2470	68.2	3900	2370	95.4						99.4	<input checked="" type="checkbox"/>	14	<input type="checkbox"/>
End 2							4720	3700	68.2	4810	2940	95.4						99.5	<input checked="" type="checkbox"/>	13	<input type="checkbox"/>
A-7510	4840	4620	100.0	4860	4520	100.0	4840	4640	68.2	4860	4570	95.4		100.8				99.8	<input checked="" type="checkbox"/>	12	<input type="checkbox"/>
A-6401	7480	7300	100.0	7580	7170	100.0	7480	7290	68.2	7580	7170	95.4		99.8				99.7	<input checked="" type="checkbox"/>	11	<input type="checkbox"/>
A-7511	9250	9000	100.0	9300	8810	100.0	9200	8810	68.2	9270	8770	95.4		97.7				99.7	<input checked="" type="checkbox"/>	10	<input type="checkbox"/>
▲ 2																			<input checked="" type="checkbox"/>	9	<input type="checkbox"/>
Start 2							9340	9060	68.2	9480	8880	95.4						99.9	<input checked="" type="checkbox"/>	8	<input type="checkbox"/>
End 1							9500	9270	68.2	9570	9070	95.4						99.8	<input checked="" type="checkbox"/>	7	<input type="checkbox"/>
GX-15417	9690	9000	100.0	10240	8540	100.0	9570	9320	68.2	9830	9160	95.4		133.1				99.8	<input checked="" type="checkbox"/>	6	<input type="checkbox"/>
A-7513	9540	9420	100.0	9560	9270	100.0	9550	9400	68.2	9660	9260	95.4		106.6				99.8	<input checked="" type="checkbox"/>	5	<input type="checkbox"/>
▲ 1																			<input checked="" type="checkbox"/>	4	<input type="checkbox"/>
Start 1							9730	9370	68.2	10330	9210	95.4						96.7	<input checked="" type="checkbox"/>	3	<input type="checkbox"/>
▲																			<input checked="" type="checkbox"/>	2	<input type="checkbox"/>

OxCal v4.1.7 Bronk Ramsey (2010); r.5 intr Atmospheric data from Reimer et al (2009);




A&M Unit I

Name	Unmodelled (BP)						Select	Page break
	from	to	%	from	to	%		
R_Date A&M Unit I	21450	20600	68.2	21500	20540	95.4	<input checked="" type="checkbox"/> 2	<input type="checkbox"/>



Royalty Paleosol

Name	Unmodelled (BP)							Select	Page break
	from	to	%	from	to	%	m		
Show all Show structure								All Visible	
R_Date GX-15762	9410	9120	68.2	9470	9010	95.4	9240	<input checked="" type="checkbox"/> 2	<input type="checkbox"/>




APPENDIX V: Trinity River Basin OxCal Results


Upper Extent Trinity River assays (Ferring 1995a)

Name	Unmodelled (BP)						Modelled (BP)						Indices				Select	Page break	
	from	to	%	from	to	%	from	to	%	from	to	%	A _{model=99}	A _{overall=100}	L	P			C
End 3							520	280	68.2	570	30	95.4					99	<input checked="" type="checkbox"/> 68	<input type="checkbox"/>
Beta-14963	630	500	100.0	660	330	100.0	630	500	68.2	660	420	95.4	105.9				99.9	<input checked="" type="checkbox"/> 67	<input type="checkbox"/>
Beta-32522	630	510	100.0	660	480	100.0	630	520	68.2	660	480	95.4	101.7				99.9	<input checked="" type="checkbox"/> 66	<input type="checkbox"/>
Beta-22488	670	510	100.0	700	480	100.0	660	530	68.2	730	480	95.4	103.3				99.8	<input checked="" type="checkbox"/> 65	<input type="checkbox"/>
TX-4001	790	510	100.0	980	300	100.0	870	550	68.2	1060	430	95.4	106.8				99.8	<input checked="" type="checkbox"/> 64	<input type="checkbox"/>
Beta-22487	680	560	100.0	690	550	100.0	670	570	68.2	710	550	95.4	99.5				99.9	<input checked="" type="checkbox"/> 63	<input type="checkbox"/>
Beta-14907	730	570	100.0	800	540	100.0	750	590	68.2	880	550	95.4	100.1				99.9	<input checked="" type="checkbox"/> 62	<input type="checkbox"/>
Beta-32521	740	660	100.0	800	560	100.0	770	660	68.2	890	560	95.4	99.9				99.9	<input checked="" type="checkbox"/> 61	<input type="checkbox"/>
Beta-32529	740	660	100.0	800	560	100.0	770	660	68.2	890	560	95.4	100				99.9	<input checked="" type="checkbox"/> 60	<input type="checkbox"/>
Beta-16416	940	790	100.0	980	730	100.0	930	800	68.2	1000	740	95.4	99.9				99.9	<input checked="" type="checkbox"/> 59	<input type="checkbox"/>
Beta-22028	960	790	100.0	1060	730	100.0	960	800	68.2	1050	740	95.4	99.9				99.9	<input checked="" type="checkbox"/> 58	<input type="checkbox"/>
Beta-13962	1180	920	100.0	1290	780	100.0	1160	910	68.2	1260	780	95.4	99.9				99.7	<input checked="" type="checkbox"/> 57	<input type="checkbox"/>
Beta-32523	1290	1090	100.0	1310	1010	100.0	1270	1100	68.2	1300	1010	95.4	99.9				99.9	<input checked="" type="checkbox"/> 56	<input type="checkbox"/>
Beta-32986	1300	1070	100.0	1350	980	100.0	1280	1090	68.2	1350	990	95.4	100				99.8	<input checked="" type="checkbox"/> 55	<input type="checkbox"/>
Beta-32530	1300	1070	100.0	1350	980	100.0	1280	1090	68.2	1350	990	95.4	99.9				99.8	<input checked="" type="checkbox"/> 54	<input type="checkbox"/>
Beta-14908	1300	1170	100.0	1310	1060	100.0	1280	1150	68.2	1320	1070	95.4	99.8				99.9	<input checked="" type="checkbox"/> 53	<input type="checkbox"/>
GX-18282	1360	1170	100.0	1520	980	100.0	1380	1120	68.2	1510	1000	95.4	100				99.8	<input checked="" type="checkbox"/> 52	<input type="checkbox"/>
Beta-14904	1400	1270	100.0	1530	1140	100.0	1430	1220	68.2	1530	1090	95.4	100				99.8	<input checked="" type="checkbox"/> 51	<input type="checkbox"/>
Beta-32524	1380	1280	100.0	1520	1170	100.0	1400	1260	68.2	1510	1160	95.4	100				99.8	<input checked="" type="checkbox"/> 50	<input type="checkbox"/>
Beta-32982	1700	1520	100.0	1740	1400	100.0	1680	1490	68.2	1780	1410	95.4	99.9				99.9	<input checked="" type="checkbox"/> 49	<input type="checkbox"/>
SMU-2404	1710	1560	100.0	1820	1520	100.0	1710	1570	68.2	1800	1530	95.4	100				99.9	<input checked="" type="checkbox"/> 48	<input type="checkbox"/>
Beta-16417	1780	1540	100.0	1880	1410	100.0	1780	1550	68.2	1870	1430	95.4	100.1				99.9	<input checked="" type="checkbox"/> 47	<input type="checkbox"/>
Beta-26737	1870	1630	100.0	1930	1560	100.0	1840	1650	68.2	1920	1570	95.4	100				99.9	<input checked="" type="checkbox"/> 46	<input type="checkbox"/>
Beta-32513	2000	1870	100.0	2130	1810	100.0	2040	1860	68.2	2130	1760	95.4	100				99.8	<input checked="" type="checkbox"/> 45	<input type="checkbox"/>
SMU-2403	2150	1940	100.0	2310	1880	100.0	2150	1960	68.2	2300	1890	95.4	100				99.8	<input checked="" type="checkbox"/> 44	<input type="checkbox"/>
Beta-32516	2360	2310	100.0	2680	2150	100.0	2470	2210	68.2	2680	2150	95.4	99.7				99.8	<input checked="" type="checkbox"/> 43	<input type="checkbox"/>
Beta-32515	2760	2720	100.0	2790	2540	100.0	2760	2580	68.2	2830	2500	95.4	99.5				99.9	<input checked="" type="checkbox"/> 42	<input type="checkbox"/>
Beta-32514	2930	2770	100.0	3000	2750	100.0	2930	2790	68.2	3020	2760	95.4	99.9				99.9	<input checked="" type="checkbox"/> 41	<input type="checkbox"/>
Beta-16418	3010	2780	100.0	3210	2740	100.0	3060	2820	68.2	3200	2760	95.4	99.9				99.8	<input checked="" type="checkbox"/> 40	<input type="checkbox"/>
Beta-16526	3390	2760	100.0	3690	2360	100.0	3380	2800	68.2	3700	2450	95.4	100				99.5	<input checked="" type="checkbox"/> 39	<input type="checkbox"/>
SMU-2402	3570	3390	100.0	3680	3350	100.0	3580	3410	68.2	3670	3360	95.4	100				99.9	<input checked="" type="checkbox"/> 38	<input type="checkbox"/>
SMU-2401	4580	4420	100.0	4810	4400	100.0	4580	4430	68.2	4770	4310	95.4	107.3				99.9	<input checked="" type="checkbox"/> 37	<input type="checkbox"/>
Beta-26738	4840	4520	100.0	4880	4410	100.0	4710	4460	68.2	4820	4340	95.4	89.2				99.8	<input checked="" type="checkbox"/> 36	<input type="checkbox"/>
▲ 3																	<input checked="" type="checkbox"/> 35	<input type="checkbox"/>	
Start 3							4900	4570	68.2	5190	4480	95.4					99.5	<input checked="" type="checkbox"/> 34	<input type="checkbox"/>
End 2							6430	5640	68.2	6580	5000	95.4					99.6	<input checked="" type="checkbox"/> 33	<input type="checkbox"/>
Beta-34051	6560	6390	100.0	6670	6300	100.0	6610	6400	68.2	6680	6320	95.4	97.6				99.8	<input checked="" type="checkbox"/> 32	<input type="checkbox"/>
Beta-46151	7430	7250	100.0	7510	7160	100.0	7420	7230	68.2	7510	7070	95.4	99.9				99.8	<input checked="" type="checkbox"/> 31	<input type="checkbox"/>
Beta-34049	7440	7260	100.0	7570	7160	100.0	7460	7260	68.2	7560	7100	95.4	100				99.8	<input checked="" type="checkbox"/> 30	<input type="checkbox"/>
SMU-2339	7940	7320	100.0	8200	6990	100.0	7930	7320	68.2	8270	6960	95.4	100				99.5	<input checked="" type="checkbox"/> 29	<input type="checkbox"/>
SMU-2400	8380	8180	100.0	8410	8160	100.0	8350	8200	68.2	8390	8090	95.4	100				99.9	<input checked="" type="checkbox"/> 28	<input type="checkbox"/>
Beta-46150	9550	9430	100.0	9730	9270	100.0	9640	9360	68.2	9870	9180	95.4	100.1				99.8	<input checked="" type="checkbox"/> 27	<input type="checkbox"/>
Beta-14905	10250	9700	100.0	10550	9530	100.0	10260	9750	68.2	10530	9570	95.4	100				99.6	<input checked="" type="checkbox"/> 26	<input type="checkbox"/>
GX-18281	11190	10290	100.0	11400	10170	100.0	11120	10410	68.2	11580	10110	95.4	100				99.6	<input checked="" type="checkbox"/> 25	<input type="checkbox"/>
SMU-2399	11180	10690	100.0	11240	10510	100.0	11100	10710	68.2	11210	10530	95.4	100				99.7	<input checked="" type="checkbox"/> 24	<input type="checkbox"/>


OxCal v4.1.8 Bronk Ramsey (2010); r.5 intr whole Atmospheric data from Reimer et al (2009)



OxCal v4.1.6 Bronk Ramsey (2010); r.5 intr. whole Atmospheric data from Reimer et al (2009)



OxCal v4.1.8 Bronk Ramsey (2010); r.6 inr whole Atmospheric data from Reimer et al. (2009);



APPENDIX VI-Categorized Bibliography

BRAZOS RIVER DRAINAGE BASIN (UPPER EXTENT)

Blum et al. 1992
Ferring 1995
Hall 1990a
Haas et al. 1986
Holliday 1985, 1988, 1995, 1997, 2000, 2009
Holliday and Johnson 1981, 1983, 1986
Holliday et al. 1983, 1985
Holliday et al. 1999
Johnson and Holliday 1980
Mandel 1992
Stafford 1981, 1983
Stricklen 1961

BRAZOS RIVER DRAINAGE BASIN (MIDDLE-LOWER EXTENT)

Abbott 2001, 2003
Alexander 2008
Bernard et al. 1962
Bernard et al. 1970
Bongino 2007
Bousman 1998
Bronaugh 1950
Brotherton 1978
Brownlow 2003
Campbell and Johnson 2004
Collins 1998
Epps 1973
Fields 1990

Fields et al. 1991
Hilliard 1997,2000
Gadus et al. 2006
Gibson 1997
Goldberg and Holliday 1998
Hill 1901
Husain 1998
Mahoney and Tomka 2001
Mahoney et al. 2003
Mehalchick 2000
Nordt 1983, 1986, 1992, 1993, 1995, 1996, 2001b, 2003, 2004
Nordt et al. 1994
Nordt et al. 1998
Patton 1987
Pearl 1997
Prochnow 2001
Ricklis 2001
Stafford 1998
Sylvia and Galloway 2006
Taha and Anderson 2008
Tharp 1988
Urista 2009
Voellinger 1990
Waters and Nordt 1995, 1996
Wilkinson and Basse 1978
Woolly 1985

COASTAL AREAS

Anderson et al. 2004
Aten 1983
Blum 1993

Blum and Aslan 2006
Blum et al. 2001
Blum and Price 1994, 1998
Blum and Törnqvist 2006
Frazier 1974
Leeder and Stewart 1996
Lowman 1949
Otvos 2004
Paine 1991
Ricklis and Blum 1997
Schumm 1998
Simms et al. 2006
Simms et al. 2007
Sylvia and Galloway 2006
Wilkinson and Basse 1978
Winker 1979

COLORADO RIVER DRAINAGE BASIN (UPPER EXTENT)

Blum and Lintz 1993
Blum and Valastro 1992
Frederick 1996
Frederick and Boutton 1996
Lintz et al. 1993
Lintz et al. 1991
Nordt and Bousman 2001
Quigg and Peck 1995
Quigg et al. 1996

COLORADO RIVER DRAINAGE BASIN (MIDDLE-LOWER EXTENT)

Abbott 1994
Baker and Penteado-Orellana 1977, 1978
Blum 1987, 1992, 1993

Blum et al. 1989
Blum et al. 1994
Blum and Valastro 1989, 1992, 1994
Caran and Baker 1986
Crawford and Frederick 2006
Fiore 1976
Frederick 1987
Kastning 1983
Largent 1991
Looney 1977
Mandel 1980
Ricklis and Collins 1995
Tinkler 1971
Urbanec 1963
Wallis 1976
Weber 1968
Weeks 1945

GUADALUPE RIVER DRAINAGE BASIN

Abbott 2008
Aiuvalasit 2006, 2007
Brown 2006
Cooke 2005
Cooke et al. 2007
Cooke et al. 2003
Frederick 2008
Houk et al. 2008
Nickels and Bousman 2010
Oksanen 2008
Ringstaff 2000
Schroeder and Oksanen 2002

Toomey 1993

Toomey et al. 1993

NECHES RIVER DRAINAGE BASIN (EAST TEXAS)

Phillips and Marion 2001

NUECES RIVER DRAINAGE BASIN

Baskin and Cornish 1989

Brown et al. 1982

Bunker 1982

Cornish and Baskin 1995

Decker et al. 2000

Durbin 1999

Gustavson 1978

Hall et al. 1986

Highley 1986

Holliday 1995

Johnson 1933

Lukowski 1987

Mear 1953, 1990, 1995, 1998

Prewitt and Paine 1987

Paine 1991

Price 1933

Ricklis 1988, 1993, 2004

Ricklis and Blum 1997

Ricklis and Cox 1998

Sayles 1935

Scott and Fox 1982

Simms 2005

Taylor and Highley 1995

Taylor 1995

Weeks 1933, 1945

RIO GRANDE-PECOS RIVER DRAINAGE BASIN

Buck 1996

Gustavson and Collins 1998

Huffington and Albritton 1941

Kochel 1988

Patton and Dibble 1982

Thomas 1972

Vierra 1998

Quigg 2000

Young et al. 1999

SABINE RIVER DRAINAGE BASIN (EAST TEXAS)

Alford and Holmes 1985

SAN ANTONIO RIVER DRAINAGE BASIN

Black et al. 1998

Collins et al. 2003

Frederick 2001

Houk and Nickels 1997

Hudler 2000

Johnson 1995

Johnson and Goode 1994

Mandel et al. 2007

Nickels et al. 2001

Nordt et al. 2002

Nordt 2001a

Osburn and Kuehn 2006

Tennis and Hard 1995

Tennis 1996

SAN JACINTO-(HOUSTON AREA) RIVER DRAINAGE BASIN

Abbott 2001

Frederick 2007

Voellinger et al. 1987

SOUTHERN HIGH PLAINS-LLANO ESTACADO REGION

Brown 1991

Caran and Baumgardner 1990

Caran 1991

Gustavson 1986a, 1986b

Gustavson et al. 1991

Holliday 1985, 1995, 1997, 2009

Holliday et al. 1994

Holliday and Mandel 2006

Holliday et al. 2008

Madole et al. 1991

Meltzer 1991

Stafford 1981

SULPHUR RIVER DRAINAGE BASIN (EAST TEXAS)

Bousman et al. 1988

Bousman and Skinner 2007

Darwin et al. 1990

Fields et al. 1993

Ferring 1995

Gadus et al. 1992

Gadus et al. 1991

Jacobs 1981

Journey et al. 1993

Rainey 1974

TRINITY RIVER DRAINAGE BASIN

Aten 1983

Bernard 1962

Bernard et al. 1970
Byers 2007
Fields 1990
Fields et al. 1991
Ferring 1990a, 1990b, 1991, 1994, 1995a, 1995b, 2000, 2001
Ferring and Yates 1997
Garvin 2008
Humphrey and Ferring 1994
McEwen 1963
Prikryl 1990
Prikryl and Yates 1987
Shuler 1935
Stovall and McAnulty 1950
Williams 2004

METHODS RADIOCARBON-STATISTICS

Acabado 2009
Banning 2000
Bayliss and Ramsey 2004
Blackwell and Buck 2008
Blackwell et al. 2006
Blockley et al. 2007
Bowman 1990
Breyer 2009
Buck et al. 2006
Buck and Blackwell 2004
Buck et al. 1996
Buck et al. 1994
Buck et al. 1992
Buck et al. 1991
Buck 2004

Chiverrell et al. 2009
Christian 1994
Everitt 2002
Fiedel 1999
Fowler et al. 1986
Grimm et al. 2009
Heaton et al. 2009
Hedman 2007
Hua 2009
Huntley 1985
Ilan 2001
Jacobson et al. 2003
Klein et al. 1982
Libby 1955
Long and Rippeteau 1974
Lowe and Walker 1997
Martin and Johnson 1995
Matthews 1985
Michael and Klein 1979
McCormac and Baillie 1993
Mook and Waterbolk 1985
Nash 1999
Ralph et al. 1973
Ramsey 1995, 1998, 2001, 2008, 2009
Reimer et al. 2009
Steier et al. 2001
Stuiver and Braziunas 1993
Stuiver and Polach 1977
Stuiver and Suess 1966
Talma and Vogel 1993
Taylor 1996

Taylor 1987, 1997, 2009
Taylor et al. 1996
Upton and Cook 2006
Van den berg 1995.
Vandenberghe 1995
Van der Plicht and Hogg 2006
Walker 2005
Zhou and Chafertz 2010

PALEOENVIRONMENT-PALEOCLIMATE-EUSTASY

Abdulah et al. 2004
Anderson et al. 2004
Baker et al. 2001
Baker et al. 2000
Baker et al. 1998
Banfield and Anderson 2004
Barber et al. 1999
Bartlein et al. 1995
Blum 1993, 2007
Blum and Aslan 2006
Blum et al. 2001
Blum and Price 1994, 1998
Blum and Straffin 2001
Blum and Törnqvist 2006
Bousman 1998a, 1998b
Bousman and Oksanen in press
Brown 1998
Bryant and Holloway 1985
Bryant and Shafer 1977
Bull 1991, 2000
Butzer et al. 2008

Caran 1998
Chiverrell et al. 2009
Clark et al. 2002
Collins 1995, 2004
Cooke 2005
Cooke et al. 2003
Daniels and Knox 2005
Dillehay 1974
Durbin 1999
Etheridge et al. 1998
Fairbanks 1989
Frederick 1995
González 2008
González and Törnqvist 2009
Hall 1990a, 1990b
Hodell et al. 2001
Howard et al. 2009
Hu et al. 1999
Huckleberry and Duff 2008
Johnson and Goode 1994
Johnson and Logan 1990
Johnstone et al. 2006
Knighton 1998
Knox 2000
Leeder and Stewart 1996
Lewin et al. 2005
Macklin et al. 2006
Macklin et al. 2002
Macklin et al. 2010
Macklin et al. 2005
Macklin and Lewin 2008

Mallouf 1981
Mayewski et al. 2004
Mock and Bartlein 1995
Musgrove 2000
Musgrove et al. 2001
Neck 1983, 1987
Nordt et al. 2007
Otvos 2004
Paine 1991, 1993
Rogers 1984
Schmieder 2009
Schumm 1993, 2003
Shuman et al. 2009
Simms et al. 2007
Smith 1992
Snow 1998
Stine 1994
Story 1990
Sylvia and Galloway 2006
Taha and Anderson 2008
Thomas and Anderson 1994
Toomey et al. 1993
Törnqvist et al. 2004
Van de Plassche et al. 1998
Wang et al. 1996
Wohlfarth 1996
Yu and Wright 2001
Zaitlin et al. 1994

REFERENCES

Abbott, James T.

- 1994 Geomorphic Context of the Barton Site (41HY202) and the Mustang Branch Site (41HY209). In *Archaic and Late Prehistoric Human Ecology in the Middle Onion Creek Valley, Hays County, Texas*. by R. A. Ricklis and M. B. Collins, 2:353–79. 2 vols. Studies in Archeology No. 19. Austin: Texas Archeological Research Laboratory, University of Texas at Austin.
- 2001 *Houston Area Geoarcheology: A Framework for Archeological Investigation, Interpretation, and Cultural Resource Management in the Houston Highway District*. Texas Department of Transportation, Environmental Affairs Division, Archeological Studies Program, Report 27. Austin, Texas.
- 2003 Appendix K: Stratigraphic and Geomorphic Evaluation of Brushy Creek at FM 973. In *Archeological Investigations at 41WM815: A Blackland Prairie Site, Williamson County, Texas*, by Russell K. Brownlow, pp. 180–191. Texas Department of Transportation, Environmental Affairs Division, Archeological Studies Program, Report 23. Austin, Texas.
- 2008 Stratigraphic and Geoarchaeological Investigations of the Guadalupe River: Interstate 35, Comal County, and Spur 98, Kerr County, Texas. In *The Gatlin Site (41KR621): Investigating Archaic Lifeways on the Southern Edwards Plateau of Central Texas*. Brett A. Houk, Kevin A. Miller, and Eric R. Oksanen, Appendix A:1–28. SWCA Cultural Resource Report No. 08-149, SWCA, Inc. Austin, Texas.

Abdulah, Kenneth C., John B. Anderson, Jennifer N. Snow, and Lynette Holdford-Jack

- 2004 The Late Quaternary Brazos and Colorado Deltas, Offshore Texas, U.S.A.—Their Evolution and the Factors That Controlled Their Deposition. In: *Late Quaternary Stratigraphic Evolution of the Northern Gulf of Mexico Margin* by John B. Anderson and Richard H. Fillon (editors), pp. 237–269. SEPM (Society for Sedimentary Geology), Special Publication No. 79, Tulsa, OK.

Acabado, S.

- 2009 A Bayesian Approach to Dating Agricultural Terraces: A Case From the Philippines. *Antiquity* 83:801–814.

Aiuvalasit, M. J.

- 2006 *Geoarchaeological Investigations at the McNeill-Gonzales Site (41VT141), Victoria County, Texas*. Unpublished Master's Thesis. Department of Anthropology. Texas A & M University, College Station, Texas.
- 2007 Geoarcheology of the McNeill Ranch Site: Implications for Paleoindian Studies of the Gulf Coastal Plain of Texas. *Bulletin of the Texas Archeological Society* 78:47–64.

- Alexander, Dawn Aileen Joyce
 2008 Geoarchaeological Investigation of Natural Formation Processes to Evaluate Context of the Clovis Component at the Gault Site (41BL323), Bell County, Texas. Unpublished Master's Thesis. Texas A & M University, College Station, Texas.
- Anderson, John B. Antonio Rodriguez, Kenneth C. Abdulah, Richard H. Fillon, Laura A. Banfield, Heather A. McKeown, and Julia S. Wellner
 2004 Late Quaternary Stratigraphic Evolution of the Northern Gulf of Mexico Margin: A Synthesis. In: *Late Quaternary Stratigraphic Evolution of the Northern Gulf of Mexico Margin* by John B. Anderson and Richard H. Fillon (editors), pp. 1–23. SEPM (Society for Sedimentary Geology), Special Publication No. 79, Tulsa, OK.
- Aten, Lawrence E.
 1983 *Indians of the Upper Texas Coast*. Academic Press, New York and London.
- Baker, Richard G., E. A. Bettis III, R. F. Dennison, L. A. Gonzalez
 2001 Plant Remains, Alluvial Chronology, and Cave Speleothem Isotopes Indicate Abrupt Holocene Climate Change at 6 ka in Midwestern USA. *Global and Planetary Change* 28:285–291.
- Baker, Richard G., Glen G. Fredlund, Rolfe D. Mandel, and E. A. Bettis III
 2000 Holocene Environments of the Central Great Plains: Multi-proxy Evidence from Alluvial Sequences, Southeastern Nebraska. *Quaternary International* 67:75–88.
- Baker, R. G., L. A. Gonzalez, M. Raymo, E. A. Bettis III, M. K. Reagan, and J. A. Dorale
 1998 Comparison of Multiple Proxy Records of Holocene Environments in the Midwestern United States. *Geology* 26(12):1131–1134.
- Baker, Victor R. and Margarida M. Penteado-Orellana
 1977 Adjustment to Quaternary Climatic Change by the Colorado River in Central Texas. *Journal of Geology* 85:395–422.
- 1978 Fluvial Sedimentation Conditioned by Quaternary Climatic Change in Central Texas. *Journal of Sedimentary Petrology* 48(2):433–451.
- Banfield, Laura A. and John B. Anderson
 2004 Late Quaternary Evolution of the Rio Grande Delta: Complex Response to Eustasy and Climate Change. In: *Late Quaternary Stratigraphic Evolution of the Northern Gulf of Mexico Margin* by John B. Anderson and Richard H. Fillon (editors), pp. 289–306. SEPM (Society for Sedimentary Geology), Special Publication No. 79, Tulsa, OK.
- Banning, J.
 2000 Interpreting Radiocarbon Results. In: *The Archaeologist's Laboratory: The Analysis of Archaeological Data*. 15:265–275. Kluwer Academic/ Plenum Publishers, New York, Boston, Dordrecht, London, Moscow.

- Baskin, Jon A. and Frank G. Cornish
 1989 Late Quaternary Fluvial Deposits and Vertebrate Paleontology, Nueces River Valley, Gulf Coastal Plain, South Texas. In *South Texas Clastic Depositional Environments*, by J. A. Baskin and J. S. Pouty (editors), pp. 23–30. Gulf Coast Association Geological Society 1989 Field Trip Guidebook, Corpus Christi Geological Society, Corpus Christi, Texas.
- Barber, D. C., A. Dyke, C. Hillaire-Marcel, A. E. Jennings, J. T. Andrews, M. W. Kerwin, G. Bilodeau, R. McNeely, J. Southons, M. D. Morehead, and J.M. Gagnon
 1999 Forcing of the Cold Event of 8,200 Years Ago by Catastrophic Drainage of Laurentide Lakes. *Nature* 400:344–348.
- Bartlein, Patrick J., Mary E. Edwards, Sarah L. Shafer, and Edward D. Barker, Jr.
 1995 Calibration of Radiocarbon Ages and the Interpretation of Paleoenvironmental Records. *Quaternary Research* 44:417–424.
- Bayliss, A. and C. B. Ramsey
 2004 Pragmatic Bayesians: a Decade of Integrating Radiocarbon Dates into Chronological Models. In: *Tools for Constructing Chronologies: Crossing Disciplinary Boundaries*, by C. E. Buck and A. R. Millard (editors), Chapter 2 pp. 25–41. Springer publishers. London, Berlin, Heidelberg, New York.
- Bernard, H. A., R. J. LeBlanc, and C. F. Major
 1962 *Recent and Pleistocene Geology of Southeast Texas*. Field Excursion No. 3, Houston Geological Society, Houston.
- Bernard, H. A., C. F. Major, Jr., B.S. Parrott, and R. J. Leblanc, Jr.
 1970 *Recent Sediments of Southeast Texas: A Field Guide to the Brazos Alluvial and Deltaic Plains and the Galveston Barrier Island Complex*. Guidebook 11, Bureau of Economic Geology, The University of Texas, Austin, Texas.
- Black, Stephen, James W. Karbula, Charles D. Frederick, and Charles G. Mear
 1998 *Archaeology Along the Wurzbach Parkway Module 5: Testing and Geoarchaeological Evaluation at the Number-6 Site (41BX996) and Other Sites*. Studies in Archeology 29. Texas Archeological Research Laboratory. The University of Texas, Austin, Texas.
- Black, Stephen and Darrell Creel
 1997 The Central Texas Burned Rock Midden Reconsidered. In: *Hot Rock Cooking on the Greater Edwards Plateau: Four Burned Rock Midden Sites in West Central Texas*. Volume I. Stephen Black, Linda Ellis, Darrell Creel, and Glen Goode (editors). Studies in Archeology 22, Texas Archeological Research Laboratory. The University of Texas, Austin, Texas
- Blackwell, P. G. and C. E. Buck
 2008 Estimating Radiocarbon Calibration Curves. *Bayesian Analysis* 3(2):225–248.
- Blackwell, P. G., C. E. Buck, and P. J. Reimer
 2006 Important Features of the New Radiocarbon Calibration Curves. *Quaternary Science Reviews* 25(5):408–413.

- Blockley, S. P. E., M. Blaauw, C. Bronk Ramsey, and J. van der Plicht
2007 Building and Testing Age Models for Radiocarbon Dates in Lateglacial and Early Holocene Sediments. *Quaternary Science Reviews* 26:1915–1926.
- Blum, Michael D.
1987 *Late Quaternary Sedimentation in the Upper Pedernales River, Texas*. Unpublished Master's Thesis, Department of Geology, University of Texas, Austin.
1992 *Modern Depositional Environments and Recent Alluvial History of the Lower Colorado River, Gulf Coastal Plain of Texas*. Ph.D. Dissertation, Department of Geology, University of Texas, Austin.
1993 Genesis and Architecture of Incised Valley Fill Sequences: A Late Quaternary Example from the Colorado River, Gulf Coastal Plain of Texas. In: *Siliclastic Sequence Stratigraphy: Recent Developments and Applications*. (P. Weimer and H. Posamentier editors) 10:259–283. AAPG (American Association of Petroleum Geologists), Memoir No. 58, Tulsa, OK.
2007 Large River Systems and Climate Change. In: *Large Rivers: Geomorphology and Management* by A. Gupta (editor), pp. 627–659. John Wiley & Sons, Ltd. New York, Chichester, Brisbane, Toronto, Singapore.
- Blum, Michael D., James T. Abbott, and Salvatore Valastro, Jr.
1992 Evolution of Landscapes on the Double Mountain Fork of the Brazos River, West Texas: Implications for Preservation and Visibility of the Archaeological Record. *Geoarchaeology* 7(4):339–370.
- Blum, Michael D. and Andres Aslan
2006 Signatures of Climate vs. Sea-level Change within Incised Valley-fill Successions: Quaternary Examples from the Texas Gulf Coast. *Sedimentary Geology* 190:177–211.
- Blum, Michael D. and Christopher W. Lintz
1993 Late Quaternary Geology in the Reservoir Basin. In: *Cultural Resource Investigations in the O. H. Ivie Reservoir, Concho, Coleman, and Runnels Counties, Texas. Volume I: Project Introduction, Setting and Methods*. by C. Lintz, N. Trierweiler, F. Oglesby, M. Blum, P. O'Neill, J. Kuhl, R. Holloway, L. Scott-Cummings, and D. Scurlock 5.2:280–314. Technical Report No. 346-I. Mariah Associates Inc. Austin, Texas.
- Blum, Michael D., Tamara J. Misner, Eric S. Collins, David B. Scott, Robert A. Morton, and Andres Aslan
2001 Middle Holocene Sea-Level Rise and Highstand at +2m, Central Texas Coast. *Journal of Sedimentary Research* 71(4):581–588.
- Blum, Michael D., J. F. Petersen, and R. S. Toomey, III
1989 *Geomorphology, Quaternary Stratigraphy, and Paleoecology of Central Texas*. Friends of the Pleistocene, Central Texas. South Central Cell, April 7-9, 1989, 7th Annual Meeting.

- Blum, Michael D. and David M. Price
 1994 Glacio-Eustatic and Climatic Controls on Quaternary Alluvial Plain Deposition, Texas Coastal Plain. *Transactions of the Gulf Coast Association of Geological Societies* 44:85–92.
- 1998 Quaternary Alluvial Plain Construction In Response To Glacio-Eustatic And Climatic Controls, Texas Gulf Coastal Plain. In: *Relative Role of Eustasy, Climate, and Tectonism in Continental Rocks* by K. W. Shanley and P. J. McCabe (editors), pp. 31–48. SEPM (Society for Sedimentary Geology), Special Publication No. 59, Tulsa, OK.
- Blum, Michael D. and Eric C. Straffin
 2001 Fluvial Responses to External Forcing: Examples from the French Massif Central, the Texas Coastal Plain (USA), the Sahara of Tunisia, and the Lower Mississippi Valley (USA). In *River Basin Sediment Systems: Archives of Environmental Change*. (D. Maddy, M. A. Macklin, and J. C. Woodward editors) pp. 195–228. Balkema Press, Netherlands.
- Blum, Michael D., Richard S. Toomey III, and Salvatore Valastro, Jr.
 1994 Fluvial Response to Late Quaternary Climatic and Environmental Change, Edwards Plateau, Texas. *Palaeogeography, Palaeoclimatology, Palaeoecology* 108:1–21.
- Blum, Michael D. and Torbjörn E. Törnqvist
 2006 Fluvial Responses to Climate and Sea-Level Change: A Review and Look Forward. *Sedimentology* 47 (Suppl. 1):2–48.
- Blum, Michael D. and Salvatore Valastro, Jr.
 1989 Fluvial Response to Late Quaternary Climatic and Environmental Change, Edwards Plateau, Texas. *Annals of American Geographers* 79:435–456.
- 1992 Quaternary Stratigraphy and Geoarchaeology of the Colorado and Concho Rivers. *Geoarchaeology* 7(5):419–448.
- 1994 Late Quaternary Sedimentation, Lower Colorado River, Gulf Coastal Plain of Texas. *Geological Society of America Bulletin* 106:1002–1016.
- Bongino, J. D.
 2007 *Late-Quaternary History of the Waco Mammoth Site: Environment Reconstruction and Determining the Cause of Death*. Unpublished Master's Thesis Department of Geology, Baylor University, Waco, Texas.
- Bousman, C. Britt
 1998 Chapter 8: Late Paleoindian Archeology. In: *Wilson-Leonard: An 11,000-year Archeological Record of Hunter-Gatherers in Central Texas. Volume I: Introduction, Background, and Syntheses*. M. Collins (editor), pp. 161–202. Studies in Archeology 31, Texas Archeological Research Laboratory. The University of Texas, Austin, Texas.
- 1998a Paleoenvironmental Change in Central Texas: The Palynological Evidence. *Plains Anthropologist* 43(164):201–219.

- 1998b Chapter 8: Late Paleoindian Archeology. In: *Wilson-Leonard: An 11,000-year Archeological Record of Hunter-Gatherers in Central Texas. Volume I: Introduction, Background, and Syntheses*. M. Collins (editor), pp. 161–202. *Studies in Archeology* 31, Texas Archeological Research Laboratory. The University of Texas, Austin, Texas.
- Bousman, C. Britt and Eric Oksanen
 In press The Paleoindian-Archaic Transition in Central Texas and Surrounding Areas. In *On the Brink*, by C. B. Bousman and B. J. Vierra (editors), pp. 250–279. Texas A&M University Press, College Station.
- Bousman, C. Brit, Michael Collins, and Timothy Pertulla
 1988 *Quaternary Geomorphology at Cooper Basin: A Framework for Archaeological Inquiry, Delta and Hopkins Counties, Texas*. Research Report No. 55. Prewitt & Associates, Inc. Austin, Texas.
- Bousman, C. B. and S. A. Skinner
 2007 The Search for Late Pleistocene Pre-Clovis Archeology in Texas: Problems and Potentials. *Bulletin of the Texas Archeological Society* 78:1–76.
- Bowman, S.
 1990 *Interpreting The Past: Radiocarbon Dating*. University of California Press, Berkeley and Los Angeles.
- Breyer, Laird
 2009 *Laird Breyer's Markov Chain Monte Carlo (MCMC) Pages*, <http://www.lbreyer.com/classic.html> (accessed February 14, 2010).
- Bronaugh, Richmond L.
 1950 *Geology of Brazos River Terraces in McLennan County, Texas*. Unpublished Master's Thesis Department of Geology, University of Texas, Austin, Texas.
- Brotherton, Mark A.
 1978 *Geomorphologic Evolution of the North Bosque River*. Unpublished Master's Thesis Department of Geology, Baylor University, Waco, Texas.
- Brownlow, R. K.
 2003 *Archeological Investigations at 41WM815: A Blackland Prairie Site, Williamson County, Texas*. Texas Department of Transportation, Environmental Affairs Division, Archeological Studies Program, Report 23. Austin, Texas.
- Brown, David O.
 1998 Late Holocene Climates of North-Central Texas. *Plains Anthropologist* 43(164):157–172.
- Brown, Kenneth Martin
 2006 *The Bench Deposits at Berger Bluff: Early Holocene-Late Pleistocene Depositional and Climatic History*. Unpublished Doctoral Dissertation. University of Texas, Austin.

- Brown, Kenneth M., Daniel R. Potter, Grant D. Hall, and Stephen L. Black
 1982 *Excavations at 41LK67 A Prehistoric Site in the Choke Canyon Reservoir, South Texas*. Choke Canyon Series Volume 7, Center for Archaeological Research, University of Texas at San Antonio.
- Buck, Caitlin E., Delil G. P. Aguilar, Cliff D. Litton, and Anthony O'Hagan
 2006 Bayesian Nonparametric Estimation of the Radiocarbon Calibration Curve. *Bayesian Analysis* 1(2):265–288.
- Buck, C. E. and P. G. Blackwell
 2004 Formal Statistical Models For Estimating Radiocarbon Calibration Curves. *Radiocarbon* 46(3):1093–1102.
- Buck, C. E., W. G. Cavanaugh, and C. D. Litton
 1996 *Bayesian Approach to Interpreting Archaeological Data*. John Wiley & Sons, Chichester, New York, Brisbane, Toronto, Singapore.
- Buck, C. E., C. D. Litton, and E. M. Scott
 1994 Making the Most of Radiocarbon Dating: Some Statistical Considerations. *Antiquity* 68:252–263.
- Buck, C. E., C. D. Litton, and A. F. M. Smith
 1992 Calibration of Radiocarbon Results Pertaining to Related Archaeological Events. *Journal of Archaeological Science* 19:497–512.
- Buck, C. E., J. B. Kenworthy, C. D. Litton, and A. F.M. Smith
 1991 Combining Archaeological and Radiocarbon Information: A Bayesian Approach to Calibration. *Antiquity* 65:808–821.
- Buck, C. E.
 2004 Bayesian Chronological Data Interpretation: Where Now? In: *Tools for Constructing Chronologies: Crossing Disciplinary Boundaries*, by C. E. Buck and A. R. Millard (editors), Chapter 1 pp. 1–24. Springer publishers. London, Berlin, Heidelberg, New York.
- Bull, William B.
 1991 *Geomorphic Responses to Climatic Change*. Oxford University Press, New York, Oxford.
- 2000 Correlation of Fluvial Aggradation Events to Times of Global Climate Change. In *Quaternary Geochronology: Methods and Applications*. (J. S. Noller, J. M. Sowers, and W. R. Lettis editors) pp. 456–464. AGU Reference Shelf 4, American Geophysical Union (AGU), Washington, D. C.
- Bunker, Russell C.
 1982 The Geomorphic Framework of Human Occupation in the Frio River Valley, Choke Canyon Dam Reservoir, Live Oak and McMullen Counties, Texas. In *Archaeological Investigations at Choke Canyon Reservoir, South Texas: The Phase I Findings*, by G. D. Hall, S. L. Black, and C. Graves Appendix II. pp. 498–517 Choke Canyon Series Volume 5, Center for Archaeological Research, University of Texas at San Antonio.

- Butzer, Karl W., James T. Abbott, Charles D. Frederick, Paul H. Lehman, Carlos E. Cordova, and John F. Oswalt
 2008 Soil-Geomorphology and “Wet” Cycles in the Holocene Record of North-Central Mexico. *Geomorphology* 101:237–277.
- Bryant, Jr., Vaughn M. and Richard G. Holloway
 1985 A Late-Quaternary Paleoenvironmental Record of Texas: An Overview of the Pollen Evidence. In *Pollen Records of Late-Quaternary North American Sediments*, by Vaughn M. Bryant, Jr. and Richard G. Holloway (editors) pp. 39–70. American Association of Stratigraphic Palynologists Foundation, Austin, Texas.
- Bryant, Jr., Vaughn M. and Harry J. Shafer
 1977 The Late Quaternary Paleoenvironment of Texas: A Model for the Archaeologist. *Bulletin of the Texas Archeological Society* 48:1–25.
- Byers, J. A.
 2007 *The Geologic and Archaeological History of the Dickie Carr Site 41PR26*. Unpublished Master’s Thesis, University of North Texas, Denton, Texas.
- Campbell, J. S. and W. C. Johnson
 2004 *Temporal Predictive Model for Fort Hood, Texas: A Pilot Study in Cowhouse Creek Drainage*. CEMML Project 532496, Department of Geography, University of Kansas.
- Caran, S. Christopher
 1998 Quaternary Paleoenvironmental and Paleoclimatic Reconstruction: A Discussion and Critique, with Examples from the Southern High Plains. *Plains Anthropologist* 43(164):111–124.
- Caran, S. Christopher and Baker, Victor R.
 1986 Flooding along the Balcones Escarpment, Central Texas. In *The Balcones Escarpment: Geology, Hydrology, Ecology, and Social Development in Central Texas*, by Abbott, P. L. and Woodruff, C. M. (editors), pp. 1–14. Comet Reproduction Services. Santa Fe Springs, California.
- Chiverrell, R. C., G. C. Foster, G. S. P. Thomas, P. Marshall, and D. Hamilton
 2009 Robust Chronologies for Landform Development. *Earth Surface Processes and Landforms* 34:319–328.
- Christian, José A.
 1994 *Bayesian Interpretation of Radiocarbon Results*. Unpublished Doctoral Dissertation. The University of Nottingham, England.
- Clark, J. S., E. C. Grimm, J. J. Donovan, S. C. Fritz, D. R. Engstrom, and J. E. Almendinger
 2002 Drought Cycles and Landscape Responses to Past Aridity on Prairies of the Northern Great Plains, USA. *Ecology* 83 (3):595–601.

- Collins, Michael B.
 1995 Forty Years of Archeology in Central Texas. *Bulletin of the Texas Archeological Society* 66:361–400.
- 1998 *Wilson-Leonard: An 11,000-year Archeological Record of Hunter-Gatherers in Central Texas. Volume I: Introduction, Background, and Syntheses*. Studies in Archeology 31, Texas Archeological Research Laboratory. The University of Texas, Austin, Texas.
- 2004 Archeology in Central Texas. In *The Prehistory of Texas*. Edited by Timothy K. Pertulla, pp. 101–126. Texas A&M University Press, College Station.
- Collins, Michael B., Dale B. Hudler, and Steve L. Black
 2003 *Pavo Real (41BX52): A Paleoindian and Archaic Camp and Work-Shop on the Balcones Escarpment, South-Central Texas*. Studies in Archeology No. 41, Austin: Texas Archeological Research Laboratory, University of Texas and Texas Department of Transportation, Environmental Affairs Division, Archeology Studies Program, Report 50, Austin, Texas.
- Comer, Clay and Diana J. Kleiner
 2009 *Handbook of Texas Online*, s.v. “Colorado River”, <http://www.tshaonline.org/handbook/online/articles/CC/rnc10.html> (accessed June 2, 2009).
- Cooke, Mary Jennifer
 2005 *Soil Formation and Erosion in Central Texas: Insights From Relict Soils and Cave Deposits*. Unpublished Ph.D. Dissertation, Department of Geology, University of Texas, Austin.
- Cooke, M. Jennifer, Libby A. Stern, Jay L. Banner, and Lawrence E. Mack
 2007 Evidence for the Silicate Source of Relict Soils on the Edwards Plateau, Central Texas. *Quaternary Research* 67:275–285.
- Cooke, M. Jennifer, Libby A. Stern, Jay L. Banner, Lawrence E. Mack, Thomas W. Stafford Jr., and Rickard S. Toomey III
 2003 Precise Timing and Rate of Massive Late Quaternary Soil Denudation. *Geology* 31(10):853–856.
- Cornish, Frank G. and Jon A. Baskin
 1995 Late Quaternary Sedimentation, Lower Nueces River, South Texas. *Texas Journal of Science* 47:191–202.
- Crawford, C. and C. F. Frederick
 2006 Geomorphology and Geoarchaeology. In *The McKinney Roughs Site 41BP627: A Stratified Late Archaic II Site on the Colorado River Terraces Bastrop County, Texas*, by Steve Carpenter, Michael Chavez, Kevin Miller, and Ken Lawrence, pp. 61–64. SWCA Cultural Resource Report No. 02-313, SWCA, Inc. Austin, Texas.
- Daniels, J. Michael and James C. Knox
 2005 Alluvial Stratigraphic Evidence for Channel Incision during the Mediaeval Warm Period on the Central Great Plains, USA. *The Holocene* 15(5):736–747.

- Darwin, Robert L., C. Reid Ferring, and Brooks B. Ellwood
1990 Geoelectric Stratigraphy and Subsurface Sediments at the Cooper Basin, NE Texas. *Geoarchaeology* 5(1):53–79.
- Decker, Susan, Stephen L. Black, and Thomas Gustavson
2000 *The Woodrow Heard Site, 41UV88 A Holocene Terrace Site in the Western Balcones Canyonlands of Southwestern Texas*. Studies in Archeology 33. Texas Archeological Research Laboratory, The University of Texas, Austin, Texas.
- Dillehay, Tom D.
1974 Late Quaternary Bison Population Changes on the Southern Plains. *Plains Anthropologist* 19(65):180–196.
- Doneker, Frances
2010 *Handbook of Texas Online*, s.v. “San Antonio River”, <http://www.tshaonline.org/handbook/online/articles/SS/rns6.html> (accessed April 17, 2010).
- Durbin, James M.
1999 *Geomorphic Response to Late Quaternary Climate and Sea-level Change, Lower Nueces River, Texas*. Unpublished Doctoral Dissertation. Department of Geology. University of Nebraska-Lincoln, Lincoln, Nebraska.
- Epps, Lawrence W.
1973 *The Geologic History of the Brazos River*. Baylor Geological Studies, Bulletin No. 24. Baylor University, Waco, Texas.
- Etheridge, Frank G., Lesli J. Wood, and S. A. Schumm
1998 Cyclic Variables Controlling Fluvial Sequence Development: Problems and Perspectives. In: *Relative Role of Eustasy, Climate, and Tectonism in Continental Rocks* by K. W. Shanley and P. J. McCabe (editors), pp. 17–29. SEPM (Society for Sedimentary Geology), Special Publication No. 59, Tulsa, OK.
- Everitt, B. S.
2002 *The Cambridge Dictionary of Statistics*. Second Edition. Cambridge University Press, Cambridge.
- Fairbanks, Richard G.
1989 A 17,000-year Glacio-eustatic Sea Level Record: Influence of Glacial Melting Rates on the Younger Dryas Event and Deep-Ocean Circulation. *Nature* 342:637–642.
- Fields, R.C. (editor)
1990 *Excavations at the Charles Cox, Lambs Creek Knoll, and Buffalo Branch Sites, Jewett Mine Project, Leon and Freestone Counties, Texas*. Reports of Investigations, Number 70. Prewitt & Associates, Inc. Austin, Texas.
- Fields, R.C., L. W. Klement, C. B. Bousman, S. A. Tomka, E. F. Gadus, and M. Howard
1991 Excavations at the Bottoms, Rena Branch, and Moccasin Springs Sites, Jewett Mine Project, Freestone and Leon Counties, Texas. Reports of Investigations, Number 82. Prewitt & Associates, Inc. Austin, Texas.

Fiore, Richard N.

- 1976 Geology and Geomorphology of the Clear Creek Drainage Basin, Western Burnet County, Texas. Unpublished Master's Thesis, University of Texas, Austin, Texas.

Ferring, Reid C.

- 1990b *Late Quaternary Geology and Geoarchaeology of the Upper Trinity River Drainage Basin, Texas*. Geological Society of America (GSA) Field Trip Guidebook #11, Dallas, Texas.
- 1991 Upper Trinity River Drainage Basin. In *Quaternary Nonglacial Geology: Conterminous U.S.*, by Roger B. Morrison editor, pp. 526–5531. The Geology of North America Volume K-2, The Geological Society of America, Inc. publishers. Boulder, Colorado.
- 1992 Alluvial Pedology and Geoarchaeological Research. In *Soils in Archaeology: Landscape Evolution and Human Occupation*, by Vance T. Holliday editor, pp. 1–39. Smithsonian Institution Press, Washington, D.C.
- 1994 *Late Quaternary Geology of the Upper Trinity River Basin*. Unpublished Doctoral Dissertation, University of Texas at Dallas, Dallas, Texas.
- 1995a The Late Quaternary Geology and Archaeology of the Aubrey Clovis Site, Texas: A Preliminary Report. In: *Ancient Peoples and Landscapes*. (E. Johnson editor) pp. 273–281. Museum of Texas Tech University, Lubbock, Texas.
- 1995b Middle Holocene Environments, Geology, and Archaeology in the Southern Plains. In: *Archaeological Geology of the Archaic Period in North America*. (E. A. Bettis III editor) 2:21–35. The Geological Society of America, Inc. publishers. Boulder, Colorado. Special Paper 297.
- 2000 Geoarchaeology in the Southern Osage Plains. In *Geoarchaeology in the Great Plains*, by Rolfe D. Mandel editor, pp. 44–78. University of Oklahoma Press, Norman.
- 2001 *The Archaeology and Paleoecology of the Aubrey Clovis Site (41DN479) Denton County, Texas*. Center for Environmental Archaeology, University of North Texas. Denton, Texas.

Ferring, C. Reid and Bonnie C. Yates

- 1997 *Holocene Geoarchaeology and Prehistory of the Ray Roberts Lake Area, North Central Texas*. Institute of Applied Sciences, North Texas State University. Denton, Texas.

Frederick, Charles D.

- 1987 *An Investigation into the Paleoenvironmental History of the Austin Mastodon Site*. Unpublished Master's Thesis, Department of Geology, University of Texas, Austin.
- 1995 *Fluvial Response to Late Quaternary Climate Change and Land Use in Central Mexico*. Unpublished Doctoral Dissertation. The University of Texas, Austin.

- 1996 Geomorphic Investigations. In: *Early Archaic Use of the Concho River Terraces: Cultural Resource Investigation 41TG307 and 41TG309, Tom Green County, San Angelo, Texas*. J. M. Quigg, J. Peck, C. Lintz, A. C. Treece, C. D. Frederick, R. Clem, G. L. Ellis, P. Schuchert, and J. T. Abbott, 5:85–108. Technical Report No. 11058. TRC Mariah Associates Inc. Austin, Texas.
- 2001 Geoarchaeological Investigations. In *Test Excavations at the Culebra Creek Site, 41BX126, Bexar County, Texas*. Nickels, David L., C. Britt Bousman, Jeff D. Leach, and Diane A. Cargill. Appendix A:254–265. Archaeological Report, No. 265, Center for Archaeological Research, University of Texas at San Antonio and Texas Department of Transportation, Environmental Affairs Division, Archeological Studies Program, Report 3. Austin, Texas.
- 2008 Paleoenvironment and Site Formation Processes. In *The Gatlin Site (41KR621): Investigating Archaic Lifeways on the Southern Edwards Plateau of Central Texas*. Brett A. Houk, Kevin A. Miller, and Eric R. Oksanen, 6:1–20. SWCA Cultural Resource Report No. 08-149, SWCA, Inc. Austin, Texas.
- Frederick, Charles D. and Thomas W. Boutton
- 1996 Appendix S Stable Isotopic Evidence of Holocene Paleoenvironmental Change in West-Central Texas. In: *Early Archaic Use of the Concho River Terraces: Cultural Resource Investigation 41TG307 and 41TG309, Tom Green County, San Angelo, Texas*. J. M. Quigg, J. Peck, C. Lintz, A. C. Treece, C. D. Frederick, R. Clem, G. L. Ellis, P. Schuchert, and J. T. Abbott, pp. S1–S18. Technical Report No. 11058. TRC Mariah Associates Inc. Austin, Texas.
- Fowler, A. J., R. Gillespie, and R. E. M. Hedges
- 1986 Radiocarbon Dating of Sediments. *Radiocarbon* 28 (2A):441–450.
- Gadus, E. Frances, Ross C. Fields, and Karl W. Kibler
- 2006 *Data Recovery Excavations at the J. B. White Site (41MM341), Milam County, Texas*. Research Report No. 145. Prewitt & Associates, Inc. Austin, Texas.
- Gard, Wayne
- 2010 *Handbook of Texas Online*, s.v. “Trinity River”, <http://www.tshaonline.org/handbook/online/articles/TT/rnr2.html> (accessed April 17, 2010).
- Garvin, Matthew G.
- 2008 *Late Quaternary Geochronologic, Stratigraphic, and Sedimentologic Framework of the Trinity River Incised Valley: East Texas Coast*. Unpublished Master’s Thesis, Department of Geology and Geophysics, Louisiana State University, Baton Rouge, Louisiana.
- Gibson, B. D.
- 1997 *The Geoarchaeology of Buttermilk Creek, Bell County, Texas*. Unpublished Master’s Thesis. Department of Anthropology. Texas A & M University, College Station, Texas.

- Goldberg, Paul and Vance T. Holliday
 1998 Chapter 6: Geology and Stratigraphy. In: *Wilson-Leonard: An 11,000-year Archeological Record of Hunter-Gatherers in Central Texas. Volume I: Introduction, Background, and Syntheses*. M. Collins (editor), pp. 77–119. Studies in Archeology 31, Texas Archeological Research Laboratory. The University of Texas, Austin, Texas.
- González, Juan L.
 2008 *Exploring the Late Holocene Sedimentary Record of the Mississippi Delta for Climate Sea-Level Links*. Unpublished Doctoral Dissertation. Tulane University, Louisiana.
- González, Juan L. and Torbjörn E. Törnqvist
 2009 A New Late Holocene Sea-Level Record from the Mississippi Delta: Evidence for a Climate/Sea Level Connection? *Quaternary Science Reviews* 28:1737–1749.
- Grimm, E. C., L. J. Maher, Jr., and D. M. Nelson
 2009 The Magnitude of Error in Conventional Bulk-Sediment Radiocarbon Dates From Central North America. *Quaternary Research* 72:301–308.
- Gustavson, Thomas C.
 1978 Bed Forms and Stratification Types of Modern Gravel Meander Lobes, Nueces River, Texas. *Sedimentology* 25:401–426.
 2000 Formation of the Holocene Terrace. In: *The Woodrow Heard Site, 41UV88 A Holocene Terrace Site in the Western Balcones Canyonlands of Southwestern Texas*. by Susan Decker, Stephen L. Black, and Thomas Gustavson editors, pp. 111–119. Studies in Archeology 33. Texas Archeological Research Laboratory, The University of Texas, Austin, Texas.
- Haas, Herbert, Vance Holliday, and Robert Stuckenrath
 1986 Dating of the Holocene Stratigraphy with Soluble and Insoluble Organic Fractions at the Lubbock Lake Archaeological Site, Texas. *Radiocarbon* 28(2A):473–485.
- Hall, Grant D.
 2010 *Texas Beyond History*, s.v. “Choke Canyon”, <http://www.texasbeyondhistory.net/choke/index.html> (accessed May 25, 2010).
- Hall, Grant D., Thomas R. Hester, and Stephen L. Black
 1986 *The Prehistoric Sites at Choke Canyon Reservoir, Southern Texas: Results of Phase II Archaeological Investigations*. Choke Canyon Series Volume 10, Center for Archaeological Research, University of Texas at San Antonio.
- Hall, Stephen A.
 1990a Channel Trenching and Climatic Change in the Southern U.S. Great Plains. *Geology* 18:342–345.

- 1990b Holocene Landscapes of the San Juan Basin, New Mexico; Geomorphic, Climatic, and Cultural Dynamics. In *Archaeological Geology of North America.*, by N. P. Lasca and J. Donahue editors, pp. 323–334. Geological Society of America, Centennial Special Volume 4. Boulder, Colorado.
- Heaton, T. J., P. G. Blackwell, and C. E. Buck
 2009 A Bayesian Approach to the Estimating of Radiocarbon Calibration Curves: The IntCal09 Methodology. *Radiocarbon* 51(4):1151–1164.
- Hedman, M.
 2007 *The Age of Everything*. University of Chicago Press, Chicago and London.
- Hendrickson Jr., Kenneth E.
 2010 *Handbook of Texas Online*, s.v. “Brazos River”, <http://www.tshaonline.org/handbook/online/articles/BB/rnb7.html> (accessed April 17, 2010).
- Hill, Robert T.
 1901 *Geography and Geology of the Black and Grand Prairies, Texas*. U. S. Geological Survey, Twenty-first Annual Report, Part VII-Texas, Washington, D. C.
- Hilliard, K. Lee
 1997 *Late-Quaternary Geology of the Waco Mammoth Site, Waco, Texas*. Unpublished Master’s Thesis Department of Geology, Baylor University, Waco, Texas.
 2000 *GIS-Based Geoarchaeological Analysis of the Owl Creek Basin, Fort Hood Military Reservation, Texas*. Unpublished Doctoral Dissertation. Department of Geology, Baylor University, Waco, Texas.
- Highley, Cheryl Lynn
 1986 *Archaeological Investigations at 41LK201, Choke Canyon Reservoir, Southern Texas*. Choke Canyon Series Volume 11, Center for Archaeological Research, University of Texas at San Antonio.
- Hodell, David A., Mark Brenner, Jason H. Curtis, and Thomas Guilderson
 2001 Solar Forcing of Drought Frequency in the Maya Lowlands. *Science* 292(5520):1367–1370.
- Holliday, Vance T.
 1985 Archaeological Geology of the Lubbock Lake Site, Southern High Plains of Texas. *Geological Society of America Bulletin* 96:1483–1492.
 1988 Genesis of a Late Holocene Soil Chronosequence at the Lubbock Lake Archaeological Site, Texas. *Annals of the Association of American Geographers* 78 (4):594–610.
 1990 Pedology in Archaeology. In *Archaeological Geology of North America.*, by N. P. Lasca and J. Donahue editors, pp. 525–540. Geological Society of America, Centennial Special Volume 4. Boulder, Colorado.

- 1995a Geoaerchology. In: *Archeological Investigations at the Loma Sandia Site (41LK28): A Prehistoric Cemetery and Campsite in Live Oak County, Texas. Volume 1.* by A. J. Taylor and C. L. Highley (editors). Chapter 3. pp. 19–30 Studies in Archeology 20. Texas Archeological Research Laboratory, The University of Texas, Austin, Texas.
- 1995b Stratigraphy and Paleoenvironments of Late Quaternary Valley Fills on the Southern High Plains. *Geological Society of America Memoir 186.*
- 1997 *Paleoindian Geoarchaeology of the Southern High Plains.* University of Texas Press, Austin.
- 2000 Vance Haynes and Paleoindian Geoarchaeology and Geochronology of the Great Plains. *Geoarchaeology* 15(6):511–522.
- 2009 Geoarchaeology and the Search for the First Americans. *Catena* 78:310–322.
- Holliday, Vance T. and Eileen Johnson
- 1981 An Update on the Plainview Occupation at the Lubbock Lake Site. *Plains Anthropologist* 26(1):251–253.
- 1983 Comments on Stafford’s “Alluvial Geology and Archaeological Potential of the Texas Southern High Plains”. *American Antiquity* 48(1):151–154.
- 1986 Re-Evaluation of the First Radiocarbon Age For the Folsom Culture. *American Antiquity* 51(2):332–338.
- Holliday, Vance T., Eileen Johnson, Herbert Haas, and Robert Stuckenrath
- 1983 Radiocarbon ages from the Lubbock Lake site, 1950–1980: Frame work for cultural and ecological change on the Southern High Plains. *Plains Anthropologist* 28:165–182.
- 1985 Radiocarbon ages from the Lubbock Lake site, 1981–1984.. *Plains Anthropologist* 30:277–291.
- Holliday, Vance T., Eileen Johnson, and Thomas W. Stafford, Jr.
- 1999 AMS Radiocarbon Dating of the Type Plainview and Firstview (Paleoindian) Assemblages: the Agony and the Ecstasy. *American Antiquity* 64(3):444–454.
- Houk, Brett A., Kevin A. Miller, and Eric A. Oksanen
- 2008 The Gatlin Site (41KR621): Investigating Archaic Lifeways on the Southern Edwards Plateau of Central Texas. SWCA Cultural Resource Report No. 08-149, SWCA, Inc. Austin, Texas.
- Houk, Brett A. and David L. Nickels
- 1997 *Phase II Archaeological Investigations at Lackland Air Force Base, San Antonio, Texas.* Archaeological Report, No. 264, Center for Archaeological Research, University of Texas at San Antonio.
- Howard, A. J., B. R. Gearey, T. Hill, W. Fletcher, and P. Marshall
- 2009 Fluvial Sediments, Correlations and Paleoenvironmental Reconstruction: The Development of Robust Radiocarbon Chronologies. *Journal of Archaeological Science* 36:2680–2688.

- Hu, F. S., D. Slawinski, H. E. Wright Jr., E. Ito, R. G. Johnson, K. R. Kelts, R. F. McEwan, and A. Boedigheimer
 1999 Abrupt Changes in North American Climate During Early Holocene Times. *Nature* 400:437–440.
- Hua, Quan
 2009 Radiocarbon: A Chronological Tool for the Recent Past. *Quaternary Geochronology* 4:378–390.
- Huckleberry, Gary and Andrew I. Duff
 2008 Alluvial Cycles, Climate, and Puebloan Settlement Shifts near Zuni Salt Lake, New Mexico, USA. *Geoarchaeology* 23(1):107–130
- Hudler, D. B.
 2000 *Modeling Paleolandscapes in Central Texas*. Unpublished Doctoral Dissertation. Department of Anthropology. University of Texas, Austin, Texas.
- Humphrey, John D. and C. Reid Ferring
 1994 Stable Isotopic Evidence for Latest Pleistocene and Holocene Climatic Change in North-Central Texas. *Quaternary Research* 41:200–213.
- Huntley, D. J.
 1985 *An Introduction to Dating, Analysis, and Location in Archaeology*. Simon Fraser University, Vancouver, B.C.
- Husain, Syed Raziuddin
 1998 *Stratigraphy of the Quaternary Terraces of the Brazos River in the Texas A&M University-Riverside Campus Area, Brazos County, Texas*. Unpublished Doctoral Dissertation. Texas A & M University, College Station, Texas.
- Ilan, Sharon
 2001 ‘Transition Dating’—A Heuristic Mathematical Approach to the Collation of Radiocarbon Dates From Stratified Sequences. *Radiocarbon* 43(2A):345–354.
- Jacobson, Robert B., James E. O’Connor, and Takashi Oguchi
 2003 Surficial Geologic Tools in Fluvial Geomorphology. In *Tools in Fluvial Geomorphology*, by Kondolf, G. Mathias and Hervé Piégay (editors), 2:25–57. John Wiley & Sons Ltd, Chichester, England.
- Johnson, E. L.
 1933 Stratigraphy of Oakville, Lagarto, and Reynosa Formations Between Nueces and Guadalupe Rivers, Texas. *Bulletin of the American Association of Petroleum Geologists*. 17(5):523–525.
- Johnson, Eileen and Holliday, Vance T.
 1980 A Plainview Kill/ Butchering Locale on the Llano Estacado—Lubbock Lake Site *Plains Anthropologist* 88(1):89–111.
- Johnson, L., Jr.,
 1995 *Past Cultures and Climates at the Jonas Terrace Site, 41ME29 of Medina County, Texas*. Office of the State Archeologist, Report No. 40: Texas Department of Transportation, Austin, Texas.

- Johnson, L., Jr., and G. T. Goode
 1994 A New Try at Dating and Characterizing Holocene Climates, as Well as Archeological Periods, on the Eastern Edwards Plateau. *Bulletin of the Texas Archeological Society* 65:1–51.
- Johnson, William C. and Brad Logan
 1990 Geoarchaeology of the Kansas River basin, central Great Plains. In *Archaeological Geology of North America.*, by N. P. Lasca and J. Donahue editors, pp. 267–299. Geological Society of America, Centennial Special Volume 4. Boulder, Colorado.
- Johnstone, E., M. G. Macklin, and J. Lewin
 2006 The Development and Application of a Database of Radiocarbon-dated Holocene Fluvial Deposits in Great Britain. *Catena* 66:14–23.
- Kemp, Leonard E.
 2008 *Buffaloed By Bison: Modeling Bison Abundance In Prehistoric Central Texas.* Unpublished Master's Thesis. University of Texas at San Antonio.
- Klein, Jeffery, J. C. Lerman, P. E. Damon, and E. K. Ralph
 1982 Calibration of Radiocarbon Dates: Tables Based on the Consensus Data of the Workshop on Calibrating the Radiocarbon Time Scale. *Radiocarbon* 24 (2):103–150.
- Knighton, David
 1998 *Fluvial Forms and Processes: A New Perspective.* Oxford University Press, Inc., New York.
- Knox, James C.
 2000 Sensitivity of Modern and Holocene Floods to Climate Change. *Quaternary Science Reviews* 19:439–457.
- Largent, Jr. F. B.
 1991 *The Prehistory of the Little Pin Oak Creek Site (41FY53), Fayette County, Texas.* Vol. II, Unpublished Master's Thesis Department of Geography, Texas A & M University, College Station, Texas.
- Leeder, M. R. and M. D. Stewart
 1996 Fluvial Incision and Sequence Stratigraphy: Alluvial Responses to Relative Sea-Level Fall and Their Detection in the Geological Record. In *Sequence Stratigraphy in British Geology.* (S. P. Hesselbo and D. N. Parkinson editors) pp. 25–39. Geological Society Special Publication No. 103, Oxford and Boston.
- Lewin, J., M. G. Macklin, and E. Johnstone
 2005 Interpreting Alluvial Archives: Sedimentological Factors in the British Holocene Fluvial Record. *Quaternary Science Reviews* 24:1873–1889.
- Libby, W. F.
 1955 *Radiocarbon Dating.* University of Chicago Press, Chicago and London.

- Lintz, Christopher, W. Nicholas Trierweiler, Fred M. Oglesby, Michael Blum, Patrick L. O'Neill, John Kuhl, Richard Holloway, Linda Scott-Cummings, and Dan Scurlock
 1993 *Cultural Resource Investigations in the O. H. Ivie Reservoir, Concho, Coleman, and Runnels Counties, Texas. Volume I: Project Introduction, Setting and Methods*. Technical Report No. 346-I. Mariah Associates Inc. Austin, Texas.
- Lintz, Christopher, W. Nicholas Trierweiler, Fred M. Oglesby, Patrick L. O'Neill, William Doering, and Michael McFaul
 1991 *A Cultural Resource Survey at Mitchell County Reservoir, Texas*. Technical Report No. 433. Mariah Associates Inc. Austin, Texas.
- Lohse, Jon C.
 2010 Personal Communication (September 14, 2010), Director Center for Archaeological Studies, Texas State University, San Marcos, Texas.
- Looney, Robert M.
 1977 *Late Quaternary Geomorphic Evolution of the Colorado River, Bastrop and Fayette Counties, Texas*. Unpublished Master's Thesis The University of Texas, Austin, Texas.
- Lukowski, Paul D.
 1987 *Archaeological Investigations Along the Leona River Watershed, Uvalde County, Texas*. Archaeological Survey Report No. 132, Center for Archaeological Research, University of Texas at San Antonio.
- Macklin, M. G., G. Benito, K. J. Gregory, E. Johnstone, J. Lewin, D. J. Michczyńska, R. Soja, L. Starkel, V. R. Thorndycraft
 2006 Past Hydrological Events Reflected in the Holocene Fluvial Record of Europe. *Catena* 66:145–154.
- Macklin, M. G., I. C. Fuller, J. Lewin, G. S. Maas, D. G. Passmore, J. Rose, J. C. Woodward, S. Black, R. H. B. Hamlin, and J. S. Rowan
 2002 Correlation of Fluvial Sequence in the Mediterranean Basin Over the Last 200 ka and their Relationship to Climate Change. *Quaternary Science Reviews* 21:1633–1641.
- Macklin, Mark G., Anna F. Jones, and John Lewin
 2010 River Response to Rapid Holocene Environmental Change: Evidence and Explanation in British Catchments. *Quaternary Science Reviews* 29(13-14):1555–1576.
- Macklin, M. G., E. Johnstone, and J. Lewin
 2005 Pervasive and Long-term Forcing of Holocene River Instability and Flooding in Great Britain by Centennial-scale Climate Change. *The Holocene* 15(7):937–943.
- Macklin, M. G. and J. Lewin
 2008 Alluvial Responses to the Changing Earth System. *Earth Surface Processes and Landforms* 33:1374–1395.

Mahoney, R. and S. Tomka

- 2001 *National Register Eligibility Testing of 41MM340 and 41MM341, along Little River, Milam County, Texas*. Archaeological Survey Report, No. 343, Center for Archaeological Research, University of Texas at San Antonio and Texas Department of Transportation, Environmental Affairs Division, Archeology Studies Program, Report 30. San Antonio.

Mahoney, Richard B., Steve A. Tomka, Raymond P. Mauldin, Harry J. Shafer, Lee C. Nordt, Russell D. Greaves, and Rebecca R. Galdeano

- 2003 *Data Recovery Excavations at 41MM340: A Late Archaic Site along Little River in Milam County, Texas*. Archaeological Survey Report, No. 340, Center for Archaeological Research, University of Texas at San Antonio and Texas Department of Transportation, Environmental Affairs Division, Archeology Studies Program, Report 54. San Antonio.

Mallouf, Robert J.

- 1981 Observations Concerning Environmental and Cultural Interactions During the Terminal Pleistocene and Early Holocene in the Big Bend of Texas and Adjoining Regions. *Bulletin of the Texas Archeological Society* 52:121–146.

Mandel, Rolfe

- 1980 *Climatic and Vegetative Change Inferred From Alluvial Paleosols in Central and South-Central Texas*. Unpublished Master's Thesis, Department of Geology, University of Texas, Austin.
- 1992 Geomorphology. In: *An Archaeological Survey of the Proposed South Bend Reservoir Area: Young, Stephens, and Throckmorton Counties, Texas*. Joe W. Saunders, C. S. Mueller-White, and David L. Carlson editors). Texas A & M Archaeological Survey Reports No. 6. College Station, Texas.

Mandel, Rolfe D., John S. Jacob, and Lee C. Nordt

- 2007 Geomorphic Investigations. In *Archaeological and Paleoecological Investigations at the Richard Beene Site South-Central Texas Volume I: Paleoecological Studies, Cultural Contexts, and Excavation Studies*. Alston V. Thoms and Rolfe D. Mandel (editors). 3:27–60. 2 vols. Reports of Investigations 8. Center for Ecological Archaeology. Texas A&M University, College Station.

Martin, C. W. and W. C. Johnson

- 1995 Variation in Radiocarbon Ages of Soil Organic Matter Fractions from Late Quaternary Buried Soils. *Quaternary Research* 43:232–237.

Matthews, J. A.

- 1985 Radiocarbon Dating of Surface and Buried Soils: Principles, Problems, and Prospects. In: *Geomorphology and Soils*. (K. S. Richards, R. R. Arnett, and S. Ellis editors) 14:269–288. George Allen & Unwin Ltd, London, Boston, and Sydney.

- Mauldin, Raymond P. and David L. Nickels
 2001 *An Archaeological Survey of Twin Buttes Reservoir, Tom Green County, Texas*. Volumes I–III. Archaeological Survey Report No. 300. Center for Archaeological Research. San Antonio, Texas.
- Mayewski, Paul A., Eelco E. Rohling, J. Curt Stager, Wibjörn Karlén, Kirk A. Maasch, L. David Meeker, Eric A. Meyerson, Françoise Gasse, Shirley van Kreveld, Karin Holmgren, Julia Lee-Thorp, Gunhild Rosqvist, Frank Rack, Michael Staubwasser, Ralph R. Schneider, and Eric J. Steig
 2004 Holocene Climate Variability. *Quaternary Research* 62:243–255.
- McCormac, F. G. and M. G. Baillie
 1993 Radiocarbon to Calendar Date Conversion: Calendrical Band Widths as a Function of Radiocarbon Precision. *Radiocarbon* 35(2):311–316.
- McEwen, Michael C.
 1963 *Sedimentary Facies of the Trinity River Delta, Texas*. Unpublished Doctoral Dissertation, Rice University, Houston, Texas.
- Mear, Charles E.
 1953 *Quaternary Geology of the Upper Sabinal River Valley, Uvalde and Bandera Counties, Texas*. Unpublished Master's Thesis. The University of Texas, Austin, Texas.
 1990 *Quaternary [sic] Geology of the Upper Sabinal River Valley, Uvalde and Bandera Counties, Texas*. Southern Texas Archaeological Association Special Publication Number Six. The University of Texas, Austin, Texas.
 1995 Quaternary Geology of the Upper Sabinal River Valley, Uvalde and Bandera Counties, Texas. *Geoarchaeology* 10(6):457–480.
 1998 Terrace Deposits and Late Quaternary Climate, South-Central Edwards Plateau, Texas. *Bulletin of the Texas Archeological Society* 69:79–88.
- McKinney, Wilson W.
 1981 Early Holocene Adaptations in Central and Southern Texas: The Problem of the Paleoindian-Archaic Transition. *Bulletin of the Texas Archeological Society* 52:91–120.
- Mehalchick, G., K. Kleinbach, D. K. Boyd, and K. W. Kibler
 2000 Geoarchaeological Investigations and National Register Testing of 52 Prehistoric Archeological Sites on Fort Hood, Texas: The 1997 Season. United States Army Fort Hood, Archeological Resource Management Series, Research Report No. 39. Prewitt & Associates, Inc. Austin, Texas.
- Michael, Henry C. and Jeffery Klein
 1979 An International Calibration for Radiocarbon Dates. *MASCA Journal* 1(2):56–57.
- Mock, Cary J. and Patrick J. Bartlein
 1995 Spatial Variability of Late-Quaternary Paleoclimates in the Western United States. *Quaternary Research* 44:425–433.

- Mook, W. G., and H. T. Waterbolk
 1985 *Handbooks for Archaeologists, No. 3 Radiocarbon Dating*. European Science Foundation, Strasbourg.
- Musgrove, MaryLynn
 2000 *Temporal Links Between Climate and Hydrology: Insights from Central Texas Cave Deposits and Groundwater*. Unpublished Doctoral Dissertation. The University of Texas, Austin.
- Musgrove, Marylynn, Jay L. Banner, Larry E. Mack, Deanna M. Combs, Eric W. James, Hai Cheng, and R. Lawrence Edwards
 2001 Geochronology of Late Pleistocene to Holocene Speleothems from Central Texas: Implications for Regional Paleoclimate. *Geological Society of America Bulletin* 113(12):1532–1543.
- Nash, Stephen E.
 1999 *Time, Trees, and Prehistory: Tree-Ring Dating and the Development of North American Archaeology 1914–1950*. The University of Utah Press, Salt Lake City, UT.
- Neck, Raymond W.
 1983 Paleoenvironmental Significance of a Nonmarine Pleistocene Molluscan Fauna From Southern Texas. *Texas Journal of Science* Vol. XXXV 2:147–154.
 1987 Changing Holocene Snail Faunas and Environments along the Eastern Caprock Escarpment of Texas. *Quaternary Research* 27:312–322.
- Nickels, David L. and C. Britt Bousman
 2010 *Archaeological Testing at San Marcos Springs (41HY160) for the Texas Rivers Center, Texas, Hays County, Texas*. Archaeological Studies Report No. 13, Center for Archaeological Studies, Texas State University, San Marcos.
- Nickels, David L., C. Britt Bousman, Jeff D. Leach, and Diane A. Cargill
 2001 *Test Excavations at the Culebra Creek Site, 41BX126, Bexar County, Texas*. Archaeological Report, No. 265, Center for Archaeological Research, University of Texas at San Antonio and Texas Department of Transportation, Environmental Affairs Division, Archeological Studies Program, Report 3. Austin, Texas.
- Nordt, Lee
 1983 *Soils-Geomorphology of the Brazos River Terraces Brazos County, Texas*. Unpublished Master's Thesis. Texas A & M University, College Station, Texas.
 1986 Regional Correlation of Terraces and Soils of the Brazos, Trinity, and Navasota Rivers of the Coastal Plain of Texas. *Soil Survey Horizons* 27(1):29–35.
 1992 *Archaeological Geology of the Ft. Hood Military Reservation, Fort Hood, Texas*. United States Army Fort Hood, Archeological Resource Management Series, Research Report No. 25. Prewitt & Associates, Inc. Austin, Texas.

- 1993 *Additional Geoarchaeological Investigations at the Ft. Hood Military Reservation, Fort Hood, Texas*. United States Army Fort Hood, Archeological Resource Management Series, Research Report No. 28. Prewitt & Associates, Inc. Austin, Texas.
- 1995 Geoarchaeology of Henson Creek: A Low-Order tributary in Central Texas. *Geoarchaeology* 10:205–221.
- 1996a Geoarchaeology of 41BX47, Upper Leon Creek Basin. In *Archaic Land Use of Upper Leon Creek Terraces: Archaeological Testing in Northern, Bexar County, Texas*. Archaeological Report, No. 234, Cynthia L. Tennis (editor), 3:11–19. Center for Archaeological Research, University of Texas at San Antonio.
- 1996b *Development of a Soil Chronosequence in Late Quaternary Alluvium in Central Texas: A Stable C Isotope and Mass Balance Approach*. Unpublished Doctoral Dissertation. Texas A & M University, College Station, Texas.
- 2001a Geoarchaeology of 41BX126. In *Test Excavations at the Culebra Creek Site, 41BX126, Bexar County, Texas*. Nickels, David L., C. Britt Bousman, Jeff D. Leach, and Diane A. Cargill. 5:36–44. Archaeological Report, No. 265, Center for Archaeological Research, University of Texas at San Antonio and Texas Department of Transportation, Environmental Affairs Division, Archeological Studies Program, Report 3. Austin, Texas.
- 2001b Geomorphology. In: *National Register Eligibility Testing of 41MM340 and 41MM341 along Little River Milam County, Texas*. R. Mahoney and S. Tomka (editors). Archaeological Survey Report No. 343. Center for Archaeological Research. San Antonio, Texas.
- 2003 Geomorphology. In: *Data Recovery Excavations at 41MM340: A Late Archaic Site Along Little River in Milam County, Texas*. R. Mahoney, S. Tomka, H. Shafer, L. Nordt, R. Greaves, and R. Galdeano (editors). Archaeological Survey Report No. 340. Center for Archaeological Research. San Antonio, Texas.
- 2004 Late Quaternary Alluvial Stratigraphy of a Low Order Tributary in Central Texas and its Response to Climate and Sediment Supply. *Quaternary Research* 62:289–300.
- Nordt, Lee C. and C. Britt Bousman
- 2001 Geoarchaeology of the Twin Buttes Project Area. In: *An Archaeological Survey of Twin Buttes Reservoir, Tom Green County, Texas*. Raymond P. Mauldin and David L. Nickels (editors) 4:39–54. Archaeological Survey Report No. 300. Center for Archaeological Research. San Antonio, Texas.
- Nordt, Lee C., Thomas W. Boutton, John S. Jacob, and Rolfe D. Mandel
- 2002 C₄ Plant Productivity and Climate-CO₂ Variations in South-Central Texas during the Late Quaternary. *Quaternary Research* 58:182–188.

- Nordt, Lee, Thomas W. Boutton, Charles T. Hallmark, and Michael R. Waters
1994 Late Quaternary Vegetation and Climate Changes in Central Texas Based on the Isotopic Composition of Organic Carbon. *Quaternary Research* 41:109–120.
- Nordt, Lee, Thomas W. Boutton, Charles T. Hallmark, and Michael R. Waters
1998 Quantifying Pedogenic Carbonate Accumulations Using Stable Carbon Isotopes. *Geoderma* 82:115–136.
- Nordt, Lee, Joseph von Fischer, and Larry Tieszen
2007 Late Quaternary Temperature Record from Buried Soils of the North American Great Plains. *Geology* 35 (2):159–162.
- North American Commission on Stratigraphic Nomenclature (NACSC)
2005 North American Stratigraphic Code. *American Association of Petroleum Geologists Bulletin*. 89(11):1547–1591.
- Oksanen, Eric O.
2008 *Archaeological Investigations at the Icehouse Site, 41HY161: A Reevaluation of Early Archaic Technology, Subsistence, and Settlement along the Balcones Escarpment and Central Texas*. Unpublished Master's Thesis. Texas State University, San Marcos.
- Osburn, Tiffany and David D. Kuehn
2006 *Evaluation of Site 41BX1623 For the SAWS Blue Wing Recycle Pipeline, Bexar County, Texas*. Miscellaneous Report of Investigations Number 366. Geo-Marine Inc. Plano, Texas.
- Otvos, Ervin G.
2004 Holocene Gulf Levels: Recognition Issues and an Updated Sea-Level Curve. *Journal of Coastal Research* 20(3):680–699.
- Paine, Jeffery G.
1991 *Late Quaternary Depositional Units, Sea Level, and Vertical Movement Along the Central Texas Coast*. Unpublished Ph.D. Dissertation, Department of Geology, University of Texas, Austin.
1993 Subsidence of the Texas Coast: Inference from Historical and late Pleistocene Sea Levels. *Tectonophysics* 222(3/4):445–453.
- Patton, Peter C.
1987 Geomorphology. In *Archeology at Aquilla Lake 1978–1982 Investigations Vol I*. by David O. Brown (editor), I:3-1–3-13. 3 vols. Texas Archeological Survey Research Report 81. University of Texas at Austin.
- Pearl, F. B.
1997 *Geoarchaeological Investigations of the Upper Lampasas River, Texas*. Unpublished Master's Thesis. Department of Anthropology. Texas A & M University, College Station, Texas.
- Phillips, Jonathan D. and Daniel A. Marion
2001 Residence Times of Alluvium in an East Texas Stream as Indicated by Stream Color. *Catena* 45:49–71.

- Prewitt, Elton R. and Jeffery Paine
 1987 The Swan Lake Site (41AS16) On Copano Bay, Aransas County, Texas: Settlement, Subsistence, and Sea Level. *Bulletin of the Texas Archeological Society* 58:147–174.
- Price, W. A.
 1933 The Role of Diastrophism in Topography of the Corpus Christi Area, South Texas. *Bulletin of the American Association of Petroleum Geologists*. 17(5):907–962.
- Prikryl, Daniel J.
 1990 *Lower Elm Fork Prehistory: A Redefinition of Cultural Concepts and Chronologies along the Trinity River, North-Central Texas*. Office of the State Archeologist Report Series, No. 37. Austin, Texas.
- Prikryl, Daniel J. and Bonnie C. Yates (editors)
 1987 *Test Excavations at 41CO141 Ray Roberts Reservoir, Cooke County, Texas*. Contributions in Archaeology No. 4, Institute of Applied Sciences, North Texas State University. Denton, Texas.
- Prochnow, Shane J.
 2001 *Paleohydrology and Geoarchaeology at the South End of the Horn Shelter Number 2 along the Brazos River, Central Texas*. Unpublished Master's Thesis Department of Geology, Baylor University, Waco, Texas.
- Quigg, J. Michael and Jay Peck
 1995 *The Rush Site (41TG346): A Stratified Late Prehistoric Locale in Tom Green County, Texas*. Technical Report No. 816C. TRC Mariah Associates Inc. Austin, Texas.
- Quigg, J. Michael, Jay Peck, Christopher Lintz, Abby C. Treece, Charles D. Frederick, Roman Clem, G. Lain Ellis, Paul Schubert, and James T. Abbott
 1996 *Early Archaic Use of the Concho River Terraces: Cultural Resource Investigation 41TG307 and 41TG309, Tom Green County, San Angelo, Texas*. Technical Report No. 11058. TRC Mariah Associates Inc. Austin, Texas.
- Rainey, Mary
 1974 *The Quaternary Stratigraphy of the North Sulphur River*. Unpublished Master's Thesis. Department of Geology, Southern Methodist University, Dallas, Texas.
- Ralph, E. K., H. N. Michael, and M. C. Han
 1973 Radiocarbon and Reality. *MASCA Newsletter* 9(1):1–4.
- Ramsey, C. B.
 1995 Radiocarbon Calibrations and the Analysis of Stratigraphy. *Radiocarbon* 37(2):425–430.
 1998 Probability and Dating. *Radiocarbon* 40(1):461–474.
 2001 Development of the Radiocarbon Calibration Program. *Radiocarbon* 43(2A):355–363.

- 2008 Deposition Models for Chronological Records. *Quaternary Science Reviews* 27:42–60.
- 2008 Radiocarbon Dating: Revolutions in Understanding. *Archaeometry* 50(2):249–275.
- 2009 Bayesian Analysis of Radiocarbon Dates. *Radiocarbon* 51(1):337–360.
- Reimer, P. J., M. G. L. Baillie, E. Bard, A. Bayliss, J. W. Beck, P. G. Blackwell, C. B. Ramsey, C. E. Buck, G. S. Burr, R. L. Edwards, M. Friedrich, P. M. Grootes, T. P. Guilderson, I. Hajdas, T. J. Heaton, A. G. Hogg, K. A. Hughen, K. F. Kaiser, B. Kromer, F. G. McCormac, S. W. Manning, R. W. Reimer, D. A. Richards, J. R. Southon, S. Talamo, J. van der Plicht, and C. E. Weyhenmeyer
- 2009 IntCal09 and Marine09 Radiocarbon Age Calibration Curves, 0-50,000 Years Cal BP. *Radiocarbon* 51(4):1111–1150.
- Rickliss, Robert A.
- 1988 Archeological Investigations at the McKinzie Site (41NU221), Nueces County, Texas. *Bulletin of the Texas Archeological Society* 58:–76.
- 1993 *A Model of Holocene Environmental and Human Adaptive Change on the Central Texas Coast: Geoarchaeological Investigations at White's Point, Nueces Bay, and Surrounding Area*. Coastal Archaeological Studies, Inc., Corpus Christi, Texas.
- 2001 *National Register Eligibility Testing at 41LE177, Alcoa Sandow Mine, Lee County, Texas: Archeological, Geoarchaeological, and Paleoenvironmental Assessment of an Upland Sandy Mantle Site*. Coastal Environments, Inc., Baton Rouge.
- 2004 Prehistoric Occupation of the Central and Lower Texas Coast. In *The Prehistory of Texas*. Edited by Timothy K. Pertulla, pp. 155–180. Texas A&M University Press, College Station.
- Rickliss, Robert A. and Michael D. Blum
- 1997 The Geoarchaeological Record of Holocene Sea Level Change and Human Occupation of the Texas Coast. *Geoarchaeology* 12(4):287–314.
- Rickliss, Robert A. and Michael B. Collins
- 1995 *Archaic and Late Prehistoric Human Ecology in the Middle Onion Creek Valley, Hays County, Texas*. 2 vols. Studies in Archeology No. 19. Austin: Texas Archeological Research Laboratory, University of Texas at Austin.
- Rickliss, Robert A. and Kim A. Cox
- 1998 Holocene Climatic and Environmental Change in the Texas Coastal Zone: Some Geoarchaeological and Ecofactual Indicators. *Plains Anthropologist* 43(164):125–136.
- Ringstaff, Christopher W.
- 2000 *A Study of Landform Evolution and Archaeological Preservation at Site 41HY165, San Marcos, Texas*. Unpublished Master's Thesis. Department of Geography, Southwest Texas State University, San Marcos.

- Rogers, R.
1984 *Kansas Prehistory: An Alluvial Geomorphological Perspective*. Unpublished Ph.D. Dissertation Department of Geography, University of Kansas, Lawrence.
- Sayles, E. B.
1935 *An Archaeological Survey of Texas*. Medallion Papers XVII, Globe, Arizona.
- Scott IV, Robert F. and Daniel E. Fox
1982 *Excavations at Sites 41LK32/32 and 41LK202 in the Choke Canyon Reservoir, South Texas*. Choke Canyon Series Volume 8, Center for Archaeological Research, University of Texas at San Antonio.
- Schaetzl, Randall and Sharon Anderson
2005 *Soils: Genesis and Geomorphology*. Cambridge University Press, Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, Sao Paulo.
- Schmieder, Jens
2009 *The Nebraska Sand Hills—Drought Variation and Landscape Stability in the Mid- to Late-Holocene Based on High-Resolution Lake Sediment Records*. Unpublished Doctoral Dissertation, University of Nebraska, Lincoln.
- Schroeder, Eric A. and Eric R. Oksanen
2002 *Data Recovery at the Armstrong Site (41CW54) Caldwell County, Texas. Volume I: Background, Methods, and site Context*. Paul Price Associates, Inc., Austin, Texas.
- Schumm, S. A.
1993 River Response to Baselevel Change: Implications for Sequence Stratigraphy. *Journal of Geology* 101:279–294.
2003 *The Fluvial System*. Blackburn Press, New Jersey.
- Shuler, Ellis H.
1935 Terraces of the Trinity River, Dallas County, Texas. *Field and Laboratory* 3:44–53.
- Shuman, Bryan N., Paige Newby, and Jeffery P. Donnelly
2009 Abrupt Climate Change as an Important Agent of Ecological Change in the Northeast U.S. Throughout the Past 15,000 Years. *Quaternary Science Reviews* 28:1693–1709.
- Simms, Alexander R.
2005 *Late Quaternary/Holocene Evolution of the Nueces Incised Valley, Central Texas*. Unpublished Doctoral Dissertation. Rice University, Houston, Texas.
- Simms, Alexander R., Kurt Lambeck, Anthony Purcell, John B. Anderson, and Antonio B. Rodriguez
2007 Sea-level History of the Gulf of Mexico Since the Last Glacial Maximum with Implications for the Melting History of the Laurentide Ice Sheet. *Quaternary Science Reviews* 26:920–940.

- Smith, A. M.
1992 Holocene Paleoclimatic Trends from Palaeoflood Analysis. *Palaeogeography, Palaeoclimatology, Palaeoecology* 97:235–240.
- Snow, Jennifer N.
1998 *Late Quaternary Highstand and Transgressive Deltas of the Ancestral Colorado River: Eustatic and Climatic Controls on Deposition*. Unpublished Master's Thesis. Rice University, Houston.
- Smyrl, Vivian Elizabeth
2010 *Handbook of Texas Online*, s.v. "Guadalupe River", <http://www.tshaonline.org/handbook/online/articles/GG/rng1.html> (accessed April 17, 2010).
- Spearing, Darwin
1991 *Roadside Geology Series: Roadside Geology of Texas*. Mountain Press Publishing Company. Missoula, MT.
- Stafford, Jr. T.
1981 Alluvial Geology and Archaeological Potential of the Texas Southern High Plains. *American Antiquity* 46(3):548–565.
1983 Geoarchaeology of the Texas Southern Plains: A Reply to Holliday and Johnson. *American Antiquity* 48(1):155–157.
1998 Chapter 25: Radiocarbon Chronostratigraphy. In: *Wilson-Leonard: An 11,000-year Archeological Record of Hunter-Gatherers in Central Texas. Volume IV: Archeological Features and Technical Analyses*. M. Collins (editor), pp. 1039–1066. Studies in Archeology 31, Texas Archeological Research Laboratory. The University of Texas, Austin, Texas.
- Steier, P., W. Rom, and S. Puchegger
2001 New Methods and Critical Aspects in Bayesian Mathematics for ¹⁴C Calibration. *Radiocarbon* 43(2A):1–15.
- Stine, Scott
1994 Extreme and Persistent Drought in California and Patagonia during Mediaeval Time. *Nature* 369:546–549.
- Stovall, J. Willis and William N. McAnulty
1950 The Vertebrate Fauna and Geologic Age of Trinity River Terraces in Henderson County, Texas. *The American Naturalist* 44(1):211–250.
- Story, D. A.
1990 Environmental Setting. In *The Archeology and Bioarcheology of the Gulf Coastal Plain: Volume I*. Dee Ann Story, Janice Guy, Barbara Burnett, Martha Doty Freeman, Jerome Rose, D. Gentry Steele, Ben Olive, and Karl Reinhard. 2:5–26. Arkansas Archeological Survey Research Series No. 38, Fayetteville, Arkansas.
- Stricklen, Jr. Fred L.
1961 *Degradational Stream Deposits of the Brazos River, Central Texas*. Geological Society of America Bulletin 72:19–36.

- Stuiver, Minze and Thomas F. Braziunas
1993 Modeling Atmospheric ^{14}C Influences and ^{14}C Ages of Marine Samples to 10,000 BC. *Radiocarbon* 35 (1):137–189.
- Stuiver, Minze and Henry A. Polach
1977 Discussion: Reporting of ^{14}C Data. *Radiocarbon* 19 (3):355–363.
- Stuiver, M. and H. S. Suess
1966 On the Relationship Between Radiocarbon Dates and True Sample Ages. *Radiocarbon* 8:534–540.
- Sylvia, Dennis A., and William E. Galloway
2006 Morphology and Stratigraphy of the Late Quaternary Lower Brazos Valley: Implications for Paleo-climate, Discharge, and Sediment Delivery. *Sedimentary Geology* 190:159–175.
- Taha, Z. Patrick and John B. Anderson
2008 The Influence of Valley Aggradation and Listric Normal Faulting on Styles of River Avulsion: A Case Study of the Brazos River, Texas, USA. *Geomorphology* 95:429–448.
- Talma, A. S. and J. C. Vogel
1993 A Simplified Approach to Calibrating ^{14}C Dates. *Radiocarbon* 35(2):317–322.
- Taylor, Anna Jean and Cheryl Lynn Highley
1995 *Archeological Investigations at the Loma Sandia Site (41LK28): A Prehistoric Cemetery and Campsite in Live Oak County, Texas. Volume 1.* Studies in Archeology 20. Texas Archeological Research Laboratory, The University of Texas, Austin, Texas.
- Taylor, Anna J.
1995 Radiocarbon Assays. In: *Archeological Investigations at the Loma Sandia Site (41LK28): A Prehistoric Cemetery and Campsite in Live Oak County, Texas. Volume 1.* by A. J. Taylor and C. L. Highley (editors). Appendix F. pp. 840–846. Studies in Archeology 20. Texas Archeological Research Laboratory, The University of Texas, Austin, Texas.
- Taylor, R. E.
2009 Six Decades of Radiocarbon Dating in New World Archaeology. *Radiocarbon* 51(1):173–212.
- 1997 Radiocarbon Dating. In: *Chronometric Dating in Archaeology: Advances in Archaeological and Museum Science Vol. 2.* (R. E. Taylor and M. J. Aitken editors) 3:65–96. Plenum Press, New York and London.
- 1987 *Radiocarbon Dating: An Archaeological Perspective.* Academic Press, Inc., Orlando and London.
- Taylor, R. E., C. V. Haynes Jr., and M. Stuiver
1996 Clovis and Folsom Age Estimates: Stratigraphic Context and Radiocarbon Calibration. *Antiquity* 70:515–525.

- Taylor, R. E., M. Stuiver, and P. J. Reimer
1996 Development and Extension of the Calibration of the Radiocarbon Time Scale: Archaeological Applications. *Quaternary Science Reviews* 15:655–668.
- Tennis, Cynthia L. and Robert J. Hard
1995 *Archaeological Survey of Upper Leon Creek Terraces, Bexar County, Texas*. Archaeological Report, No. 233, Center for Archaeological Research, University of Texas at San Antonio.
- Tennis, Cynthia L.
1996 *Archaic Land Use of Upper Leon Creek Terraces: Archaeological Testing in Northern Bexar County, Texas*. Archaeological Report, No. 234, Center for Archaeological Research, University of Texas at San Antonio.
- Texas Parks and Wildlife Department (TPWD)
2009 Natural Regions of Texas, http://www.tpwd.state.tx.us/landwater/land/maps/gis/map_downloads/ (accessed March 2009).
- Tharp, Tommy Lee
1988 *Aspects of the Leon River Drainage History with Implications to Other Central Texas Streams*. Unpublished Master's Thesis. Department of Geology, Baylor University, Waco, Texas.
- Thomas, Mark A. and John B. Anderson
1994 Sea-Level Controls on the Facies Architecture of the Trinity/Sabine Incised-Valley System, Texas Continental Shelf. In: *Incised-valley Systems: Origins and Sedimentary Sequences* by Robert W. Dalrymple, Ron Boyd, and Brian Zaitlin (editors), pp. 63–82. SEPM (Society for Sedimentary Geology), Special Publication No. 51, Tulsa, OK.
- Thoms, Alston V. and Ben W. Olive
1993 Archaeological Data and Late Prehistoric Period Population Dynamics. In: *Archaeological Survey at Fort Hood, Texas Fiscal Years 1991 and 1992: The Cantonment and Belton Lake Periphery Areas*. by Alston V. Thoms (editor), pp. 43–60. United States Army Fort Hood, Archeological Resource Management Series, Research Report No. 27.
- Tinkler, K. J.
1971 Active Valley Meanders in South-Central Texas and Their Wider Implications. *Geological Society of America Bulletin* 82:1783–1800.
- Toomey III, Rickard S.
1993 *Late Pleistocene and Holocene Faunal and Environmental Changes at Hall's Cave, Kerr County, Texas*. Vols. 1 and 2 Unpublished Ph.D. dissertation, Department of Geology, The University of Texas at Austin.
- Toomey III, Rickard S., Michael D. Blum, and Salvatore Valastro Jr.
1993 Late Quaternary Climates and Environments of the Edwards Plateau, Texas. *Global and Planetary Change* 7:299–320.

- Törnqvist, Torbjörn E., Juan L. González, Lee A. Newsom, Klaas van der Borg, Arie F. M. Jong, and Charles W. Kurnik
2004 Deciphering Holocene Sea-level History on the Gulf Coast: A High-Resolution Record from the Mississippi Delta. *Geological Society of America Bulletin* 116(7/8):1026–1039.
- Urbanec, Don A.
1963 *Stream Terraces and Related Deposits in the Austin Area, Texas*. Unpublished Master's Thesis The University of Texas, Austin, Texas.
- Urista, Juan Carlos
2009 *Stratigraphy and Geomorphology of the Vernor Mammoth Site, Clute, Brazoria County Texas*. Unpublished Master's Thesis. Texas A & M University, College Station, Texas.
- U. S. Geological Survey (USGS)
2010 Midway group, undivided. <http://tin.er.usgs.gov/geology/state/sgmc-unit.php?unit=TXPNmi%3B0> (accessed August 17, 2010).
- Upton, G. and I. Cook
2006 *A Dictionary of Statistics*. Second Edition. Oxford University Press, Oxford and New York.
- Van den berg, Jan H.
1995 Prediction of Alluvial Channel Pattern of Perennial Rivers. *Geomorphology* 12:259–279.
- Vandenberghe, J.
1995 Timescales, Climate and River Development. *Quaternary Science Reviews* 14:631–638.
- Van de Plassche, Orson, Klaas van der Borg, and Arie F. M. de Jong
1998 Sea Level–Climate Correlation During the Past 1400 yr. *Geology* 26 (4):319–322.
- Van de Plicht, J. and A. Hogg
2006 A Note on Reporting Radiocarbon. *Quaternary Geochronology* 1:237–240.
- Voellinger, Leonard R.
1990 *The Geomorphological Analysis of the Lower Brazos River Basin, Texas Applied to the Management of Its Cultural Resources*. Unpublished Master's Thesis Department of Geography, Southwest Texas State University, San Marcos.
- Walker, M. J. C.
2005 *Quaternary Dating Methods*. John Wiley & Sons, Ltd, West Sussex, England.

- Wallis, John C.
1976 *Evidence of Climatic Change From Fluvial Geomorphology and Alluvial Soils of the Colorado River, Texas*. Unpublished Master's Thesis The University of Texas, Austin, Texas.
- Wang, Yang, Ronald Amundson, and Susan Trumbore
1996 Radiocarbon Dating of Soil Organic Matter. *Quaternary Research* 45:282–288.
- Waters, Michael R. and Lee C. Nordt
1995 Late Quaternary Alluvial History of the Brazos River in East-Central Texas. *Quaternary Research* 43:311–319.
1996 Geomorphic Reconnaissance of Selected Segments of the MIDTEXAS Pipeline. In *Archaeological Survey of the Proposed 130–Mile MIDTEXAS Pipeline, Gonzales, Dewitt, Lavaca, Colorado, Austin, and Waller Counties, Texas*. Kevin A. Miller (editor), 5:39–68. SWCA Cultural Resource Report No. 95-154, SWCA, Inc. Austin, Texas.
- Weber, Gerald E.
1968 *Geology of the Fluvial Deposits of the Colorado River Valley, Central Texas*. Unpublished Master's Thesis The University of Texas, Austin, Texas.
- Weddle, Robert S.
2010 *Handbook of Texas Online*, s.v. “Nueces River”, <http://www.tshaonline.org/handbook/online/articles/NN/rnn15.html> (accessed May 18, 2010).
- Weeks, A. W.
1933 Lissie, Reynosa, and Upland Terrace Deposits of Coastal Plain of Texas Between Brazos River and Rio Grande. *Bulletin of the American Association of Petroleum Geologists*. 17(5):453–487.
1945 Quaternary Deposits of Texas Coastal Plain Between Brazos River and Rio Grande. *Bulletin of the American Association of Petroleum Geologists*. 29(12):1693–1720.
- Wilkinson, Bruce H. and Robert A. Basse
1978 Late Holocene History of the Central Texas Coast from Galveston Island to Pass Cavallo. *Geological Society of America Bulletin* 89:1592–1600.
- Williams, Marikka L.
2004 *Interpreting Prehistoric Patterns: Site-Catchment Analysis in the Upper Trinity River Basin of North Central Texas*. Unpublished Master's Thesis, University of North Texas, Denton, Texas.
- Winans, Melissa C.
2010 Personal Communication (September 15, 2010), Database Analyst, Texas Natural Science Center, University of Texas, Austin. Personal Communication.
- Wohlfarth, Barbara
1996 The Chronology of the Last Termination: A Review of Radiocarbon-Dated, High-Resolution Terrestrial Stratigraphies. *Quaternary Science Reviews* 14(4):267–284.

Woolly, Buck W.

1985 *Channel Changes on the Brazos River Below Whitney Dam Bosque, Hill, and McLennan Counties, Texas*. Unpublished Master's Thesis. Baylor University, Waco, Texas.

Yu, Z. and H. E. Wright, Jr.

2001 Response of Interior North America to Abrupt Climate Oscillations in the North Atlantic Region During the Last Deglaciation. *Earth Science Reviews* 52:333–369.

Zaitlin, Brian A., Robert W. Dalrymple, and Ron Boyd

1994 The Stratigraphic Organization of Incised-Valley Systems Associated with Relative Sea-Level Change. In: *Incised-valley Systems: Origins and Sedimentary Sequences* by Robert W. Dalrymple, Ron Boyd, and Brian Zaitlin (editors), pp. 45–60. SEPM (Society for Sedimentary Geology), Special Publication No. 51, Tulsa, OK.

Zhou, Jie and Henry S. Chafertz

2010 Pedogenic Carbonates in Texas: Stable-Isotope Distributions and Their Implications for Reconstructing Region-Wide Paleoenvironments. *Journal of Sedimentary Research* 80:137–150.

VITA

Ken L. Lawrence was born in Wichita, Kansas, on October 7, 1968, the son of Gary and Judy Lawrence. After graduation from Yates Center High School in 1987, he attended the University of Kansas and received a Bachelor of Arts in Anthropology in 1992. During the following eight years he worked as an archaeologist for P-III & Associates and SWCA in South Dakota, Colorado, Wyoming, Oregon, New Mexico, and Arizona with an emphasis in Nevada and Utah on the Great Basin. In 2000, he moved to Austin, Texas to work with SWCA where he is currently employed. During the last 18 years of archaeological work, he has developed an interest in geoarchaeological research and geomorphic processes and is currently improving his experience and education in these areas. In the fall of 2007, he entered the Department of Anthropology of Texas State.

Permanent address: 4604 Federal Circle

Austin, Texas 78744

This thesis was typed by Ken Lawrence.