

South Dakota State University

Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange

SDSU Extension Fact Sheets

SDSU Extension

1967

Oat Production in South Dakota

Elmer E. Sanderson

Leon S. Wood

Follow this and additional works at: https://openprairie.sdstate.edu/extension_fact

Recommended Citation

Sanderson, Elmer E. and Wood, Leon S., "Oat Production in South Dakota" (1967). *SDSU Extension Fact Sheets*. 1328.

https://openprairie.sdstate.edu/extension_fact/1328

This Fact Sheet is brought to you for free and open access by the SDSU Extension at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in SDSU Extension Fact Sheets by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



For current policies and practices, contact SDSU Extension

Website: extension.sdstate.edu

Phone: 605-688-4792

Email: sdsu.extension@sdstate.edu

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture.

FS 384

Oat Production

in
South Dakota



Cooperative Extension Service
South Dakota State University
Brookings, South Dakota

Oat Production in South Dakota

By Elmer E. Sanderson, Extension agronomist—crops,
and Leon S. Wood, Extension Plant pathologist.

South Dakota is a large oat producing state, but it's acreage has been gradually declining. The largest oat acreage was in 1954 when over 4 million acres were planted to oats. For the 5-year period of 1963-67, the average acreage was about 2½ million acres with an average annual production of nearly 92 million bushels.

The oat crop has added considerably to South Dakota agriculture economy. Oats rank first in total acres among all small grain crops and is surpassed only by wheat in total valuation.

Oats is produced both as a cash crop and as a feed crop. Usually oats has an economic advantage if used for livestock feed, and especially if used for hay or silage. Experiment Station agronomists found that the total digestible nutrients produced per acre can be doubled if the entire oat crop can be harvested as forage and fed to livestock.

OATS IN THE ROTATION

Oats usually follows an intertilled or row crop that leaves the soil in condition to be prepared quickly for spring planting. The yield is greater following a row crop than following other small grain crops in the rotation. Since legumes and plant residues help maintain soil productivity, the establishment of legumes is frequently accomplished using oats as a companion crop.

RESPOND TO SOIL FERTILITY

Oats, like all small grains, develops rapidly in the early spring when soil temperatures are cold. Under these conditions, the nitrogen release is low. If available plant food is lacking, especially nitrogen, the oat plant will be yellow, short, and have few or no tillers. Under such conditions, grain quality may be good but the yield disappointing. This is especially true for early varieties which must make their growth earlier in the season than late varieties.

Oats will respond to commercial fertilizers. There is some question as to whether the use of commercial fertilizers will always pay in the lower rainfall areas in western South Dakota. A soil test should be made to accurately determine soil fertility levels and plant food needs.

SEEDBED PREPARATION

Double disking and harrowing row crop land is a common method of preparing a seedbed for oats. This method is relatively cheap, fast, and leaves 3 to 4 inches of loose, friable soil on the surface with firm soil beneath. Spring plowing may give increased yields but is slow and more costly. All plowed land should be packed either before or after planting to prevent excessive moisture evaporation. Firm seedbeds are invaluable when drought conditions prevail.

TIME OF SEEDING

Oats should be seeded as soon as the soil can be properly worked with usual farming equipment. "Mudding in" before the surface soil has a chance to dry is not a good practice.

METHOD AND RATE OF SEEDING

Seeding with a grain drill is the best method. Drilling distributes the seed evenly at a uniform depth in moist soil where conditions are favorable for germination. Drilling is especially recommended for the drier areas. Broadcasting and disking-in is a cheaper seeding practice and can be justified on small acreages. Seed at the rate of 2 to 2½ bushels per acre. Western areas may even seed less than 2 bushels. Increase the rate slightly for broadcasting.

USE GOOD SEED

First requirement of a successful crop is to use pure seed, free of weed seed and of high germinating ability. A good farmer will seed nothing else. Certified seed is your assurance of good seed.

Research tests show that fully mature, plump, high test weight seed produces stronger seedlings with more rapid emergence than light weight seed. Therefore, it is important to plant only good clean seed where plenty of wind has been used to remove light weight and diseased seeds.

SEED TREATMENT

Proper seed treatment with a recommended fungicide is often a good practice. Seed treatment controls both loose and covered smut of oats and can help control certain other seed-borne and soil-borne diseases.

More detailed information on seed treatment and recommended fungicides are given in Fact Sheet 193 "Seed Treatment."

The actual cost of seed treatment per acre is small. Except for small amounts of seed, custom treating has proved practical. For custom and home treatment, instructions on the seed treatment container as to rate and method of treatment should be followed carefully. Thorough mixing of the fungicide and seed is necessary to get good seed coverage and derive maximum benefits from the treatment.

WEED CONTROL

Under normal rainfall conditions in South Dakota, you cannot expect to raise a weed crop and oat crop on the same field in the same year. A planned cultural weed control program should be practiced throughout the cropping rotation and should be supplemented with herbicides as needed. Most oat varieties are more tolerant to MCPA than to 2,4-D. General recommendations are to use one-third pound of 2,4-D acid in an ester or one-half pound in an amine form or one-half pound of MCPA acid per acre. Apply during the tolerant periods of growth which are usually between the 5-leaf and early boot stages. For complete information on spraying to control weeds in oats, variety tolerance, and so on, consult your county agent or secure Fact Sheet "Weed Control in Small Grains."

HARVESTING AND STORING

Most of the oat acreage is harvested with a combine, either direct or from a windrow. Because of weeds, uneven ripening, shattering, and possible high moisture of the grain, the windrowing and combining method is most common in eastern oat producing counties. Highest quality grain is obtained by allowing the oats to mature and threshing as soon as the grain is dry enough for safe storage. The moisture content of the grain should be 14% or less for safe storage.

MARKETING THE OAT CROP

Much of the oat crop is fed to livestock and then marketed in the form of livestock and livestock products. This means of utilizing the oats is usually the most profitable.

A very significant amount is marketed as a cash crop. Studying the cash oat market and selling when the price tends to be the highest, can increase the net profit. Usually, the price is down at harvest time and for several months after harvest. In order to take advantage of peak oat prices at certain seasons, it means some form of grain storage, either on the farm or at custom storage facilities.

South Dakota oats is in demand as feed oats in several southern states and has become known as the "truck oat market." The market usually pays a sizeable premium and starts in mid-winter and continues until near the next harvest season.

Oat producers should not forget about the milling oat market as in some years there is a worthwhile premium paid for oats meeting the milling oat standard. Good milling oats must possess certain quality factors, namely (1) low amount of foreign material, (2) freedom from mixture with other crops and (3) plump kernels with 36 pounds test weight or better.

DISEASES

Breeding new varieties is an ever-continuing process, mainly because new disease troubles follow new varieties. Sometimes a disease may have been minor or previously unknown. On occasions, new races or varieties of well-known disease organisms or little-known races have increased and caused severe losses. Under such changing conditions, older varieties which were once popular may reappear and produce well until older diseases also return. Newer varieties generally are much more disease resistant than those of a generation ago.

Leaf Rust (Crown Rust) is usually the most serious disease of oats in South Dakota. When severe, it causes lodging, light-weight kernels and low yields. Leaf rust races now prevalent in the United States consist of two groups: the older races to which many of the recommended varieties have moderately resistant reactions, and the new races (264, 295, and 326) to which most present varieties are all susceptible. Portal is resistant to Race 264. It cannot be predicted how rapidly these newer races will increase in importance, but plant pathologists and plant breeders are hopeful of developing oat varieties resistant to these races. Attempts to stay one step ahead of the constantly changing development of leaf and stem rust races accounts for the rather rapid change in oat varieties.

Stem Rust has not been serious in South Dakota in recent years. Because stem rust has not caused severe damage to oats since 1957, there may be a tendency to overlook the importance of stem rust resistance for successful oat production in South Dakota. Stem rust could become important again if newly discovered subraces become widely distributed. Races 6A, and 6AF can attack all present available commercial varieties. If, or how rapidly, these races increase in importance cannot be predicted. Research workers in several states have recently found sources of resistance in some breeding lines, and this resistance will be incorporated into commercial varieties as rapidly as possible.

"Red Leaf" of oats, a Virus disease, was present in most areas of the state in 1959 and reduced yields where infection was high early in the growing season. This disease has been minor since 1959, although insect carriers of the virus have been observed early in the growing season each year. All commercial oat varieties are susceptible; however, Brave and Jaycee have shown some field tolerance.

Smut resistance is found in all recommended varieties. Some nonrecommended varieties including Ajax and Nodaway are susceptible to certain races. Smut-susceptible varieties should be seed treated for control of this disease.

Halo blight, a bacterial leaf-spot disease, frequently is found early in the season. At first spots are yellow and later turn brown. The appearance of this disease on leaves often causes much concern. But the plants tend to outgrow the disease and the effect on yields is usually minor.

SELECTING THE BEST VARIETY

Selecting the best oat variety for a farm or for a certain field is an important decision. Growing an adapted variety or varieties insures more stable production. Ignoring this principle often invites disappointments and causes fluctuations in farm income. Recommendations and variety descriptions given in this Fact Sheet should help South Dakota farmers to choose their varieties.

There is no one variety of oats that is best for all areas or for all situations. Factors determining the selection of a variety are: (1) local climatic environments such as elevation, normal expected rainfall, and temperature, (2) soil type, (3) soil fertility, (4) varietal performance, (5) market demand, and (6) crop use.

VARIETY RECOMMENDATIONS

The list of recommended varieties for South Dakota (table below) is based on reliable and impartial information obtained from Experiment Station tests conducted throughout the state. These recommendations are based not only on yield but also on maturity, disease and insect resistance, straw strength, grain quality, market need, and so forth. Variety recommendations, according to "crop adaptation areas," are

given in Fact Sheet "Field Crop Varieties in South Dakota" for the current year.

The table gives the important characteristics of the more commonly grown oat varieties in South Dakota. The recommended group represents a list of good varieties adapted in one or more areas of the state. It is recognized that other varieties may have local interest and satisfactory performance. In some cases, varieties not recommended may not be inferior to those recommended but may merely represent duplication of qualities already available.

Secure these Fact Sheets for additional information on oats production:

- Crop Variety Recommendations
- Seed Treatment
- Fertilizing Small Grains
- Weed Control in Small Grains
- Producing Milling Oats

Agronomic Characteristics and Stem and Leaf Rust Reactions

Varieties Recommended	Yielding Ability	Bushel Weight	Stem ¹ Rust	Leaf ¹ Rust	Plant Height	Lodging Resistance	Seed Color
Early-Maturity							
Dupree	Medium	Medium	S ²	S ²	Short	Medium	White-Gray
Jaycee	Medium	High	MR	MS	Short	Good	Yellowish-White
Santee	Medium	Medium	MS	MS	Medium	Good	Ivory
Tippecanoe	Medium	High	MS	MS	Short	Excellent	Yellow
Tyler	Medium	High	MS	MS	Short	Excellent	Yellow-Tan
Medium Early-Maturity							
Clintland 64	Medium	High	MR	MR	Medium	Good	Yellow
Medium-Maturity							
Brave	High	Medium	MR	MS	Medium	Medium	Yellow
Burnett	High	High	MR	MS	Med-Tall	Good	Yellow-White
Coachman	Medium	High	MR	MS	Medium	Good	Yellow
Holden	High	Medium	MR	MR	Medium	Good	Yellow
Garland	High	High	MR	MR	Med-Short	Good	Yellow
Portal	High	Medium	MR	MR	Medium	Good	Yellow
Late-Maturity							
Kelsey	Excellent	Medium	MR	MR	Med-Tall	Medium	White
Lodi	High	Medium	MR	MR	Tall	Excellent	White
Rodney	High	High	MR	MS	Tall	Good	White
Sioux	Excellent	Medium	MR	MS	Tall	Medium	White
Varieties Not Recommended							
Early-Maturity							
Andrew	High	Medium	S	S	Medium	Medium	Yellow
Minhafer	Medium	Medium	MR	MS	Medium	Good	Yellow
Neal	Medium	Medium	MS	S	Med-Short	Good	Ivory
Nodaway	Medium	Medium	MR	S	Medium	Good	White
Ransom	Medium	Medium	MS	S	Med-Short	Medium	Yellow
Medium Early-Maturity							
Clintford	Medium	High	MS	MS	Med-Short	Excellent	Yellow-White
Clintland 60	Medium	High	MR	MS	Medium	Good	Yellow
O'Brien	Medium	Medium	MR	MR	Med-Tall	Good	Yellow
Medium-Maturity							
Dodge	Medium	Medium	MR	MR	Medium	Good	Yellow
Marion	Medium	Medium	S	S	Medium	Medium	Tan-White
Stormont	Medium	Medium	MR	MS	Medium	Medium	White
Medium Late-Maturity							
Ajax	High	Low	S	S	Tall	Medium	White
Orbit	High	Low	MR	MR	Medium	Good	White
Portage	High	Medium	MS	MR	Tall	Medium	White
Sauk	High	Medium	S	MS	Tall	Medium	Yellow
Late-Maturity							
Branch	Medium	Medium	S	MS	Tall	Medium	White
Harmon	High	Low	MR	MS	Tall	Medium	White
Garry	High	Medium	MR	MS	Tall	Good	White
Ortley	High	High	MR	MS	Tall	Medium	White
Russell	Medium	Medium	MR	MR	Tall	Medium	White

¹Disease reactions to the prevalent races of rust.

²R=Resistant; MR=Moderately Resistant; MS=Moderately Susceptible; S=Susceptible.