

South Dakota State University

Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange

SDSU Extension Fact Sheets

SDSU Extension

1969

Grasshopper Control in South Dakota – 1969

B. H. Kantack

Wayne L. Berndt

P. A. Jones

Follow this and additional works at: https://openprairie.sdstate.edu/extension_fact

Recommended Citation

Kantack, B. H.; Berndt, Wayne L.; and Jones, P. A., "Grasshopper Control in South Dakota – 1969" (1969). *SDSU Extension Fact Sheets*. 1421.

https://openprairie.sdstate.edu/extension_fact/1421

This Fact Sheet is brought to you for free and open access by the SDSU Extension at Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in SDSU Extension Fact Sheets by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



For current policies and practices, contact SDSU Extension

Website: extension.sdstate.edu


Phone: 605-688-4792

Email: sdsu.extension@sdstate.edu

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture.

GRASSHOPPER CONTROL

South Dakota—1969



Cooperative Extension Service
South Dakota State University
U. S. Department of Agriculture

Grasshopper Control in South Dakota—1969

by B. H. KANTACK, extension entomologist; WAYNE L. BERNDT, extension pesticide specialist;
and P. A. JONES, experiment station entomologist

GRASSHOPPER PROBLEMS

Grasshoppers are usually prevalent in some areas of South Dakota each year. Heavy outbreaks also are likely to occur periodically where extensive acreages are infested over a wide area.

Severe infestations occur during seasons where hot and dry weather conditions prevail. Farmers and ranchers should watch for grasshoppers early in the season and initiate control measures immediately where problems exist.

APPLY CONTROL MEASURES EARLY

Two general groups of injurious grasshoppers are of economic importance in South Dakota. Species of economic importance are usually grouped into cropland and rangeland grasshoppers.

The best time to control grasshoppers is when they are small and before they migrate into cropland areas. Apply insecticides shortly after the main egg hatch is completed. There are a number of advantages in this early treatment: (1) fewer acres will have to be treated and less insecticide is necessary to obtain control; (2) injurious grasshopper species are killed before they have had the opportunity to injure the crops; (3) early treatments before the grasshoppers have reached maturity prevent egg deposition which helps reduce the potential grasshopper threat for the following crop year.

Number of Grasshopper Adults Per Square Yard

Field	Margin	Infestation or damage category
0-2	5-10	non-economic
3-7	11-20	light
8-14	21-40	moderate to abundant
15-28	41-80	threatening to severe
over 28	over 80	very severe

The above table applies both to infestations in cropland and rangeland. Insecticide control is advised when the infestation reaches threatening levels.

TREATING GRASSHOPPER INFESTATIONS IN DIFFERENT SITUATIONS

Cropland (Small Grains, etc.). Fall and spring tillage of the soil helps reduce grasshopper populations. Inspect all grain fields periodically and treat only areas where grasshoppers are found. Field margins and headlands are areas where young grasshoppers are usually present. In South Dakota, winter

wheat fields are often severely damaged each fall; border spraying of these fields can prevent considerable damage.

Legume Fields. Usually the entire field is infested and must be sprayed to obtain adequate control. Since some grasshopper species hatch later, the main hatch will not be completed until after the first crop of hay has been harvested. Delay spraying until the second crop is about 3 inches tall in legume fields. Sometimes growers leave a small trap strip from the first cutting to attract grasshoppers from the remainder of the field. This strip is then sprayed with a suitable insecticide.

Conservation Reserve (Soilbank). In fields where forage crop cover is lacking and weeds are prevalent, the entire field will often be infested with grasshoppers. In fields where grass growth is good, the grasshoppers will usually be concentrated along the margins and borders of the field. Under dry conditions considerable migration from these soilbank areas into adjacent crops usually occurs.

Roadside. Fence rows and roadsides are favorite oviposition sites for grasshoppers. Often a properly timed spray in these areas after the main grasshopper hatch will eliminate the need for a costly spray on adjacent cropland. Spray these areas before the grasshoppers begin migrating into adjacent cropland.

Pastureland. Grasshopper problems often arise in pastures that have been overgrazed or mismanaged. Heavy infestations in pastures result in considerable loss of grass and grazing capacity. Grasshoppers from pasture areas often invade adjacent field crops later in the season.

Proper pasture management and improvement methods will discourage grasshopper populations. When insecticides are needed for control of grasshoppers in pastures, it is very important that the proper insecticide is selected so that grazing is permitted either immediately or soon after application.

Rangeland. In rangeland areas, grasshopper problems frequently arise. A severe rangeland infestation can reduce the grass growth so that the carrying capacity of livestock is greatly lowered. For lands under federal domain, the responsible agency treats the infested areas when the situation warrants control measures. In areas of privately owned rangeland, co-

The U. S. Department of Agriculture will pay one-third of the cost on privately owned land and state land, the rancher must pay the remaining two-thirds. The U. S. Department of Agriculture will pay the full cost of control on government land on short term lease or leased on a permit basis, provided the privately owned land intermingled or adjacent to the government land is signed up. Government land that is under direct management and control of a governmental agency is included in this category.

Government land in long-term lease is handled the same as private land, that is, the Department of Agriculture will pay one-third and the rancher leasing the land must pay two-thirds.

The material used is a liquid spray, applied by aircraft. No spraying is done until all the grasshoppers have hatched (late June or early July) and the spraying must be completed before the grasshoppers start laying their eggs (the last few days of July or the first week of August).

See pages 3-4 for Insecticide Recommendations

PRECAUTIONS

Select the insecticide to fit the situation. Always follow the label and use recommended rates. Pay particular attention to the waiting periods required between the time of application and harvest or grazing of the treated areas. Never apply an unlabeled insecticide or allow an unlabeled insecticide to be applied on your crop. Always read and understand the label.

SPECIAL NOTE

Aldrin and dieldrin cannot be used for grasshopper control on field crops, forage or pasture lands, as the USDA label approval has been withdrawn.

CANCELLED

USDA labels for Chlordane and Heptachlor on grass and the label for Chlordane on alfalfa were cancelled in 1968.

**INSECTICIDES RECOMMENDED AND PRECAUTIONS FOR PROPER USE
FOR GRASSHOPPER CONTROL ON VARIOUS CROPS**

CROP	INSECTICIDE	DOSAGE*	REMARKS, PRECAUTIONS
Small grain	Toxaphene	1½-2 lbs./A.	Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitation on use of grain.
	Chlordane	1 lb./A.	Do not apply after heads start to form. Do not graze treated forage. No restrictions on the use of grain.
	Malathion	1 lb./A.	Do not harvest for 7 days.
	Parathion (Ethyl or Methyl)	8 oz. /A.	Experienced aerial applicators only. Do not harvest for 15 days.
	Malathion (ultra low-volume concentrate)	8 fl. oz./A.	Commercial aerial applicators only. Do not harvest for 7 days.
Corn	Carbaryl (Sevin)	1½-2 lbs./A.	No time limitation.
	Malathion	1 lb./A.	Do not harvest for 5 days.
	Malathion (ultra low-volume concentrate)	8 fl. oz./A.	Aerial applicators only. Do not harvest for 5 days.
	Toxaphene	1½-2 lbs./A.	Do not feed treated forage to dairy cattle or animals being finished for slaughter. No limitation on use of grain.
	Chlordane	1 lb./A.	Do not feed treated forage to dairy cattle or animals being finished for slaughter. No restrictions on the use of grain.
	Parathion (Ethyl or Methyl)	8 ozs./A.	Experienced aerial applicators only. Do not harvest for 12 days.
Sorghum	Carbaryl (Sevin)	1½ lbs./A.	No time limitation on corn or sorghum grown for forage. Do not apply on sorghum grown for grain within 21 days of harvest.
	Toxaphene (grain sorghum only)	1½ lbs./A.	Do not graze toxaphene treated forage with dairy animals or animals being finished for slaughter. Do not ensile treated forage. Do not apply more than once after heads start to form. Do not harvest within 28 days.
	Ethyl Parathion	8 ozs./A.	Experienced aerial applicators only. Do not harvest for 12 days.
(CAUTION: Do not spray methyl parathion or mixtures of ethyl and methyl parathion on sorghum as plant injury may occur).			
Soybeans	Carbaryl (Sevin)	1½ lbs./A.	No time limitations.
	Malathion	1 lb./A.	Do not apply within 1 day of harvest.
	Malathion (ultra low-volume concentrate)	8 fl. ozs./A.	Do not harvest for 7 days.
	Toxaphene	1½ lbs./A.	Do not feed treated forage to dairy animals or animals being finished for slaughter. No restrictions on threshed beans. Do not harvest for 21 days.
	Chlordane	1 lb./A.	Do not feed to dairy animals or animals being finished for slaughter. No restrictions on threshed beans.
	Parathion (Ethyl)	8 ozs./A.	Experienced aerial applicators only. Do not harvest for 15 days.
	Parathion (Methyl)	8 oz./A.	Experienced aerial applicators only. Do not harvest for 20 days.
	Guthion	8 oz./A.	Experienced aerial applicators only. Do not harvest for 21 days.

CROP	INSECTICIDE	DOSAGE*	REMARKS, PRECAUTIONS
Alfalfa, clover-forage and seed	Malathion	1 lb./A.	No time limitations.
	Carbaryl (Sevin)	1½ lbs./A.	No time limitations.
	Diazinon	1 lb./A.	Do not graze livestock within 2 days or cut for hay within 10 days of application. Do not spray alfalfa in bloom.
	Dibrom (Naled)	1 lb./A.	Do not apply within 4 days of cutting for hay.
	Cygon	4 ozs./A.	Do not apply more than once per season or within 28 days of harvest.
	Parathion (Ethyl or Methyl)	8 oz./A.	Experienced aerial applicators only. Do not apply within 15 days of harvest
	Guthion	8 ozs./A.	Experienced aerial applicators only. Do not harvest for 16 days.
	Malathion (ultra low-volume concentrate)	8 fl. ozs./A.	No waiting period. Do not spray alfalfa or clover in bloom.

NOTE: Malathion ultra low-volume concentrate is not recommended for grasshopper control after alfalfa is over eight inches tall or where very dense foliage is present. Where dense foliage is encountered and ULV malathion is being used, a five to six foot boom height with a 50-foot swath width is recommended.

To protect bee population, spray before 7:00 a.m. or after 7:00 p.m. Never spray alfalfa in bloom. Notify local beekeepers before spraying.

SPECIAL NOTE: Never spray aldrin, dieldrin, heptachlor or chlordane on alfalfa or clover. The registration for dieldrin and heptachlor was cancelled on alfalfa and clover in 1964. Foliar applications of aldrin and dieldrin were cancelled on a number of other field crops early in 1966. The chlordane label for alfalfa was cancelled in 1968.


Pastures and ranges, roadsides, field margins, ditches and other lands	Malathion	1 lb./A.	No time limitation. Dairy cattle must be removed from the pasture during application of the sprays.
	Malathion (ultra low-volume concentrate)	8 fl. oz./A.	Aerial applicators only. May be harvested or grazed on day of application.
	Carbaryl (Sevin)	1½ lbs./A.	No time limitation.
	Toxaphene	1¼-2 lbs./A.	Apply only one application per person. Do not graze meat animals in treated fields within six weeks of slaughter. Do not apply to forage to be sold commercially or shipped interstate.

CAUTION: All dairy animals must be removed from areas being treated when malathion or Sevin are being applied but can be turned into areas after spraying operation is completed. Avoid applying insecticides near streams, farm ponds, etc. to avoid any possibility of injuring fish and wildlife.

*In terms of actual toxicant per acre

Use of a trade name does not indicate endorsement of one product over another.

1969 GRASSHOPPER OUTLOOK FOR SOUTH DAKOTA



Should ACCIDENTAL POISONING occur, direct contact can be made with:

South Dakota Poison Information Center
Vermillion, South Dakota
Telephone (605)624-3432

COOPERATIVE EXTENSION SERVICE
U. S. DEPARTMENT OF AGRICULTURE
SOUTH DAKOTA STATE UNIVERSITY
Brookings, S. D. 57006
OFFICIAL BUSINESS
10M-3-69-8862

Postage and Fees Paid
U. S. Department of Agriculture



Cooperative Extension Service
South Dakota State University
U. S. Department of Agriculture

**GRASSHOPPER
CONTROL**
South Dakota-1969