

Chapman University

Chapman University Digital Commons

ESI Working Papers

Economic Science Institute

5-11-2020

Age Appropriate Wisdom? Ethnobiological Knowledge Ontogeny in Pastoralist Mexican Choyeros

Eric Schniter

Chapman University, schniter@chapman.edu

Shane J. Macfarlan

University of Utah

Juan J. Garcia

University of Utah

Gorgonio Ruiz-Campos

Universidad Autónoma de Baja California

Diego Guevara Beltran

University of Arizona

See next page for additional authors

Follow this and additional works at: https://digitalcommons.chapman.edu/esi_working_papers



Part of the [Economic Theory Commons](#), and the [Other Economics Commons](#)

Recommended Citation

Schniter, E., Macfarlan, S. J., Garcia, J. J., et al. (2020). Age appropriate wisdom? Ethnobiological knowledge ontogeny in pastoralist Mexican Choyeros. *ESI Working Paper 20-31*.

https://digitalcommons.chapman.edu/esi_working_papers/326/

This Article is brought to you for free and open access by the Economic Science Institute at Chapman University Digital Commons. It has been accepted for inclusion in ESI Working Papers by an authorized administrator of Chapman University Digital Commons. For more information, please contact laughtin@chapman.edu.

Age Appropriate Wisdom? Ethnobiological Knowledge Ontogeny in Pastoralist Mexican Choyeros

Comments

ESI Working Paper 20-31

Authors

Eric Schniter, Shane J. Macfarlan, Juan J. Garcia, Gorgonio Ruiz-Campos, Diego Guevara Beltran, Brenda B. Bowen, and Jory C. Lerbak



Click here to view linked References

1
2
3
4 **Age Appropriate Wisdom?**

5
6 **Ethnobiological Knowledge Ontogeny in Pastoralist Mexican Choyeros**

7
8 Eric Schniter^{1*}, Shane J. Macfarlan^{2,3,4}, Juan J. Garcia⁵, Gorgonio Ruiz-Campos⁶, Diego Guevara Beltran⁷, Brenda
9 B. Bowen^{4,8}, Jory C. Lerback⁸

10
11 ¹Economic Science Institute, Chapman University, Orange, California, USA

12 ²Department of Anthropology, University of Utah, Salt Lake City, Utah, USA

13 ³Center for Latin American Studies, University of Utah, Salt Lake City, Utah, USA

14 ⁴Global Change and Sustainability Center, University of Utah, Salt Lake City, Utah, USA

15 ⁵Department of Education, Culture, & Society, University of Utah, Salt Lake City, Utah, USA

16 ⁶Departamento de Biología, Universidad Autónoma de Baja California, Ensenada, Baja California, México

17 ⁷Department of Psychology, University of Arizona, Tucson, Arizona, USA

18 ⁸Department of Geology & Geophysics, University of Utah, Salt Lake City, Utah, USA

19
20
21 *Corresponding author: Eric Schniter, eschniter@gmail.com (email), 714-628-7272 (telephone). ORCID:
22 <https://orcid.org/0000-0002-2381-3616>
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **Abstract**
5

6 We investigate whether age profiles of ethnobiological knowledge development are consistent with predictions
7 derived from life history theory about the timing of productivity and reproduction. Life history models predict
8 complementary knowledge profiles developing across the lifespan for women and men as they experience changes
9 in embodied capital and the needs of dependent offspring. We evaluate these predictions using an ethnobiological
10 knowledge assessment tool developed for an off-grid pastoralist population, known as *Choyeros*, from Baja
11 California Sur, Mexico. Our results indicate that while individuals acquire knowledge of most dangerous items and
12 edible resources by early adulthood, knowledge of plants and animals relevant to the age and sex divided labor
13 domains and ecologies (e.g., women’s house gardens, men’s herding activities in the wilderness) continues to
14 develop into middle adulthood but to different degrees and at different rates for men and women. As the demands of
15 offspring accumulate for parents with age, reproductive aged adults continue to develop their knowledge to meet
16 their children’s needs. After controlling for vision, post-reproductive adults’ show the greatest ethnobiological
17 knowledge. These findings extend our understanding of the evolved human life history by illustrating how changes
18 in embodied capital and the needs of dependent offspring predict the development of men’s and women’s
19 ethnobiological knowledge across the lifespan.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 *Keywords:* Traditional ecological knowledge; ethnobiological knowledge; learning; embodied capital; life history
39 theory; Baja California Sur, México
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

Life history models predict knowledge profiles developing across the lifespan as organisms learn, grow, reproduce, invest (e.g. nutritionally, educationally) in dependent offspring, and senesce. Debates in anthropology have centered on alternative interpretations of what shapes these age-specific knowledge profiles should take across the lifespan. One view, inspired by general life-history models applied to primate foraging (e.g., Schuppli et al., 2012) suggests that effort allocated to productivity (which includes investments in acquiring knowledge) should increase up until physical maturation and plateau for the remainder of the life course (e.g. see discussion in Kramer & Ellison, 2010). This “wise adults” view suggests that the wisdom of younger adults should be equally developed as that of older adults. A different view, held widely among social scientists is that vocabulary and worldly knowledge, which relies on experience, (what Cattell (1963) called "crystallized intelligence") continues to accumulate past the age of physical maturation, resulting in “wisdom of elders” (Berlin, 1992; Gurven et al., 2017; Koster et al., 2016). Heterogeneity in aging and the experience of stochastic shocks (e.g. injury, illness) contribute to unequal abilities among elders to develop this wisdom, challenging the universality of wise elders claim. Previous studies have reported peak ages for ethnobiological knowledge through a wide range of adult ages: around 30-80 years of age for Mayangna and Miskito forager-horticulturalists of Nicaragua (Koster et al., 2016), around 30-40 years for Djerma farmers, and 40-50 years for Peulh herdsman of Nigeria (Ayantunde et al., 2008).¹

The embodied capital perspective (Kaplan, 1996) considers (from a biological fitness maximization perspective) how individuals and their parents optimally invest in growing and maintaining the body’s physical capital (e.g. bones, muscles, neurons) and functional capital (e.g. strength and coordination, skill, knowledge). According to embodied capital theory, extreme investments in cognitive capital have led to a human life history pattern of complementary “age appropriate” knowledge profiles developing across the lifespan for women and men as they experience changes in their stocks of embodied capital and demand of their dependent offspring (Del Giudice et al., 2016). This perspective predicts that levels of productivity and childrearing increase through reproductive parent ages when offspring provisioning demands (for both calories and knowledge/experience) are growing (peaking approximately 35-45 years of age), and should remain high until impacted by senescence (e.g. of physical mobility, visual, and hearing systems). We use “age appropriate” to refer to the nuanced view that an older

¹ Others also report ethnobiological knowledge variation at different ages and across generations (Hewlett et al., 2011; Kline et al., 2013; Ohmagari & Berkes, 1997; Schniter et al., 2015; Zarger & Stepp, 2004; Zent, 2001).

1
2
3
4 adults' ability to demonstrate knowledge is constrained by their particular abilities (e.g. to see, to move about, to
5 remember), and that there is increasing heterogeneity in these abilities with age.
6
7

8 To gauge ethnobiological knowledge, we evaluate responses to a set of images showing culturally salient
9 plants and animals that we presented to Mexican *Choyeros*, an off-grid arid land pastoralist population whose
10 livelihood has been closely tethered to arid land springs and open rangelands of the Sierra de la Giganta mountains
11 of Baja California Sur for more than 300 years (Cariño Olvera, 2014; Koster et al., 2019). From an evolved life
12 history strategy perspective, we predict how age-appropriate complementary knowledge profiles develop across the
13 lifespan for *Choyero* men and women as they experience changes in embodied capital and the needs of dependent
14 offspring. We report age profiles of *Choyero* ethnobiological knowledge development that are consistent with
15 predictions derived from life history theory. This paper sheds new light on the distribution of ethnobiological
16 knowledge maintained by a relatively isolated and small-scale subsistence society.
17
18
19
20
21
22
23
24
25

26 **Hypotheses and predictions**

27 We test predictions based on our overarching hypothesis that the time-path of ethnobiological knowledge
28 development corresponds with functional “life goals” and embodied capital constraints specific to each of three
29 stages of the human life course. In the *pre-reproductive life stage* the goals are to: (1) acquire basic skills and
30 knowledge (e.g., ability to recognize and name edible and dangerous plants and animals) to support one’s self; and
31 (2) attract and secure a mate, which often involves demonstrating basic competence consistent with the first goal.
32 The goals of the *reproductive life stage* are to (3) reproduce, optimally developing and using skills and knowledge to
33 support reproduction, such that (4) offspring are successfully raised to be able to accomplish pre-reproductive and
34 reproductive goals 1, 2, 3, and 4 (this entails transferring both material resources and *knowledge* to dependents).
35 After reproduction ceases with menopause, additional years of life are needed for the last-born offspring to fully
36 achieve goals 1, 2, 3, and 4. During transition to the *post-reproductive life stage*, adults (5) continue making
37 contributions towards fulfilling kin’s goals 1, 2, 3, and 4, despite (and in compensation for limitations related to)
38 age-related senescence. Below we derive several predictions concerning ethnobiological knowledge development
39 across the life course.
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 **Pre-reproductive life stage.** The pre-reproductive life stage includes gestation, infancy, childhood,
55 adolescence, and ends at first reproduction. During this period, young adults are generally protected, remain
56 relatively free of responsibilities that require great exertion or risk, and are supported by the surplus food production
57
58
59
60
61
62
63
64
65

1
2
3
4 and knowledge provided by older adults (Kaplan, 1994; Kaplan et al., 2000; Lee, 2013; Lee & Kramer, 2002).
5
6 Social learning of conceptual elements and procedures needed for survival in a subsistence society is well underway
7
8 during childhood (Hewlett et al., 2011; Kline et al., 2013; Ruddle & Chesterfield, 1977; Stieglitz et al., 2013).
9

10 Cross- culturally people name and classify plants and animals - a key to rapid and effective learning
11
12 (Atran, 1990; Berlin, 1992; Mithen, 2006). What is even more striking is the ease with which some of this
13
14 knowledge is acquired at young ages – human minds appear to be pre- tuned for acquiring and processing
15
16 information about certain classes of animals and plants relevant to adaptive challenges in early life stages throughout
17
18 our evolutionary past (Atran, 1990; Barrett et al., 2016). Below we consider the reasons and evidence for
19
20 preferential knowledge of plants and animals that are potentially dangerous and/or food.
21

22 It is important for youngsters to prioritize acquiring information about dangerous plants and animals
23
24 without experiencing morbid and mortal hazards first-hand. It appears that children are born with an innate
25
26 understanding of what distinguishes living from inanimate things and begin learning about plants and animals at the
27
28 youngest ages (Atran, 1990; Barrett & Behne, 2005; Carey & Spelke, 1994; Keil, 1987, 1994). Infants in the first
29
30 and second year of life selectively identify plants over artifacts as food sources, but also show a greater reluctance
31
32 manipulating plants due to the dangers they can pose (Wertz & Wynn, 2014a, 2014b). As an alternative to direct
33
34 hands-on exploration, infants prefer to learn about plants indirectly by watching or receiving cues from adults
35
36 (Elsner & Wertz, 2019; Oña et al., 2019). Likewise, cross-cultural evidence shows that children show preferential
37
38 attention to, learning of, and ability to remember culturally transmitted information about dangerous animals
39
40 (Barrett & Broesch, 2012).
41

42 Over the course of evolutionary history, it was also likely important for youngsters to identify plants and
43
44 animals as natural sources of food or as competitors for food, the latter of which could present danger (Barrett, 2005;
45
46 Brantingham, 1998). Though dangerous hazards prevent child foraging opportunities in many ecologies, some
47
48 children as young as young as 3-5 years of age forage with other children or while accompanying their mothers on
49
50 foraging trips providing rich opportunities to gain important information about plant and animal dangers (Bird &
51
52 Bird, 2002; Crittenden et al., 2009; Hewlett, 2017). Brown (1985) has suggested that, given the hazards of
53
54 agricultural crop and livestock failures, an interest in wild plant and animals that are potential food sources may
55
56 have been crucial to the survival of small-scale agriculture groups. Others have noted that edible plants and animals
57
58 are more salient than non-edible things in the minds of horticultural people (e.g., Berlin, 1992; Malinowski, 1925).
59
60
61
62
63
64
65

1
2
3
4 During the pre-reproductive life stage ethnobiological knowledge is transmitted via specific social vectors.
5
6 Vertical transmission of important knowledge and subsistence skills during childhood often comes from same-sex
7
8 parental figures (Borgerhoff Mulder et al., 2019; Garfield et al., 2016; Reyes-García et al., 2009; Schniter et al.,
9
10 2015). This vertical transmission pattern promotes information stability over generations, even in the face of
11
12 sociopolitical, economic, and environmental change (Hewlett & Cavalli-Sforza, 1986; Ohmagari & Berkes, 1997;
13
14 Zarger & Stepp, 2004). The necessary strength and size needed for coping with dangers, independent performance,
15
16 and dedicated practice of difficult and physically demanding skills are often undeveloped until after the adolescent
17
18 growth spurt (Bogin & Smith, 1996), when adult stature is nearly attained and most subsistence skills are finally
19
20 mastered (e.g., Cain, 1977; Draper, 1976; Ruddle & Chesterfield, 1977). As learning-by-doing continues during
21
22 adolescence and early adulthood, people often have greater opportunities to work and play alongside peers, learn
23
24 horizontally from peers, and socialize beyond the nuclear family (Gallois et al., 2017; Hewlett & Cavalli-Sforza,
25
26 1986; Reyes-García et al., 2016). Oblique transmission (from older relatives other than parents), often same-sex, is
27
28 reported to be a dominant mode of ethnobotanical knowledge learning and subsistence skill development after early
29
30 childhood (Garfield et al., 2016; Reyes-García et al., 2009; Schniter, 2009; Setalaphruk & Price, 2007). A pattern of
31
32 knowledge acquisitions that includes delays due to physical strength development and due to delayed social
33
34 opportunities for learning suggests the possibility of continued knowledge development into adulthood (consistent
35
36 with wisdom of elders and embodied capital model predictions).
37

38
39 *Prediction 1. Knowledge of plants and animals will develop monotonically in the pre-reproductive life*
40
41 *stage and children will develop more knowledge of plants and animals that are potentially food and/or dangerous.*
42

43 **Reproductive life stage.** The reproductive life stage for females is defined by fertility, spanning from the
44
45 onset of reproduction to its cessation prior to menopause. During the reproductive stage, a primary goal is
46
47 maintaining sufficient food production, education, and care to meet the waxing and waning demands of a family of
48
49 dependents accumulated under a fertility schedule. The embodied capital model proposes that parental and
50
51 grandparental childcare and support allow children the learning and on-the-job training opportunities in areas of
52
53 food production, craft production, and childrearing (Kaplan, 1996). Parental costs of provisioning and caring for
54
55 young are offset by offspring's increasing surplus returns over the long reproductive and post-reproductive lifespan
56
57 (Kaplan et al., 2000; Robson & Kaplan, 2003).
58
59
60
61
62
63
64
65

1
2
3
4 Maintaining proximity to helpful kin may be important because one's ability to be economically self-
5 sufficient and fully support dependents is often achieved later than sexual maturity (Hill & Hurtado, 2009; Howell,
6 2010; R. D. Lee, 2013; Marlowe, 2007; Walker et al., 2013). Delayed productivity and downward resource flows
7 among adults means parenting and grand parenting roles will often blend together towards the end of this stage, with
8 adults possibly contributing support to two generations of dependents prior to cessation of the reproductive stage.
9

10
11 During fertile adult years, as the number and net caloric demands of dependents increase within a family,
12 parents and grandparents are faced with greater provisioning and caregiving demands. Children's daily caloric net
13 consumption requirements almost double between early childhood and teenage years (Institute of Medicine, 2002).
14

15
16 When the net caloric demand of dependents is at its highest, adults who rely on subsistence skills produce
17 the greatest amount of food of their life (Gurven & Walker, 2006). During these times of high productivity, adults
18 will also be actively engaged with relevant knowledge. Others have proposed that ethnobiological knowledge
19 appears to develop as a function of time allocated to related tasks (Guest, 2002; Koster et al., 2016; Reyes-García et
20 al., 2013). In general, we expect that performance with knowledge skills related to subsistence (like plant and animal
21 identification and naming) is most needed, most practiced, and thus at highest levels around the time of greatest
22 offspring demand.
23
24

25
26 With a focus on labor substitution within the household, Bock (2002a, 2002b) suggested that family
27 members make optimal contributions to fitness given their strength and knowledge as a function of age, and
28 differences in energetics of reproduction. From this a sexual division of labor is expected where men and women
29 invest in household and childrearing contributions that complement one another. Over the course of human
30 evolution and across cultures today, starting in early childhood, females typically move through smaller home
31 ranges while males range over significantly larger areas (Gaulin & Hoffman, 1988). In subsistence societies we see
32 evidence of a clear sexual division of labor that affects men's and women's activities in and experience of the
33 environment (Gurven et al., 2012; Kaplan et al., 2009). Plants may be differentially distributed (e.g. near human
34 habitation vs. in the wilderness) and encountered differently by men and women who range across and frequent
35 different environments.² Sex differences in men's and women's activities in their respective economic domains
36 could affect reproductive aged adults' exposure to and knowledge of the plants.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

59 ² Like plants, animals may also be differentially distributed across environments (e.g. human inhabited vs
60 uninhabited), but potentially less so due to their mobility and movements across environments.
61

1
2
3
4 *Prediction 2. Ethnobiological knowledge should continue to develop into reproductive adulthood as*
5
6 *parents continue to meet their offspring's burgeoning needs*
7

8 *Prediction 3. During the reproductive life stage, ethnobiological plant knowledge relevant to gender*
9
10 *associated domains develops to different degrees and at different rates for women and men.*
11

12 **Post-reproductive life stage.** The post-fertility period that we call the “post-reproductive life stage” is
13
14 mostly an extension of the reproductive career. Physical strength that contributes to productive subsistence work is
15
16 highest earlier in adulthood and declines into later adulthood, a human aging process referred to as sarcopenia
17
18 (Rosenberg, 1997; Walker et al., 2002; Walker & Hill, 2003). By the seventh decade of life when one’s children are
19
20 all economically self-sufficient, caloric production among subsistence people declines (Amoss & Harrell, 1981;
21
22 Gurven et al., 2012; Gurven & Kaplan, 2006; Walker & Hill, 2003), food transfers to kin declines (Hooper, 2011;
23
24 Kaplan et al., 2000), and functional disability increases (Kaplan et al., 2010; Stieglitz et al., 2015). Despite their
25
26 functional deficits, older adults are often capable of transferring important information to and helping younger kin
27
28 (Burke & Mackay, 1997; Schniter, 2014). Delayed productivity, extending beyond peak ages of offspring
29
30 dependency, is consistent with both embodied capital model and grandmother hypothesis (Hawkes, 2003; Hawkes et
31
32 al., 1989) expectations. While the grandmother hypothesis is a one-sex model of life history evolution that focuses
33
34 on selection for older females who are grandmothers, the embodied capital model is a two-sex model that predicts
35
36 both sexes will make important complementary post-reproductive contributions to fitness. For example, by sharing
37
38 their late life wisdom with close kin.
39

40
41 Late life wisdom is also consistent with a cognitive reserve hypothesis arguing that encephalization of brain
42
43 tissue, particularly in the cerebral cortex, which is specialized in the storage, retrieval, and processing of
44
45 experiences, represents an increased investment in the capacity to transfer important information (Parker &
46
47 McKinney, 2012). Returns from accumulated “cognitive capital”, for example via transfers of ethnobiological
48
49 knowledge (e.g. via applied and communicated knowledge and experience), may show stability with advancing
50
51 adult age or even increase until the final decade of life (Park et al., 2002; Salthouse, 1993; Schaie, 1994). Thus,
52
53 conceptual and procedural learning may continue in the post-reproductive life stage, producing wise elders decades
54
55 after the expected age of peak transfers needed to meet dependents’ needs.
56

57 Specific physical and cognitive abilities (e.g., strength, hearing, seeing) may decline with age and
58
59 negatively impact older adults’ ability to develop or recall knowledge stored in memory. Overall memory
60
61

1
2
3
4 performance is traditionally understood in terms of the concerted functioning of three stages – encoding, storage,
5 and retrieval (Craik & Rose, 2012). The encoding and storage of plant and animal information tends to take place at
6 early ages when people are first exposed to information about them. Retrieval of stored information based on
7 stimulus prompt (e.g. recognizing a seen item which triggers the retrieval of its name) is improved with practice, a
8 process sometimes called “rehearsal” but also consistent with practice and “general use models” predicting
9 knowledge. Age related decrements in ability to correctly name recognized items may reflect inefficiencies of
10 retrieval operations involved in memory but may also reflect age-related changes in activity and experience that
11 affect how older adults interact with memory prompts in their environment. Age-related physical and cognitive
12 declines affect older adults’ interaction with the stimulus that facilitates memory.
13
14
15
16
17
18
19
20
21

22 Finally, visual acuity directly affects how carefully one can recognize encountered items. Recognition
23 affects rehearsal and recall ability. If people are not seeing well, they might not be recognizing things as clearly as
24 when they saw better and therefore have less reason to recall and rehearse their memories (e.g. details and names).
25 Perceptual deterioration reduces recall of specific detail from the encoded record, leading to degraded memory
26 (Craik & Rose, 2012; Schneider & Pichora-Fuller, 2000).
27
28
29
30
31

32 As heterogeneity in physical (Maddox & Clark, 1992; Nelson & Dannefer, 1992) and mental abilities
33 (Ardila, 2007; Christensen et al., 1999) increases at later ages, it reinforces greater specialization in the post-
34 reproductive life stage, with older adults increasingly focusing on skills best suited to their abilities and contribution
35 opportunities. Heterogeneous patterns of aging suggest that while a “wisdom of elders” pattern may not emerge for
36 most or on average, the highest levels of knowledge demonstrated among adults may be differentially represented by
37 a few wise individuals in their post-reproductive years.
38
39
40
41
42
43

44 *Prediction 4. The highest level of ethnobiological knowledge will be observed among a subset of adults in*
45 *their post-reproductive years.*
46
47

48 *Prediction 5. Knowledge in gender associated economic domains among post-reproductive adults is*
49 *explained by visual acuity variation and the different male and female activity patterns in their gender associated*
50 *economic domains.*
51
52
53
54

55 **Methods**

56
57
58
59
60
61
62
63
64
65

1
2
3
4 Below we present information about our study sample, their biogeographical setting, way of life, and
5
6 ethnobiological knowledge breadth. We also describe research we conducted that contributed to our knowledge
7
8 about our sample including their fertility schedules, family budgets, and ethnobiological knowledge.
9

10 **Study sample and ecology**

11
12 Our study was conducted across four ranching communities (Santa Maria de Toris, San Pedro de la Presa,
13
14 La Higuera, and La Soledad) in the southern Sierra de La Giganta mountain range within the municipality of La Paz,
15
16 Baja California Sur, México (see Figure 1).
17

18 <Insert Figure 1 about here>
19

20
21 **Choyeros.** Approximately 4000 people known as *Choyeros* currently occupy the Sierra de La Giganta
22
23 (hereafter “*Giganta*”) mountain range of Baja California Sur, one of the least populated areas in one of the least
24
25 inhabited states in Mexico (Comisión Nacional de Areas Naturales Protegidas, 2020). *Choyeros* refer to themselves
26
27 by this name which roughly means people of the local cholla cactus (*Cylindropuntia cholla*) desert. With the
28
29 *Giganta* covering approximately 7,400 square kilometers (Leon de La Luz et al. 2008), Choyero population density
30
31 is about 1 person per 2 square kilometers. These *Choyeros* reside on working ranches distributed over the many
32
33 drainages found throughout the *Giganta* and most are descended through consanguineal and/or affinal ties from the
34
35 original Euro-American families who colonized the region over the last 300 years (Koster et al., 2019; Macfarlan et
36
37 al., 2020). *Choyeros* share culture history with other ranching populations found throughout the peninsula and have
38
39 been referred to as *rancheros*, *Californios*, or *Oasiana-Rancheros* in the academic literature (Crosby, 2015; Cariño
40
41 Olvera, 2014).
42

43
44 **Biogeographical setting.** Biogeographically, the *Giganta* is characterized as Sonoran Desert (Shreve &
45
46 Wiggins, 1964) with sacrocaulescent scrubland vegetation. It has a hot arid climate with most of its rainfall
47
48 generated by tropical storms and hurricanes occurring during the late summer and early fall (August-October)
49
50 (INEGI, 2016). The *Giganta* is Baja California Sur’s largest mountain range, spanning approximately 150 km along
51
52 a NW-SE axis (León De La Luz & Coria Benet, 2018). The mountain range is situated in a geological complex
53
54 known as the Comondú Formation, which is composed of clastic sandstones, volcanic rock, and conglomerates.
55
56 Although the range is situated more closely to the Gulf of California along its precipitous eastern escarpment, the
57
58 majority of its surface slopes towards the Magdalena plain in the west, producing a number of intermittent-stream
59
60 drainages that terminate at the Pacific Ocean. It is along these drainages that a variety of perennial wetlands emerge,
61

1
2
3
4 either as perched water tables, springs or hanging gardens (Maya et al., 1997; Ruiz-Campos et al., 2014). These
5
6 wetlands are species rich and diverse (León De La Luz & Domínguez Cadena, 2006; Riemann & Ezcurra, 2005;
7
8 Ruiz-Campos et al., 2002). The wetlands act as waypoints for resident and migratory species (Erickson et al., 2008;
9
10 Rodríguez-Estrella, 2005), provide forage and pasture for livestock, and have been making human life possible in
11
12 this harsh and unforgiving landscape for at least the last 4000 years (Henrickson, 2014; Macfarlan & Henrickson,
13
14 2010).

15
16 **Cultural adaptation to arid land pastoralism in the Giganta.** *Choyeros* have accommodated the
17
18 *Giganta*'s hot arid climate and sparse rainfall over the last three centuries by culturally adapting to the perennial
19
20 springs scattered along mountain drainages (Cariño Olvera, 2001; CONANP, 2014). Any spring able to water a
21
22 suitable home site creates potential for a *Choyero* ranch. The population maintains many of the productive traditions
23
24 that were first brought to the region during the Jesuit mission era by the colonists, such as leather working, metal
25
26 smithing, heritage crop production, and ranching (de Grenade & Nabhan, 2013a, 2013b; Nabhan et al., 2010).
27
28 Historically and still today, *Choyeros* remain relatively isolated from the larger Mexican society: they are outside the
29
30 reach of market places, paved roads, mail service, electrical grid, cell-phone networks, potable water systems,
31
32 sewage, and waste disposal systems (Cariño Olvera, 2001; CONANP, 2014; Crosby, 2015).

33
34
35 The primary subsistence mode is goat and cattle ranching, with an emphasis on meat and cheese production
36
37 for export by roaming merchants to local and regional markets. The average ranch has four or more people, mostly
38
39 related as kin extended across multi-generational families. Households may also maintain sheep, pigs, turkeys,
40
41 chickens, and rabbits for domestic consumption, as well as horses, donkeys, and mules for transportation.
42
43 Households supplement their diet through the governmental sponsored food program CONASUPO, as well as with
44
45 food purchased from urban markets found in Las Pocitas, Ciudad Constitución, and La Paz. Near their homes, most
46
47 ranches maintain *huertas* and/or *jardines*. *Huertas* are typically tended by adult male and female family members
48
49 and provide edible foods for the household, as well as feed for domestic livestock (e.g. alfalfa, sorghum). Beyond
50
51 their collaboration in the *huertas*, *Choyeros* have sex divided labor domains and ecologies that affect the genders'
52
53 differential exposure to plants (encounters of which vary by ecology). *Jardines* are most often tended by female
54
55 heads of household and serve important household functions such as providing shade, decorative flowers, culinary
56
57 herbs, and medicine (e.g. for herbal infusion into teas). When not tending to livestock near the ranch, men are most
58
59
60
61
62
63
64
65

1
2
3
4 often tending to livestock in the *campo* or the open-range desert wilderness of the *Giganta*. While *Choyero* children
5
6 can assist parents with household chores, they do not engage in substantial ranch work outside of the house.
7

8 Because they are places where fertile soils and fresh water come together, both *huertas* and *jardines* serve
9
10 as attractive oases for a variety of plant and animal species. Many *huertas* that were quite important for ranch-level
11
12 subsistence throughout Baja California's mountains a few generations ago are now less productive, as ranchers have
13
14 increasingly focused on milk, cheese, and meat production for sale to the market (Cariño Olvera, 2001; Crosby,
15
16 2015; de Grenade & Nabhan, 2013a). Aside from market sales and occasional itinerant wage labor, *Choyeros* remain
17
18 relatively unintegrated into the larger market economy and subsist on yearly incomes that lie below the poverty line
19
20 (CONAPO, 2010). According to government statistics, *Choyero* ranches are classified as "*muy bajo*" - the indicator
21
22 of being in Mexico's lowest (poorest and most underdeveloped) socio-economic stratum (SEDESOL, 2010).
23

24 Open range lands like those used by *Choyeros* are distinguished by their rugged landscapes and the vast
25
26 distances traversed, due to the far-reaching mobility of livestock (Huntsinger, 2016). Cattle and goats often travel
27
28 long distances (>10 miles a day) between elevations, to water and food, and away from harsh weather or to avoid
29
30 predators. Cattle and goats are able to make use of a wide variety of rangeland plants of low nutritional quality and
31
32 that may present dangers to other animals and humans. Because food and water are scarce in the arid ranges of the
33
34 *Giganta*, the ability for livestock to find scarce resources is key to the rancher's success. The movement of livestock
35
36 is also sometimes under the control of the herder (who depends on wayfinding knowledge to traverse the vast
37
38 landscape), such as when rounded up and brought back to an enclosure or when moved to greener pasture after a
39
40 rain. As such, pastoralists who depend on livestock in harsh environments have developed culturally specific
41
42 subsistence traditions and bodies of knowledge that allow them to both support and control the grazing and
43
44 reproduction of livestock in otherwise unproductive landscapes (Fernandez-Gimenez & Febre, 2006). For example,
45
46 *Choyeros* make use of horses, donkeys, and mules that facilitate human travel over these ranges whether ridden or
47
48 loaded with packs, and herding dogs - that when well-trained can effectively herd livestock without the guidance or
49
50 instructions of human herders. *Choyeros* have a keen sense of when and where resources can be found the
51
52 landscape, and when hazards need to be avoided (e.g. high predation rates by coyote and mountain lions during the
53
54 dry season, flood and storm risks during the summer monsoonal season). Currently, no information exists
55
56 concerning *Choyero* ecological knowledge variation.
57
58
59
60
61
62
63
64
65

1
2
3
4 ***Choyero ethnobiological knowledge and school education.*** To meet the challenges of becoming
5
6 competent off-grid ranchers in the Giganta, *Choyeros* acquire, develop, maintain, and transmit a rich body of
7
8 knowledge about the local ecology associated with springs, gardens, and wild rangeland. This includes knowledge
9
10 over 1) predators (e.g. coyotes, pumas), 2) vectors for infectious disease (e.g. racoons, skunks), 3) venomous
11
12 animals (e.g. rattlesnakes, scorpions), 4) poisonous and dangerous plants (e.g., arrowpoison plant, cacti), 5) wild and
13
14 domesticated plants and animals eaten by humans or livestock and 6) plants and animals for construction, shade,
15
16 cooking fuel, medicine, ritual, and artisanal crafts. *Choyero* knowledge may also be supplemented by or
17
18 complemented by education and social exposure gained with schooling.
19

20
21 Educational achievement has been on the rise for the last several decades (INEGI, 2004). Nevertheless, due
22
23 to economic hardship, costs of transportation and school supplies prevent some *Choyeros* from maximal exposure to
24
25 education. Families who wish their children to pursue a formal education must send them to public schools for
26
27 primary, middle-school and high school education. Due to the challenge of transporting children from remote
28
29 ranches to schoolhouses on a daily basis, a unique feature of the educational system in Baja California Sur is the
30
31 placement of schools, cafeterias, dormitories, teachers, and social welfare officers in rural locations through the
32
33 *Coordinación Estatal de Albergues Escolares*. Most children attending school are brought to the school for five days
34
35 and four overnights a week and then return home every weekend. The *albergues escolares* program established rural
36
37 schools in La Soledad in 1969 and in Santa Maria de Toris in 1980.

38
39 Local history and culture of Baja California Sur is neglected in the school curriculum, which is taught by
40
41 non-local teachers and features the history and culture of mainland Mexico. Knowledge about the local ecology and
42
43 about subsistence skills is also not gained directly from a school education. As such, like many of the ranching
44
45 cultures of Baja California Sur, *Choyeros* are expected to acquire most of their ethnobiological education informally
46
47 in the social and work contexts of ranch life. This kind of practical ranching knowledge is typically learned from
48
49 same-sex peers and adults (Cariño Olvera, 2001; Crosby, 2015).

50
51 While ethnobiological information is not taught in the schools, there are reasons that educational
52
53 achievement and ethnobiological knowledge might be related. One possibility is that formal education and
54
55 experience of nature and subsistence activities are mutually exclusive opportunities for children, such that those who
56
57 choose to develop their knowledge through scholastic studies do so at the expense of developing their
58
59 ethnobiological knowledge (Nabhan & St Antoine, 1993; Zent, 2001). However, a mix of both kinds of learning
60
61
62
63
64
65

1
2
3
4 should be possible for *Choyero* children because formal education is coming from non-*Choyeros* in a schoolhouse
5 setting, while ethnobiological education is likely coming from a combination of older kin at home on the ranch and
6 near aged peers (who may be encountered at school). As adolescents and young adults, *Choyeros* might also learn
7 ethnobiological information from *Choyeros* living on other ranches, especially neighbors nearby whom they might
8 see more often. As such, a positive correlation between school education and ethnobiological knowledge may exist
9 if more capable individuals self-select into both achieving more years of education and acquiring more
10 ethnobiological knowledge from others.

11 **Choyero demographics and fertility**

12 Based on community censuses, we implemented interviews and demographic surveys with 143 heads of
13 household from across 70 ranches in four communities across five field seasons (2014-2018). We used information
14 from these interviews and surveys to determine characteristics of *Choyero* demographics, fertility schedules, and
15 childrearing demands on parents.

16 **Fertility schedules.** Based on a sample of 46 women across 4 communities (born between 1930 and 2000),
17 the mean female age of first reproduction among *Choyeros* is 22.6 (SD 5). On average, *Choyero* men become fathers
18 by the age of 27 (the average husband is 4.4 (SD 7) years older than his wife). The average fertility schedule of a
19 *Choyero* man is tied to that of his wife, with an interbirth interval (IBI) of 3.9 years and a total fertility rate (TFR) of
20 3.7 surviving offspring. As such, *Choyero* fertility schedules extend roughly from age 16 to about age 46. See Table
21 1 for additional details about *Choyero* fertility.

22 <Insert Table 1 about here>

23 **Childrearing demands on parents.** We estimated the parent ages at which dependent offspring impose
24 peak dependency (for food and essential information) on parents based on a cumulative model of relative costs from
25 generalizable features of human nutritional demand schedules across the pre-reproductive life stage (per Institute of
26 Medicine, 2002) and estimated fertility schedules based on the *Choyero* IBIs and TFR reported in Table 1.
27 According to this model, the average *Choyero* mother aged 40-45 (and her husband of expected age 44.4-49.4) is at
28 peak dependency of 3.7 expected offspring younger than age 20 (two of them adolescents and one a very young
29 adult).

30 Among *Choyeros*, offspring that remain at natal ranches maintain net dependency (consuming more than
31 they produce) until they marry and start living more independently - typically after age 20. *Choyero* last-borns will

1
2
3
4 often remain dependent until their mother is in her sixties. Young *Choyero* parents tend to live in close proximity to
5
6 and are affiliated with older kin from whom they receive support (Crosby, 2015)
7

8 Consistent with the estimates above for parent ages at peak offspring dependency, our sampling at the
9
10 household level from across 70 ranches indicates that the number of co-resident dependent offspring (around an
11
12 average of 2) relying on a head of household is greatest for heads of household in the 35-45 year-old age range (see
13
14 Figure 2). The number of dependent offspring observed for heads of households of a given age is always less than
15
16 the number of expected offspring because a large proportion of our sample included adults with no coresident
17
18 dependents.
19

20 <Insert Figure 2 about here>
21

22 **Choyero ethnobotanical lexicon**

23
24 In 2015, two authors (SJM & JJG) asked 46 heads of households from 28 ranches (n-males=29; n-
25
26 females=17; Mean Age=52 years) within one community (Santa Maria de Toris) to free-list plants and animals they
27
28 associated with *Giganta* spring ecosystems to develop an initial lexicon for discussing ecosystem knowledge. This
29
30 free-list task identified an initial set of 218 animals and 201 plants. Following the free-list task, the researchers,
31
32 along with a focus group of 10 adults (seven males, three females; Mean Age=43 years; Min/Max Age=21/74 years)
33
34 from four ranches, linked the emic plant and animal terms with the Linnaean classification system (i.e. Family,
35
36 genus, and species) using photos and descriptions from botanical and faunal field guides (i.e., Grismer, 2002;
37
38 Howell & Webb, 1995; Kays & Wilson, 2009; León De La Luz & Coria Benet, 2018; Rebman et al., 2012; Rodd et
39
40 al., 2007; Wiggins, 1980). Furthermore, this focus group clarified that certain plants and animals were referred to by
41
42 multiple emic terms. Based on this information, we obtained from this focus group, using a process of consensus,
43
44 the term that was most likely to be used amongst the ranching community. We noted and consolidated these terms.
45
46 Occasionally, a plant description could not be linked to a field guidebook photo. As such, we asked the focus group
47
48 to find examples of the plant for us to photograph. These photos were then sent to botanical experts at the Red Butte
49
50 Garden at the University of Utah, the San Diego Natural History Museum, and from the Departamento de Biología
51
52 at the Universidad Autónoma de Baja California, who assisted with linking the photo to its associated Linnaean
53
54 classification. Three plant species could not be reconciled, resulting in a final set of 162 plant and 125 animal
55
56 species. Of these 287 species, 72 plants and 50 animals were included in our ethnobiological assessment task
57
58 (described below).
59
60
61
62
63
64
65

1
2
3
4 **Plant and animal categorizations**
5

6 In May 2016, two key informants (1 male and 1 female) suggested to two authors (SJM and JJG) that the
7 consolidated list of 162 plants mentioned above could be categorized according to where the plants are found:
8 *jardines*, *huertas*, or in the *campo*. Using a process of consensus, the two key informants provided their assessments
9 (45 *jardin* plants; 18 *huerta* plants; 99 *campo* plants).
10
11

12
13
14 In June 2018, four authors (SJM, ES, JJG, and DGB) employed surveys about plant and animal uses with
15 heads of households from 38 ranches from three communities (Santa Maria de Toris, La Higuera, & San Pedro de
16 La Presa), as part of our larger project on *Choyero* ethnobiological knowledge. In these surveys we asked
17 participants to enumerate plants and animals which conform to a number of use categories, two of which we report
18 on here (“human plant comestibles”(food), “dangerous plants and animals”(danger)). For our purposes, a plant item
19 was considered “food” if a single respondent identified it as such and provided its preparation for consumption.
20 Respondents provided 210 unique plant concepts, 60 of which were identified as human food. We then linked these
21 reports of food uses to our initial plant database. Of the 87 plants shown in our ethnobiological assessment task
22 (described below), 27 were categorized as “food” (see Appendix A). Of the 50 animals shown in the ethnobiological
23 assessment task, we also categorized five (i.e., three species of birds, a jack rabbit, and a fish) as “food”, based on
24 reports and evidence that these were eaten. Among a subset of these ranches (n-ranches=3) we also asked
25 participants to enumerate dangerous plants and animals. Respondents provided examples of 10 dangerous plants and
26 5 dangerous animals along with common reasons for their danger: poisonous sap, sharp thorns, dangerous and
27 potentially disease transmitting bite. Based on these dangerous features we categorized 28 items as dangerous: 7
28 cacti, 7 shrubs and trees with thorns or toxicity, 14 predatory or disease transmitting animals (see Appendix A).
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **Ethnobiological knowledge assessment task**
46

47 In 2017 and 2018, seventy-one individuals (33 females and 38 males) aged 5 to 86 years old from 40
48 ranches across three communities (Santa Maria de Toris: n = 55; La Higuera: n = 5; and La Soledad: n = 11) were
49 presented an ethnobiological knowledge assessment task (see Table 2 for sample characteristics). As a first step of
50 the task, individuals took a vision acuity test using “tumbling E” eye charts presented on a laptop computer, a robust
51 and easy to use diagnostic tool that is practical for populations with innumerate or analphabetic participants
52 (Messina & Evans, 2006). From this test we derived a visual acuity score (ranging from 1 to 11). A higher score
53 indicates better visual acuity. Visual acuity across the sample was 10.6 on average (SD=0.77, min=6 max=11) on
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 the 1-11 scale (Table 2). The ethnobiological knowledge assessment task presented a sequence of 137 slides on a
5
6 laptop computer featuring images of 87 plants and 50 animals (Appendix A). The set of plants and animals used in
7
8 our ethnobiological knowledge assessment task was informed by the ethnobiological research methods described
9
10 above and selected based on the availability of clear and representative photographs.

11
12 <Insert Table 2 about here>
13

14 The order of items presented was varied with about half of participants seeing items in reversed order.
15
16 Assessments of ethnobiological knowledge were conducted in Spanish in the privacy of individual homes and
17
18 occurred as part of a larger set of household interviews regarding ranching demography and lifestyles. These
19
20 household interviews additionally informed us of individuals' educational achievement, their ranch affiliation, and
21
22 community membership. For each of 137 plant and animal images presented in sequence, the researcher (ES) asked
23
24 the participant whether they recognize the item in the image. If the participant answered affirmatively, the researcher
25
26 asked the follow-up question about what the item is called. Responses were recorded and coded "correctly named" if
27
28 matching any of the locally appropriate culturally correct names used for species identification (see Macfarlan et al.,
29
30 2020).
31

32 **Analytic models**

33
34 Our analytic models are performed using STATA/IC 16.1 (Rabe-Hesketh & Skrondal, 2012). We
35
36 investigate ethnobiological knowledge with two-stage logistic regression models. In the first stage of these models
37
38 we predict an individual's ability to recognize R_i a plant or animal item shown to them (R_i equals 1 if item i is
39
40 recognized and zero otherwise). In the second stage, we predict an individual's ability to correctly name C_i a
41
42 recognized plant or animal item (C_i equals 1 if item i is named correctly and zero otherwise).
43
44

45 Multiple individuals may reside at the same ranch, so we nest the outcome variable at both the level of the
46
47 individual and the ranch. As such, we employ multi-level logistic regressions, whereby we use an exchangeable
48
49 variance-covariance structure at the level of the individual and ranch. Furthermore, we employ robust standard
50
51 errors for all fixed effects in all models. We assess 6 classes of models of ethnobiological knowledge: 1) a model
52
53 that examines knowledge across all life stages; 2) a model that examines knowledge of food and or dangerous items
54
55 (relative to all other investigated items) across all life stages; 3) a model that exams sex differences in knowledge of
56
57 *jardin* plants only; 4) a model that exams sex differences in knowledge of *campo* plants only; 5) a model of sex
58
59 differences in *jardin* plant knowledge for post-reproductive adults only; 6) a model of sex differences in *campo* plant
60
61

1
2
3
4 knowledge for post-reproductive adults only. For the age ranges of *pre-reproductive* (< 23 years), *reproductive* (23
5 to 45 years), and *post-reproductive* (> 45 years) life stages demarcated in or bounding the models, we used the
6 observations of *Choyero* life stage tendencies detailed in the **Fertility schedules** subsection above.
7
8

9
10 Where appropriate for each class of model, we specify the results of the following co-variates: age (in
11 years), age squared, sex (1=male, 0=female), visual acuity or *vision* (scored 1 to 11), educational achievement or
12 *education* (in years), and whether the item is potential food and/or danger or not (1=food or danger; 0=Other). All
13 models presented in this study can be replicated using the associated STATA DO and Data files provided as
14 electronic supplementary materials.
15
16
17
18

19 20 **Results**

21
22 Original data of this study are available at Mendeley Data (Schniter et al., 2020). Although the scope of our
23 research is to understand how factors associated with specific life stage moments affect ethnobiological knowledge,
24 we first consider the variables generated by our ethnobiological knowledge assessment task. Across all ages, most
25 participants recognized the majority of plant ($M = 71.0\%$) and animal ($M = 85.9\%$) items presented, but fewer were
26 able to correctly name those plant ($M = 42.6\%$) and animal ($M = 59.0\%$) items according to local nomenclature.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

99 100 **What explains knowledge across all life stages?**

101
102 We fit a two-stage model of ethnobiological knowledge across all life stages. This allows our results to be
103 compared to previous work on the topic (e.g., Koster et al., 2016) and portrays the overall development of
104 knowledge across the life course within our sample. We find that reported recognition of plant and animal species
105 takes a quadratic, concave shape over the life course (Age: Odds Ratio = 1.1; $p < .001$; Age Squared: Odds Ratio =
106 0.99; $p < .001$) and that better visual acuity is associated with improved ability to recognize (Odds Ratio = 1.50; $p <$
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2
3
4 .001) (Table 3 Model (1)). Educational achievement has no effect on recognition (Odds Ratio = 1.02; $p = .41$) and
5
6 the difference between males' and females' recognition is not significant (Odds Ratio = 1.17; $p = .19$) (Table 3
7
8 Model (1)). We find that the ability to correctly name recognized species takes a quadratic, concave shape over the
9
10 life course (Age: Odds Ratio = 1.1; $p < .001$; Age Squared: Odds Ratio = 0.99; $p < .001$) and that better visual acuity
11
12 is associated with improved ability to correctly name (Odds Ratio = 1.20; $p = .032$) (Table 3 Model (1), Figure 3).
13
14 Educational achievement has no effect on correct naming (Odds Ratio = 1.00; $p = .87$) and males correctly name
15
16 recognized species more than females (Odds Ratio = 1.20; $p = .03$) (Table 3 Model (1)).
17

18 <Insert Table 3 about here>

19
20 <Insert Figure 3 about here>

21 22 **What explains ethnobiological knowledge in the pre-reproductive life stage?**

23
24 To evaluate whether the rates of ethnobiological knowledge development in pre-reproductive individuals
25
26 are continuously increasing yet different for species that are potentially dangerous and food (compared to those that
27
28 are not), we employ a two stage model of ethnobiological knowledge across all life stages. Despite predicted effects
29
30 in our models, we observed no variation in vision for participants in the pre-reproductive life stage. We find that
31
32 recognition of plant and animal species continuously increases during the pre-reproductive life stage (Age: Odds
33
34 Ratio = 1.11; $p < .001$; Age Squared: Odds Ratio = 0.99; $p < .001$) and that recognition of dangerous and/or food
35
36 species increases at more than twice the rate than for non-dangerous and non-food species (Odds Ratio = 2.05, $p <$
37
38 $.001$) (Table 3 Model (2)).
39

40
41 We also find that the ability to correctly name recognized species continuously increases during the pre-
42
43 reproductive life stage (Age: Odds Ratio = 1.11; $p < .001$; Age Squared: Odds Ratio = 0.99; $p < .001$) and that
44
45 correct naming of dangerous and/or food species increases by 80 percent compared to non-dangerous and non-food
46
47 species (Odds Ratio = 1.80, $p < .001$) (Table 3 Model (2), Figure 4). On average, knowledge of dangerous and/or
48
49 food species is greater during the pre-reproductive life stage, with 42.65% of these species correctly named
50
51 compared to the 20.78% of other species. While rates of this knowledge acquisition change across the lifespan, the
52
53 greater knowledge of food and dangerous species than for other species studied is evident across all life stages
54
55 (Appendix B, Figure 4).
56

57 <Insert Figure 4 about here>

58 59 **What explains ethnobiological knowledge in the reproductive life stage?**

1
2
3
4 Results of our estimated family budgets suggest that parents face growing offspring needs at least until the
5
6 end of the reproductive lifestage when offspring demands may be highest, and that those demands fade but can
7
8 continue for decades after, stretching into parents' post-reproductive years. To evaluate whether the rates of
9
10 ethnobiological knowledge development in reproductive adults are continuously increasing as parents continue to
11
12 meet their offspring's burgeoning needs, we first focus on trends in the reproductive lifestage revealed by our
13
14 models of ethnobiological knowledge development across the lifespan (Table 3 Models (1) and (2)). Despite
15
16 predicted effects in our models, we observed no variation in vision for participants in the reproductive life stage.
17
18 These models both reveal that correct naming of plant and animal species continuously increases with younger adult
19
20 ages such that the highest levels of correctly named species occur at the end of the reproductive life stage (Figures 3
21
22 and 4), when offspring demands are highest.

23
24 During the reproductive life stage, females correctly name 61.4% of *jardin* plants compared to males who
25
26 correctly name 44.6%, while males correctly name 49.4% of *campo* plants compared to females who correctly name
27
28 36.0% (Appendix B). To assess the role of age and gender on *jardin* and *campo* plant knowledge in the reproductive
29
30 lifespan we use two multi-level two-stage logistic regression models. For brevity, we report only results of the
31
32 second stage of these models (3) and (4) in text, concerning the ability to correctly name recognized *jardin* plants
33
34 and *campo* plants, respectively (see Table 4 for complete model results). Our first model shows that males correctly
35
36 name substantially fewer *jardin* plants than females do (Odds Ratio = 0.55; $p < .001$) (Table 4, Model (3)). Our
37
38 second model shows that males correctly name more *campo* plants than females do (Odds Ratio = 1.5; $< .001$)
39
40 (Table 4, Model (4)). These models show that, overall and relative to their opposite gender, being a reproductive
41
42 aged female improves the probability of knowing plants associated with house gardens by 45%, while being a
43
44 reproductive aged male improves the probability of knowing plants associated with herding activities by 50%
45
46 (Figures 5 and 6).

47
48 <Insert Table 4 about here>

49
50 <Insert Figures 5 and 6 about here>

51 52 53 **What explains ethnobiological knowledge in the post-reproductive life stage?**

54
55 Post-reproductive adults are most representative of the top decile of ethnobiological knowledge in our
56
57 sample, occupying seven out of eight of the highest ranks for recognizing and correctly naming plants and animals.
58
59 The ages of these most expert adults in the top decile ranged from 46 to 63 with a mean age of 53. On average, post-
60
61

1
2
3
4 reproductive adults correctly name 49.0% of plants and animals, while reproductive adults correctly name 52.7%
5
6 (Appendix B).

7
8 During the post-reproductive life stage, females correctly name 50.0% of *jardin* plants compared to males
9
10 who correctly name 35.4%, while males correctly name 44.3% of *campo* plants compared to females who correctly
11
12 name 35.5% (Appendix B). Compared to earlier lifestages in our sample that have no variation in visual acuity, post
13
14 reproductive adults show the greatest variation in visual acuity (min=6, max=11).

15
16 To evaluate whether the rates of ethnobiological knowledge in post-reproductive adults are predicted by
17
18 visual acuity, age, sex, and gender-associated economic domains, we employ a pair of multi-level two-stage logistic
19
20 regression models that consider only adults in the post-reproductive lifespan (Table 5 Models (5) and (6)). These
21
22 models examine male and female knowledge of *jardin* and *campo* plants. For brevity, we report only results of the
23
24 second stage of these models (5) and (6) in text, concerning male and female abilities to correctly name recognized
25
26 *jardin* plants and *campo* plants (see Table 5 for complete model results).

27
28 We find that sex (Odds Ratio = 0.53; $p = .001$) is a significant predictor of correctly naming *jardin* plants,
29
30 even after controlling for age (Odds Ratio = 0.96; $p = .01$) and visual acuity (Odds Ratio = 1.46; $p = .023$), with
31
32 post-reproductive females able to name more house garden plants than post-reproductive males (Table 5 Model (5)).
33
34 Among post-reproductive adults, we do not find an effect of sex (Odds Ratio = 1.27; $p = .09$) on correctly naming
35
36 *campo* plants, after controlling for age (Odds Ratio = 0.97; $p = .01$) and visual acuity (Odds Ratio = 1.38; $p = .01$)
37
38 (Table 5 Model (6)). While the average proportion of *campo* plants correctly named by women does not change
39
40 between reproductive and post-reproductive life stages, the portion of *campo* plants correctly named by males
41
42 declines, with post-reproductive aged adults naming 10.3% fewer *campo* plants than younger reproductive aged
43
44 adults (calculated from life stage proportions reported in Appendix B).

45
46
47 <Insert Table 5 about here>

48 49 Discussion

50
51 We investigated whether age profiles of *Choyero* ethnobiological knowledge development are consistent
52
53 with predictions derived from life history theory about the timing of productivity and reproduction. We see evidence
54
55 of age-appropriate complementary knowledge profiles developing for *Choyero* men and women as they experience
56
57 changes in embodied capital and needs of dependent offspring. We find that while individuals acquire knowledge of
58
59 most dangerous items and edible resources by early adulthood, knowledge of plants and animals relevant to the age
60
61

1
2
3
4 and sex divided labor domains and ecologies (e.g., women’s house gardens, men’s herding activities in the
5 wilderness) develops into middle adulthood but to different degrees and at different rates for women and men.
6
7 Across our models, we see peak predicted ethnobiological knowledge for men and women aged 5-60, near the
8
9 ages at which parents experience the greatest demand from dependent offspring, bearing support for a “wisdom of
10
11 adults” view of ethnobiological knowledge development.
12
13

14 A widespread “wisdom of elders” view has been that highest levels of ethnobiological knowledge should
15
16 be found among post-reproductive older adults generally (Koster et al., 2016). When we consider the top decile of
17
18 most knowledge *Choyeros* sampled, we find that almost all of these experts are post-reproductive aged adults
19
20 ranging in age from 46 to 63 – bearing support for a nuanced “wisdom of elders” view of ethnobiological knowledge
21
22 development. On the other hand, average rates of recognition and correct naming for reproductive adults are higher
23
24 than for post-reproductive adults, challenging a “wisdom of elders” view. Our study resolves this apparent paradox
25
26 by showing the effects of vision on ethnobiological knowledge, and the possible effects of reduced activity by males
27
28 in the *campo* on their knowledge of *campo* plants (contributing to ethnobiological knowledge). The effects of
29
30 sensory acuity (vision and hearing) on intelligence among older adults are large; after controlling for age and
31
32 hearing, Lindenberger and Baltes (1994) found that vision accounts for 43% of the variance in intelligence among a
33
34 sample of older adults. Additional cognitive and physiological measures affecting men and women’s activities in
35
36 their economic domains might explain additional variance in ethnobiological knowledge and should be considered
37
38 for future research on older adult knowledge. If individuals are unable (e.g. due to vision limitations and perhaps
39
40 other limitations like difficulty hearing or immobility) to effectively recognize and talk about specific plant or
41
42 animal examples being referred to with a visual examples, they are severely incapacitated in their ability to reliably
43
44 transmit knowledge about those things. While these age-associated limitations may prevent many older adults from
45
46 reliably transmitting ethnobiological knowledge, a small subset of older adults appear to be differentially advantaged
47
48 with better abilities to function as ethnobiological experts. Experts can be of great cultural value and of great value
49
50 with conservation efforts, as we discuss below.
51
52

53 Conservation biologists (e.g., León de la Luz & Domínguez Cadena, 2006; Riemann & Ezcurra, 2005;
54
55 Rodríguez-Estrella, 2005) and non-governmental agencies (e.g. World Wildlife Fund) have implicated rancher
56
57 lifestyles as primary vectors for *Giganta* ecosystem decline, suggesting ignorance of the local ecology, but those
58
59 conclusions are typically drawn from non-representative samples in areas of relatively high population densities or
60
61
62
63
64
65

1
2
3
4 vehicular traffic that tend to be more depleted of biodiversity. Elsewhere, in natural protected areas of Baja
5
6 California Sur, conservation biologists have suggested that local ranchers' environmental traditional knowledge is of
7
8 value for developing conservation, sustainable use, and local economic development policies (Pío-León et al., 2017).
9
10 Our own partnerships leveraging local *Choyeros* ethnobiological has led to novel findings regarding species range
11
12 extensions (MacFarlan et al., 2019). Our research shows that, as a group, *Choyero* ranchers embody a great
13
14 distributed body of ethnobiological knowledge: they generated for us a vast inventory of 287 known plants and
15
16 animals and provided detail about various names used to refer to those species and species traits and uses that are
17
18 important to their way of life. While there is substantial variability in ecological knowledge among *Choyero*
19
20 ranchers, most are knowledgeable of most items in our ethnobiological knowledge assessment tool by adulthood –
21
22 meaning they have, at minimum, the necessary visual recognition ability (to trigger memory of having seen the plant
23
24 or animal item before) and semantic memory (to recall the appropriate lexicon by which people refer to the item
25
26 specifically) to indicate knowledge of the species. Furthermore, a few experts are highly knowledgeable of almost
27
28 all items in our tool's set. In aggregate, our ethnobiological lexicon results, species classification results, and the
29
30 analytic results of models predicting knowledge of species included in the ethnobiological knowledge assessment
31
32 task are important evidence of the rich ethnobiological knowledge that *Choyeros* maintain. As primary stakeholders
33
34 whose economic and cultural livelihoods have been tethered to arid land springs over the last 300 years, it is
35
36 possible that *Choyero* knowledge and social institutions either support or can be leveraged to improve the goal of
37
38 long-term *Giganta* ecosystem sustainability. For example, their detailed understanding of the biota and its natural
39
40 occurrence can facilitate cost-effective biological inventory programs seeking to identifying species presence-
41
42 absence, abundance, distributions, range extensions, or introductions (Berkes et al., 2000).
43
44

45 That reproductively and post-reproductively aged *Choyero* women and men maintain separate but
46
47 complementary ethnobiological knowledge fits both the literature on the evolution of the sexual division of labor in
48
49 humans generally (e.g. Gurven et al., 2012; Macfarlan, 2016), as well as the human ecology of the *Giganta*,
50
51 specifically. In the *Giganta*, ranches are typically established in valley bottoms on level ground near springs.
52
53 Because springs and flat terrain occur in a sporadic fashion throughout this region, ranches can be separated by large
54
55 distances from one another. This has caused households to become highly independent (Crosby 2015). In response,
56
57 women and men have developed distinct divisions of labor and associated knowledge to support autarkic household
58
59 functioning. Although this produces strongly gendered economic domains, there is a complementarity to gender
60
61
62
63
64
65

1
2
3
4 dynamics within *Choyero* communities, whereby people explicitly recognize the value of each gender's role in
5 household economics. Among *Choyeros*, men's greater knowledge of *campo* plants compliments women's greater
6 knowledge of *jardin* plants during the reproductive life stage, and while women retain their relative advantage in
7 *jardin* knowledge into post-reproductive years, men lose a relative advantage in *campo* knowledge – possibly
8 associated with declining exposure to species in the campo with advancing post-reproductive years.
9

14 **Limitations and future prospects**

16 We discuss limitations to our study method and results. Our methods to capture a portrait of
17 ethnobiological knowledge development across the life course are based on cross-sectional sample design, rather
18 than longitudinal study. Cross-sectional sample design may not correctly reveal the path of knowledge development
19 if secular changes might differentially affect the life stage cohorts (e.g. educational opportunity has improved since
20 at least 1980). We controlled for possible effects of educational achievement on ethnobiological knowledge across
21 our full sample and found no effect. Our analyses also reveal that substantial differences in ethnobiological
22 knowledge exists at the ranch level, however ranches have also changed over the past century. Future research on
23 pastoralist ethnobiological knowledge should carefully investigate whether ranch-level differences may be
24 associated with economic differentiation between households, distance from neighbors, and different family
25 histories in the *Giganta*.
26
27
28
29
30
31
32
33
34
35

36 Our set of items presented for visual identification may not provide the most appropriate representations of
37 the actual items and may not be adequately characterizing the breadth of species knowledge essential for ranching.
38 The color photographs we presented of items were assembled from images opportunistically gathered from a variety
39 of personal and publicly available plant photo archives. An alternative approach for visual identification is to present
40 participants controlled transects or a collection of specimens (both of which present practical challenges for
41 permitting, and for controlling quality and reliable stability of over time) (e.g., see Begossi, 1996; Zarger & Stepp,
42 2004). Visual identification and naming tasks are commonly used methods for investigating knowledge but there are
43 others (some of which we also used) that reveal dimensions of ethnobiological knowledge including questionnaire
44 (pile sort, multiple choice, free-listing), semi-structured interview, and observation (Reyes-García et al., 2007).
45 Missing from this report on *Choyero* ethnobiology is related knowledge about insects, fungi, and knowledge about
46 agriculture, animal husbandry, weather, hydrology (e.g. of local springs), and geology.
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Last, interpretation of survey results from older adult respondents as evidence of their ethnobiological
5
6 “knowledge” necessitates careful consideration of the difficulty in lexical access that many older adults experience,
7
8 which may be distinct from their ability to see and recognize, recall familiarity with, and produce knowledge of the
9
10 species example (Craik, 1998; Grady & Craik, 2000). Word-finding failures increase with age (Burke et al., 1991),
11
12 and almost all elderly people experience increased difficulty in retrieving proper names for recognized things
13
14 (Cohen & Burke, 1993). While we rely on knowledge revealed by participants in response to questions on our
15
16 survey, we have no measure of how people view one another’s contributions or expertise. We also have no
17
18 independent measure of subsistence ability to directly correlate with knowledge as others did (Koster et al., 2016;
19
20 Reyes-García et al., 2007; Schniter et al., 2018). While we fall short of providing a comprehensive measure of
21
22 ethnobiological knowledge based on a full range of complimentary methods, we leverage a mix of ethnobiological
23
24 research methods that contributed to a detailed study of the ability to name recognized plants and animals across the
25
26 lifecourse. From this study we can draw several conclusions about characteristics of the evolved human life history.
27

28 **Conclusion**

29
30 This research represents an important contribution to the literature on ethnobiological knowledge among a
31
32 small-scale subsistence population, since pastoralism (practiced by *Choyeros*) is the subsistence mode least
33
34 represented in this literature (Reyes-García et al., 2007). Furthermore, the research highlights the ethnobiological
35
36 knowledge of an understudied culture that is feared to be in rapid decline (Crosby 2015). With this research we also
37
38 provide a new test of the embodied capital model hypothesis that the life course of ethnobiological knowledge
39
40 development is shaped by the changing risks, capabilities, and wisdom of individuals with age and experience (Del
41
42 Giudice et al., 2016), and the changing needs of the family budget (Bock, 2002a, 2002b; Crittenden et al., 2013;
43
44 Gurven & Kaplan, 2006; Schniter et al., 2015). As adults become grandparents and advance through their post-
45
46 reproductive years, a number of physical and sensory declines reduce the profitability of various types of economic
47
48 activities. In response to these declines, older adults are expected to shift their efforts towards low-strength yet
49
50 knowledge-intensive service niches (Schniter et al., 2018). The knowledge required to know what plants and
51
52 animals look like (to recognize them) and what recognized plants and animals are called (according to culturally
53
54 correct names) bears characteristics making it optimal for continued development over the life course: these forms
55
56 of knowledge require extensive learning and facilitate communication for teaching about features of the local
57
58
59
60
61
62
63
64
65

1
2
3
4 ecology, supporting the childrearing, subsistence goals, and conservation efforts distributed within and between
5
6 multi-generational familial groups.
7
8

9 **Declarations**

10
11 Funding: This research project was supported by National Geographic Society (Research & Exploration
12 Award# HJ-099R-17), National Science Foundation (IBSS-L Award# 1743019), Funding Incentive Seed Grant
13 (University of Utah), Center for Latin American Studies (University of Utah), Society, Water, & Climate Seed Grant
14 (University of Utah), and NEXUS Pilot Grant (University of Utah), Economic Science Institute (Chapman
15 University), Division of Anthropology (California State University Fullerton).
16
17

18 Conflicts of interest/Competing interests: We have no conflicts of interest to declare.
19

20 Availability of data and material (data transparency): Raw data for this study is available at *Mendeley Data*
21 <https://doi.org/10.17632/hgd8zdyf2c.1>
22

23 Ethics approval: All procedures were in accordance with the ethical standards of the Declaration of Helsinki and
24 the University of Utah Human Research Ethics Committee (approval number IRB 00083096)
25

26 Consent to participate: Informed consent was obtained from all adult individuals included in this study. Both child
27 assent and parental permission were obtained for all child participants included in this study.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- Amoss, P. T., & Harrell, S. (1981). *Other Ways of Growing Old: Anthropological Perspectives*. Stanford University Press.
- Ardila, A. (2007). Normal aging increases cognitive heterogeneity: Analysis of dispersion in WAIS-III scores across age. *Archives of Clinical Neuropsychology*, 22(8), 1003–1011. <https://doi.org/10.1016/j.acn.2007.08.004>
- Atran, S. (1990). *Cognitive Foundations of Natural History: Towards an Anthropology of Science*. Cambridge University Press.
- Ayantunde, A. A., Briejer, M., Hiernaux, P., Udo, H. M. J., & Tabo, R. (2008). Botanical Knowledge and its Differentiation by Age, Gender and Ethnicity in Southwestern Niger. *Human Ecology*, 36(6), 881–889. <https://doi.org/10.1007/s10745-008-9200-7>
- Barrett, H. C. (2005). Cognitive Development and the Understanding of Animal Behavior. In *Origins of the social mind: Evolutionary psychology and child development*, (pp. 438–467).
- Barrett, H. C., & Behne, T. (2005). Children’s understanding of death as the cessation of agency: A test using sleep versus death. *Cognition*, 96(2), 93–108. <https://doi.org/10.1016/j.cognition.2004.05.004>
- Barrett, H. C., & Broesch, J. (2012). Prepared social learning about dangerous animals in children. *Evolution and Human Behavior*, 33(5), 499–508. <https://doi.org/10.1016/j.evolhumbehav.2012.01.003>
- Barrett, H. C., Peterson, C. D., & Frankenhuis, W. E. (2016). Mapping the Cultural Learnability Landscape of Danger. *Child Development*, 87(3), 770–781. <https://doi.org/10.1111/cdev.12495>
- Begossi, A. (1996). Use of ecological methods in ethnobotany: Diversity indices. *Economic Botany*, 50(3), 280. <https://doi.org/10.1007/BF02907333>
- Berkes, F., Colding, J., & Folke, C. (2000). Rediscovery of Traditional Ecological Knowledge as Adaptive Management. *Ecological Applications*, 10(5), 1251–1262. [https://doi.org/10.1890/1051-0761\(2000\)010\[1251:ROTEKA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[1251:ROTEKA]2.0.CO;2)
- Berlin, B. (1992). *Ethnobiological classification: Principles of categorization of plants and animals in traditional societies*. Princeton University Press.
- Bird, R. B., & Bird, D. W. (2002). Constraints of knowing or constraints of growing? *Human Nature*, 13(2), 239–267. <https://doi.org/10.1007/s12110-002-1009-2>
- Bock, J. (2002a). Evolutionary demography and intrahousehold time allocation: School attendance and child labor among the Okavango Delta Peoples of Botswana. *American Journal of Human Biology*, 14(2), 206–221. <https://doi.org/10.1002/ajhb.10040>
- Bock, J. (2002b). Learning, life history, and productivity. *Human Nature*, 13(2), 161–197. <https://doi.org/10.1007/s12110-002-1007-4>
- Borgerhoff Mulder, M., Towner, M. C., Baldini, R., Beheim, B. A., Bowles, S., Colleran, H., Gurven, M., Kramer, K. L., Mattison, S. M., Nolin, D. A., Scelza, B. A., Schniter, E., Sear, R., Shenk, M. K., Voland, E., & Ziker, J. (2019). Differences between sons and daughters in the intergenerational transmission of wealth.

- 1
2
3
4 *Philosophical Transactions of the Royal Society B: Biological Sciences*, 374(1780), 20180076.
5 <https://doi.org/10.1098/rstb.2018.0076>
6
- 7 Brantingham, P. J. (1998). Hominid–Carnivore Coevolution and Invasion of the Predatory Guild. *Journal of*
8 *Anthropological Archaeology*, 17(4), 327–353. <https://doi.org/10.1006/jaar.1998.0326>
9
- 10 Brown, C. H. (1985). Polysemy, overt marking, and function words. *Language Sciences*, 7(2), 283–332.
11 [https://doi.org/10.1016/S0388-0001\(85\)80003-6](https://doi.org/10.1016/S0388-0001(85)80003-6)
12
- 13 Burke, D. M., & Mackay, D. G. (1997). Memory, language, and ageing. *Philosophical Transactions of the Royal*
14 *Society of London. Series B: Biological Sciences*, 352(1363), 1845–1856.
15 <https://doi.org/10.1098/rstb.1997.0170>
16
- 17 Burke, D. M., MacKay, D. G., Worthley, J. S., & Wade, E. (1991). On the tip of the tongue: What causes word
18 finding failures in young and older adults? *Journal of Memory and Language*, 30(5), 542–579.
19 [https://doi.org/10.1016/0749-596X\(91\)90026-G](https://doi.org/10.1016/0749-596X(91)90026-G)
20
- 21 Cain, M. T. (1977). The Economic Activities of Children in a Village in Bangladesh. *Population and Development*
22 *Review*, 3(3), 201–227. <https://doi.org/10.2307/1971889>
23
- 24 Carey, & Spelke, E. S. (1994). *Mapping the mind: Domain specificity in cognition and culture*. Cambridge
25 University Press.
26
- 27 Cariño Olvera, M. (2001). La oasisidad: Núcleo de la cultura sudcaliforniana. *Gaceta Ecológica*, 60, 57–68.
28
- 29 Cariño Olvera, M. (2014). Oasisidad: Identidad geográfica sudcaliforniana y expresión local de la sustentabilidad.
30 *Oasis Sudcalifornianos: Para Un Rescate de La Sustentabilidad Local*. (M. Cariño Olvera and A. Ortega,
31 Eds.), 73–106.
32
- 33 Cattell, R. B. (1963). Theory of fluid and crystallized intelligence: A critical experiment. *Journal of Educational*
34 *Psychology*, 54(1), 1–22. <https://doi.org/10.1037/h0046743>
35
- 36 Christensen, H., Mackinnon, A. J., Korten, A. E., Jorm, A. F., Henderson, A. S., Jacomb, P., & Rodgers, B. (1999).
37 An analysis of diversity in the cognitive performance of elderly community dwellers: Individual differences
38 in change scores as a function of age. *Psychology and Aging*, 14(3), 365–379. [https://doi.org/10.1037/0882-](https://doi.org/10.1037/0882-7974.14.3.365)
39 [7974.14.3.365](https://doi.org/10.1037/0882-7974.14.3.365)
40
- 41 Cohen, G., & Burke, D. M. (1993). Memory for proper names: A review. *Memory*, 1(4), 249–263.
42 <https://doi.org/10.1080/09658219308258237>
43
- 44 Comision Nacional de Areas Naturales Protegidas. (2020). *Regiones de áreas naturales protegidas de México*.
45 http://sig.conanp.gob.mx/website/pagsig/datos_anp.htm
46
- 47 CONANP. (2014). *Reserva de La Biósfera Sierras La Giganta y Guadalupe* (p. 296) SEMARNAT Secretario de
48 Medio Ambiente y Recursos Naturales.
49
- 50 CONAPO. (2010). *Índice de marginación por localidad 2010*.
51 http://www.conapo.gob.mx/es/CONAPO/Indice_de_Marginacion_por_Localidad_2010
52
- 53 Craik, F. I. M. (1998, September 28). *Memory, aging, and survey measurement*. Cognition, Aging and Self-Reports.
54 <https://doi.org/10.4324/9780203345115-12>
55
- 56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Craik, F. I. M., & Rose, N. S. (2012). Memory encoding and aging: A neurocognitive perspective. *Neuroscience &*
5 *Biobehavioral Reviews*, 36(7), 1729–1739. <https://doi.org/10.1016/j.neubiorev.2011.11.007>
6
7 Crittenden, A. N., Conklin-Brittain, N. L., Marlowe, F. W., Schoeninger, M. J., & Wrangham, R. (2009). Foraging
8 strategies and diet composition of Hadza children. *American Journal of Physical Anthropology*.
9 <https://doi.org/10.1002/ajpa.21030>
10
11 Crittenden, A. N., Conklin-Brittain, N. L., Zes, D. A., Schoeninger, M. J., & Marlowe, F. W. (2013). Juvenile
12 foraging among the Hadza: Implications for human life history. *Evolution and Human Behavior*, 34(4),
13 299–304. <https://doi.org/10.1016/j.evolhumbehav.2013.04.004>
14
15 Crosby, H. W. (2015). *Californio Portraits: Baja California's Vanishing Culture*. University of Oklahoma Press.
16
17 de Grenade, R., & Nabhan, G. P. (2013a). Baja California peninsula oases: An agro-biodiversity of isolation
18 and integration. *Applied Geography*, 41, 24–35. <https://doi.org/10.1016/j.apgeog.2013.03.008>
19
20 de Grenade, R., & Nabhan, G. P. (2013b). Agrobiodiversity in an Oasis Archipelago. *Journal of Ethnobiology*,
21 33(2), 203–236. <https://doi.org/10.2993/0278-0771-33.2.203>
22
23 Del Giudice, M., Gangestad, S. W., & Kaplan, H. S. (2016). Life history theory and evolutionary psychology. In *The*
24 *handbook of evolutionary psychology: Foundations, Vol. 1, 2nd ed* (pp. 88–114). John Wiley & Sons Inc.
25
26 Draper, P. (1976). 9 Social and Economic Constraints on Child Life among the !Kung. In R. B. Lee & I. DeVore
27 (Eds.), *Kalahari Hunter-Gatherers: Studie of !Kung San and Their Neighbors*. Harvard University Press.
28 <https://doi.org/10.4159/harvard.9780674430600.c16>
29
30
31
32 Elsner, C., & Wertz, A. E. (2019). The seeds of social learning: Infants exhibit more social looking for plants than
33 other object types. *Cognition*, 183, 244–255. <https://doi.org/10.1016/j.cognition.2018.09.016>
34
35 Erickson, R. A., Hamilton, R. A., & Mlodinow, S. G. (2008). Status review of Belding's Yellowthroat Geothlypis
36 beldingi, and implications for its conservation. *Bird Conservation International*, 18(3), 219–228.
37 <https://doi.org/10.1017/S0959270908000221>
38
39 Fernandez-Gimenez, M. E., & Febre, S. L. (2006). Mobility in pastoral systems: Dynamic flux or downward trend?
40 *International Journal of Sustainable Development & World Ecology*, 13(5), 341–362.
41 <https://doi.org/10.1080/13504500609469685>
42
43 Gallois, S., Duda, R., & Reyes-García, V. (2017). Local ecological knowledge among Baka children: A case of
44 “children's culture”? *Journal of Ethnobiology*, 37(1), 60–80.
45
46
47 Garfield, Z. H., Garfield, M. J., & Hewlett, B. S. (2016). A Cross-Cultural Analysis of Hunter-Gatherer Social
48 Learning. In H. Terashima & B. S. Hewlett (Eds.), *Social Learning and Innovation in Contemporary*
49 *Hunter-Gatherers: Evolutionary and Ethnographic Perspectives* (pp. 19–34). Springer Japan.
50 https://doi.org/10.1007/978-4-431-55997-9_2
51
52 Gaulin, S., & Hoffman, H. (1988). Evolution and development of sex differences in spatial ability. In *Human*
53 *reproductive behaviour: A Darwinian perspective* (pp. 129–152).
54
55 Grady, C. L., & Craik, F. I. (2000). Changes in memory processing with age. *Current Opinion in Neurobiology*,
56 10(2), 224–231. [https://doi.org/10.1016/S0959-4388\(00\)00073-8](https://doi.org/10.1016/S0959-4388(00)00073-8)
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Grismer, L. L. (2002). *Amphibians and Reptiles of Baja California, Including Its Pacific Islands and the Islands in*
5 *the Sea of Cortés*. University of California Press.
6
7 Guest, G. (2002). Market Integration and the Distribution of Ecological Knowledge within an Ecuadorian Fishing
8 Community. *Journal of Ecological Anthropology*, 6(1), 38–49. <https://doi.org/10.5038/2162-4593.6.1.3>
9
10 Gurven, M., Fuerstenberg, E., Trumble, B., Stieglitz, J., Beheim, B., Davis, H., & Kaplan, H. (2017). Cognitive
11 performance across the life course of Bolivian forager-farmers with limited schooling. *Developmental*
12 *Psychology*, 53(1), 160–176. <https://doi.org/10.1037/dev0000175>
13
14 Gurven, M., & Kaplan, H. (2006). Determinants of time allocation across the lifespan. *Human Nature*, 17(1), 1–49.
15 <https://doi.org/10.1007/s12110-006-1019-6>
16
17 Gurven, M., Stieglitz, J., Hooper, P. L., Gomes, C., & Kaplan, H. (2012). From the womb to the tomb: The role of
18 transfers in shaping the evolved human life history. *Experimental Gerontology*, 47(10), 807–813.
19 <https://doi.org/10.1016/j.exger.2012.05.006>
20
21 Gurven, M., & Walker, R. (2006). Energetic demand of multiple dependents and the evolution of slow human
22 growth. *Proceedings of the Royal Society B: Biological Sciences*, 273(1588), 835–841.
23 <https://doi.org/10.1098/rspb.2005.3380>
24
25 Hawkes, K. (2003). Grandmothers and the evolution of human longevity. *American Journal of Human Biology*,
26 15(3), 380–400. <https://doi.org/10.1002/ajhb.10156>
27
28 Hawkes, K., O'Connell, J. F., & Blurton Jones, N. G. (1989). Hardworking Hadza grandmothers. *Comparative*
29 *Socioecology: The Behavioral Ecology of Mammals and Man*, 341-366.
30
31 Henrickson, C. N. (2014). *The Archaeology of Cueva Santa Rita: A Late Holocene Rockshelter in the Sierra de La*
32 *Giganta of Baja California Sur, Mexico. Unpublished Doctoral Dissertation*. University of California
33 Berkeley.
34
35 Hewlett, B. S. (2017). *Hunter-Gatherer Childhoods: Evolutionary, Developmental, and Cultural Perspectives*.
36 Routledge.
37
38 Hewlett, B. S., & Cavalli-Sforza, L. L. (1986). Cultural Transmission among Aka Pygmies. *American*
39 *Anthropologist*, 88(4), 922–934. JSTOR.
40
41 Hewlett, B. S., Fouts, H. N., Boyette, A. H., & Hewlett, B. L. (2011). Social learning among Congo Basin hunter–
42 gatherers. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 366(1567), 1168–1178.
43 <https://doi.org/10.1098/rstb.2010.0373>
44
45 Hill, K., & Hurtado, A. M. (2009). Cooperative breeding in South American hunter–gatherers. *Proceedings of the*
46 *Royal Society B: Biological Sciences*, 276(1674), 3863–3870. <https://doi.org/10.1098/rspb.2009.1061>
47
48 Hooper, P. (2011). *The structure of energy production and redistribution among Tsimane' forager-horticulturalists*
49 [University of New Mexico]. https://digitalrepository.unm.edu/anth_etds/33
50
51 Howell, N. (2010). *Life Histories of the Dobe !Kung: Food, Fatness, and Well-being Over the Life Span*. University
52 of California Press.
53
54 Howell, S. N. G., & Webb, S. (1995). *A Guide to the Birds of Mexico and Northern Central America*. OUP Oxford.
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Huntsinger, L. (2016). Enabling sustainable pastoral landscapes: Building social capital to restore natural capital. In
5 *Mountain pastures and livestock farming facing uncertainty: Environmental, technical and socio-economic*
6 *challenges*. Casasús I. (ed.), Lombardi G. (ed.): Vol. n. 116 (pp. 315–325). CIHEAM.
7
8 INEGI. (2004). *El rezago educativo en la población mexicana*.
9 <http://en.www.inegi.org.mx/app/biblioteca/ficha.html?upc=702825497538>
10
11 INEGI. (2016). *Anuario estadístico y geográfico de Baja California*.
12 [http://internet.contenidos.inegi.org.mx/contenidos/Productos/prod_serv/contenidos/espanol/bvinegi/product](http://internet.contenidos.inegi.org.mx/contenidos/Productos/prod_serv/contenidos/espanol/bvinegi/productos/nueva_estruc/anuarios_2016/702825083663.pdf)
13 [os/nueva_estruc/anuarios_2016/702825083663.pdf](http://internet.contenidos.inegi.org.mx/contenidos/Productos/prod_serv/contenidos/espanol/bvinegi/productos/nueva_estruc/anuarios_2016/702825083663.pdf)
14
15 Institute of Medicine. (2002). *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids,*
16 *Cholesterol, Protein, and Amino Acids*.
17
18 Kaplan, H. (1994). Evolutionary and Wealth Flows Theories of Fertility: Empirical Tests and New Models.
19 *Population and Development Review*, 20(4), 753–791. JSTOR. <https://doi.org/10.2307/2137661>
20
21 Kaplan, H. (1996). A theory of fertility and parental investment in traditional and modern human societies.
22 *American Journal of Physical Anthropology*, 101(S23), 91–135. [https://doi.org/10.1002/\(SICI\)1096-](https://doi.org/10.1002/(SICI)1096-8644(1996)23+<91::AID-AJPA4>3.0.CO;2-C)
23 [8644\(1996\)23+<91::AID-AJPA4>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1096-8644(1996)23+<91::AID-AJPA4>3.0.CO;2-C)
24
25 Kaplan, H., Gurven, M., Winking, J., Hooper, P. L., & Stieglitz, J. (2010). Learning, menopause, and the human
26 adaptive complex. *Annals of the New York Academy of Sciences*, 1204(1), 30–42.
27 <https://doi.org/10.1111/j.1749-6632.2010.05528.x>
28
29 Kaplan, H., Hill, K., Lancaster, J., & Hurtado, A. M. (2000). A theory of human life history evolution: Diet,
30 intelligence, and longevity. *Evolutionary Anthropology: Issues, News, and Reviews*, 9(4), 156–185.
31 [https://doi.org/10.1002/1520-6505\(2000\)9:4<156::AID-EVAN5>3.0.CO;2-7](https://doi.org/10.1002/1520-6505(2000)9:4<156::AID-EVAN5>3.0.CO;2-7)
32
33 Kaplan, H., Hooper, P. L., & Gurven, M. (2009). The evolutionary and ecological roots of human social
34 organization. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1533), 3289–
35 3299. <https://doi.org/10.1098/rstb.2009.0115>
36
37 Kays, R. W., & Wilson, D. E. (2009). *Mammals of North America: Second Edition*. Princeton University Press.
38
39 Keil, F. C. (1987). Conceptual development and category structure. In *Concepts and conceptual development:*
40 *Ecological and intellectual factors in categorization* (pp. 175–200). Cambridge University Press.
41
42 Keil, F. C. (1994). The birth and nurturance of concepts by domains: The origins of concepts of living things. In
43 *Mapping the mind: Domain specificity in cognition and culture* (pp. 234–254). Cambridge University
44 Press. <https://doi.org/10.1017/CBO9780511752902.010>
45
46 Kline, M. A., Boyd, R., & Henrich, J. (2013). Teaching and the Life History of Cultural Transmission in Fijian
47 Villages. *Human Nature*, 24(4), 351–374. <https://doi.org/10.1007/s12110-013-9180-1>
48
49 Koster, J., Bruno, O., & Burns, J. L. (2016). Wisdom of the Elders? Ethnobiological Knowledge across the Lifespan.
50 *Current Anthropology*, 57(1), 113–121. <https://doi.org/10.1086/684645>
51
52 Koster, J., Lukas, D., Nolin, D., Power, E., Alvergne, A., Mace, R., Ross, C. T., Kramer, K., Greaves, R., Caudell,
53 M., MacFarlan, S., Schniter, E., Quinlan, R., Mattison, S., Reynolds, A., Yi-Sum, C., & Massengill, E.
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 (2019). Kinship ties across the lifespan in human communities. *Philosophical Transactions of the Royal*
5 *Society B*. <https://royalsocietypublishing.org/doi/abs/10.1098/rstb.2018.0069>
6
7 Kramer, K. L., & Ellison, P. T. (2010). Pooled energy budgets: Resituating human energy -allocation trade-offs.
8 *Evolutionary Anthropology: Issues, News, and Reviews*, 19(4), 136–147.
9 <https://doi.org/10.1002/evan.20265>
10
11 Lee, R. D. (2013). Intergenerational Transfers, the Biological Life Cycle, and Human Society. *Population and*
12 *Development Review*, 38(Suppl 1), 23–35. <https://doi.org/10.1111/j.1728-4457.2013.00549.x>
13
14 Lee, R. D., & Kramer, K. L. (2002). Children’s Economic Roles in the Maya Family Life Cycle: Cain, Caldwell,
15 and Chayanov Revisited. *Population and Development Review*, 28(3), 475–499.
16 <https://doi.org/10.1111/j.1728-4457.2002.00475.x>
17
18 León De La Luz, J. L., & Coria Benet, R. D. C. (2018). *Flora Iconográfica de Baja California Sur*.
19 <http://dspace.cibnor.mx:8080/handle/123456789/2131>
20
21 León De La Luz, J. L., & Domínguez Cadena, R. (2006). Hydrophytes of the oases in the Sierra de la Giganta of
22 Central Baja California Sur, Mexico: Floristic composition and conservation status. *Journal of Arid*
23 *Environments*, 67(4), 553–565. <https://doi.org/10.1016/j.jaridenv.2006.03.012>
24
25 Lindenberger, U., & Baltes, P. B. (1994). Aging and intelligence. In *Encyclopedia of human intelligence* (pp. 52–
26 66). Macmillan.
27
28 MacFarlan, S., Garcia, J. J., Schniter, E., Guevara Beltran, D., Amador Bibo, J., & Ruiz-Campos, G. (2019).
29 Geographic Distribution: Elgaria velazquezi (Central Baja California Alligator Lizard). *Herpetological*
30 *Review*, 50(1), 10.
31
32 Macfarlan, S. J., Garcia, J. J., & Schniter, E. (2020). Choyero identified plants and animals. *Mendeley Data*, 1.
33 <https://doi.org/10.17632/kjds8jztzv.1>
34
35 Macfarlan, S. J., & Henrickson, C. N. (2010). Inferring Relationships Between Indigenous Baja California Sur and
36 Seri/Comcáac Populations Through Cultural Traits. *Journal of California and Great Basin Anthropology*,
37 30(1), 51–67. JSTOR.
38
39 Macfarlan, S. J., Schacht, R., Foley, C., Cahoon, S., Osusky, G., Tayler, E., Henrickson, C. N., & Schniter, E. (n.d.).
40 Marriage dynamics in old Lower California: Ecological constraints and reproductive value in an arid
41 peninsular frontier. *Biodemography and Social Biology, Revise and Resubmit*.
42
43 Maddox, G. L., & Clark, D. O. (1992). Trajectories of Functional Impairment in Later Life. *Journal of Health and*
44 *Social Behavior*, 33(2), 114–125. JSTOR. <https://doi.org/10.2307/2137250>
45
46 Malinowski, B. K. (1925). *Magic, Science, Religion*. Souvenir Press.
47
48 Marlowe, F. W. (2007). Hunting and Gathering: The Human Sexual Division of Foraging Labor. *Cross-Cultural*
49 *Research*, 41(2), 170–195. <https://doi.org/10.1177/1069397106297529>
50
51 Maya, Y., Coria, R., & Domínguez, R. (1997). Caracterización de los oasis. In *Los oasis de la península de Baja*
52 *California* (pp. 5–25).
53
54 Messina, E., & Evans, J. M. (2006). *Standards for visual acuity*.
55 <https://pdfs.semanticscholar.org/7925/68f1947235eeb3eae2186a3adff4f67c6fae.pdf>
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Mithen, S. (2006). Ethnobiology and the evolution of the human mind. *Journal of the Royal Anthropological*
5 *Institute*, 12(s1), S45–S61. <https://doi.org/10.1111/j.1467-9655.2006.00272.x>
6
7 Nabhan, G. P., Garcia, J., Routson, R., Routson, K., & Cariño-Olvera, M. (2010). Desert oases as genetic refugia of
8 heritage crops: Persistence of forgotten fruits in the mission orchards of Baja California, Mexico.
9 *International Journal of Biodiversity and Conservation*, 2(4), 56–69.
10
11 Nabhan, G. P., & St Antoine, S. (1993). The loss of floral and faunal story: The extinction of experience. In *The*
12 *Biophilia Hypothesis*. Edited by Stephen R. Kellert, Edward O. Wilson (pp. 229–250). Island Press.
13
14 Nelson, E. A., & Dannefer, D. (1992). Aged Heterogeneity: Fact or Fiction? The Fate of Diversity in Gerontological
15 Research. *The Gerontologist*, 32(1), 17–23. <https://doi.org/10.1093/geront/32.1.17>
16
17 Ohmagari, K., & Berkes, F. (1997). Transmission of Indigenous Knowledge and Bush Skills Among the Western
18 James Bay Cree Women of Subarctic Canada. *Human Ecology*, 25(2), 197–222.
19 <https://doi.org/10.1023/A:1021922105740>
20
21 Oña, L., Oña, L. S., & Wertz, A. E. (2019). The evolution of plant social learning through error minimization.
22 *Evolution and Human Behavior*, 40(5), 447–456. <https://doi.org/10.1016/j.evolhumbehav.2019.05.009>
23
24 Park, D. C., Lautenschlager, G., Hedden, T., Davidson, N. S., Smith, A. D., & Smith, P. K. (2002). Models of
25 visuospatial and verbal memory across the adult life span. *Psychology and Aging*, 17(2), 299–320.
26 <https://doi.org/10.1037/0882-7974.17.2.299>
27
28 Parker, S. T., & McKinney, M. L. (2012). *Origins of Intelligence: The Evolution of Cognitive Development in*
29 *Monkeys, Apes, and Humans*. JHU Press.
30
31 Pío-León, J. F., Delgado-Vargas, F., Murillo-Amador, B., León-de-la-Luz, J. L., Vega-Aviña, R., Nieto-Garibay, A.,
32 Córdoba-Matson, M., & Ortega-Rubio, A. (2017). Environmental traditional knowledge in a natural
33 protected area as the basis for management and conservation policies. *Journal of Environmental*
34 *Management*, 201, 63–71. <https://doi.org/10.1016/j.jenvman.2017.06.032>
35
36 Rabe-Hesketh, S., & Skrondal, A. (2012). *Multilevel and Longitudinal Modeling using Stata-Volume II: Categorical*
37 *Responses Counts and Survival*. 3rd Edn. Stata Press.
38
39 Rebman, J., Roberts, N., & Ezcurra, E. (2012). *Baja California Plant field guide*. San Diego Natural History
40 Museum.
41
42 Reyes-García, V., Broesch, J., Calvet-Mir, L., Fuentes-Peláez, N., McDade, T. W., Parsa, S., Tanner, S., Huanca, T.,
43 Leonard, W. R., & Martínez-Rodríguez, M. R. (2009). Cultural transmission of ethnobotanical knowledge
44 and skills: An empirical analysis from an Amerindian society. *Evolution and Human Behavior*, 30(4), 274–
45 285. <https://doi.org/10.1016/j.evolhumbehav.2009.02.001>
46
47 Reyes-García, V., Gallois, S., & Demps, K. (2016). A Multistage Learning Model for Cultural Transmission:
48 Evidence from Three Indigenous Societies. In H. Terashima & B. S. Hewlett (Eds.), *Social Learning and*
49 *Innovation in Contemporary Hunter-Gatherers: Evolutionary and Ethnographic Perspectives* (pp. 47–60).
50 Springer Japan. https://doi.org/10.1007/978-4-431-55997-9_4
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Reyes-García, V., Guèze, M., Luz, A. C., Paneque-Gálvez, J., Macía, M. J., Orta-Martínez, M., Pino, J., & Rubio-
5 Campillo, X. (2013). Evidence of traditional knowledge loss among a contemporary indigenous society.
6 *Evolution and Human Behavior*, 34(4), 249–257. <https://doi.org/10.1016/j.evolhumbehav.2013.03.002>
7
8 Reyes-García, V., Martí, N., McDADE, T., Tanner, S., & Vadez, V. (2007). Concepts and methods in studies
9 measuring individual ethnobotanical knowledge. *Journal of Ethnobiology*, 27(2), 182–203.
10 [https://doi.org/10.2993/0278-0771\(2007\)27\[182:CAMISM\]2.0.CO;2](https://doi.org/10.2993/0278-0771(2007)27[182:CAMISM]2.0.CO;2)
11
12 Riemann, H., & Ezcurra, E. (2005). Plant endemism and natural protected areas in the peninsula of Baja California,
13 Mexico. *Biological Conservation*, 122(1), 141–150. <https://doi.org/10.1016/j.biocon.2004.07.008>
14
15 Robson, A. J., & Kaplan, H. S. (2003). The Evolution of Human Life Expectancy and Intelligence in Hunter-
16 Gatherer Economies. *American Economic Review*, 93(1), 150–169.
17
18 Rodd, T., Bryant, G., & Barnard, L. (2007). *The Plant Finder*. Firefly Books. [http://agris.fao.org/agris-](http://agris.fao.org/agris-search/search.do?recordID=US201300120161)
19 [search/search.do?recordID=US201300120161](http://agris.fao.org/agris-search/search.do?recordID=US201300120161)
20
21 Rodríguez-Estrella, R. (2005). Terrestrial birds and conservation priorities in Baja California Peninsula. In *Bird*
22 *Conservation Implementation and Integration in the Americas: Proceedings of the Third International*
23 *Partners in Flight Conference, March 20-24, 2002, Asilomar, California* (Vol. 1, p. 115). U.S. Department
24 of Agriculture, Forest Service, Pacific Southwest Research Station.
25
26 Rosenberg, I. H. (1997). Sarcopenia: Origins and Clinical Relevance. *The Journal of Nutrition*, 127(5), 990S-991S.
27 <https://doi.org/10.1093/jn/127.5.990S>
28
29 Ruddle, K., & Chesterfield, R. (1977). *Education for Traditional Food Procurement in the Orinoco Delta*.
30 University of California Press.
31
32 Ruiz-Campos, G., Camarena-Rosales, F., González-Acosta, A. F., Maeda-Martinez, A. M., García de León, F. J.,
33 Varela-Romero, A., & Andreu-Soler, A. (2014). Estatus actual de conservación de seis especies de peces
34 dulceacuícolas de la península de Baja California, México. *Revista Mexicana de Biodiversidad*, 85(4),
35 1235–1248. <https://doi.org/10.7550/rmb.43747>
36
37 Ruiz-Campos, G., Castro-Aguirre, J. L., Contreras-Balderas, S., de Lourdes Lozano-Vilano, M., González-Acosta,
38 A. F., & Sánchez-González, S. (2002). An annotated distributional checklist of the freshwater fish from
39 Baja California Sur, México. *Reviews in Fish Biology and Fisheries*, 12(2), 143–155.
40 <https://doi.org/10.1023/A:1025067202218>
41
42 Salthouse, T. A. (1993). Speed and Knowledge as Determinants of Adult Age Differences in Verbal Tasks. *Journal*
43 *of Gerontology*, 48(1), P29–P36. <https://doi.org/10.1093/geronj/48.1.P29>
44
45 Schaie, K. W. (1994). The course of adult intellectual development. *American Psychologist*, 49(4), 304–313.
46 <https://doi.org/10.1037/0003-066X.49.4.304>
47
48 Schneider, B. A., & Pichora-Fuller, M. K. (2000). Implications of perceptual deterioration for cognitive aging
49 research. In *The handbook of aging and cognition*, 2nd ed (pp. 155–219). Lawrence Erlbaum Associates
50 Publishers.
51
52 Schniter, E. (2009). *Why Old Age: Non-Material Contributions and Patterns of Aging Among Older Adult Tsimane'*
53 [Dissertation]. University of California, Santa Barbara.
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 Schniter, E. (2014). Older Adults' Contributions to the Tsimane Forager-Farmer Economy. *Anthropology & Aging*,
5 35(1), 56–58. <https://doi.org/10.5195/aa.2014.62>
6
- 7 Schniter, E., Gurven, M., Kaplan, H. S., Wilcox, N. T., & Hooper, P. L. (2015). Skill ontogeny among Tsimane
8 forager-horticulturalists. *American Journal of Physical Anthropology*, 158(1), 3–18.
9 <https://doi.org/10.1002/ajpa.22757>
10
- 11 Schniter, E., Macfarlan, S. J., & Garcia, J. J. (2020). Choyero ethnobiological knowledge survey. *Mendeley Data*, 1.
12 <https://doi.org/10.17632/hgd8zdyf2c.1>
13
- 14 Schniter, E., Wilcox, N. T., Beheim, B. A., Kaplan, H. S., & Gurven, M. (2018). Information transmission and the
15 oral tradition: Evidence of a late-life service niche for Tsimane Amerindians. *Evolution and Human*
16 *Behavior*, 39(1), 94–105. <https://doi.org/10.1016/j.evolhumbehav.2017.10.006>
17
- 18 Schuppli, C., Isler, K., & van Schaik, C. P. (2012). How to explain the unusually late age at skill competence among
19 humans. *Journal of Human Evolution*, 63(6), 843–850. <https://doi.org/10.1016/j.jhevol.2012.08.009>
20
- 21 SEDESOL. (2010). *Catalago de Localidades: Resumen de Localidades del municipio* (Sistema de Apoyo Para La
22 Planeación Del PDZP).
23 <http://www.microrregiones.gob.mx/catloc/LocdeMun.aspx?tipo=clave&campo=loc&ent=03&mun=003>
24
- 25 Setalaphruk, C., & Price, L. L. (2007). Children's traditional ecological knowledge of wild food resources: A case
26 study in a rural village in Northeast Thailand. *Journal of Ethnobiology and Ethnomedicine*, 3(1), 33.
27 <https://doi.org/10.1186/1746-4269-3-33>
28
- 29 Shreve, F., & Wiggins, I. L. (1964). *Vegetation and Flora of the Sonoran Desert*. Stanford University Press.
30
- 31 Stieglitz, J., Gurven, M., Kaplan, H., & Hooper, P. L. (2013). Household Task Delegation among High-Fertility
32 Forager-Horticulturalists of Lowland Bolivia. *Current Anthropology*, 54(2), 232–241.
33 <https://doi.org/10.1086/669708>
34
- 35 Stieglitz, J., Schniter, E., von Rueden, C., Kaplan, H., & Gurven, M. (2015). Functional Disability and Social
36 Conflict Increase Risk of Depression in Older Adulthood Among Bolivian Forager-Farmers. *The Journals*
37 *of Gerontology: Series B*, 70(6), 948–956. <https://doi.org/10.1093/geronb/gbu080>
38
- 39 Walker, R., Beckerman, S., Flinn, M. V., Gurven, M., von Rueden, C. R., Kramer, K. L., Greaves, R. D., Córdoba,
40 L., Villar, D., Hagen, E. H., Koster, J. M., Sugiyama, L., Hunter, T. E., & Hill, K. R. (2013). Living with
41 Kin in Lowland Horticultural Societies. *Current Anthropology*, 54(1), 96–103.
42 <https://doi.org/10.1086/668867>
43
- 44 Walker, R., & Hill, K. (2003). Modeling growth and senescence in physical performance among the ache of eastern
45 paraguay. *American Journal of Human Biology*, 15(2), 196–208. <https://doi.org/10.1002/ajhb.10135>
46
- 47 Walker, R., Hill, K., Kaplan, H., & McMillan, G. (2002). Age-dependency in hunting ability among the Ache of
48 Eastern Paraguay. *Journal of Human Evolution*, 42(6), 639–657. <https://doi.org/10.1006/jhev.2001.0541>
49
- 50 Wertz, A. E., & Wynn, K. (2014a). Thyme to touch: Infants possess strategies that protect them from dangers posed
51 by plants. *Cognition*, 130(1), 44–49. <https://doi.org/10.1016/j.cognition.2013.09.002>
52
- 53 Wertz, A. E., & Wynn, K. (2014b). Selective Social Learning of Plant Edibility in 6- and 18-Month-Old Infants.
54 *Psychological Science*, 25(4), 874–882. <https://doi.org/10.1177/0956797613516145>
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Wiggins, I. L. (1980). *Flora of Baja California*. Stanford University Press.

<https://www.cabdirect.org/cabdirect/abstract/19810668708>

Zarger, R. K., & Stepp, J. R. (2004). Persistence of Botanical Knowledge among Tzeltal Maya Children. *Current Anthropology*, 45(3), 413–418. <https://doi.org/10.1086/420908>

Zent, S. (2001). Acculturation and ethnobotanical knowledge loss among the Piaroa of Venezuela: Demonstration of a quantitative method for the empirical study of TEK change. In *On biocultural diversity: Linking Language, Knowledge, and the Environment*, L. Maffi (Ed.) (pp. 190–211). Harper Collins.

Table 1. *Choyero* reproductive demographics.

	N	Mean (SD)	Median	Min, Max
Age at first birth (years) among women	53	22.6 (5)	22	15, 42
IBI* between offspring 1 & 2 (days)	43	1335(761)	1190	224/2985
IBI between offspring 2 & 3 (days)	33	1565(1108)	1307	244/4204
IBI between offspring 3 & 4 (days)	19	1719(1688)	896	367/6768
IBI between offspring 4 & 5 (days)	7	864(728)	661	314/2466
IBI between offspring 5 & 6 (days)	6	1530(1736)	649	377/4822
IBI between offspring 6+ (days)	4	809(538)	656	342/1586
IBI All (days)	112	1430(1125)	1044	224/6768
Total fertility rate among women ¹	27	3.7(2)	3	1/9

*IBI: Interbirth intervals.

¹Total fertility was based on our sample of women who were older than 45 years.

Table 2. *Choyero* ethnobiological knowledge sample characteristics for full sample (all), females, and males.

	N	Mean (SD)	Median	Min, Max
Full sample	71			
Age	71	46.6 (17.73)	48	5, 86
Visual acuity	71	10.6 (0.77)	11	6, 11
Educational achievement (years)	71	5.0 (3.71)	6	0, 12
Females	33			
Age	33	47.3 (18.33)	49	5, 79
Visual acuity	33	10.5 (0.97)	11	6, 11
Educational achievement (years)	33	4.6 (3.39)	6	0, 10
Males	38			
Age	38	45.8 (17.42)	45.5	16, 86
Visual acuity	38	10.7 (0.53)	11	9, 11
Educational achievement (years)	38	5.3 (3.98)	6	0, 12

Table 3. Two stage multi-level logit regression model coefficients associated with *Choyero* ethnobiological knowledge.

	Model							
	(1)				(2)			
	B (RSE) ¹	OR (RSE) ²	z	p	B (RSE) ¹	OR (RSE) ²	z	p
DV: Recognized^{3,4}								
Age (years)	0.097(0.02)	1.10(0.02)	4.9	<.001	0.100(0.02)	1.11(0.02)	4.9	<.001
Age Squared (years)	-0.001(0.0002)	0.99(0.0002)	-4.3	<.001	-0.001(0.0002)	0.99(0.0002)	-4.7	<.001
Visual Acuity	0.410(0.11)	1.50(0.17)	3.7	<.001	0.460(0.12)	1.59(0.2)	3.9	<.001
Education (years)	0.023(0.028)	1.02(0.03)	0.8	.41				
Sex Male (=1)	0.160(0.12)	1.17(0.14)	1.3	.19				
Food or Danger (=1)					0.720(0.05)	2.05(0.1)	13.6	<.001
Constant	-5.400(1.4)	0.005(0.006)	-4.0	<.001	-6.100(1.5)	0.01(0.003)	-4.1	<.001
DV: Named^{5,6}								
Age (years)	.0930(0.016)	1.10(0.02)	5.9	<.001	0.097(0.01)	1.11(0.02)	7.1	<.001
Age Squared (years)	-0.001(0.00019)	0.99(0.0002)	-4.9	<.001	-0.001(0.0001)	0.99(0.0001)	-6.5	<.001
Visual Acuity	0.150(0.07)	1.20(0.08)	2.1	.032	0.190(0.06)	1.20(0.07)	3.1	.002
Education (years)	0.003(0.018)	1.00(0.02)	0.2	.87				
Sex Male (=1)	0.160(0.07)	1.20(0.08)	2.2	.03				
Food or Danger (=1)					0.580(0.05)	1.80(0.09)	11.7	<.001
Constant	-3.190(0.089)	0.04(0.04)	-3.6	<.001	-3.900(0.8)	0.02(0.02)	-4.7	<.001

¹OR(RSE): Odds Ratio (Robust Standard Errors); ²B(RSE): Unstandardized coefficient (Robust Standard Errors)

³Model 1 Wald $X^2 = 46.6$; n observations = 9,727; n ranches = 40; N individuals = 71; $p < .0001$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.09(0.09) and Ego = 0.4(0.2)

⁴Model 2 Wald $X^2 = 266.4$; n observations = 9,727; n ranches = 40; N individuals = 71; $p < .0001$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.01(0.01) and Ego = 0.4(0.2)

⁵Model 1 Wald $X^2 = 113.1$; n observations = 7,435; n ranches = 40; N individuals = 71; $p < .0001$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.03(0.03) and Ego = 0.8(0.3)

⁶Model 2 Wald $X^2 = 181.7$; n observations = 7,435; n ranches = 40; N individuals = 71; $p < .0001$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.05(0.03) and Ego = 0.8(0.3)

Table 4. Two stage multi-level logit regression model coefficients associated with *Choyero* ethnobiological plant knowledge.

	Model							
	(3 Jardin Plants)				(4 Campo Plants)			
	B (RSE) ¹	OR (RSE) ²	z	p	B (RSE) ¹	OR (RSE) ²	z	p
DV: Recognized^{3,4}								
Age (years)	0.09 (0.02)	1.09(0.03)	3.6	<.001	0.09(0.02)	1.1(0.02)	4.2	<.001
Age Squared (years)	-0.001(0.0003)	0.99(0.0003)	-3.5	<.001	-0.001(0.0002)	0.99(0.0002)	-3.9	<.001
Visual Acuity	0.410(0.11)	1.74(0.3)	3.7	<.001	0.5(0.1)	1.6(0.02)	3.7	<.001
Sex Male (=1)	-0.6 (0.20)	0.55(0.09)	-3.7	<.001	0.3 (0.2)	1.4(0.2)	2.1	.04
Constant	-6.2 (1.8)	0.002(0.004)	-3.4	.001	-6.4(1.6)	0.002(0.003)	-3.9	<.001
DV: Named^{5,6}								
Age (years)	.1(0.03)	1.13(0.03)	4.7	<.001	0.1(0.02)	1.11(0.02)	5.5	<.001
Age Squared (years)	-0.001(0.0003)	0.99(0.0003)	-4.6	<.001	-0.001(0.0002)	0.99(0.0002)	-4.7	<.001
Visual Acuity	0.3(0.1)	1.37(0.2)	2.9	.004	0.2(0.1)	1.2(0.1)	4.2	.08
Sex Male (=1)	-0.6(0.1)	0.55(0.07)	-4.4	<.001	0.4(0.1)	1.5(0.1)	4.2	<.001
Constant	-5.1(1.5)	0.006(0.009)	-3.5	<.001	-4.2(1.2)	0.02(0.02)	-3.5	<.001

¹OR(RSE): Odds Ratio (Robust Standard Errors); ²B(RSE): Unstandardized coefficient (Robust Standard Errors)

³Model 3 Wald $X^2 = 30.1$; n observations = 2,556; n ranches = 40; N individuals = 71; $p < .0001$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.2(0.09) and Ego = 0.3(0.1)

⁴Model 4 Wald $X^2 = 31.7$; n observations = 3,7673; n ranches = 40; N individuals = 71; $p < .0001$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.05(0.01) and Ego = 0.5(0.2)

⁵Model 3 Wald $X^2 = 31.1$; n observations = 1,893; n ranches = 40; N individuals = 71; $p < .0001$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.1(0.06) and Ego = 0.1(0.06)

⁶Model 4 Wald $X^2 = 78.5$; n observations = 2,575; n ranches = 40; N individuals = 71; $p < .0001$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 7.84e-32(5.47e-31) and Ego = 0.2(0.06)

Table 5. Two stage multi-level logit regression model coefficients associated with *Choyero* post-reproductive ethnobiological plant knowledge.

	Model							
	(5 Jardin Plants)				(6 Campo Plants)			
	B (RSE) ¹	OR (RSE) ²	z	p	B (RSE) ¹	OR (RSE) ²	z	p
DV: Recognized^{3,4}								
Age (years)	-0.03 (0.02)	0.97(0.01)	-2.2	.027	-0.03 (0.01)	0.97(0.01)	-2.2	.03
Visual Acuity	0.6(0.2)	1.8(0.3)	3.5	<.001	0.5(0.1)	1.69(0.2)	3.7	<.001
Sex Male (=1)	-0.7 (0.3)	0.48(0.1)	-2.5	.012	0.2 (0.3)	1.21(0.3)	0.7	.048
Constant	-2.5 (1.7)	0.08(0.1)	-1.5	.15	-2.8 (1.6)	0.068(0.1)	-1.8	.08
DV: Named^{5,6}								
Age (years)	-0.04 (0.01)	0.96(0.01)	-2.6	.01	-0.03 (0.01)	0.97(0.01)	-2.6	.01
Visual Acuity	0.4(0.2)	1.46(0.2)	2.3	.023	0.3(0.1)	1.38(0.2)	2.5	.01
Sex Male (=1)	-0.6 (0.3)	0.53(0.1)	-2.4	.015	0.2 (0.1)	1.27(0.2)	1.7	.09
Constant	-1.0 (1.4)	0.4(0.5)	-0.7	.47	-1.4 (1.0)	0.26(0.3)	-1.3	.19

¹OR(RSE): Odds Ratio (Robust Standard Errors); ²B(RSE): Unstandardized coefficient (Robust Standard Errors)


³Model 5 Wald $X^2 = 18.1$; n observations = 1,368; n ranches = 26; N individuals = 38; $p = .0004$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.17(0.1) and Ego = 0.4(0.1)

⁴ Model 6 Wald $X^2 = 19.4$; n observations = 2014; n ranches = 26; N individuals = 38; $p = .0002$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.1(0.01) and Ego = 0.5(0.2)

⁵ Model 5 Wald $X^2 = 12.2$; n observations = 972; n ranches = 26; N individuals = 38; $p = .00067$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.3(0.3) and Ego = 0.2(0.3)

⁶ Model 6 Wald $X^2 = 18.5$; n observations = 1348; n ranches = 26; N individuals = 38; $p = .0003$; Random effects estimates of estimated standard deviation in the intercept on the logit scale (and its standard error) include Ranch = 0.2(0.09) and Ego = 0.03(0.05)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 2. Choyero ranches and communities in the southern Sierra de La Giganta of Baja California Sur, México




Figure 3. The predicted *Choyero* ability to correctly name plant and animal species is extracted from the regression model and plotted as a function of age, age square, sex, visual acuity, and education, with random effects for data nested at the level of ranches and individuals.




Figure 4. The predicted *Choyero* ability to correctly name plant and animal species is extracted from the regression model and plotted as a function of age, age square, visual acuity, and whether or not the species were potentially food and/or dangerous, with random effects for data nested at the level of ranches and individuals.




Figure 5. The predicted *Choyero* ability to correctly name house garden (*jardin*) plants is extracted from the regression model and plotted as a function of age, age square, sex, and visual acuity, with random effects for data nested at the level of ranches and individuals.




Figure 6. The predicted *Choyero* ability to correctly name wild (*campo*) plants is extracted from the regression model and plotted as a function of age, age square, sex, and visual acuity, with random effects for data nested at the level of ranches and individuals.

Appendix A.

Note: For the “tumbling E” eye chart used to assess visual acuity and the original stimulus in the Ethnobiological Knowledge Assessment Tool consisting of 137 slides showing the images of plants and animals listed in the table below, see [Schniter, E., Macfarlan, S. J., & Garcia, J. J. \(2020\). Choyero ethnobiological knowledge survey. *Mendeley Data*, 1. <https://doi.org/10.17632/hgd8zdyf2c.1>](#)

Table 7. Plants and animals in the Ethnobiological Knowledge Assessment Tool.

Item #	Family	Genus	species	Choyero common names	Plant=1 Animal=0	Food =1	Danger =1	Jardin =1	Huerta =1	Campo =1
1	Arecaceae	<i>Phoenix</i>	<i>dactylifera</i>	Datil, Mata de Datil; Datiles; Palma de Datil	1	1	0	1	1	0
2	Moraceae	<i>Ficus</i>	<i>carica</i>	Higo, Higuera	1	1	0	1	1	0
3	Myrtaceae	<i>Psidium</i>	<i>guajava</i>	Guayabo(a)	1	1	0	1	1	0
4	Fabaceae	<i>Tamarindus</i>	<i>sp.</i>	Tamarindo	1	1	0	1	1	0
5	Rosaceae	<i>Prunus</i>	<i>persica</i>	Durazno	1	1	0	1	1	0
6	Rutaceae	<i>Casimiroa</i>	<i>edulis</i>	Zapote (Amarillo o Blanco)	1	1	0	1	1	0
7	Rutaceae	<i>Citrus</i>	<i>sinensis</i>	Naranja(o)	1	1	0	1	1	0
8	Rutaceae	<i>Citrus</i>	<i>aurantium</i>	Naranja amarga (agria)	1	1	0	1	1	0
9	Rutaceae	<i>Citrus</i>	<i>limon</i>	Limón real, Limón	1	1	0	1	1	0
10	Rutaceae	<i>Citrus</i>	<i>medica</i>	Cidra, Limón agrio	1	1	0	1	1	0
11	Oleaceae	<i>Olea</i>	<i>europaea</i>	Olivo, Aceituna	1	1	0	1	1	0
12	Passifloraceae	<i>Passiflora</i>	<i>ligularis</i>	Granadilla, Granada china, Rosal de pasion, Maracuya	1	1	0	1	1	0
13	Rosaceae	<i>Prunu</i>	<i>armeniaca</i>	Chabacano, Alberechigo, Albericoque, Manzano	1	1	0	1	1	0
14	Fabaceae	<i>Pithecellobium</i>	<i>dulce</i>	Guamuchil, Mesquite con pechita dulce, Roscas	1	1	0	1	1	0
15	Fabaceae	<i>Prosopis</i>	<i>palmeri</i>	Palo fiero; Palo Hierro	1	0	0	0	0	1
16	Fabaceae	<i>Olneya</i>	<i>tesota</i>	Uña de gato	1	0	1	0	0	1
17	Fouquieriaceae	<i>Fouquieria</i>	<i>diguetii</i>	Palo Adan	1	0	1	0	0	1
18	Euphorbiaceae	<i>Jatropha</i>	<i>cuneata</i>	Matacora	1	0	0	0	0	1

19	Acanthaceae	<i>Ruellia</i>	<i>californica peninsularis</i>	Rama parda	1	0	0	0	0	1
20	Burseraceae	<i>Bursera</i>	<i>epinnata</i>	Copal Blanco	1	0	0	0	0	1
21	Fabaceae	<i>Acacia</i>	<i>farnesiana</i>	Huizache	1	0	1	0	0	1
22	Cactaceae	<i>Stenocereus</i>	<i>gummosus</i>	Pitaya agria	1	1	1	0	0	1
23	Fabaceae	<i>Mimosa</i>	<i>distachya</i>	Garabatillo	1	0	1	0	0	1
24	Fabaceae	<i>Parkinsonia</i>	<i>microphylla</i>	Lipua	1	0	1	0	0	1
25	Fabaceae	<i>Parkinsonia</i>	<i>praecox</i>	Palo brea	1	0	1	0	0	1
26	Rhamnaceae	<i>Colubrina</i>	<i>viridis</i>	Palo colorado	1	0	0	0	0	1
27	Anacardiaceae	<i>Cyrtocarpa</i>	<i>edulis</i>	Ciruela de campo/monte	1	1	0	0	0	1
28	Cochlospermaceae	<i>Amoreuxia</i>	<i>palmatifida</i>	Saya	1	1	0	0	0	1
29	Anacardiaceae	<i>Pachycormus</i>	<i>discolor</i>	Copalquin	1	0	0	0	0	1
30	Asteraceae	<i>Baccharis</i>	<i>salicifolia</i>	Guatamote	1	0	0	0	0	1
31	Zygophyllaceae	<i>Larrea</i>	<i>tridentata</i>	Gobernadora	1	0	0	0	0	1
32	Polygonaceae	<i>Antigonon</i>	<i>leptopus</i>	San Miguel	1	0	0	0	0	1
33	Anacardiaceae	<i>Mangifera</i>	<i>indica</i>	Mango	1	1	0	1	1	0
34	Convolvulaceae	<i>Merremia</i>	<i>aurea</i>	Yuca	1	0	0	0	0	1
35	Moraceae	<i>Ficus</i>	<i>palmeri</i>	Higuera cimaron, Higuera silvestre, Zalate	1	1	0	0	0	1
36	Euphorbiaceae	<i>Sebastiania</i>	<i>bilocularis</i>	Hierba de flecha	1	0	1	0	0	1
37	Fabaceae	<i>Calliandra</i>	<i>californica</i>	Tabardillo	1	0	0	0	0	1
38	Cactaceae	<i>Cylindropuntia</i>	<i>molesta var. clavellina</i>	Ciribe, Caribe, Clavellina, Clavin	1	0	1	0	0	1
39	Euphorbiaceae	<i>Adelia</i>	<i>brandegeei</i>	Pimientilla	1	0	0	0	0	1
40	Asteraceae	<i>Ambrosia</i>	<i>ambrosioides</i>	Chicura	1	0	0	0	0	1
41	Geraniaceae	<i>Geranium</i>	<i>spp.</i>	Geranio	1	0	0	1	0	0
42	Asteraceae	<i>Amauria</i>	<i>rotundifolia</i>	Manzanilla	1	0	0	1	0	0
43	Fabaceae	<i>Aeschynomene</i>	<i>nivea Brandegeei</i>	Vara prieta	1	0	0	0	0	1
44	Cactaceae	<i>Mammillaria</i>	<i>albicans</i>	Viejito, "Chiltipin"	1	1	1	0	0	1
45	Apocynaceae	<i>Vallesia</i>	<i>glabra</i>	Huitatave, Hutatave	1	1	0	0	0	1

46	Malvaceae	<i>Hibiscus</i>	<i>spp.</i>	Obelisco	1	1	0	1	0	0
47	Typhaceae	<i>Typha</i>	<i>domingensis</i>	Tule	1	0	0	0	0	1
48	Portulacaceae	<i>Portulaca</i>	<i>grandiflora</i>	Amore, Hielitos	1	0	0	1	0	0
49	Lamiaceae	<i>Coleus</i>	<i>blumei</i>	Brocado	1	0	0	1	0	0
50	Simmondsiaceae	<i>Simmondsia</i>	<i>chinensis</i>	Jojoba	1	0	0	0	0	1
51	Solanaceae	<i>Solanum</i>	<i>hindsianum</i>	Mariola	1	0	0	0	0	1
52	Rubiaceae	<i>Randia</i>	<i>capitata</i>	Palo santo, papache	1	1	0	0	0	1
53	Nolinaceae	<i>Nolina</i>	<i>palmeri var. brandegeei</i>	Datilillo	1	0	0	0	0	1
54	Krameriaceae	<i>Krameria</i>	<i>bicolor</i>	Mesquitillo	1	0	0	0	0	1
55	Rutaceae	<i>Esenbeckia</i>	<i>flava</i>	Palo amarillo	1	0	0	0	0	1
56	Fabaceae	<i>Delonix</i>	<i>regia</i>	Arbol del fuego, Tabachin, Tabachina	1	0	0	1	0	0
57	Saururaceae	<i>Anemopsis</i>	<i>californica</i>	Hierba de manso	1	0	0	0	0	1
58	Stixaceae	<i>Forchhammeria</i>	<i>watsonii</i>	Palo San Juan	1	0	0	0	0	1
59	Asteraceae	<i>Tagetes</i>	<i>spp.</i>	Cenpasuchil, Cempasuchil, Tenpasuchil, Tempasuchil	1	0	0	1	0	0
60	Sapotaceae	<i>Sideroxylon</i>	<i>occidentale</i>	Bebelama	1	0	0	0	0	1
61	Euphorbiaceae	<i>Jatropha</i>	<i>vernica</i>	Lomboy sangregado, Lomboy rojo	1	0	0	0	0	1
62	Euphorbiaceae	<i>Euphorbia</i>	<i>lomelii</i>	Candelilla, Liga, Periquito de Amor, Periquito, Perrico	1	0	0	0	0	1
63	Euphorbiaceae	<i>Euphorbia</i>	<i>pulcherrima</i>	Noche buena	1	0	0	1	0	0
64	Anacardiaceae	<i>Schinus</i>	<i>molle</i>	Pirul	1	0	0	0	0	1
65	Malpighiaceae	<i>Callaeum</i>	<i>macropterum</i>	Gallineta	1	0	0	0	0	1
66	Cactaceae	<i>Echinocereus</i>	<i>brandegeei</i>	Pitayita, Casa de Ratas	1	0	1	0	0	1
67	Asteraceae	<i>Gazania</i>	<i>spp.</i>	Solecito, Novia del Sol, Muchacho del Sol	1	0	0	1	0	0
68	Fabaceae	<i>Senna</i>	<i>covesii</i>	Hojasen	1	0	0	0	0	1
69	Apocynaceae	<i>Catharanthus</i>	<i>roseus</i>	Terecita	1	0	0	1	0	0
70	Asteraceae	<i>Encelia</i>	<i>farinosa</i>	Incensio, Incienso	1	0	0	0	0	1

71	Oleaceae or Fabaceae	<i>Ligustrum</i> or <i>Clitoria</i>	<i>japonicum</i> or <i>ternatea</i>	Manga de niño	1	0	0	1	0	0
72	Liliaceae	<i>Lilium</i>	<i>spp.</i>	Lirio, Asuzena	1	0	0	1	0	0
73	Cannaceae	<i>Canna</i>	<i>spp.</i>	Platanillo, Mariposa	1	0	0	1	0	0
74	Aizoaceae	<i>Trianthema</i>	<i>portulacastrum</i>	Verdolaga, Verdolaga de cochi	1	1	0	0	0	1
75	Asteraceae	<i>Ambrosia</i>	<i>camphorata</i>	Estafiete, Estafiate	1	0	0	0	0	1
76	Convolvulaceae	<i>Ipomoea</i>	<i>batatas</i>	Camote	1	1	0	0	1	0
77	Moraceae	<i>Ficus</i>	<i>benjamina</i>	Benjasmin	1	0	0	1	0	0
78	Combretaceae	<i>Combretum</i>	<i>indicum</i>	Madre selva	1	0	0	1	0	0
79	Fabaceae	Leucaena	leucocephala	Guaje, Guiachi, Guajil, Bombita/Bombito	1	1	0	1	0	1
80	Cactaceae	Stenocereus	eruca	Chirinola	1	0	1	0	0	1
81	Arecaceae	Brahea	brandegeei	Palmilla	1	0	0	1	0	1
82	Cactaceae	<i>Peniocereus</i>	<i>striatus</i>	Cardoncillo, Matraca	1	0	1	0	0	1
83	Fabaceae	<i>Caesalpinia</i>	<i>placida</i>	Cresta de gallo, Navajia	1	0	0	1	0	1
84	Cactaceae	<i>Myrtillocactus</i>	<i>cochal</i>	Gualama, Pilita, Pila	1	1	1	0	0	1
85	Asteraceae	<i>Zinnia</i>	<i>spp.</i>	Zinia	1	0	0	1	0	0
86	Malvaceae	<i>Gossypium</i>	<i>davidsonii</i>	Algodon	1	0	0	1	0	0
87	Lamiaceae	<i>Ocimum</i>	<i>basilicum</i>	Albacar, Albaca, Albahaca	1	0	0	1	0	0
88	Columbidae	<i>Zenaida</i>	<i>asiatica</i>	Paloma	0	1	0	0	0	0
89	Canidae	<i>Canis</i>	<i>latrans</i>	Coyote	0	0	1	0	0	0
90	Cathartidae	<i>Cathartes</i>	<i>aura teter</i>	Aura, Zopilote, Buitre	0	0	1	0	0	0
91	Cardinalidae	<i>Cardinalis</i>	<i>cardinalis</i>	Cardinal, Carindal Pardo	0	0	0	0	0	0
92	Felidae	<i>Lynx</i>	<i>rufus peninsularis</i>	Gato montes, Leoncillo, Linxe	0	0	1	0	0	0
93	Odontophoridae	<i>Callipepla</i>	<i>californica</i>	Chacuaca, Godorniz	0	1	0	0	0	0
94	Accipitridae	<i>Accipiter</i>	<i>cooperii</i>	Gavilansillo; Gavilan pollero	0	0	1	0	0	0
95	Leporidae	<i>Lepus</i>	<i>californicus</i>	Liebre	0	1	0	0	0	0
96	Icteridae	<i>Icterus</i>	<i>cucullatus</i>	Calandria, Caliandra	0	0	0	0	0	0
97	Phrynosomatidae	<i>Urosaurus</i>	<i>lahtelai</i> and	Cachora	0	0	0	0	0	0

<i>nigricaudus</i>										
98	Procyonidae	<i>Procyon</i>	<i>lotor</i>	Mapache	0	0	1	0	0	0
99	Strigidae	<i>Bubo</i>	<i>virginianus</i>	Tecolote	0	0	1	0	0	0
100	Falconidae	<i>Caracara</i>	<i>plancus auduboni</i>	Quelele, Cara cara	0	0	1	0	0	0
101	Sciuridae	<i>Otospermophilus</i>	<i>atricapillus</i>	Ardilla, Ardillon	0	0	0	0	0	0
102	Columbidae	<i>Columbina</i>	<i>passerina</i>	Torcasa, Tortulita, Zacatera, Paloma Zacatera	0	1	0	0	0	0
103	Troglodytidae	<i>Campylorhynchus</i>	<i>brunneicapillus</i>	Urraca	0	0	0	0	0	0
104	Accipitridae	<i>Parabuteo</i>	<i>unicinctus</i>	Halcón	0	0	1	0	0	0
105	Colubridae	<i>Pituophis</i>	<i>vertebralis</i>	Coralillo, Alicante	0	0	0	0	0	0
106	Mimidae	<i>Mimus</i>	<i>polyglottos</i>	Centzontle	0	0	0	0	0	0
107	Mimidae	<i>Toxostoma</i>	<i>cinereum</i>	Huirigo, Huirico, Urico, Huiri, Huitacoche, Huitlacoche	0	0	0	0	0	0
108	Poliophtilidae	<i>Poliophtila</i>	<i>caerulea</i>	Chivirito	0	0	0	0	0	0
109	Mustelidae	<i>Taxidea</i>	<i>taxus</i>	Tejon	0	0	1	0	0	0
110	Procyonidae	<i>Bassariscus</i>	<i>astutus palmarius</i>	Babisuri	0	0	0	0	0	0
111	Heteromyidae	<i>Chaetodipus</i>	<i>spinatus</i>	Ratón	0	0	1	0	0	0
112	Corvidae	<i>Aphelocoma</i>	<i>californica</i>	Pajaro azul	0	0	0	0	0	0
113	Strigidae	<i>Megascops</i>	<i>kennicotti</i>	Lechuza	0	0	1	0	0	0
114	Geomyidae	<i>Thomomys</i>	<i>umbrinus alticolus</i>	Tuza, Topo ciego, Topo	0	0	0	0	0	0
115	Ardeidae	<i>Ardea</i>	<i>herodias</i>	Garza (morena, azul, gris)	0	0	0	0	0	0
116	Ptiliognatidae	<i>Phainopepla</i>	<i>nitens</i>	Cardenal negro, Cardenal prieto, Cardenal parda	0	0	0	0	0	0
117	Iguanidae	<i>Ctenosaura</i>	<i>hemilopha</i>	Iguana de arbol, Iguana de palo, Iguana de mesquite	0	0	0	0	0	0
118	Phrynosomatidae	<i>Callisaurus</i>	<i>draconoides</i>	Cachimba	0	0	0	0	0	0
119	Rallidae	<i>Fulica</i>	<i>americana</i>	Gallineta	0	0	0	0	0	0
120	Phyllodactylidae	<i>Phyllodactylus</i>	<i>xanti</i>	Salamanques(a)	0	0	0	0	0	0
121	Phrynosomatidae	<i>Phrynosoma</i>	<i>coronatum</i>	Camaleon	0	0	0	0	0	0
122	Heteromyidae	<i>Dipodomys</i>	<i>merriami</i>	Rata canguro	0	0	1	0	0	0

123	Charadriidae	<i>Charadrius</i>	<i>vociferus vociferus</i>	Tildillo	0	0	0	0	0	0
124	Falconidae	<i>Falco</i>	<i>sparverius peninsularis</i>	Aguililla	0	0	1	0	0	0
125	Cardinalidae	<i>Pheucticus</i>	<i>melanocephalus</i>	Pintillo, Pajaro chichareo / chicarero, Pajaro trigero	0	0	0	0	0	0
126	Sciuridae	<i>Ammospermophilus</i>	<i>leucurus</i>	Juancito, Juanito, Chichimoco	0	0	0	0	0	0
127	Dytiscidae	over 160 genera	about 4,000 species	Cucaracha (del agua), la vieja de los chubasco	0	0	0	0	0	0
128	Caprimulgidae	<i>Phalaenoptilus</i>	<i>nuttallii</i>	Tapa camino, Caudebe	0	0	0	0	0	0
129	Elateridae	<i>Chalcolepidius</i>	<i>rubripennis</i>	Tronadores	0	0	0	0	0	0
130	Colubridae	<i>Thamnophis</i>	<i>hammondii</i>	Culebra de agua/de pozo	0	0	0	0	0	0
131	Icteridae	<i>Icterus</i>	<i>parisorum</i>	Serrano	0	0	0	0	0	0
132	Accipitridae	<i>Buteo</i>	<i>jamaicensis calurus</i>	Gavilan, Aguililla cola roja	0	0	1	0	0	0
133	Laniidae	<i>Lanius</i>	<i>ludovicianus</i>	Pajaro cabezon, Pajaro lelo, Cenzontle tapojero/entapojado/atapojado	0	0	0	0	0	0
134	Troglodytidae	<i>Salpinctes</i>	<i>obsoletus</i>	Piedredero, Catalina, Pajaro de Cantil/de los Cantiles	0	0	0	0	0	0
135	Eleotridae	<i>Dormitator</i>	<i>latrifons</i>	Puyeki	0	1	0	0	0	0
136	Tyrannidae	<i>Sayornis</i>	<i>nigricans</i>	Albañil, Pajaro lodero, Lodorito, Cantilerito, Pajaro de las posas, Pajaro de lodo	0	0	0	0	0	0
137	Scolopacidae	<i>Calidris</i>	<i>pusilla</i>	Sarapico, Gallito	0	0	0	0	0	0

Appendix B.

Table 8. Choyero ethnobiological knowledge for full sample (all), females, and males.

Item category	Items <i>i</i> (n)	Recognized $= \sum_{i=1}^n R_i$				Correctly Named $= \sum_{i=1}^n C_i$			
		Mean% (SD)	Median%	Min%,	Max%	Mean% (SD)	Median%	Min%,	Max%
All (N=71)									
all plants & animal:	137	76.4 (14.53)	80.3	29.2,	99.3	48.6 (15.37)	48.9	7.3,	91.2
dangerous	28	84.7 (13.20)	89.3	39.3,	100	54.0 (16.31)	53.6	14.3,	89.3
food items	32	82.1 (13.35)	84.4	31.3,	100	62.3 (17.15)	65.6	6.3,	96.9
animal	50	85.9 (13.95)	90.0	34.0,	100	59.0 (16.50)	58.0	12.0,	96.0
plant	87	71.0 (16.95)	74.7	17.2,	98.9	42.6 (16.73)	42.5	2.3,	88.5
<i>jardin</i> plants	36	74.1 (17.85)	77.8	16.7,	100	44.6 (19.80)	47.2	0.0,	91.7
<i>huerta</i> plants	16	80.2 (16.64)	87.5	25.0,	100	60.7 (20.11)	62.5	0.0,	100
<i>campo</i> plants	53	68.4 (18.22)	69.8	15.1,	100	39.5 (17.52)	41.5	1.9,	88.7
Females (N=33)									
all plants & animal:	137	73.1 (16.23)	75.2	29.2,	97.1	44.6 (15.98)	44.5	7.3,	75.9
dangerous	28	80.5 (15.80)	85.7	39.3,	100	45.7 (14.17)	46.4	14.3,	78.6
food items	32	80.1 (15.62)	81.3	31.3,	96.9	59.4 (18.78)	62.5	12.5,	90.6
animal	50	80.1 (16.14)	84.0	34.0,	100	49.9 (14.57)	52.0	12.0,	84.0
plant	87	69.1 (18.27)	72.4	17.2,	95.4	41.5 (18.11)	39.1	3.4,	78.2
<i>jardin</i> plants	36	77.1 (18.58)	83.3	16.7,	100	50.8 (21.91)	50.0	5.6,	91.7
<i>huerta</i> plants	16	78.4 (18.95)	81.3	25.0,	100	59.1 (21.03)	62.5	12.5,	93.8
<i>campo</i> plants	53	63.1 (19.13)	66.0	15.1,	94.3	33.0 (16.64)	32.1	1.9,	67.9
Males (N=38)									
all plants & animal:	137	79.3 (12.37)	81.8	50.4,	99.3	52.0 (14.12)	53.3	13.9,	91.2
dangerous	28	88.3 (9.20)	89.3	67.9,	100	61.2 (14.65)	62.5	21.4,	89.3
food items	32	83.9 (10.94)	84.4	56.3,	100	64.9 (15.39)	65.6	6.3,	96.9
animal	50	91.1 (9.24)	93.0	66.0,	100	66.8 (13.95)	68.0	30.0,	96.0
plant	87	72.6 (15.79)	74.7	33.3,	98.9	43.5 (15.61)	43.7	2.3,	88.5
<i>jardin</i> plants	36	71.4 (17.00)	72.2	27.8,	97.2	39.3 (16.26)	38.9	0.0,	88.9
<i>huerta</i> plants	16	81.7 (14.41)	87.5	37.5,	100	62.2 (19.44)	62.5	0.0,	100
<i>campo</i> plants	53	73.1 (16.23)	72.6	32.1,	100	45.2 (16.44)	47.2	3.8,	88.7

Note: Variables R_i and C_i are equal to 1 when item i is recognized or named correctly, respectively, and zero otherwise.

Appendix B.

Table 9. Choyero ethnobiological knowledge in pre-reproductive lifestage (<23 years old) for full sample (all), females, and males.

Item category	Items <i>i</i> (n)	Recognized = $\sum_{i=1}^n R_i$				Correctly Named = $\sum_{i=1}^n C_i$			
		Mean% (SD)	Median%	Min%,	Max%	Mean% (SD)	Median%	Min%,	Max%
All (N=7)									
all plants & animal:	137	63.2 (20.22)	70.1	32.1	81.8	30.8 (17.67)	29.9	7.3	49.6
dangerous	28	73.0 (19.87)	82.1	39.3	89.3	37.8 (16.73)	35.7	14.3	60.7
food items	32	70.5 (17.01)	81.3	40.7	84.4	46.9 (22.75)	50.0	12.5	71.9
animal	50	69.7 (22.20)	78.0	34.0	92.0	40.6 (22.27)	44.0	12.0	66.0
plant	87	59.4 (19.16)	65.5	31.0	78.2	25.1 (15.65)	24.1	3.5	43.7
<i>jardin</i> plants	36	64.7 (18.75)	72.3	38.9	91.7	29.0 (17.34)	38.9	5.6	47.2
<i>huerta</i> plants	16	68.8 (20.09)	75.0	37.5	87.5	45.5 (21.56)	50.0	12.5	68.8
<i>campo</i> plants	53	56.3 (21.00)	56.6	26.4	81.1	21.8 (17.63)	20.8	1.9	47.2
Females (N=3)									
all plants & animal:	137	47.7 (19.87)	40.9	32.1	70.1	15.6 (12.48)	9.5	7.3	29.9
dangerous	28	57.1 (22.30)	50.0	39.3	82.1	22.6 (8.99)	21.4	14.3	32.1
food items	32	58.3 (20.81)	53.1	40.6	81.2	27.1 (20.09)	18.8	12.5	50.0
animal	50	52.6 (22.74)	46.0	34.0	78.0	23.3 (17.93)	14.0	12.0	44.0
plant	87	44.8 (18.25)	37.9	31.0	65.5	11.1 (9.57)	8.0	3.5	21.8
<i>jardin</i> plants	36	55.6 (20.03)	50.0	38.9	77.8	16.7 (19.25)	5.6	5.6	38.9
<i>huerta</i> plants	16	52.1 (20.09)	43.8	37.5	75.0	27.1 (20.09)	18.8	12.5	50.0
<i>campo</i> plants	53	38.4 (16.05)	32.1	26.4	56.6	6.3 (3.93)	7.5	1.9	9.4
Males (N=4)									
all plants & animal:	137	74.9 (11.56)	79.9	57.7	81.8	42.2 (10.79)	46.4	26.3	49.6
dangerous	28	84.8 (4.49)	85.7	78.6	89.3	49.1 (10.26)	50.0	35.7	60.7
food items	32	79.7 (5.42)	81.3	71.88	84.4	61.7 (8.98)	62.5	50.0	71.9
animal	50	82.5 (11.47)	86.0	66.0	92.0	53.5 (16.03)	59.0	30.0	66.0
plant	87	70.4 (11.77)	75.3	52.9	78.2	35.6 (9.24)	37.4	24.1	43.7
<i>jardin</i> plants	36	71.5 (17.03)	72.2	50.0	91.7	38.2 (9.45)	40.3	25.0	47.2
<i>huerta</i> plants	16	81.3 (7.22)	81.3	75.0	87.5	59.4 (8.07)	59.4	50.0	68.8
<i>campo</i> plants	53	69.8 (11.93)	71.7	54.7	81.1	33.5 (13.73)	33.0	20.8	47.2

Note: Variables R_i and C_i are equal to 1 if item i is recognized or named correctly, respectively, and zero otherwise.

Table 10. Choyero ethnobiological knowledge in reproductive lifestage (= 23 to 45 years old) for for full sample (all), females, and males.

Item category	Items <i>i</i> (n)	Recognized $= \sum_{i=1}^n R_i$				Correctly Named $= \sum_{i=1}^n C_i$			
		Mean% (SD)	Median%	Min%,	Max%	Mean% (SD)	Median%	Min%,	Max%
All (N=26)									
all plants & animal:	137	81.4 (9.35)	82.1	59.1	98.5	52.7 (8.72)	54.0	32.1	73.0
dangerous	28	88.5 (11.13)	92.9	50.0	100.0	56.3 (12.75)	57.1	25.0	85.7
food items	32	88.3 (6.76)	87.5	71.9	100.0	70.8 (7.75)	71.9	56.3	84.4
animal	50	89.0 (10.71)	92.0	48.0	100.0	60.5 (12.59)	60.0	30.0	86.0
plant	87	77.1 (10.75)	77.6	54.0	97.7	48.1 (9.31)	48.3	32.2	65.5
<i>5jardin</i> plants	36	81.0 (11.38)	83.3	52.8	94.4	51.7 (13.34)	50.0	30.6	80.6
<i>huerta</i> plants	16	88.5 (7.44)	87.5	75.0	100.0	72.1 (10.19)	71.9	50.0	87.5
<i>campo</i> plants	53	73.9 (13.96)	71.7	45.3	100.0	43.8 (12.09)	45.3	20.8	64.2
Females (N=11)									
all plants & animal:	137	78.0 (10.08)	80.3	59.1	89.1	48.7 (8.83)	51.8	32.1	57.7
dangerous	28	83.1 (13.47)	89.3	50.0	92.9	46.4 (9.58)	46.4	25.0	60.7
food items	32	87.2 (6.91)	87.5	71.9	96.9	69.0 (8.55)	68.8	56.3	81.3
animal	50	82.9 (12.60)	86.0	48.0	92.0	50.2 (9.31)	52.0	30.0	62.0
plant	87	75.2 (10.53)	78.2	54.0	87.4	47.9 (9.55)	48.3	33.3	60.9
<i>jardin</i> plants	36	85.3 (7.57)	86.1	66.7	94.4	61.4 (11.14)	61.1	47.2	80.6
<i>huerta</i> plants	16	88.6 (7.30)	87.0	75.0	100.0	72.2 (9.83)	68.8	62.5	87.5
<i>campo</i> plants	53	67.6 (14.51)	67.9	45.3	84.9	36.0 (10.59)	34.0	20.8	50.9
Males (N=15)									
all plants & animal:	137	83.9 (8.23)	82.5	72.3	98.5	55.6 (7.65)	54.7	44.5	73.0
dangerous	28	92.4 (7.25)	92.9	75.0	100.0	63.6 (9.56)	64.3	46.4	85.7
food items	32	89.2 (6.77)	87.5	78.1	100.0	72.1 (7.13)	75.0	59.4	84.4
animal	50	93.5 (6.39)	94.0	78.0	100.0	68.1 (8.67)	68.0	54.0	86.0
plant	87	78.5 (11.06)	77.0	59.8	97.7	48.4 (9.45)	48.3	32.2	65.5
<i>jardin</i> plants	36	77.8 (12.82)	75.0	52.8	94.4	44.6 (10.10)	41.7	30.6	66.7
<i>huerta</i> plants	16	88.3 (7.79)	87.5	75.0	100.0	72.1 (10.80)	75.0	50.0	87.5
<i>campo</i> plants	53	78.5 (11.99)	77.4	56.6	100.0	49.4 (9.96)	52.8	30.2	64.2

Note: Variables R_i and C_i are equal to 1 if item i is recognized or named correctly, respectively, and zero otherwise.

Table 11. Choyero ethnobiological knowledge in post-reproductive lifestage (46+ years old) for for full sample (all), females, and males.

Item category	Items <i>i</i> (n)	Recognized $= \sum_{i=1}^n R_i$				Correctly Named $= \sum_{i=1}^n C_i$			
		Mean% (SD)	Median%	Min%, Max%	Mean% (SD)	Median%	Min%, Max		
All (N=38)									
all plants & animal	137	75.5 (14.90)	77.0	29.2 99.3	49.0 (16.52)	47.1	13.9 91.2		
dangerous	28	84.3 (12.08)	87.5	42.9 100.0	55.4 (17.09)	51.8	21.4 89.3		
food items	32	80.0 (14.23)	79.7	31.3 100.0	59.4 (18.09)	59.4	6.3 96.9		
animal	50	86.8 (12.40)	90.0	50.0 100.0	61.3 (16.00)	60.0	30.0 96.0		
plant	87	68.9 (18.72)	72.4	17.2 98.9	42.0 (18.74)	39.1	2.3 88.5		
<i>jardin</i> plants	36	71.1 (19.97)	72.2	16.7 100.0	42.7 (22.08)	38.9	0.0 91.7		
<i>huerta</i> plants	16	76.6 (18.30)	75.0	25.0 100.0	55.8 (21.42)	56.3	0.0 100.0		
<i>campo</i> plants	53	66.9 (19.34)	67.9	15.1 100.0	39.9 (18.97)	38.7	3.8 88.7		
Females (N=19)									
all plants & animal	137	74.3 (15.62)	75.2	29.2 97.1	46.8 (15.42)	43.8	16.8 75.9		
dangerous	28	82.7 (13.69)	85.7	42.9 100.0	48.9 (14.04)	46.4	25.0 78.6		
food items	32	79.4 (15.77)	78.1	31.3 96.9	58.9 (17.62)	59.4	15.6 90.6		
animal	50	82.7 (13.40)	84.0	50.0 100.0	54.0 (12.58)	52.0	30.0 84.0		
plant	87	69.4 (19.33)	73.6	17.2 95.4	42.6 (18.50)	39.1	9.2 78.2		
<i>jardin</i> plants	36	75.7 (20.41)	80.6	16.7 100.0	50.0 (21.91)	50.0	8.3 91.7		
<i>huerta</i> plants	16	76.6 (19.53)	75.0	25.0 100.0	56.6 (20.14)	56.3	12.5 93.8		
<i>campo</i> plants	53	64.3 (19.72)	66.0	15.1 94.3	35.5 (17.24)	32.1	7.6 67.9		
Males (N=19)									
all plants & animal	137	76.6 (14.47)	81.0	50.4 99.3	51.3 (17.66)	51.1	13.9 91.2		
dangerous	28	85.9 (10.34)	89.3	67.9 100.0	61.8 (17.73)	67.9	21.4 89.3		
food items	32	80.6 (12.91)	81.3	56.3 100.0	59.9 (19.02)	59.4	6.3 96.9		
animal	50	90.9 (10.03)	96.0	66.0 100.0	68.6 (15.96)	70.0	34.0 96.0		
plant	87	68.4 (18.59)	70.1	33.3 98.9	41.4 (19.47)	40.2	2.3 88.5		
<i>jardin</i> plants	36	66.4 (18.91)	63.9	27.8 97.2	35.4 (20.22)	30.6	0.0 88.9		
<i>huerta</i> plants	16	76.6 (17.53)	75.0	37.5 100.0	54.9 (23.16)	56.3	0.0 100.00		
<i>campo</i> plants	53	69.5 (19.12)	71.7	32.1 100.0	44.3 (20.02)	45.3	3.8 88.7		

Note: Variables R_i and C_i are equal to 1 if item i is recognized or named correctly, respectively, and zero otherwise.