

***C. elegans* LRP-2 functions in vulval precursor cell polarity**

Paul J Minor^{1,2} and Paul W Sternberg^{1§}

¹Division of Biology and Biological Engineering, Caltech, Pasadena, CA 91125

²Department of Biology, Hopkins Marine Station of Stanford University, Pacific Grove, CA 93950

[§]To whom correspondence should be addressed: pws@caltech.edu

Relevant Genotype	% P-Rv1	n	p value
N2	0	100	
<i>lin-17(n671)</i>	74	100	
<i>lin-18(e620)</i>	31	100	
<i>lin-17(n671); lin-18(e620)</i>	100	40	
<i>egl-20(hu120)</i>	0	66	
<i>lin-17(n671); egl-20(hu120)</i>	6	52	<0.0001
<i>cam-1(gm122)</i>	0	54	
<i>vang-1(ok1142)</i>	0	58	
<i>lrp-2(gk272)</i>	0	40	
<i>lrp-2(gk292)</i>	0	40	
<i>lin-17(n671); cam-1(gm122)</i>	50	54	0.0042
<i>lin-17(n671); vang-1(ok1142)</i>	48	60	0.0013
<i>lin-17(n671); lrp-2(gk272)</i>	55	40	0.0430
<i>lin-17(n671); lrp-2(gk292)</i>	54	55	0.0198
<i>lin-17(n671); lrp-2(gk272); cam-1(gm122)</i>	49	39	

Table 1. LRP-2 functions in ground polarity: P values in comparison to *lin-17(n671)* single mutants using Fisher's Exact Test.

Description

The *C. elegans* vulva is formed from divisions of three vulval precursor cells (VPCs) – P5.p, P6.p, and P7.p – arranged along the anteroposterior axis in the ventral epithelium (Sulston and Horvitz, 1977). Previous analyses show the orientation of P5.p and P7.p descendants is determined by the interaction of multiple Wnt signals. Specifically, in the absence of all Wnts, the VPCs display a randomized orientation, which is likely the default (Green *et al.*, 2008; Minor *et al.* 2013). Two separate Wnts from the anchor cell, LIN-44 and MOM-2 acting through receptors LIN-17/Frizzled and LIN-18/Ryk, respectively, regulate P7.p orientation (Ferguson *et al.*, 1987; Sternberg and Horvitz, 1988; Sawa *et al.*, 1996; Inoue *et al.*, 2004; Gleason *et al.*, 2006). In the absence of these signals the orientation of the progeny of P7.p mimic those of P5.p and face toward the posterior of the worm, a phenotype referred to as posterior-reversed vulval

lineage (P-Rvl). This posterior orientation is dependent on the instructive signal EGL-20, a Wnt expressed in the tail acting through CAM-1/ROR and VANG-1/Van Gogh, and is referred to as “ground polarity” (Green *et al.*, 2008).

Here we examine the role of a low-density lipoprotein receptor, *lrp-2*, and its role in controlling the orientation of P7.p daughter cells. To investigate this interaction double mutants were constructed with both alleles of *lrp-2* and *lin-17(n671)* (Table 1). Much like *cam-1(gm122)* and *vang-1(ok1142)*, both alleles of *lrp-2* suppress the *lin-17(n671)* phenotype from 74 to approximately 50% P-Rvl leading us to hypothesize that *lrp-2* functions in the same pathway as *cam-1* and *vang-1*. Furthering this hypothesis we have shown that, like *cam-1* and *vang-1*, *lrp-2* controls the localization of SYS-1/b-catenin (Minor and Sternberg, 2019). To ensure that this phenotype was a result of loss of *lrp-2* function as opposed to background effects we injected a fosmid (WRM0617cA02) containing the full-length sequence of *lrp-2* and found that it does rescue the double mutant phenotype of *lin-17(n671); lrp-2(gk272)* from 55 to 73%. In order to better test this hypothesis a triple mutant was constructed between *lin-17(n671)*, *lrp-2(gk272)*, and *cam-1(gm122)* (Table 1). This triple mutant displays the same P-Rvl penetrance as both the *lin-17(n671); lrp-2(gk272)* and *lin-17(n671); cam-1(gm122)* double mutants confirming that *lrp-2* functions in the same pathway as *cam-1*.

Methods

Strains were constructed by standard procedures (Ferguson and Horvitz, 1985).

Reagents

Strains:

N2

MT1306: *lin-17(n671)* (Ferguson and Horvitz, 1985)

PS4818: *lin-18(e620)*

PS3976: *lin-17(n671)/hT2 [qIs48 (myo-2::gfp etc.)]; +/- hT2 [qIs48]; lin-18(e620)*

PS5401: *egl-20(hu120)*

PS5530: *lin-17(n671); egl-20(hu120)*

NG2615: *cam-1(gm122)*

RB1125: *vang-1(ok1142)*

VC543: *lrp-2(gk272)*

VC621: *lrp-2(gk292)*

PS4809: *lin-17(n671); cam-1(gm122)*

PS5651: *lin-17(n671); vang-1(ok1142)*

lin-17(n671); lin-18(e620) animal awere segregated from **PS3976**.

The *lin-17(n671); lrp-2(gk292)* double mutant was constructed by crossing *lrp-2(gk292)* males with strain **MT1488:** *lin-17(n671); unc-13(e1091)* hermaphrodites.

The *lin-17(n671); lrp-2(gk272)* double mutant constructed by crossing *lrp-2(gk272)* males with strain **MT1488:** *lin-17(n671); unc-13(e1091)* hermaphrodites.

lin-17(n671); lrp-2(gk272); cam-1(gm122) Triple mutant constructed by crossing *lin-17(n671); lrp-2(gk272)* hermaphrodites with *cam-1(gm122)* and sequencing for homozygous triples.

PS6581: *qIs95[pSYS-1::VENUS::SYS-1 with ptx-3::dsRed as coninjection marker];*

PS6576: *lin-18(e620) sem-5(n1779)*

PS6580: *qIs95; sem-5(n1779)*

PS6579: *Pegl-17::CWN-1::GFP (unnamed construct) in cwn-1(ok546); lin-18(e620)*

References

Ferguson, E. L. and Horvitz, H. R. (1985). Identification and characterization of 22 genes that affect the vulval cell lineages of the nematode *Caenorhabditis elegans*. *Genetics* 110, 17-72. PMID: 3996896.

Ferguson, E. L., Sternberg, P. W. and Horvitz, H. R. (1987). A genetic pathway for the specification of the vulval cell lineages of *Caenorhabditis elegans*. *Nature* 326, 259-267. PMID: 2881214.

- Gleason, J. E., Szyleyko, E. A. and Eisenmann, D. M. (2006). Multiple redundant Wnt signaling components function in two processes during *C. elegans* vulval development. *Developmental biology* 298, 442-457. PMID: 16930586.
- Green, J. L., Inoue, T. and Sternberg, P. W. (2008). Opposing Wnt pathways orient cell polarity during organogenesis. *Cell* 134, 646-656. PMID: 18724937.
- Inoue, T., Oz, H. S., Wiland, D., Gharib, S., Deshpande, R., Hill, R. J., Katz, W. S. and Sternberg, P. W. (2004). *C. elegans* LIN-18 is a Ryk ortholog and functions in parallel to LIN-17/Frizzled in Wnt signaling. *Cell* 118, 795-806. PMID: 15369677.
- Minor P.J., He T.F., Sohn C.H., Asthagiri A.R., and Sternberg P.W. (2013). FGF signaling regulates Wnt ligand expression to control vulval cell lineage polarity in *C. elegans*. *Development* 140(18):3882-91. PMID: 23946444.
- Minor, P.J. and Sternberg, P.W. (2019). LRP-2 control the localization of *C. elegans* SYS-1/Beta-catenin. *microPublication Biology*. 10.17912/micropub.biology.000151 DOI: 10.17912/micropub.biology.000151
- Mizumoto, K. and Sawa, H. (2007). Two betas or not two betas: regulation of asymmetric division by beta-catenin. *Trends in cell biology* 17, 465-473. PMID: 17919911.
- Phillips, B. T., Kidd, A. R., 3rd, King, R., Hardin, J. and Kimble, J. (2007). Reciprocal asymmetry of SYS-1/beta-catenin and POP-1/TCF controls asymmetric divisions in *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences of the United States of America* 104, 3231-3236. PMID: 17296929.
- Sawa, H., Lobel, L. and Horvitz, H. R. (1996). The *Caenorhabditis elegans* gene *lin-17*, which is required for certain asymmetric cell divisions, encodes a putative seven-transmembrane protein similar to the *Drosophila* frizzled protein. *Genes & development* 10, 2189-2197. PMID: 8804313.
- Sternberg, P. W. and Horvitz, H. R. (1988). *lin-17* mutations of *Caenorhabditis elegans* disrupt certain asymmetric cell divisions. *Developmental biology* 130, 67-73. PMID: 3181641.
- Sulston, J. E. and Horvitz, H. R. (1977). Post-embryonic cell lineages of the nematode, *Caenorhabditis elegans*. *Developmental biology* 56, 110-156. PMID: 2578115.

Funding: Howard Hughes Medical Institute, with whom PWS was an Investigator. The National Institute of Neurological Disorders and Stroke of the National Institutes of Health under award number 1F32NS098658-01A1 awarded to PJM.

Reviewed By: Hitoshi Sawa

History: Received July 29, 2019 Accepted August 6, 2019 Published August 27, 2019

Copyright: © 2019 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Minor, PJ; Sternberg, PW (2019). *C. elegans* LRP-2 functions in vulval precursor cell polarity. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000152>